

Genomic organization, intragenic tandem duplication, and expression analysis of chicken *TGFBR2* gene

Bolin Ning,^{*,†,‡} Jiaxin Huang,^{*,†,‡} Haidong Xu,^{*,†,‡} Yuqi Lou,^{*,†,‡} Weishi Wang,^{*,†,‡} Fang Mu,^{*,†,‡}
Xiaohong Yan,^{*,†,‡} Hui Li,^{*,†,‡} and Ning Wang^{*,†,‡,1}

^{*}Key Laboratory of Chicken Genetics and Breeding, Ministry of Agriculture and Rural Affairs, Harbin 150030, China; [†]Key Laboratory of Animal Genetics, Breeding and Reproduction, Education Department of Heilongjiang Province, Harbin 150030, China; and [‡]College of Animal Science and Technology, Northeast Agricultural University, Harbin 150030, China

ABSTRACT Transforming growth factor beta receptor II (*TGFBR2*), a core member of the transforming growth factor- β (*TGF- β*) signaling pathway. To date, chicken *TGFBR2* (*cTGFBR2*) genomic structure has not been fully explored. Here, the complete sequences of *cTGFBR2* transcript isoforms were determined by 5' and 3' rapid amplification of cDNA ends (5' & 3' **RACE**) and reverse transcription polymerase chain reaction (**RT-PCR**); the tissue expression profiling of *cTGFBR2* transcript isoforms was performed using quantitative real-time polymerase chain reaction (**qRT-PCR**). The results showed that *cTGFBR2* gene produced 3 transcript isoforms through alternative transcription initiation, splicing, and polyadenylation, which were designated as *cTGFBR2-1*, *cTGFBR2-2*, and *cTGFBR2-3*, respectively. These 3 *cTGFBR2* transcript isoforms encoded 3 protein isoforms: *cTGFBR2-1*, *cTGFBR2-2*, and *cTGFBR2-3*. Duplication analysis revealed that, unlike other animal species, *cTGFBR2* gene harbored a 5.5-kb intragenic tandem duplication. Tissue expression profiling in the 4-wk-old Arbor Acres (**AA**) broiler chickens showed that

cTGFBR2-1 was ubiquitously expressed, with high expression in abdominal fat, subcutaneous fat, lung, gizzard, and muscle; *cTGFBR2-2* was highly expressed in heart, kidney, gizzard, and muscle; *cTGFBR2-3* was weakly expressed in all the tested chicken tissues. Tissue expression profiling in the 7-wk-old broiler chickens of the fat and lean lines of Northeast Agricultural University broiler lines divergently selected for abdominal fat content (**NEAUHLF**) showed that *cTGFBR2-1* was significantly differentially expressed in all the tested tissues except heart, *cTGFBR2-2* was significantly differentially expressed in all the tested tissues except subcutaneous fat and liver, and *cTGFBR2-3* was significantly differentially expressed in all the tested tissues between the lean and fat lines. Intriguingly, in the fat line, the 3 *cTGFBR2* transcript isoforms were expressed to varying degrees in all the 3 tested fat tissues, while in the lean line, only *cTGFBR2-1* was expressed in all the 3 tested fat tissues. This is the first report of intragenic tandem duplication within *TGFBR2* gene. Our findings pave the way for further studies on the functions and regulation of *cTGFBR2* gene.

Key words: chicken, *TGFBR2*, transcript isoform, intragenic tandem duplication, gene expression

2022 Poultry Science 101:102169
<https://doi.org/10.1016/j.psj.2022.102169>

INTRODUCTION

Transforming growth factor- β (*TGF- β*) signaling pathway regulates a variety of biological processes throughout embryonic development and postnatal life, as well as during the presentation of human disease (Lönn et al., 2009; Moustakas and Heldin, 2009; Pardali et al., 2010; Xu et al., 2012). There are 3 *TGF- β* forms

(*TGF- β 1*, 2, and 3) and 3 *TGF- β* receptors (*TGFBR1*, 2, and 3) (Vander Ark et al., 2018). *TGFBR2* plays an important role in the *TGF- β* signaling pathway. Upon ligand binding, *TGFBR2* forms a signaling complex with *TGFBR1* and activates *TGFBR1* via phosphorylation. The activated *TGFBR1* in turn phosphorylates its downstream receptor-regulated SMADs (**R-SMADs**, **SMAD2/3**). Subsequently, the activated R-SMADs bind to SMAD4, migrate to the nucleus, and regulate *TGF- β* target gene transcription.

TGFBR2 serves as an initial regulator of the *TGF- β* signaling pathway and is involved in many cellular processes (Moustakas and Heldin, 2009). Loss or reduction of its expression can lead to uncontrolled cell growth (Yuan et al., 2020). Defects by *TGFBR2* knockout involve many

© 2022 The Authors. Published by Elsevier Inc. on behalf of Poultry Science Association Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Received April 4, 2022.

Accepted September 2, 2022.

¹Corresponding author: wangning@neau.edu.cn

phylogenetic and major organ systems, including developmental defects of the kidney, skeleton, and lymphatic network (James et al., 2013; Peters et al., 2017; Dumbrava et al., 2021). During the development of chicken embryos, *TGFBR2* shows dynamically and frequently overlapping expression patterns in numerous embryonic cell layers and structures, and is involved in chicken-specific developmental processes including somitogenesis, cardiogenesis, and vasculogenesis (Cooley et al., 2014).

TGFBR2 genomic structure has been intensively studied in mammals. Both human and mouse *TGFBR2* genes are composed of 8 exons and 7 introns. Due to alternative splicing, *TGFBR2* gene generates 2 transcript isoforms in humans and mice, which encode 2 protein isoforms, named as T β R II and T β R II -B, respectively (Hirai and Fijita, 1996; Krishnaveni et al., 2006). Recently, a novel human *TGFBR2* transcript isoform named as T β R II -SE was discovered, which encodes a soluble protein (Bertolio et al., 2021). To date, chicken *TGFBR2* (*cTGFBR2*) genomic structure has not been fully explored, and whether the *cTGFBR2* gene encodes multiple transcript and protein isoforms is unclear. In the present study, we for the first time characterized *cTGFBR2* transcript isoforms, dissected the *cTGFBR2* genomic structure and identified a 5.5-kb intragenic tandem duplication.

MATERIALS AND METHODS

Experimental Animals and Tissue Collection

Our animal experiments were conducted according to the guidelines for the care and use of experimental animals established by the Ministry of Science and Technology of the People's Republic of China (approval no. 2006-398) and were approved by the Laboratory Animal Management Committee of Northeast Agricultural University

(Harbin, Heilongjiang, PRC). All experimental chickens were fed under the same environmental conditions with free intake of food and water. Abdominal fat, subcutaneous fat, gizzard fat, heart, liver, spleen, lung, kidney, gizzard, proventriculus, pancreas, and muscle tissue were collected from 4-wk-old Arbor Acres (AA) broiler chickens. All of the above tissues except the lung were collected from 7-wk-old broiler chickens of fat and lean lines of Northeast Agricultural University broiler lines divergently selected for abdominal fat content (NEAUHLF) (24th generation). All samples were washed with a solution containing 0.75% NaCl, snap frozen in liquid nitrogen, and stored at -80°C until RNA isolation.

RNA Isolation and Reverse Transcription

The frozen tissue samples were ground into powder in liquid nitrogen by using a mortar and pestle. Total RNAs were isolated using Trizol reagent (Invitrogen, Carlsbad, CA) according to the manufacturer's protocol. RNA quality was assessed by denaturing formaldehyde agarose gel electrophoresis and quantified by using a spectrophotometer. The extracted RNA was reverse transcribed using the PrimeScriptTM RT reagent Kit (Takara, Dalian, China) according to the manufacturer's protocol.

5' and 3' Rapid Amplification of cDNA Ends

The 5' and 3' rapid amplification of cDNA ends (5' & 3' RACE) of *cTGFBR2* was performed using the SMARTer RACE 5'/3' Kit (Takara, Dalian, China) according to the manufacturer's protocol. The gene-specific primers for the 5' and 3' RACE were designed by Primer 6.0 based on *cTGFBR2* mRNA (NM_205428.1). The primer sequences are listed in Table 1. The RACE products were separated by agarose gel electrophoresis

Table 1. The primers used in the present study.

Type	Primer name	Primer sequence	
RACE primer	5'RACE-GSP1	R: ACAGCGAGATGTCATTTCCAGAG	
	5'RACE-GSP2	F: TTCCCAGAGGACCAAAGC	
	3'RACE-GSP1	F: CTCATGGCTTTGGTCTCTG	
	3'RACE-GSP2	F: CTCGCTGTAATGGTGTGG	
RT-PCR primer	TGFBR2-1-F/TGFBR2-1-R	F: TGCAGCGCCGAAGTGAAGTTTTTC R: TACATCTCCCTGCCAGAGCCAC	
	TGFBR2-1-F/ TGFBR2-2-R	F: TGCAGCGCCGAAGTGAAGTTTTTC R: GAACAGAGTCAGGCTGTGGTATG	
	TGFBR2-1-F/ TGFBR2-3-R	F: TGCAGCGCCGAAGTGAAGTTTTTC R: GTCCGTAAGAGCCACTCAACATA	
	TGFBR2-2-F/ TGFBR2-1-R	F: GGGGAGAAGGAAGTGAAGTTTTTC R: TACATCTCCCTGCCAGAGCCAC	
	TGFBR2-2-F/ TGFBR2-2-R	F: GGGGAGAAGGAAGTGAAGTTTTTC R: GAACAGAGTCAGGCTGTGGTATG	
	TGFBR2-2-F/ TGFBR2-3-R	F: GGGGAGAAGGAAGTGAAGTTTTTC R: GTCCGTAAGAGCCACTCAACATA	
	PCR primer	Genome- F/R	F: AGAAGGATGATGGGGACTGA R: TGGTTTGAGCACGTTGTTGC
	qRT-PCR primer	<i>cTGFBR2</i> -total-F/R	F: CTAAGAGACAGAGGGCGACC R: ACAGCTTCTCCTGAGAGCT
<i>cTGFBR2</i> -1-F/R		F: TAGAAGCAAGGAAAATGG R: CTGGCTTATGGGGATCAA	
<i>cTGFBR2</i> -3-F/R		F: TTGGTCTCTGGGAAATG R: AGATGGGCTTTGTAGTGC	
TBP-F/R		F: GCGTTTGTCTGTGTTATTATGAG R: TCCTTGCTGCCAGTCTGGAC	

Abbreviations: TGFBR2, transforming growth factor beta receptor II; TBP, TATA-box binding protein.

and recovered using DNA Gel Recovery Kit (Axygen, Union City, CA) following the manufacturer's protocol. All recovered products were cloned using pEASY blunt simple cloning vector (TransGen, Beijing, China) according to the manufacturer's protocol, and transformed into *Escherichia coli* (*E. coli*). Single colonies were picked for PCR identification and sent to Genewiz biological company (Suzhou, China) for Sanger sequencing.

Reverse Transcription Polymerase Chain Reaction and Genomic PCR

Reverse transcription polymerase chain reaction (RT-PCR) was used to identify *cTGFBR2* transcript isoforms on the cDNAs generated from various chicken tissues. The primers are shown in Table 1. Thermal cycling was performed as follows: 5 min for the initial denaturation at 95°C, followed by 40 cycles of 15 s at 95°C, 15 s at 60°C, and 180 s at 72°C, and ending with a final extension time of 10 min at 72°C in a Master cycler nexus GSX1 (Eppendorf, Hamburg, Germany). Genomic PCR was used to detect the intragenic tandem duplication of *cTGFBR2* gene on blood genomic DNAs extracted from 4 different chicken breeds (Northeast Agricultural University F2 Resource Line, NEAUHLF, AA broiler chickens, and Lidian chickens), which were stored in our laboratory. The primers are shown in Table 1. The forward primer was located in exon 3 of *cTGFBR2* gene and the reverse primer was located in exon 2'. If there is an intragenic tandem duplication comprising exons 2 and 3 and intron 2 in the *cTGFBR2* gene, genomic PCR amplification with this primer pair generates a 2,000-bp product, otherwise, no PCR products would be obtained. Thermal cycling was performed as follows: 5 min for the initial denaturation at 95°C, followed by 40 cycles of 30 s at 95°C, 30 s at 63°C, 150 s at 72°C, and ending with a final extension time of 10 min at 72°C in a Master cycler nexus GSX1 (Eppendorf, Hamburg, Germany). PCR products were purified, cloned into pEASY blunt simple cloning vector (TransGen, Beijing, China), and sent to Genewiz biological company (Suzhou, China) for Sanger sequencing.

Quantitative Real-Time Polymerase Chain Reaction

The relative expression of *cTGFBR2-1*, *cTGFBR2-2*, and *cTGFBR2-3* were determined by quantitative real-time polymerase chain reaction (qRT-PCR) using FastStart Universal SYBR Green Master [Rox] (Roche, Basel, Switzerland). The primers are shown in Table 1. The expression levels of *cTGFBR2-1* and *cTGFBR2-3* were detected with primer pairs *cTGFBR2-1-F/R* and *cTGFBR2-3-F/R*, respectively. The expression level of *cTGFBR2-2* was calculated by the total expression of *cTGFBR2-1* plus *cTGFBR2-2* detected with the primer pair *cTGFBR2-total-F/R* minus the *cTGFBR2-1* expression detected with the primer pair *cTGFBR2-1-F/R*. The qRT-PCR was performed on ABI Prism 7500 sequence

detection system (Applied Biosystems, Foster City, CA). The cycling conditions were used as follows: 30 s at 95°C, followed by 40 cycles of 5 s at 95°C and 30 s at 60°C. TATA-box binding protein (*TBP*) was used as an internal reference, and the relative gene expression was calculated using the $2^{-\Delta C_t}$ method.

Western Blotting Analysis

The frozen tissue samples were ground into powder in liquid nitrogen by using a mortar and pestle and lysed using RIPA buffer containing 1% PMSF (Beyotime, Shanghai, China). Proteins were separated by 12% SDS-PAGE and transferred onto nitrocellulose membranes (Millipore, Bedford, MA). Then the membranes were blocked for 120 min and incubated overnight at 4°C with primary antibodies against TGFBR2 (1:1,000, Santa, Dallas, Texas) and β -actin (1:1,000, Beyotime, Shanghai, China). After the blots were rinsed with PBST 3 times, they were incubated with HRP-conjugated anti-mouse secondary antibody (1:3,000, Beyotime, Shanghai, China) for 60 min at room temperature. The blots were observed with an ECL Plus detection kit (Beyotime, Shanghai, China).

Bioinformatics Analysis

cTGFBR2 genomic sequence was downloaded from the National Center for Biotechnology Information (NCBI) website (https://www.ncbi.nlm.nih.gov/nucleotide/NC_052533.1?report=fasta&from=39709134&to=39773208). Intragenic tandem duplication analysis was performed using the DNASTAR software. Open reading frame (ORF) analysis and sequence identity analysis were performed using the DNAMAN software. Sequence alignment was performed using the basic local alignment search tool (BLAST) (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>). Protein domain analysis was performed using SMART (<http://smart.embl-heidelberg.de/>). Polyadenylation signal (PAS) analysis was performed using the ITB tools (<http://itbtools.ba.itb.cnr.it/utrscan>).

Statistical Analysis

Results are expressed as means \pm SD. Comparisons between groups were performed using the multiple *t* tests. Statistical differences were considered significant when *P* < 0.05.

RESULTS

Identification of the 5' and 3' Terminal Sequences of *cTGFBR2* mRNA

To obtain the full-length *cTGFBR2* mRNA, we performed the 5' and 3' RACE on the pooled cDNA samples. The 5' RACE and 3' RACE PCR products were analyzed by electrophoresis on 2% agarose gels. The results showed that there were 2 different-sized 5'




Figure 1. Identification of the 5' and 3' terminal regions of *cTGFB2* transcript isoforms. (A) 5' RACE analysis of *cTGFB2* transcript isoforms. M, marker. (B) 3' RACE analysis of *cTGFB2* transcript isoforms. M, marker. (C) Schematic structure of the 5'-terminal region of *cTGFB2* transcript isoforms. (D) Schematic structure of the 3'-terminal regions of *cTGFB2* transcript isoforms. Exons are numbered inside the boxes.

RACE PCR products (Figure 1A) and 3 different-sized 3' RACE PCR products (Figure 1B). All 5' and 3' RACE PCR products were purified, cloned and sequenced. A total of 36 recombinant plasmids containing the 5' RACE PCR products were sequenced. The sequencing analysis identified 2 distinct 5'-terminal sequences of *cTGFB2*: 1,493 (11 colonies) and 1,918 bp (25 colonies) in size, which were designated as Type a and Type b, respectively (Figure 1C). A total of 82 recombinant plasmids containing the 3' RACE PCR products were sequenced. The sequencing analysis identified 3 distinct 3'-terminal sequences of *cTGFB2*: 913 (39 colonies), 2,560 (26 colonies), and 1,640 bp (17 colonies), which were designated as Type A, Type B, and Type C, respectively (Figure 1D).

Identification of *cTGFB2* Transcript Isoforms

Based on the 5' and 3' RACE analysis results, we speculated that the *cTGFB2* gene may generate at most 6 transcript isoforms (Figure 2A). To identify *cTGFB2* transcript isoforms, we designed the transcript isoform-specific primer sets for the 6 possible transcript isoforms (Table 1 and Figure 2A) and performed RT-PCR on the pooled cDNA. The expected RT-PCR product sizes for 6 possible *cTGFB2* transcript isoforms were 1,953, 2,904, 2,360, 2,028, 3,321, and 2,777 bp, respectively. The RT-PCR products were analyzed by electrophoresis on 2% agarose gels. The RT-PCR with primer sets TGFBR2-1-F/TGFBR2-1-R, TGFBR2-2-F/TGFBR2-

2-R, and TGFBR2-2-F/TGFBR2-3-R yielded the expected RT-PCR products, but RT-PCR with other 3 primer sets (TGFBR2-1-F/TGFBR2-2-R, TGFBR2-1-F/TGFBR2-3-R, and TGFBR2-2-F/TGFBR2-1-R) yielded no RT-PCR products (Figure 2B). The 3 expected RT-PCR products were cloned and a total of 180 recombinant plasmids were sequenced. The sequencing results showed, as expected, the sizes of these 3 RT-PCR products were 1,953 (52 colonies), 3,321 (49 colonies), and 2,777 bp (79 colonies), respectively. Collectively, these results suggest that the *cTGFB2* gene produces 3 transcript isoforms, which were designated as *cTGFB2-1*, *cTGFB2-2*, and *cTGFB2-3*. Their nucleotide sequences were deposited in GenBank database under accession numbers ON164837, ON164838, and ON164839, respectively.

There are 2 *cTGFB2* transcript sequences (NM_205428.2 and XM_015281321.4) in GenBank database. Sequence identity analysis using DNAMAN software showed that *cTGFB2-1* shared 57.88 and 50.36% nucleotide identities, respectively, with the 2 reported *cTGFB2* transcript sequences; *cTGFB2-2* shared 86.08 and 97.18% nucleotide identities, respectively, and *cTGFB2-3* shared 46.71 and 57.88% nucleotide identities, respectively. The 5' UTR sizes of the 3 *cTGFB2* transcript isoforms were 140, 223, and 223 bp, respectively. The 3' UTR sizes of the 3 *cTGFB2* transcript isoforms were 535, 2,196, and 1,413 bp, respectively. The coding region sequence of *cTGFB2-1* and *cTGFB2-2* were completely identical to that of NM_205428.2 and XM_015281321.4, respectively. The *cTGFB2-1* contained an additional 23




Figure 2. Identification of *cTGFBR2* transcript isoforms. (A) Schematic structure of 6 possible *cTGFBR2* transcript isoforms and schematic location of the 6 transcript isoform-specific primer sets used for the identification of *cTGFBR2* transcript isoforms. (B) RT-PCR identification of *cTGFBR2* transcript isoforms. RT-PCR was performed on pooled cDNA to identify *cTGFBR2* transcript isoforms with the 6 transcript-specific primer sets. RT-PCR products were analyzed by 2% agarose gel electrophoresis. M, marker. Lane 1, RT-PCR with primer set TGFBR2-1-F and TGFBR2-1-R; Lane 2, RT-PCR with primer set TGFBR2-1-F and TGFBR2-2-R; Lane 3, RT-PCR with primer set TGFBR2-1-F and TGFBR2-3-R; Lane 4, RT-PCR with primer set TGFBR2-2-F and TGFBR2-1-R; Lane 5, RT-PCR with primer set TGFBR2-2-F and TGFBR2-2-R; Lane 6, RT-PCR with primer set TGFBR2-2-F and TGFBR2-3-R.

nucleotides in 5' UTR and lacked 1,663 nucleotides in 3' UTR, compared with the NM_205428.2. The *cTGFBR2-2* contained an additional 106 nucleotides in 5' UTR and lacked 2 nucleotides in 3' UTR, compared with the XM_015281321.4. The *cTGFBR2-3* contained an additional 83 nucleotides in 5' UTR, and an additional 878 nucleotides in 3' UTR, compared with the *cTGFBR2-1*. The *cTGFBR2-3* possessed the same 5' UTR as *cTGFBR2-2* but lacked 783 nucleotides in 3' UTR relative to the *cTGFBR2-2*.

ORF analysis using DNAMAN software showed that these 3 identified *cTGFBR2* transcript isoforms could encode 3 protein isoforms: *cTGFBR2-1* (557 amino acids), *cTGFBR2-2* (671 amino acids), and *cTGFBR2-3* (628 amino acids; **Figure 3A**). Comparison of amino acid sequences of *cTGFBR2-1*, *cTGFBR2-2*, *cTGFBR2-3* with the published chicken TGFBR2 (**cTGFBR2**) revealed that, *cTGFBR2-1* shared 100% identity with NP_990759.1 (encoded by NM_205428.2), *cTGFBR2-2*

shared 100% identity with XP_015136807.1 (encoded by XM_015281321.4), but *cTGFBR2-3* shared 74.52 and 91.51% identity with NP_990759.1 and XP_015136807.1, respectively. By comparison, *cTGFBR2-3* contained an additional 114 amino acids at its N-terminal, compared with the *cTGFBR2-1*, while lacked the C-terminal 43 amino acids, compared with the *cTGFBR2-1* and *cTGFBR2-2* (**Figure S1**).

The protein structural domain analysis using the SMART database showed that, similar to human and mouse TGFBR2, the 3 *cTGFBR2* isoforms consisted of 4 primary structural domains: the signal peptide (**SP**), transforming growth factor beta receptor 2 ectodomain (**ecTbetaR2**), transmembrane region (**TR**), and serine/threonine-protein kinases (**S_TKc**; **Figure 3A**). By comparison, *cTGFBR2-2* and *cTGFBR2-3* have 1 more ecTbetaR2 domain than *cTGFBR2-1* (**Figure 3A**). To confirm the presence of these 3 *cTGFBR2* isoforms, we performed western blotting using TGFBR2 monoclonal antibody (sc-



**B**

Figure 3. Structural analysis and detection of *cTGFBFR2* protein isoforms. (A) The domains of *cTGFBFR2* isoforms were predicted using SMART (<http://smart.embl-heidelberg.de/>). SP, signal peptide; ecTbetaR2, Transforming growth factor beta receptor 2 ectodomain; TR, Transmembrane region; S_TKc, Serine/threonine-protein kinases. (B) Western blotting analysis of *cTGFBFR2* protein isoform expression in the pooled tissue protein lysate (abdominal fat, heart, liver, kidney, gizzard, and muscle). The pooled tissue protein lysate was generated from the 4-wk-old AA broilers and immunoblotted with the indicated antibodies.

17799, Santa, Dallas, Texas) to detect *cTGFBFR2* in the pooled protein sample consisting of abdominal fat, heart, liver, kidney, gizzard, muscle in equal proportions. Western blotting detected the predicted 63, 76, and 71 kDa protein bands of *cTGFBFR2* isoforms (Figure 3B), suggesting that *cTGFBFR2* gene indeed encodes 3 protein isoforms.

Characterization of *cTGFBFR2* Genomic Structure

To determine the genomic structure of the *cTGFBFR2* gene, we aligned these 3 *cTGFBFR2* transcript isoform

sequences against chicken genome sequence (GRCg7b) using the BLAST at the NCBI. As expected, these 3 *cTGFBFR2* transcript isoforms were mapped to the *cTGFBFR2* locus on chromosome 2 in the Gallus gallus genome database (GRCg7b). The *cTGFBFR2* gene spanned a region of approximately 64-kb (Chr.2: 39,709,028-39,773,208). The nucleotide lengths of each exon and intron of *cTGFBFR2* gene are as follows: exon 1a/1 (83/222 bp), intron 1 (26,763 bp), exon 2 (151 bp), intron 2 (1,517 bp), exon 3 (191 bp), intron 3 (1,747 bp), exon 2' (151 bp), intron 4 (1,622 bp), exon 3' (191 bp), intron 5 (11,718 bp), exon 4 (800 bp), intron 6 (4,245 bp), exon 5 (142 bp), intron 7 (8,713 bp), exons 6/6a




Figure 4. Schematic structure of *cTGFBR2* genomic and transcript isoforms. Abbreviations: ATG, start codon; TAA and TAG, stop codon; TSS, transcription start site; PAS, polyadenylation signal; CR-PAS, coding region PAS; UTR-PAS, untranslated region PAS.

(128/1,464 bp), intron 8 (1,955 bp), and exons 7/7a (715/1,661 bp; [Figure 4](#)). The *cTGFBR2-1* consisted of 7 exons (exons 1-7), *cTGFBR2-2* consisted of 9 exons (exon 1a/1, exons 2-6, exons 2' and 3', exons 7/7a), and *cTGFBR2-3* consisted of 8 exons (exon 1a/1, exons 2-5, exons 2' and 3', exons 6/6a; [Figure 4](#)). All the intron/exon splice junctions followed the GT-AG rule.

Furthermore, BLAST analysis revealed that the *cTGFBR2* gene had 2 alternative transcription start sites (TSS1 and TSS2), which were 82 bp apart. The *cTGFBR2-2* and *cTGFBR2-3* shared TSS1 and *cTGFBR2-1* employed TSS2 ([Figure 4](#)). These 3 *cTGFBR2* transcript isoforms shared the same start codon (ATG) on exon 1, the *cTGFBR2-1* and *cTGFBR2-2* shared the same stop codon (TAG) located in exon 7, whereas *cTGFBR2-3* positioned its stop codon (TAA) in exon 6a ([Figure 4](#)). Sequence analysis showed that *cTGFBR2* gene had 3 alternative 3' UTRs (535, 2,196, and 1,413 bp, respectively), indicating that the *cTGFBR2* gene undergoes alternative polyadenylation (APA). We used ITB tools to identify the PAses for the 3 alternative 3' UTRs. The results showed that as expected, *cTGFBR2* gene had 3 PAses (AATAAA), one of the PAses, located in exon 6a, was 6 bp upstream of the poly (A) site of *cTGFBR2-3* ([Figure 4](#)). The other 2 PAses, located in the exons 7 and 7a, were 9 and 16 bp, respectively, upstream of the poly (A) sites of *cTGFBR2-1* and *cTGFBR2-2* ([Figure 4](#)). The PAS in exon 6a was a coding region PAS (CR-PAS), which led to the production of the novel protein isoform *cTGFBR2-3*. The other 2 PAses were the untranslated region PAS (UTR-PAS), and did not affect the coding capacity of *cTGFBR2*, but caused the different 3' UTR lengths between *cTGFBR2-1* and *cTGFBR2-2*.

Intragenic Duplication of *cTGFBR2* Gene

The structural domain analysis showed that the *cTGFBR2-1* protein isoform had 1 ecTbetaR2 domain, while both *cTGFBR2-2* and *cTGFBR2-3* protein isoforms had 2 tandem ecTbetaR2 domains, which had high amino acid sequence identity (90.99%; [Figure 5A](#)). Using DNAMAN software, we analyzed the mRNA sequences coding the 2 tandem ecTbetaR2 domains in *cTGFBR2-2* and *cTGFBR2-3*. The results showed that the coding sequences for the 2 tandem ecTbetaR2 domains (324 and 330 bp, respectively) shared 97.08% nucleotide identity ([Figure 5B](#)).

Protein domain repeats derive from the intragenic tandem duplication ([Björklund et al., 2010](#)). To verify the presence of intragenic tandem duplication in the *cTGFBR2* gene locus, we performed duplication analysis of the *cTGFBR2* genomic sequence (Chr.2: 39,709,028-39,773,208, 64-kb) using DNASTAR software. The results displayed that the *cTGFBR2* genomic sequence had an intragenic tandem duplication of 5.5-kb (Chr.2: 39,736,096-39,741,665). Duplication region 1 (Chr.2: 39,736,096-39,737,954) was 1,859 bp in length and consisted of exons 2 and 3, and intron 2. Duplication region 2 (Chr.2: 39,739,702-39,741,665) was 1,964 bp in length and consisted of exons 2' and 3', and intron 4. These 2 intragenic duplication regions shared 87.68% nucleotide identity ([Figure 5C](#)), and the nucleotide identities between exon 2 and exon 2', exon 3 and exon 3', intron 2 and intron 4, were 93.38, 100.00, and 85.71%, respectively. Duplication region 1 encoded the first ecTbetaR2 domain of *cTGFBR2-2* and *cTGFBR2-3*, and duplication region 2 encoded the second ecTbetaR2 domain of *cTGFBR2-2* and *cTGFBR2-3*.

To test whether this *cTGFBR2* intragenic tandem duplication is commonly present in chickens, we




Figure 5. Schematic of *cTGFB2* intragenic duplication. (A) Schematic of *cTGFB2* with a tandem repeat of 2 ecTbetaR2 domains. (B) Schematic of *cTGFB2* transcript isoforms with a tandem duplication of exons 2 and 3. (C) Schematic of *cTGFB2* genomic intragenic duplication. The black square represents the region of intragenic duplication.

performed the duplication-specific genomic PCR to detect the intragenic tandem duplication in 4 chicken breeds: Northeast Agricultural University F2 Resource Line, NEAUHLF, AA broiler chickens, and Lindian chickens (a local Chinese chicken breed). The genomic PCR results showed that a single band of the expected size (2,000-bp) was amplified using the duplication-specific primer pair Genome-F/R in all the 4 tested chicken breeds (Figures 6A and 6B), and Sanger sequencing further confirmed that *cTGFB2* gene harbored the

intragenic tandem duplication in all the tested chickens. To test whether this intragenic tandem duplication is chicken-specific, we performed bioinformatics analysis of *TGFB2* genomic, mRNA, and protein sequences in various animal species including goose, duck, horse, penguin, eagle, human, and mouse. The bioinformatics analysis showed that there was no intragenic tandem duplication in *TGFB2* genomic and mRNA sequences, and no tandem domain repeats in TGFB2 protein in these tested animal species. Taken together, these




Figure 6. PCR identification of *cTGFB2* intragenic duplication. (A) Schematic locations of PCR primers used for the identification of *cTGFB2* intragenic duplication. (B) Genomic PCR identification of the intragenic duplication of *cTGFB2* in various chicken breeds. M, marker; Control, water as the negative control; Lanes 1 and 2, Northeast Agricultural University F2 Resource Line; Lanes 3 and 4, Lindian chicken breed; Lanes 5 and 6, NEAUHLF; Lanes 7 and 8, AA broiler chicken.

results suggest that the tandem domain repeat and intragenic tandem duplication of the *TGFBR2* gene is chicken-specific.

Tissue Expression Patterns of *cTGFBR2* Transcript Isoforms

To understand the functions of these 3 *cTGFBR2* transcript isoforms, we performed tissue expression profiling of the 3 *cTGFBR2* transcript isoforms in various tissues of the 4-wk-old AA broiler chickens using qRT-PCR with the primer pairs (Table 1, Figure 7A). The results showed that the *cTGFBR2-1* was ubiquitously expressed in all the tested chicken tissues, with high expression in abdominal fat, subcutaneous fat, lung, gizzard, and muscle, but low expression in gizzard fat, heart, liver, spleen, kidney, proventriculus, and pancreas. The *cTGFBR2-2* was highly expressed in the heart, kidney, gizzard, and muscle, whereas low or no expression was detected in abdominal fat, subcutaneous fat, gizzard fat, liver, spleen, lung, proventriculus, and pancreas. The *cTGFBR2-3* was weakly expressed in all the tested tissues, with no expression in abdominal fat, subcutaneous fat, gizzard fat, spleen, lung, proventriculus, and pancreas (Figure 7B). These results suggest that the 3 *cTGFBR2* transcript isoforms may play different roles in different tissues and organs.

To confirm the tissue expression profiling results, we also performed tissue expression profiling of the 3 *cTGFBR2* transcript isoforms in the 7-wk-old chickens of NEAUHLF. In both lean and fat lines, only *cTGFBR2-1* was ubiquitously expressed in all the tested tissues (Figure 7C). Comparison of gene expression

between the lean and fat lines revealed that *cTGFBR2-1* was significantly differentially expressed in all the tested tissues except the heart, and *cTGFBR2-2* was significantly differentially expressed in all the tested tissues except the subcutaneous fat and liver, and *cTGFBR2-3* was significantly differentially expressed in all the tested tissues (Figure 7C, $P < 0.05$).

It is worth noting that, in the fat line, the 3 *cTGFBR2* transcript isoforms were expressed to varying degrees in all the 3 tested fat tissues (Figure 7C). However, in the lean line, only *cTGFBR2-1* was expressed in all the 3 tested fat tissues, *cTGFBR2-2* and *cTGFBR2-3* were only expressed in subcutaneous fat (Figure 7C). These results suggest that the expression of the 3 *cTGFBR2* transcript isoforms is differentially regulated in adipose tissues at different anatomic sites.

DISCUSSION

In the present study, we demonstrated that *cTGFBR2* gene produces 3 transcript isoforms (*cTGFBR2-1*, *cTGFBR2-2*, *cTGFBR2-3*) through alternative transcription initiation, splicing, and polyadenylation and these transcript isoforms encode 3 cTGFBR2 protein isoforms (cTGFBR2-1, cTGFBR2-2, cTGFBR2-3). Among these 3 *cTGFBR2* transcript isoforms, *cTGFBR2-3* was a novel *cTGFBR2* transcript isoform, which encodes a novel cTGFBR2 protein isoform.

Intragenic duplication plays an important role in the evolution, inheritance and variation of life, and involves gene expression, transcription regulation, chromosome construction and physiological metabolism (Brahmachari




Figure 7. Tissue expression profiling of *cTGFBR2* transcript isoforms in various chicken tissues. (A) Schematic locations of qRT-PCR primers used in the tissue expression profiling of *cTGFBR2* transcript isoforms. (B) Expression profiling of the 3 *cTGFBR2* transcript isoforms in various tissues of the 4-wk-old AA chickens. (C) Expression comparison of the 3 *cTGFBR2* transcript isoforms in various tissues of fat and lean lines of NEAUHLF. Abbreviations: AF, abdominal fat; SF, subcutaneous fat; GF, gizzard fat; H, heart; Li, liver; Sp, spleen; Lu, lung; K, kidney; G, gizzard; P, proventriculus; Pa, pancreas; M, muscle.

et al., 1995; Saier Jr, 2016; Nava Rodrigues et al., 2019; Xu et al., 2022). For example, a recent study showed the crest phenotype in domestic chicken is caused by a 197 bp duplication in the intron of *HOXC10* (Li et al., 2021). Many eukaryotic proteins contain tandem domain repeats (Light et al., 2012), which are often caused by intragenic duplication (Björklund et al., 2010). In the present study, we demonstrated that the intragenic duplication resulted in a tandem repeat of 2 tandem ecT-betaR2 domains in *cTGFB2*. Protein domain repeats have 2 modes of evolution: 1) the conservative mode, whereby repeats occur before species differentiation and thus the same repeat structure is maintained within different species (Hughes, 2000); 2) the concerted mode, whereby repeats are homogenized within species by independent duplication after species differentiation (Thomas et al., 1997). In the present study, we demonstrated that the intragenic tandem duplication of the *TGFB2* gene is chicken-specific. We reason that intragenic tandem duplication of *cTGFB2* gene occurs after species differentiation, and has evolved in a concerted fashion in chickens.

It has been shown that there are marked species differences in TGF- β signaling pathway (Kruithof et al., 2012). For example, chickens had significantly higher TGF- β signaling activity in lower jaw skeleton than ducks (Smith et al., 2022). Chickens and mice exhibited distinct TGF- β ligand expression patterns during the epithelial to mesenchymal cell transformation (Azhar et al., 2003; Molin et al., 2003) and had contradictory requirements for the TGF- β ligands and receptors during the endocardial-to-mesenchymal cell transformation (Jiao et al., 2006; Yamagishi et al., 2012). In the present study, we provided evidence that *TGFB2*, the core receptor of the TGF- β signaling pathway, differs in structure between chickens and other species. Our finding might partially explain the differences in TGF- β signaling pathway activity between chickens and other species.

APA is one of the important post-transcriptional mechanisms and plays an important regulatory role in many biological processes (Nourse et al., 2020). PAS is divided into CR-PAS and UTR-PAS, of which UTR-PAS is the most common (Chen et al., 2017). UTR-PAS generates various transcript isoforms with different 3' UTR lengths (Jambhekar and Derisi, 2007). CR-PAS causes genes to encode different C-terminal protein isoforms, and some of the isoforms may lack functional domains (Tian and Manley, 2017). In the present study, we for the first time demonstrated that *cTGFB2* gene undergoes APA and that *cTGFB2* gene possesses both UTR-PAS and CR-PAS. Similarly, chicken growth hormone receptor (*GHR*) gene has been shown to have both CR-PAS and UTR-PAS (Lau et al., 2007). *TGFB2* is known to play vital roles in multiple developmental processes. Considering the importance of APA in posttranscriptional regulation, it is essential to explore the roles and molecular mechanisms of APA in *cTGFB2*.

Tissue expression profiling analysis showed that the *cTGFB2-1* was ubiquitously expressed in all the tested tissues of the 4-wk-old AA broiler chickens and the 7-

wk-old broiler chickens of NEAUHAL, which was similar to the tissue expression pattern of human *TGFB2* (Fagerberg et al., 2014), suggesting that *cTGFB2-1* and human *TGFB2* may have similar functions in tissue development. In the present study, our results showed that *cTGFB2-1* in 10 tested tissues, *cTGFB2-2* in 9 tested tissues, and *cTGFB2-3* in all the tested tissues, displayed significantly differential expression levels between the fat and lean lines of NEAUHAL, suggesting that these 3 *cTGFB2* transcript isoforms may contribute to various trait differences between the fat and lean lines of NEAUHAL.

Intriguingly, in the present study, we found that the expression of these *cTGFB2* transcript isoforms in the abdominal fat and gizzard fat was significantly higher in the fat line than in the lean line (Figure 7C). It has been known that activation of TGF- β signaling pathway inhibits adipogenesis, and the expression of adipogenesis-related genes such as *PPAR γ* (Feng and Derynck, 2005; Hill, 2009; Li and Wu, 2020). *TGFB2* knock-down in hASCs cells significantly increased the expression of *PPAR γ* and *C/EBP α* and promoted adipogenesis (Kim et al., 2009). Our previous study found that *PPAR γ* mRNA expression in chicken abdominal fat was significantly higher in the fat line than in the lean line at 7-wk-old broiler chickens of NEAUHAL (Sun et al., 2014). Based on our previous study, and the inhibitory role of TGF- β signaling pathway in mammalian adipogenesis and adipogenesis-related genes expression, we supposed that the expression of the 3 *cTGFB2* transcript isoforms in abdominal fat would be lower in the fat line than in the lean line. However, our expression results contradicted this supposition. This unexpected result may be explained by 2 reasons. First, it has been shown that *TGFB2* gene expression was promoted by activation of PI3K/Akt signaling pathway (Budi et al., 2015), and that PI3K/Akt pathway activity was higher in obese individuals than in normal individuals (Chi et al., 2017). Hence, PI3K/Akt signaling pathway activity in the abdominal fat may be higher in the fat line than in the lean line, leading to increased expression of *cTGFB2* transcript isoforms in the fat line relative to the lean line. Second, adipose tissue *TGFB2* expression may be differentially regulated between chickens and mammals.

In conclusion, we defined *cTGFB2* genomic structure, characterized *cTGFB2* transcript and protein isoforms, and identified an intragenic tandem duplication within *cTGFB2*. To the best of our knowledge, this is the first report of the intragenic tandem duplication within *TGFB2* gene. Our findings pave the way for further investigating the roles and regulation of *cTGFB2* gene.

ACKNOWLEDGMENTS

This work was supported by National 973 Project of China (No.2009CB941604), China National Natural Science Foundation Grant (No.31872346), and the China

Agriculture Research System of MOF and MARA (No. CARS-41).

DISCLOSURES

We declare no conflicts of interest.

SUPPLEMENTARY MATERIALS

Supplementary material associated with this article can be found in the online version at doi:10.1016/j.psj.2022.102169.

REFERENCES

- Azhar, M., J. E. J. Schultz, I. Grupp, G. W. Dorn, P. Meneton, D. G. M. Molin, A. C. Gittenberger-de Groot, and T. Doetschman. 2003. Transforming growth factor *beta* in cardiovascular development and function. *Cytokine. Growth. Factor. Rev.* 14:391–407.
- Bertolio, M. S., A. La Colla, A. Carrea, A. Romo, G. Canziani, S. M. Echarte, S. Campisano, G. P. Barletta, A. M. Monzon, T. M. Rodriguez, A. N. Chisari, and R. A. Dewey. 2021. A novel splice variant of human TGF- β type II receptor encodes a soluble protein and its Fc-Tagged version prevents liver fibrosis *in vivo*. *Front. Cell. Dev. Biol.* 9:690397.
- Björklund, A. K., S. Light, R. Sagit, and A. Elofsson. 2010. Nebulin: a study of protein repeat evolution. *J. Mol. Biol.* 402:38–51.
- Brahmachari, S. K., G. Meera, P. S. Sarkar, P. Balagurumoorthy, J. Tripathi, S. Raghavan, U. Shaligram, and S. Pataskar. 1995. Simple repetitive sequences in the genome: structure and functional significance. *Electrophoresis* 16:1705–1714.
- Budi, E. H., B. P. Muthusamy, and R. Derynck. 2015. The insulin response integrates increased TGF- β signaling through Akt-induced enhancement of cell surface delivery of TGF- β receptors. *Sci. Signal.* 8:ra96.
- Chen, W., Q. Jia, Y. Song, H. Fu, G. Wei, and T. Ni. 2017. Alternative polyadenylation: methods, findings, and impacts. *Genom. Proteom. Bioinf.* 15:287–300.
- Chi, Y., J. Li, N. Li, Z. Chen, L. Ma, W. Peng, X. Pan, M. Li, W. Yu, X. He, B. Geng, Q. Cui, Y. Liu, and J. Yang. 2017. FAM3A enhances adipogenesis of 3T3-L1 preadipocytes via activation of ATP-P2 receptor-Akt signaling pathway. *Oncotarget* 8:45862–45873.
- Cooley, J. R., T. A. Yatskevych, and P. B. Antin. 2014. Embryonic expression of the transforming growth factor beta ligand and receptor genes in chicken. *Dev. Dyn.* 243:497–508.
- Dumbrava, M. G., J. L. Lacañale, C. J. Rowan, and N. D. Rosenblum. 2021. Transforming growth factor beta signaling functions during mammalian kidney development. *Pediatr. Nephrol.* 36:1663–1672.
- Fagerberg, L., B. M. Hallstrom, P. Oksvold, C. Kampf, D. Djureinovic, J. Odeberg, M. Habuka, S. Tahmasebpour, A. Danielsson, K. Edlund, A. Asplund, E. Sjostedt, E. Lundberg, C. A. Szgyarto, M. Skogs, J. O. Takanen, H. Berling, H. Tegel, J. Mulder, P. Nilsson, J. M. Schwenk, C. Lindskog, F. Danielsson, A. Mardinoglu, A. Sivertsson, K. von Feilitzen, M. Forsberg, M. Zwahlen, I. Olsson, S. Navani, M. Huss, J. Nielsen, F. Ponten, and M. Uhlen. 2014. Analysis of the human tissue-specific expression by genome-wide integration of transcriptomics and antibody-based proteomics. *Mol. Cell. Proteomics.* 13:397–406.
- Feng, X. H., and R. Derynck. 2005. Specificity and versatility in TGF- β signaling through smads. *Annu. Rev. Cell. Dev. Biol.* 21:659–693.
- Hill, C. S. 2009. Nucleocytoplasmic shuttling of smad proteins. *Cell. Res.* 19:36–46.
- Hirai, R., and T. Fujita. 1996. A human transforming growth factor- β type II receptor that contains an insertion in the extracellular domain. *Exp. Cell. Res.* 223:135–141.
- Hughes, A. L. 2000. Modes of evolution in the protease and kringle domains of the plasminogen-prothrombin family. *Mol. Phylogenet. Evol.* 14:469–478.
- Jambhekar, A., and J. L. Derisi. 2007. *Cis*-acting determinants of asymmetric, cytoplasmic RNA transport. *RNA* 13:625–642.
- James, J. M., A. Nalbandian, and Y. S. Mukoyama. 2013. TGF β signaling is required for sprouting lymphangiogenesis during lymphatic network development in the skin. *Development* 140:3903–3914.
- Jiao, K., M. Langworthy, L. Batts, C. B. Brown, H. L. Moses, and H. S. Baldwin. 2006. Tgf β signaling is required for atrioventricular cushion mesenchyme remodeling during *in vivo* cardiac development. *Development* 133:4585–4593.
- Kim, Y. J., S. J. Hwang, Y. C. Bae, and J. S. Jung. 2009. MiR-21 regulates adipogenic differentiation through the modulation of TGF- β signaling in mesenchymal stem cells derived from human adipose tissue. *Stem. Cells.* 27:3093–3102.
- Krishnaveni, M. S., J. L. Hansen, W. Seeger, R. E. Morty, S. P. Sheikh, and O. Eickelberg. 2006. Constitutive homo- and hetero-oligomerization of T β R β II-B, an alternatively spliced variant of the mouse TGF- β type II receptor. *Biochem. Biophys. Res. Commun.* 351:651–657.
- Kruithof, B. P. T., S. N. Duim, A. T. Moerkamp, and M. J. Goumans. 2012. TGF β and BMP signaling in cardiac cushion formation: lessons from mice and chicken. *Differentiation* 84:89–102.
- Lau, J. S., C. W. Yip, K. M. Law, and F. C. Leung. 2007. Cloning and characterization of chicken growth hormone binding protein (cGHBP). *Domest. Anim. Endocrin.* 33:107–121.
- Li, J., M. O. Lee, B. W. Davis, P. Wu, S. M. Hsieh Li, C. M. Chuong, and L. Andersson. 2021. The crest phenotype in domestic chicken is caused by a 197 bp duplication in the intron of *HOXC10*. *G3 (Bethesda)* 11:jkaa048.
- Li, S. N., and J. F. Wu. 2020. TGF- β /SMAD signaling regulation of mesenchymal stem cells in adipocyte commitment. *Stem. Cell. Res. Ther.* 11:41.
- Light, S., R. Sagit, S. S. Ithychanda, J. Qin, and A. Elofsson. 2012. The evolution of filamin-A protein domain repeat perspective. *J. Struct. Biol.* 179:289–298.
- Lönn, P., A. Morén, E. Raja, M. Dahl, and A. Moustakas. 2009. Regulating the stability of TGF β receptors and smads. *Cell. Res.* 19:21–35.
- Molin, D. G., U. Bartram, K. Van der Heiden, L. Van Iperen, C. P. Speer, B. P. Hierck, R. E. Poelmann, and A. C. Gittenberger-de-Groot. 2003. Expression patterns of *Tgfb1-3* associate with myocardialisation of the outflow tract and the development of the epicardium and the fibrous heart skeleton. *Dev. Dyn.* 227:431–444.
- Moustakas, A., and C. H. Heldin. 2009. The regulation of TGF β signal transduction. *Development* 136:3699–3714.
- Nava Rodrigues, D., N. Casiraghi, A. Romanel, M. Crespo, S. Miranda, P. Rescigno, I. Figueiredo, R. Riisnaes, S. Carreira, S. Sumanasuriya, P. Gasperini, A. Sharp, J. Mateo, A. Makay, C. McNair, M. Schiewer, K. Knudsen, G. Boysen, F. Demichelis, and J. S. de Bono. 2019. RB1 heterogeneity in advanced metastatic castration-resistant prostate cancer. *Clin. Cancer. Res.* 25:687–697.
- Nourse, J., S. Spada, and S. Danckwardt. 2020. Emerging roles of RNA 3'-end cleavage and polyadenylation in pathogenesis, diagnosis and therapy of human disorders. *Biomolecules* 10:915.
- Pardali, E., M. J. Goumans, and P. ten Dijke. 2010. Signaling by members of the TGF- β family in vascular morphogenesis and disease. *Trends. Cell. Biol.* 20:556–567.
- Peters, S. B., Y. Wang, and R. Serra. 2017. Tgfb2 is required in Osterix expressing cells for postnatal skeletal development. *Bone* 97:54–64.
- Saier, M. H. Jr 2016. Transport protein evolution deduced from analysis of sequence, topology and structure. *Curr. Opin. Struct. Biol.* 38:9–17.
- Smith, S. S., D. Chu, T. Qu, J. A. Aggleton, and R. A. Schneider. 2022. Species-specific sensitivity to TGF β signaling and changes to the Mmp13 promoter underlie avian jaw development and evolution. *eLife* 11:e66005.
- Sun, Y. N., Y. Gao, S. P. Qiao, S. Z. Wang, K. Duan, Y. X. Wang, H. Li, and N. Wang. 2014. Epigenetic DNA methylation in the

- promoters of *peroxisome proliferator-activated receptor γ* in chicken lines divergently selected for fatness. *J. Anim. Sci.* 92:48–53.
- Thomas, G. H., E. C. Newbern, C. C. Korte, M. A. Bales, S. V. Muse, A. G. Clark, and D. P. Kiehart. 1997. Intragenic duplication and divergence in the spectrin superfamily of proteins. *Mol. Biol. Evol.* 14:1285–1295.
- Tian, B., and J. L. Manley. 2017. Alternative polyadenylation of mRNA precursors. *Nat. Rev. Mol. Cell. Biol.* 18:18–30.
- Vander Ark, A., J. Cao, and X. Li. 2018. TGF- β receptors: in and beyond TGF- β signaling. *Cell. Signal.* 52:112–120.
- Xu, F., E. Aref-Eshghi, J. Wu, J. Schubert, G. Wertheim, T. Bhatti, J. Pogoriler, M. Patel, K. Cao, A. Long, Z. Fan, E. H. Denenberg, E. A. Fanning, D. M. Wilmoth, M. Luo, L. K. Conlin, A. S. Dain, K. Zelle, S. Baldino, N. Balamuth, S. MacFarland, M. M. Li, and Y. Zhong. 2022. A novel *TP53* tandem duplication in a child with Li-Fraumeni syndrome. *Cold. Spring. Harb. Mol. Case. Stud.* 8: a006181.
- Xu, P., J. Liu, and R. Derynck. 2012. Post-translational regulation of TGF- β receptor and smad signaling. *FEBS. Lett.* 586:1871–1884.
- Yamagishi, T., K. Ando, H. Nakamura, and Y. Nakajima. 2012. Expression of the *Tgfb2* gene during chick embryogenesis. *Anat. Rec. (Hoboken)*. 295:257–267.
- Yuan, J., K. Yi, and L. Yang. 2020. TGFBR2 regulates hedgehog pathway and cervical cancer cell proliferation and migration by mediating SMAD4. *J. Proteome. Res.* 19:3377–3385.