

REVIEW

Open Access

# A systematic review of the role of vitamin insufficiencies and supplementation in COPD

Ioanna G Tsiligianni\*, Thys van der Molen

## Abstract

**Background:** Pulmonary inflammation, oxidants-antioxidants imbalance, as well as innate and adaptive immunity have been proposed as playing a key role in the development of COPD. The role of vitamins, as assessed either by food frequency questionnaires or measured in serum levels, have been reported to improve pulmonary function, reduce exacerbations and improve symptoms. Vitamin supplements have therefore been proposed to be a potentially useful additive to COPD therapy.

**Methods:** A systematic literature review was performed on the association of vitamins and COPD. The role of vitamin supplements in COPD was then evaluated.

**Conclusions:** The results of this review showed that various vitamins (vitamin C, D, E, A, beta and alpha carotene) are associated with improvement in features of COPD such as symptoms, exacerbations and pulmonary function. High vitamin intake would probably reduce the annual decline of FEV1. There were no studies that showed benefit from vitamin supplementation in improved symptoms, decreased hospitalization or pulmonary function.

## Introduction

COPD in all stages of severity is a very prevalent disease and a great burden for patients and society [1]. In affluent countries COPD is related to smoking over a long period of time, whereas in many other countries it is also related to indoor and outdoor air pollution [1]. The pathology of chronic obstructive pulmonary disease include pulmonary inflammation, oxidants-antioxidants imbalance, protease-antiprotease imbalance, and both innate and adaptive immunity [2,3]. Smoking cessation has been proven to be effective in stopping further deterioration of pulmonary function, reducing symptoms and improving overall health [4]. Smoking cessation however, seems to have only limited influence on the inflammatory process that is associated with COPD. This inflammatory process is probably initiated by oxidative stress and forms the basis of the pathophysiology of COPD [5-8]. Thus the inflammatory process that is associated with COPD seems to be triggered by noxious gasses such as smoking and serious indoor or outdoor air pollution. Oxidative stress caused by these noxious gasses at the level of the epithelium of the bronchial

tree might have play a key role in this inflammatory process. It is therefore possible that anti oxidant therapy or an intensive anti oxidant diet could have an influence on the inflammatory process and the progression of COPD. Over the last two decades a number of studies have suggested that COPD risk is associated with vitamins that all have antioxidant properties and with an anti oxidant diet. Low diet-intake of vitamins has been reported to reduce natural defenses and increase the possibility of airway inflammation [9]. Furthermore, a higher intake of fruits and vegetables was associated with a lower risk of COPD, lower mortality and an improvement of spirometric values [10-17].

When levels of vitamins were measured in the serum they were found to be significantly lower in COPD patients than in control subjects [18]. The association of vitamins with pulmonary diseases is further supported by a meta-analysis of 40 studies in patients with asthma. This meta-analysis revealed that relatively low dietary intake of vitamins A and C were associated with statistically significant increased odds of asthma and wheezing [19].

A large number of studies and reviews highlight an association of vitamins with lung function in healthy subjects and COPD patients [20-27]. Recently a randomized controlled trial suggested that a dietary shift to

\* Correspondence: i.tsiligianni@med.umcg.nl  
Department of General Practice, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

higher antioxidant food intake was associated with improvement in lung function [25]. Furthermore, several studies associate vitamins with a reduction in symptoms, respiratory infections and exacerbations [28-37].

Although for vitamin D the role in respiratory diseases has been clarified through its implication in immunity, for most other vitamins the mechanism of action is less clear [38-42]. Indeed we know that 1,25-dihydroxyvitamin D stimulates both innate and adaptive immunity, in addition to mineralization and calcium homeostasis. Further support on the important role of Vitamin D is given by the fact that it regulates genes that are implicated in apoptosis and cellular proliferation [39], known to be an important step in COPD pathogenesis [42]. Vitamin D has both immunomodulatory and antiinflammatory properties [43]. For vitamin A, B, C, and E, studies highlight their role in COPD risk as well as their connection with COPD outcomes such as symptoms and improvement in spirometric values, without a clear mechanism of action. This article aims to update the knowledge we have about the association between vitamin intake and COPD in outcome measures, as well as to assess the potential role of vitamin supplements.

## Methods

A systematic literature search was performed from 1989 until June 2010 in Pubmed, Embase and Cochrane Collaboration containing the following keywords: COPD in conjunction with smoking, gene polymorphisms, vitamins, FEV1, vitamin C, vitamin E, vitamin D, vitamin A, b-carotene, and vitamin supplements. Further articles were identified from the reference lists of the included articles. In order to be as accurate as possible we included in the present review only studies that have measured serum vitamin levels or used validated food frequency questionnaires to assess the role of vitamins.

Other dietary factors -cured meats, fish, whole grains and alcohol- that have been reported to be associated with the risk of chronic obstructive pulmonary disease or with an increase in symptoms were not included in this review [44-51]. Further we did not examine the influence of caloric intake on COPD. Weight loss and muscle wasting, considered complications of COPD strongly associated with diet, are also not included in the review.

## Results

### Methods of assessing vitamin status: which is best?

The literature reports two essentially different ways to measure vitamins: Serum levels and Food Frequency Questionnaires (FFQ). Both measures have their advantages and disadvantages.

#### Serum levels of vitamins

The assessment of serum levels of vitamins can have the advantage of being more objective than patient's

reported intake. However, serum level assessment of vitamins has the disadvantage that they represent the more recent intake, and for some vitamins such as vitamin C, the levels in peripheral blood are not representative of intake and do not change accordingly [19,52].

#### Food Frequency questionnaire

Food frequency questionnaires on the other hand present a large heterogeneity with differences in assessing periods (from 1 day to two years), number of items on food questionnaires (ranging from 44 to 350) and use of portion size questions [53]. Further, it has been suggested that FFQs do not always detect weak associations [54]. Regarding vitamin C intake large differences were found between FFQs that used portion size questions instead of using standard portions [53]. Another problem regarding FFQ use is that it is difficult to determine which particular vitamin is associated with COPD and if it is the vitamins in fruits and vegetables that are associated with COPD, or another confounding nutrient. Resveratrol for example, a phenolic antioxidant is present in many fruits and is associated with anti-inflammatory activity [55,56]. Therefore it is not clear if in some cases, the vitamins have the beneficial effect, or other nutrients such as resveratrol with antioxidant and anti-inflammatory properties.

Another important obstacle in FFQ is that an assessment of vitamin D from food intake would lead to an incorrect estimation in for example Mediterranean countries where there is a high skin synthesis because of the sunlight.

We found 14 references [31,57-69] that assessed the relationship between a vitamin rich diet as assessed by a FFQ and the subsequent improvement of spirometric values and symptoms. Twelve studies [32,33,35, 52,63,70-76] measured serum levels of vitamins. For the above mentioned reasons, and in order to have a more precise overview, we decided to include both FFQ and serum level studies.

### Food intake patterns: Effect of vitamins on the risk of developing COPD and associated mortality

Varraso et al in a study of 72,043 women identified 754 cases of newly diagnosed COPD [44]. In this study a healthy diet (fruit, vegetables, fish, whole-grain products) was compared with a Western diet (refined grains, cured and red meats, desserts, French fries). The healthy diet was associated with a lower risk of COPD [44]. This could be considered to be due to the overall diet, or indicate a possible positive effect of vitamins on COPD risk, as fruits are considered sources rich with vitamins. From the same author another study comparing the same patterns of diet showed the same results in 111 self-reported cases of newly diagnosed COPD in men [46]. Celik et al used a food frequency questionnaire and found that the

consumption of fruits and vegetables was significantly lower in COPD patients compared to the control group [77]. Fruit intake was related to a lower 25-year incidence of chronic bronchitis and emphysema [57] as well as spirometry improvement [78]. A recent randomized controlled trial has shown that a dietary shift to more anti-oxidant foods such as fruits and vegetables is associated with improvement in lung function [25].

These studies showed a protective role of vitamins against COPD but did not measure vitamins in serum or with a FFQ. Therefore conclusions about the special role of any vitamin in COPD could not be obtained.

### Vitamin D and COPD

Vitamin D is extremely important for the human body. It has a significant role in bone mineralization, in calcium and phosphorous absorption, and is important in the immune system [39,40]. Its most important role however is in bone structure development and bone turn over, as a low vitamin D level is directly associated with osteoporosis. The main sources of vitamin D are skin synthesis and diet. The precursor form is 7-dehydrocholesterol which with UVB is transformed to vitamin D3. Vitamin D3 is transported via the D-binding protein (DBP) to the liver where with hydroxylation reactions transforms to 25(OH)D3 and which is transported again by DBP to the kidneys where it takes its active form of 1,25(OH)2D3. 25OHD3 can also be transformed in 1,25(OH)2D3 in the immune cells [42]. Some studies showed a DBP Gc-1F allele presence that was higher in COPD patients [79,80] and Schellenberg et al found that the Gc2 homozygous genotype was protective for COPD [81]. Other polymorphisms associated with vitamin D binding protein gene are related to clinical differences in families with alpha-1-antitrypsin deficiency [82].

COPD is characterized by inflammation induced by macrophages and neutrophils (innate immunity). COPD is considered a disease where proinflammatory cytokines are increased and has a Th2 response with a predominance of CD8 lymphocytes (adaptive immunity). 1,25-dihydroxyvitamin D stimulates innate immunity probably due to activation of cathelicidin (antimicrobial peptides) to enhance the bacterial killing via Toll-like receptors [83,84]. Vitamin D receptors (VDR) are present in various cells of both innate (i.e. macrophages) and adaptive immunity (i.e. T and B cells). Vitamin D is able to modulate both types of immunity therefore minimizing inflammation [85]. Vitamin D in general is involved in modulating cellular proliferation, suppressing TH cells, [86], downregulating cytokines such as IL-2 [87], as well as in the inhibition of dendritic cells [88], all of which are known to be important in the COPD pathway. Regarding respiratory function, vitamin D plays a significant role in airway remodeling through the inhibition of TNFa and enhancement of IL-10 in immune cells [39]. Vitamin D also seems to play a role as an alternative treatment

strategy to reverse glucocorticoid resistance through its ability to restore IL-10 response [89]. This is important since glucocorticoid resistance is a pivotal barrier to the anti inflammatory treatment of COPD.

Patients with COPD have an increased prevalence of osteoporosis (from 9-69%) and osteopenia (from 27-67%) [90-93]. Malnutrition and low vitamin D levels could be a cause of this higher prevalence [91,94]. The majority of COPD patients have vitamin D deficiency [39,41,95-97] therefore vitamin D supplementation in patients with COPD has been proposed [40].

Black et al reported that higher vitamin D levels were associated with better lung function [72]. In this study that used cross-sectional data from the Third National Health and Nutrition Examination Survey 14.091 people aged >20 years were included. The mean difference between the highest and the lowest quintile of 25-hydroxyvitamin D serum concentration was 126 ml in FEV 1, and 172 ml for FVC after adjustment for factors that affect lung function (age, gender, smoking, etc) [72] (Table 1).

Vitamin D insufficiency has been reported to be associated with an increased incidence of chronic respiratory infections [29,33-35]. There are some studies that also suggest that low serum 25-hydroxy vitamin D levels are associated with upper and lower respiratory tract infection [33-35]. In one large cross sectional study with 18.883 participants, this association was stronger in COPD patients [33] (Table 1). Ebstein Barr virus infection, which is often found in COPD patients, is also associated with low levels of vitamin D [98,99]. Liou et al reported a relation between Toll-like receptors, external triggers and vitamin D-mediated innate immunity, and suggested that differences in the ability of human populations to produce vitamin D may contribute to susceptibility to microbial infections [100].

Finally, Vitamin D could play an important role as an antioxidant therapy, not only for the significant improvement in spirometric values, but also because it has been proposed as a novel treatment to cachexia and sarcopenia in COPD patients [101].

### Vitamin C and E

The role of vitamin C (also known as ascorbate or L-ascorbic acid) in the human body is essential. It has antioxidant properties, is involved in various metabolic reactions, and some studies report it also plays a role in the immune system [102,103]. It is considered important for the maintenance of the connective tissue and bone remodeling [102]. Vitamin E has antioxidant properties as well, and has been reported to have a protective role in the prevention of atherosclerosis and carcinogenesis [104].

In one study that included 3 European Countries a trend ( $P < 0.05$ ) of lower COPD mortality was observed with

**Table 1 Studies connecting spirometric values or incidence of respiratory infections with Vitamin D**

Vitamin D-Ref	No of participants	FFQ or plasma levels	Results
[33]	18,883	Plasma levels	Lower 25(OH)D levels were independently associated with recent URTI (odds ratio [OR], 1.36-1.24). The association between 25(OH)D level and URTI was stronger in patients with chronic obstructive pulmonary disease odds ratio; 2.26.
[35]	800	Plasma levels	Subjects with serum 25(OH)D concentrations < 40 nmol/L (n = 24) had significantly ( $P = 0.004$ ) more days of absence from duty due to respiratory infection (median: 4; quartile 1-quartile 3: 2-6) than did control subjects (2; 0-4; n = 628; incidence rate ratio 1.63; 95% CI: 1.15, 2.24).
[72]	14,091	Plasma levels	The mean difference between the highest and the lowest quintile of 25-hydroxyvitamin D serum concentration was 126 mL (SE:22 mL) in FEV 1, and 172 mL (SE:22 mL) for FVC.

vitamin E intake, while no trend was found with vitamin C after adjustment for age, smoking and country [16]. Higher levels of vitamin C and E in both serum and FFQ in healthy subjects were associated with an increase in FEV1 and FVC [31,32,58-66,69,70]. More details are depicted in Table 2. In one study an increase of 20 micro-mol/Lt in plasma vitamin C concentration was associated with a 13% reduction in the risk of developing obstructive airway disease OR: 0.87 (CI:0.77-0.98) [75].

Studies regarding the role of vitamin C and E in respiratory symptoms showed that low levels were associated with more wheezing, phlegm production and dyspnea [28,31,32,36,37]. Tug et al found both vitamin E and vitamin A levels were significantly lower during exacerbations of COPD than in patients with stable COPD [30].

Takkouche et al, in 1667 cases of the common cold in the general population suggested that intake of vitamin C and zinc was not related to the occurrence of common cold [105]. Nevertheless vitamin C decreases the duration of common cold symptoms which might be important in patients with COPD [106].

### Vitamin A and B

Vitamin A (retinol and carotens) plays an important role in several functions of the human body including vision, bone and skin health, and further has an innate antioxidant activity. Vitamin B is involved in various steps of metabolism and enhances immunity. High levels of vitamin A, b-carotene and/or alpha-carotene were associated with increase in FEV1 and FVC in most of the studies [31,32,63,65,66,69-71,76] although there are some exceptions [57,61,68]. More details are presented in Table 2. High serum beta carotene levels in a general population sample of 523 subjects were associated with the expression of a gene polymorphism that connected with a slower FEV1 decline [107].

Hirayama et al reported that the highest level of intake of vitamin A resulted in a 52% ( $p = 0.008$ ) reduction in COPD risk [108] while in another study the risk for COPD was associated with lower levels of plasma vitamin A ( $p < 0.01$ )[108].

Fimognari et al reported lower levels of folate and vitamin B12 in COPD patients, resulted in an increased

**Table 2 Studies connecting Vitamin C, E, A, alpha and beta-carotene with spirometric values improvement**

Vitamin	FFQ studies	Plasma levels studies	Improvement in spirometric values	No association with spirometric values
Vit C	31,58,59,60,61, 62,63,65,66	32,52, 63,69,70	<b>Serum:</b> FEV1 improvement in ml from 17-94 ml and FVC improvement from 16.4-94 ml for an SD variation <b>FFQ:</b> FEV1 improvement in ml from 37-53 ml and FVC improvement from 23.3-79 ml for an SD variation	52
Vit E	31,58,59,61, 62,64,65	32,69,70	<b>Serum:</b> An SD increase in plasma levels of vitamin E had a median range of FEV1 increase in ml from 12-59.3 ml <b>FFQ:</b> An SD increase had a median range of FEV1 increase in ml from 20.1-93 ml and for FVC from 23.1 -54 ml, respectively	31,58,61
Vit A	61,68	32,70	32,70	61,68
b-carotene	31,57, 63,65,66,69	32,69, 70, 76	<b>Serum:</b> Improvement in FEV1 ranges from 22-31.2 ml <b>FFQ:</b> Improvement in FEV1 = 60 ml, FVC= 75 ml	57
a-carotene		70,71	<b>Serum:</b> Improvement in FEV1 for one SD increase 23.7 ml <sup>70</sup> . Subjects in the fifth quintile of serum beta-carotene had a 195 ml (95% confidence interval [95% CI]: 40 to 351 ml) higher and those in the fifth quintile of alpha-carotene had a 257 ml (95% CI: 99 to 414 ml) higher FEV(1) compared with subjects in the first quintile of these carotenoids <sup>71</sup> .	

**Table 3 Vitamin supplementation and COPD outcome measured**

Reference	Supplementation	No of patients	Effect
[18]	Supplementation E and C. 10 of 21 COPD patients were given vitamin E (200 UI/day) and vitamin C (500 mg/day) for 1 month.	21 COPD and 10 controls.	The exercise time increased significantly in the 10 COPD patients who were treated (exercise time 6.4+1.8 vs 8.7+2.1 min, p = 0.01). (Bruce protocol-graded treadmill exercise test).
[36]	Supplementation alpha-tocopherol (50 mg/d) and beta-carotene (20 mg/d) supplementation, for 5-8 years.	29,133 people (Cancer prevention study)	The supplementation did not affect the recurrence or incidence of chronic cough, phlegm or dyspnea. Relative risk for the above mentioned symptoms around 1 with or without supplementation.
[111]	Vit E supplementation. 400 IU daily for 12 weeks.	30 COPD patients	Spirometric measurements. Changed not significant either on day 1 or after 12 weeks of vitamin E supplementation.
[112]	Vit E supplementation Patients were divided into two groups: group A- placebo group (n = 14), receiving only standard therapy, and group B- vitamin E-supplemented group (n = 10), receiving 400 IU of vitamin E capsules twice daily in addition to standard therapy, for 8 weeks.	24 COPD patients.	There was a similar degree of lung function and clinical improvement in both groups.
[113]	Vit C and E. Patients were randomly assigned to placebo (n = 8), 400 mg/day vitamin E (E400, n = 9), 200 mg/day vitamin E (E200, n = 9), or 250 mg/day vitamin C (C250, n = 9) for 12 weeks.	35 COPD patients	No improvement in lung function after 12 weeks of supplementation.
[114]	Vit A supplementation for 30 days. (healthy nonsmokers (n = 7), healthy smokers (n = 7), mild chronic obstructive pulmonary disease (COPD-mild) patients (n = 9), COPD-moderate-severe patients (n = 7), and COPD-moderate-severe patients with exacerbation (+ex;n = 6)	36 people-21 COPD n = 6.	Improvement in lung function mean increase for 1-s forced expiratory volume (FEV1) = 22.9% in the COPD-vitamin A group.
[115]	Supplementation 600 mg vitamin E, 250 mg vitamin C, and 20 mg β-carotene daily 5-year treatment period. All participants randomly allocated to receive vitamin supplementation or placebo.	20,536 UK adults (aged 40-80) with coronary disease, other occlusive arterial disease, or diabetes	No significant differences were observed between the treatment groups in forced expiratory volume during one second (FEV1: 2.06 L vitamin-allocated vs 2.06 L placebo-allocated; difference 0.00 L [SE 0.01]) or in forced vital capacity (FVC: 2.83 L vs 2.82 L; difference 0.01 L [SE 0.01]). Nor were significant differences observed in the numbers of participants hospitalised for chronic obstructive pulmonary disease or asthma (149 [1.5%] vs 133 [1.3%]) or for any other non-neoplastic respiratory cause (641 [6.2%] vs 642 [6.3%]).

plasma level of total homocysteine, a known cardiovascular risk factor [109].

Regarding the role of β-carotene in respiratory symptoms, two studies showed a beneficial association with cough [32,36] and one study showed no correlation with symptoms, except for wheezing [31].

#### Vitamin supplementation

Antioxidant supplementation has been proposed to be helpful in patients with COPD as a way to reduce oxidative stress and inflammation, and improve spirometric values [24]. Multivitamin supplementation has been reported to be popular for patients with COPD especially among older patients [110].

Seven studies reported the effect of vitamin supplements on several outcomes of COPD [18,36,111-115]. All these studies showed a large heterogeneity regarding which vitamins had been supplemented, the dosage and

the duration of the vitamin supplementation, ranging from 4 weeks to 5 years [111-115]. Secondly, different outcomes were measured such as spirometric values, symptoms and exercise capacity. A randomized controlled trial in high-risk individuals for cardiovascular events that received antioxidant vitamins (vitamin C, E and β-carotene) supplementation for 5 years failed to identify any improvement in 5-year mortality and in spirometric values or hospitalization due to COPD. However this study excluded patients with severe COPD [115]. More details are depicted in Table 3. Also a cohort study of 77,719 participants using multi-vitamin supplements was not related to the total mortality [116].

Little is known regarding the prevention of upper respiratory tract infections after supplementation of Vitamin D although some studies report a trend for improvement [117] but some others do not confirm that (ranges of OR = 0.77-0.95) [118,119].

## Conclusion

The results of this review show that intake of various vitamins are associated with improvement in features of COPD such as symptoms, exacerbations and pulmonary function. Increased vitamin intake could probably reduce the annual decline of FEV1. Although the mechanisms behind these effects are often not clear, this might open possibilities to develop drugs that modify or prevent COPD. Dietary interventions directed towards high vitamin intake might be an additional approach towards COPD management. Although there are many studies that associate vitamins with improvement in lung function tests, there is no clear evidence of the benefit of vitamin supplements. Most studies regarding supplements showed no benefit of multivitamin supplementation in symptoms, spirometric function or hospitalization for COPD.

This review suggests that future work is needed with prospective randomized controlled trials, that would explore the role of vitamins as well as the effectiveness of vitamin supplements on outcomes such as symptoms, spirometric values, health status, risk of development of COPD and exacerbations rates.

### Authors' contributions

Both authors (IGT, TvdM) wrote and revised the manuscript, and approved the final version.

### Competing interests

The authors declare that they have no competing interests.

Received: 18 July 2010 Accepted: 6 December 2010

Published: 6 December 2010

### References

- Global Initiative for Chronic Obstructive Lung Disease. [http://www.goldcopd.com], (assessed June 2010).
- Saetta M, Turato G, Maestrelli P, Mapp CE, Fabbri LM: Cellular and structural bases of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2001, **163**:1304-1309.
- Grashoff WF, Sont JK, Sterk PJ, Hiemstra PS, de Boer WI, Stolk J, Han J, van Krieken JM: Chronic obstructive pulmonary disease: role of bronchiolar mast cells and macrophages. *Am J Pathol* 1997, **151**:1785-1790.
- Willemsen BW, Postma DS, Timens W, ten Hacken NH: The impact of smoking cessation on respiratory symptoms, lung function, airway hyperresponsiveness and inflammation. *Eur Respir J* 2004, **23**(3):464-76.
- Rahman I, Adcock IM: Oxidative stress and redox regulation of lung inflammation in COPD. *Eur Respir J* 2006, **28**(1):219-42.
- Siafakas NM, Tzortzaki EG: Few smokers develop COPD. Why? *Respir Med* 2002, **96**(8):615-24.
- Brigham KL: Role of free radicals in lung injury. *Chest* 1986, **89**(6):859-63.
- Sanguinetti CM: Oxidant/antioxidant imbalance: role in the pathogenesis of COPD. *Respiration* 1992, **59**:Suppl 1:20-3.
- Heffner JE, Repine JE: Pulmonary strategies of antioxidant defense. *Am Rev Respir Dis* 1989, **140**(2):531-54.
- Brug J, Schols A, Mesters I: Dietary change, nutrition education and chronic obstructive pulmonary disease. *Patient Educ Couns* 2004, **52**(3):249-57.
- Watson L, Margetts B, Howarth P, Dorward M, Thompson R, Little P: The association between diet and chronic obstructive pulmonary disease in subjects selected from general practice. *Eur Respir J* 2002, **20**(2):313-8.
- Tabak C, Feskens EJ, Heederik D, Kromhout D, Menotti A, Blackburn HW: Fruit and fish consumption: a possible explanation for population differences in COPD mortality (The Seven Countries Study). *Eur J Clin Nutr* 1998, **52**(11):819-25.
- Tabak C, Arts IC, Smit HA, Heederik D, Kromhout D: Chronic obstructive pulmonary disease and intake of catechins, flavonols, and flavones: the MORGAN Study. *Am J Respir Crit Care Med* 2001, **164**(1):61-4.
- Denny SL, Thompson RL, Margetts BM: Dietary factors in the pathogenesis of asthma and chronic obstructive pulmonary disease. *Curr Allergy Asthma Rep* 2003, **3**(2):130-6.
- Carey IM, Strachan DP, Cook DG: Effects of changes in fresh fruit consumption on ventilatory function in healthy British adults. *Am J Respir Crit Care Med* 1998, **158**(3):728-33.
- Walda IC, Tabak C, Smit HA, Räsänen L, Fidanza F, Menotti A, Nissinen A, Feskens EJ, Kromhout D: Diet and 20-year chronic obstructive pulmonary disease mortality in middle-aged men from three European countries. *Eur J Clin Nutr* 2002, **56**(7):638-43.
- Strachan DP, Cox BD, Erzindoglu SW, Walters DE, Whichelow MJ: Ventilatory function and winter fresh fruit consumption in a random sample of British adults. *Thorax* 1991, **46**(9):624-9.
- Agacikten A, Basigit I, Ozden M, Yildiz F, Ural D, Maral H, Boyaci H, Ilgazli A, Komsuoglu B: The effects of antioxidants on exercise-induced lipid peroxidation in patients with COPD. *Respirology* 2004, **9**(1):38-42.
- Allen S, Britton JR, Leonardi-Bee JA: Association between antioxidant vitamins and asthma outcome measures: systematic review and meta-analysis. *Thorax* 2009, **64**(7):610-9.
- Schünemann HJ, Freudenheim JL, Grant BJ: Epidemiologic evidence linking antioxidant vitamins to pulmonary function and airway obstruction. *Epidemiol Rev* 2001, **23**(2):248-67.
- Sridhar MK: Nutrition and lung health. *Proc Nutr Soc* 1999, **58**(2):303-8.
- Smit HA: Chronic obstructive pulmonary disease, asthma and protective effects of food intake: from hypothesis to evidence? *Respir Res* 2001, **2**(5):261-4.
- Smit HA, Grievink L, Tabak C: Dietary influences on chronic obstructive lung disease and asthma: a review of the epidemiological evidence. *Proc Nutr Soc* 1999, **58**(2):309-19.
- Romieu I, Trenga C: Diet and obstructive lung diseases. *Epidemiol Rev* 2001, **23**(2):268-87.
- Keranis E, Makris D, Rodopoulou P, Martinou H, Papamakarios G, Daniil Z, Zintzaras E, Gourgoulianis KI: Impact of dietary shift to higher antioxidant foods in COPD: A Randomized Trial. *Eur Respir J* 2010.
- Schünemann HJ, Grant BJ, Freudenheim JL, Muti P, Browne RW, Drake JA, Klocke RA, Trevisan M: The relation of serum levels of antioxidant vitamins C and E, retinol and carotenoids with pulmonary function in the general population. *Am J Respir Crit Care Med* 2001, **163**(5):1246-55.
- Kelly FJ: Vitamins and respiratory disease: antioxidant micronutrients in pulmonary health and disease. *Proc Nutr Soc* 2005, **64**(4):510-26.
- Schwartz J, Weiss ST: Dietary factors and their relation to respiratory symptoms. The Second National Health and Nutrition Examination Survey. *Am J Epidemiol* 1990, **132**(1):67-76.
- Zasloff M: Fighting infections with vitamin D. *Nat Med* 2006, **12**(4):388-90.
- Tug T, Karatas F, Terzi SM: Antioxidant vitamins (A, C and E) and malondialdehyde levels in acute exacerbation and stable periods of patients with chronic obstructive pulmonary disease. *Clin Invest Med* 2004, **27**(3):123-8.
- Grievink L, Smit HA, Ocké MC, van 't Veer P, Kromhout D: Dietary intake of antioxidant (pro)-vitamins, respiratory symptoms and pulmonary function: the MORGAN study. *Thorax* 1998, **53**(3):166-71.
- Kelly Y, Sacker A, Marmot M: Nutrition and respiratory health in adults: findings from the health survey for Scotland. *Eur Respir J* 2003, **21**(4):664-71.
- Ginde AA, Mansbach JM, Camargo CA Jr: Association between serum 25-hydroxyvitamin D level and upper respiratory tract infection in the Third National Health and Nutrition Examination Survey. *Arch Intern Med* 2009, **169**(4):384-90.
- Karatekin G, Kaya A, Salihoglu O, Balci H, Nuhoglu A: Association of subclinical vitamin D deficiency in newborns with acute lower respiratory infection and their mothers. *Eur J Clin Nutr* 2009, **63**(4):473-7.
- Laaksi I, Ruohola JP, Tuohimaa P, Auvinen A, Haataja R, Pihlajamäki H, Ylikomi T: An association of serum vitamin D concentrations < 40 nmol/L with acute respiratory tract infection in young Finnish men. *Am J Clin Nutr* 2007, **86**(3):714-7.

36. Rautalahti M, Virtamo J, Haukka J, Heinonen OP, Sundvall J, Albanes D, Huttunen JK: **The effect of alpha-tocopherol and beta-carotene supplementation on COPD symptoms.** *Am J Respir Crit Care Med* 1997, **156**(5):1447-52.
37. Bodner C, Godden D, Brown K, Little J, Ross S, Seaton A: **Antioxidant intake and adult-onset wheeze: a case-control study.** Aberdeen WHEASE Study Group. *Eur Respir J* 1999, **13**(1):22-30.
38. Chishimba L, Thickett DR, Stockley RA, Wood AM: **The vitamin D axis in the lung: a key role for vitamin D-binding protein.** *Thorax* 2010, **65**(5):456-62.
39. Hughes DA, Norton R: **Vitamin D and respiratory health.** *Clin Exp Immunol* 2009, **158**(1):20-5.
40. Janssens W, Lehocka A, Carremans C, Bouillon R, Mathieu C, Decramer M: **Vitamin D beyond bones in chronic obstructive pulmonary disease: time to act.** *Am J Respir Crit Care Med* 2009, **179**(8):630-6.
41. Wright RJ: **Make no bones about it: increasing epidemiologic evidence links vitamin D to pulmonary function and COPD.** *Chest* 2005, **128**(6):3781-3.
42. Holick MF: **Vitamin D deficiency.** *N Engl J Med* 2007, **357**(3):266-81.
43. Adams JS, Hewison M: **Unexpected actions of vitamin D: new perspectives on the regulation of innate and adaptive immunity.** *Nat Clin Pract Endocrinol Metab* 2008, **4**(2):80-90.
44. Varraso R, Fung TT, Barr RG, Hu FB, Willett W, Camargo CA Jr: **Prospective study of dietary patterns and chronic obstructive pulmonary disease among US women.** *Am J Clin Nutr* 2007, **86**(2):488-95.
45. Tabak C, Smit HA, Heederik D, Ocké MC, Kromhout D: **Diet and chronic obstructive pulmonary disease: independent beneficial effects of fruits, whole grains, and alcohol (the MORGEN study).** *Clin Exp Allergy* 2001, **31**(5):747-55.
46. Varraso R, Fung TT, Hu FB, Willett W, Camargo CA: **Prospective study of dietary patterns and chronic obstructive pulmonary disease among US men.** *Thorax* 2007, **62**(9):786-91.
47. Varraso R, Jiang R, Barr RG, Willett WC, Camargo CA Jr: **Prospective study of cured meats consumption and risk of chronic obstructive pulmonary disease in men.** *Am J Epidemiol* 2007, **166**(12):1438-45.
48. Varraso R, Fung TT, Barr RG, Hu FB, Willett W, Camargo CA Jr: **Prospective study of dietary patterns and chronic obstructive pulmonary disease among US women.** *Am J Clin Nutr* 2007, **86**(2):488-95.
49. Jiang R, Paik DC, Hankinson JL, Barr RG: **Cured meat consumption, lung function, and chronic obstructive pulmonary disease among United States adults.** *Am J Respir Crit Care Med* 2007, **175**(8):798-804.
50. Butler LM, Koh WP, Lee HP, Tseng M, Yu MC, London SJ: **Singapore Chinese Health Study. Prospective study of dietary patterns and persistent cough with phlegm among Chinese Singaporeans.** *Am J Respir Crit Care Med* 2006, **173**(3):264-70.
51. Butler LM, Koh WP, Lee HP, Yu MC, London SJ: **Dietary fiber and reduced cough with phlegm: a cohort study in Singapore.** *Am J Respir Crit Care Med* 2004, **170**(3):279-87.
52. Cook DG, Carey IM, Whincup PH, Papacosta O, Chirico S, Bruckdorfer KR, Walker M: **Effect of fresh fruit consumption on lung function and wheeze in children.** *Thorax* 1997, **52**(7):628-33.
53. Molag ML, de Vries JH, Ocké MC, Dagnelie PC, van den Brandt PA, Jansen MC, van Staveren WA, van't Veer P: **Design characteristics of food frequency questionnaires in relation to their validity.** *Am J Epidemiol* 2007, **166**(12):1468-78.
54. Schatzkin A, Kipnis V, Carroll RJ, Midthune D, Subar AF, Bingham S, Schoeller DA, Troiano RP, Freedman LS: **A comparison of a food frequency questionnaire with a 24-hour recall for use in an epidemiological cohort study: results from the biomarker-based Observing Protein and Energy Nutrition (OPEN) study.** *Int J Epidemiol* 2003, **32**(6):1054-62.
55. Olas B, Wachowicz B: **Resveratrol, a phenolic antioxidant with effects on blood platelet functions.** *Platelets* 2005, **16**(5):251-60.
56. Sharafkhaneh A, Velamuri S, Badmaev V, Lan C, Hanania N: **The potential role of natural agents in treatment of airway inflammation.** *Ther Adv Respir Dis* 2007, **1**(2):105-20.
57. Miedema I, Feskens EJ, Heederik D, Kromhout D: **Dietary determinants of long-term incidence of chronic nonspecific lung diseases. The Zutphen Study.** *Am J Epidemiol* 1993, **138**(1):37-45.
58. Morabia A, Sorenson A, Kumanyika SK, Abbey H, Cohen BH, Chee E: **Vitamin A, cigarette smoking, and airway obstruction.** *Am Rev Respir Dis* 1989, **140**(5):1312-6.
59. Britton JR, Pavord ID, Richards KA, Knox AJ, Wisniewski AF, Lewis SA, Tattersfield AE, Weiss ST: **Dietary antioxidant vitamin intake and lung function in the general population.** *Am J Respir Crit Care Med* 1995, **151**(5):1383-7.
60. Ness AR, Khaw KT, Bingham S, Day NE: **Vitamin C status and respiratory function.** *Eur J Clin Nutr* 1996, **50**(9):573-9.
61. McKeever TM, Scrivener S, Broadfield E, Jones Z, Britton J, Lewis SA: **Prospective study of diet and decline in lung function in a general population.** *Am J Respir Crit Care Med* 2002, **165**(9):1299-303.
62. Dow L, Tracey M, Villar A, Coggon D, Margetts BM, Campbell MJ, Holgate ST: **Does dietary intake of vitamins C and E influence lung function in older people?** *Am J Respir Crit Care Med* 1996, **154**(5):1401-4.
63. Hu G, Zhang X, Chen J, Peto R, Campbell TC, Cassano PA: **Dietary vitamin C intake and lung function in rural China.** *Am J Epidemiol* 1998, **148**(6):594-9.
64. Butland BK, Fehily AM, Elwood PC: **Diet, lung function, and lung function decline in a cohort of 2512 middle aged men.** *Thorax* 2000, **55**(2):102-8.
65. Tabak C, Smit HA, Räsänen L, Fidanza F, Menotti A, Nissinen A, Feskens EJM, Heederik D, Kromhout D: **Dietary factors and pulmonary function: a cross-sectional study in middle aged men from three European countries.** *Thorax* 1999, **54**:1021-1026.
66. Ochs-Balcom HM, Grant BJ, Muti P, Sempos CT, Freudenheim JL, Browne RW, McCann SE, Trevisan M, Cassano PA, Iacoviello L, Schünemann HJ: **Antioxidants, oxidative stress, and pulmonary function in individuals diagnosed with asthma or COPD.** *Eur J Clin Nutr* 2006, **60**(8):991-9.
67. Schwartz J, Weiss ST: **Relationship between dietary vitamin C intake and pulmonary function in the First National Health and Nutrition Examination Survey (NHANES I).** *Am J Clin Nutr* 1994, **59**(1):110-4.
68. Shahar E, Folsom AR, Melnick SL, Tockman MS, Comstock GW, Shimakawa T, Higgins MW, Sorlie PD, Szkoł M: **Does dietary vitamin A protect against airway obstruction? The Atherosclerosis Risk in Communities (ARIC) Study Investigators.** *Am J Respir Crit Care Med* 1994, **150**(4):978-82.
69. Hu G, Cassano PA: **Antioxidant nutrients and pulmonary function: the Third National Health and Nutrition Examination Survey (NHANES III).** *Am J Epidemiol* 2000, **151**:975-981.
70. McKeever TM, Lewis SA, Smit HA, Burney P, Cassano PA, Britton J: **A multivariate analysis of serum nutrient levels and lung function.** *Respir Res* 2008, **9**:67.
71. Grievink L, de Waart FG, Schouten EG, Kok FJ: **Serum carotenoids, alpha-tocopherol, and lung function among Dutch elderly.** *Am J Respir Crit Care Med* 2000, **161**(3 Pt 1):790-5.
72. Black PN, Scragg R: **Relationship between serum 25-hydroxyvitamin d and pulmonary function in the third national health and nutrition examination survey.** *Chest* 2005, **128**(6):3792-8.
73. Pearson P, Britton J, McKeever T, Lewis SA, Weiss S, Pavord I, Fogarty A: **Lung function and blood levels of copper, selenium, vitamin C and vitamin E in the general population.** *Eur J Clin Nutr* 2005, **59**(9):1043-8.
74. Førli L, Pedersen JL, Bjørtoft O, Blomhoff R, Kofstad J, Boe J: **Vitamins A and E in serum in relation to weight and lung function in patients with advanced pulmonary disease.** *Int J Vitam Nutr Res* 2002, **72**(6):360-8.
75. Sergeant LA, Jaeckel A, Wareham NJ: **Interaction of vitamin C with the relation between smoking and obstructive airways disease in EPIC Norfolk. European Prospective Investigation into Cancer and Nutrition.** *Eur Respir J* 2000, **16**(3):397-403.
76. Grievink L, Smit HA, Veer P, Brunekreef B, Kromhout D: **Plasma concentrations of the antioxidants beta-carotene and alpha-tocopherol in relation to lung function.** *Eur J Clin Nutr* 1999, **53**(10):813-7.
77. Celik F, Topcu F: **Nutritional risk factors for the development of chronic obstructive pulmonary disease (COPD) in male smokers.** *Clin Nutr* 2006, **25**(6):955-61.
78. Strachan DP, Cox BD, Erzinclioglu SW, Walters DE, Whichelow MJ: **Ventilatory function and winter fresh fruit consumption in a random sample of British adults.** *Thorax* 1991, **46**(9):624-9.
79. Ito I, Nagai S, Hoshino Y, Muro S, Hirai T, Tsukino M, Mishima M: **Risk and severity of COPD is associated with the group-specific component of serum globulin 1F allele.** *Chest* 2004, **125**(1):63-70.
80. Ishii T, Keicho N, Teramoto S, Azuma A, Kudoh S, Fukuchi Y, Ouchi Y, Matsuse T: **Association of Gc-globulin variation with susceptibility to COPD and diffuse panbronchiolitis.** *Eur Respir J* 2001, **18**(5):753-7.

81. Schellenberg D, Paré PD, Weir TD, Spinelli JJ, Walker BA, Sandford AJ: Vitamin D binding protein variants and the risk of COPD. *Am J Respir Crit Care Med* 1998, **157**(3 Pt 1):957-61.
82. Wood AM, Needham M, Simmonds MJ, Newby PR, Gough SC, Stockley RA: Phenotypic differences in alpha 1 antitrypsin-deficient sibling pairs may relate to genetic variation. *COPD* 2008, **5**(6):353-9.
83. Yim S, Dhawan P, Ragunath C, Christakos S, Diamond G: Induction of cathelicidin in normal and CF bronchial epithelial cells by 1,25-dihydroxyvitamin D(3). *J Cyst Fibros* 2007, **6**(6):403-10.
84. Trinchieri G, Sher A: Cooperation of Toll-like receptor signals in innate immune defence. *Nat Rev Immunol* 2007, **7**(3):179-90.
85. Bhalla AK, Amento EP, Clemens TL, Holick MF, Krane SM: Specific high-affinity receptors for 1,25-dihydroxyvitamin D3 in human peripheral blood mononuclear cells: presence in monocytes and induction in T lymphocytes following activation. *J Clin Endocrinol Metab* 1983, **57**(6):1308-10.
86. Chen S, Sims GP, Chen XX, Gu YY, Chen S, Lipsky PE: Modulatory effects of 1,25-dihydroxyvitamin D3 on human B cell differentiation. *J Immunol* 2007, **179**(3):1634-47.
87. Lemire JM, Adams JS, Kermani-Arab V, Bakke AC, Sakai R, Jordan SC: 1,25-Dihydroxyvitamin D3 suppresses human T helper/inducer lymphocyte activity in vitro. *J Immunol* 1985, **134**(5):3032-5.
88. Penna G, Adorini L: 1, Alpha,25-dihydroxyvitamin D3 inhibits differentiation, maturation, activation, and survival of dendritic cells leading to impaired alloreactive T cell activation. *J Immunol* 2000, **164**(5):2405-11.
89. Barnes PJ, Adcock IM: Glucocorticoid resistance in inflammatory diseases. *Lancet* 2009, **373**(9678):1905-17.
90. Sin DD, Man JP, Man SF: The risk of osteoporosis in Caucasian men and women with obstructive airways disease. *Am J Med* 2003, **114**(1):10-4.
91. Incalzi RA, Caradonna P, Ranieri P, Basso S, Fuso L, Pagano F, Ciappi G, Pistelli R: Correlates of osteoporosis in chronic obstructive pulmonary disease. *Respir Med* 2000, **94**(11):1079-84.
92. Ferguson GT, Calverley PM, Anderson JA, Jenkins CR, Jones PW, Willits LR, Yates JC, Vestbo J, Celli B: Prevalence and Progression of Osteoporosis in Patients With COPD. Results From TORCH. *Chest* 2009, **136**(6):1456-65.
93. Graat-Verboom L, Wouters EF, Smeenk FW, van den Borne BE, Lunde R, Spruit MA: Current status of research on osteoporosis in COPD: a systematic review. *Eur Respir J* 2009, **34**(1):209-18.
94. Franco CB, Paz-Filho G, Gomes PE, Nascimento VB, Kulak CA, Boguszewski CL, Borba VZ: Chronic obstructive pulmonary disease is associated with osteoporosis and low levels of vitamin D. *Osteoporos Int* 2009, **20**(11):1881-7.
95. Riancho JA, González Macías J, Del Arco C, Amado JA, Freijanes J, Antón MA: Vertebral compression fractures and mineral metabolism in chronic obstructive lung disease. *Thorax* 1987, **42**(12):962-6.
96. Shane E, Silverberg SJ, Donovan D, Papadopoulos A, Staron RB, Addesso V, Jorgesen B, McGregor C, Schulman L: Osteoporosis in lung transplantation candidates with end-stage pulmonary disease. *Am J Med* 1996, **101**(3):262-9.
97. de Batle J, Romieu I, Antó JM, Méndez M, Rodríguez E, Balcells E, Ferrer A, Gea J, Rodriguez-Roisin R, García-Aymerich J: The PAC-COPD Study Group. Dietary habits of firstly admitted Spanish COPD patients. *Respir Med* 2009, **103**(12):1904-10.
98. McManus TE, Marley AM, Baxter N, Christie SN, Elborn JS, O'Neill HJ, Coyle PV, Kidney JC: High levels of Epstein-Barr virus in COPD. *Eur Respir J* 2008, **31**(6):1221-6.
99. Grant WB: A possible role for Epstein-Barr virus infection in COPD? *Eur Respir J* 2008, **32**(5):1412-3.
100. Liu PT, Stenger S, Li H, Wenzel L, Tan BH, Krutzik SR, Ochoa MT, Schaubert J, Wu K, Meinken C, Kamen DL, Wagner M, Bals R, Steinmeyer A, Zügel U, Gallo RL, Eisenberg D, Hewison M, Hollis BW, Adams JS, Bloom BR, Modlin RL: Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response. *Science* 2006, **311**(5768):1770-3.
101. Kung T, Springer J, Doehner W, Anker SD, von Haehling S: Novel treatment approaches to cachexia and sarcopenia: highlights from the 5th Cachexia Conference. *Expert Opin Investig Drugs* 2010, **19**(4):579-85.
102. Deruelle F, Baron B: Vitamin C: is supplementation necessary for optimal health? *J Altern Complement Med* 2008, **14**(10):1291-8.
103. Jacob RA: Assessment of human vitamin C status. *J Nutr* 1990, **120**(Suppl 11):1480-5.
104. Dutta A, Dutta SK: Vitamin E and its role in the prevention of atherosclerosis and carcinogenesis: a review. *J Am Coll Nutr* 2003, **22**(4):258-68.
105. Takkouche B, Regueira-Méndez C, García-Closas R, Figueiras A, Gestal-Otero JJ: Intake of vitamin C and zinc and risk of common cold: a cohort study. *Epidemiology* 2002, **13**(1):38-44.
106. Heimer KA, Hart AM, Martin LG, Rubio-Wallace S: Examining the evidence for the use of vitamin C in the prophylaxis and treatment of the common cold. *J Am Acad Nurse Pract* 2009, **21**(5):295-300.
107. Guenegou A, Boczkowski J, Aubier M, Neukirch F, Leynaert B: Interaction between a heme oxygenase-1 gene promoter polymorphism and serum beta-carotene levels on 8-year lung function decline in a general population: the European Community Respiratory Health Survey (France). *Am J Epidemiol* 2008, **167**(2):139-44.
108. Hirayama F, Lee AH, Binns CW, Zhao Y, Hiramatsu T, Tanikawa Y, Nishimura K, Taniguchi H: Do vegetables and fruits reduce the risk of chronic obstructive pulmonary disease? A case-control study in Japan. *Prev Med* 2009, **49**(2-3):184-9.
109. Fimognari FL, Loffredo L, Di Simone S, Sampietro F, Pastorelli R, Monaldo M, Violi F, D'Angelo A: Hyperhomocysteinaemia and poor vitamin B status in chronic obstructive pulmonary disease. *Nutr Metab Cardiovasc Dis* 2009, **19**(9):654-9.
110. Hirayama F, Lee AH, Binns CW, Taniguchi H: Dietary supplementation by Japanese patients with chronic obstructive pulmonary disease. *Complement Ther Med* 2009, **17**(1):37-43.
111. Daga MK, Chhabra R, Sharma B, Mishra TK: Effects of exogenous vitamin E supplementation on the levels of oxidants and antioxidants in chronic obstructive pulmonary disease. *J Biosci* 2003, **28**(1):7-11.
112. Nadeem A, Raj HG, Chhabra SK: Effect of vitamin E supplementation with standard treatment on oxidant-antioxidant status in chronic obstructive pulmonary disease. *Indian J Med Res* 2008, **128**(6):705-11.
113. Wu TC, Huang YC, Hsu SY, Wang YC, Yeh SL: Vitamin E and vitamin C supplementation in patients with chronic obstructive pulmonary disease. *Int J Vitam Nutr Res* 2007, **77**(4):272-9.
114. Paiva SA, Godoy I, Vannucchi H, Fávaro RM, Geraldo RR, Campana AO: Assessment of vitamin A status in chronic obstructive pulmonary disease patients and healthy smokers. *Am J Clin Nutr* 1996, **64**(6):928-34.
115. Heart Protection Study Collaborative Group. MRC/BHF Heart Protection: Study of antioxidant vitamin supplementation in 20,536 high-risk individuals: a randomised placebo-controlled trial. *Lancet* 2002, **360**(9326):23-33.
116. Pocobelli G, Peters U, Kristal AR, White E: Use of supplements of multivitamins, vitamin C, and vitamin E in relation to mortality. *Am J Epidemiol* 2009, **170**(4):472-83.
117. Aloia JF, Li-Ng M: Re: epidemic influenza and vitamin D. *Epidemiol Infect* 2007, **135**(7):1095-6; author reply 1097-8.
118. Liu BA, McGeer A, McArthur MA, Simor AE, Aghdassi E, Davis L, Allard JP: Effect of multivitamin and mineral supplementation on episodes of infection in nursing home residents: a randomized, placebo-controlled study. *J Am Geriatr Soc* 2007, **55**(1):35-42.
119. Graat JM, Schouten EG, Kok FJ: Effects of daily vitamin E and multivitamin-mineral supplementation on acute respiratory tract infections in elderly persons: a randomized controlled trial. *JAMA* 2002, **288**(6):715-721.

doi:10.1186/1465-9921-11-171

**Cite this article as:** Tsiliogianni and van der Molen: A systematic review of the role of vitamin insufficiencies and supplementation in COPD. *Respiratory Research* 2010 11:171.