



ISSN 1600-5368

OPEN ACCESS

Crystal structure of ethyl 2-([(4Z)-3,5-dioxo-1-phenylpyrazolidin-4-ylidene]methyl]amino)acetate

Shaaban K. Mohamed,^{a,b} Mehmet Akkurt,^c Joel T. Mague,^d Eman A. Ahmed^e and Mustafa R. Albayati^{f,*}

^aChemistry and Environmental Division, Manchester Metropolitan University, Manchester M1 5GD, England, ^bChemistry Department, Faculty of Science, Minia University, 61519 El-Minia, Egypt, ^cDepartment of Physics, Faculty of Sciences, Erciyes University, 38039 Kayseri, Turkey, ^dDepartment of Chemistry, Tulane University, New Orleans, LA 70118, USA, ^eChemistry Department, Faculty of Science, Sohag University, 82524 Sohag, Egypt, and ^fKirkuk University, College of Science, Department of Chemistry, Kirkuk, Iraq. *Correspondence e-mail: shaabankamel@yahoo.com

Received 18 July 2014; accepted 20 July 2014

Edited by H. Stoeckli-Evans, University of Neuchâtel, Switzerland

The title compound, C₁₄H₁₅N₃O₄, is nearly planar, the dihedral angle between the planes of the phenyl and pyrazolidine rings being 1.13 (7) Å, and that between the plane of the pyrazolidine ring and the mean plane of the side chain [C–N–C–C(=O)–O; r.m.s. deviation = 0.024 Å] being 2.52 (7)°. This is due in large part to the presence of the intramolecular N–H···O and C–H···O hydrogen bonds. In the crystal, pairwise N–H···O hydrogen bonds form inversion dimers, which are further associated into layers, lying very close to plane (1̄20), via pairwise C–H···O hydrogen bonds. The layers are then weakly connected through C–H···O hydrogen bonds, forming a three-dimensional structure.


Keywords: crystal structure; hydrogen bonding; hydrogen-bonded dimers; pyrazolidine-3,5-dione; aminoacetic acid ester.

CCDC reference: 1015152

1. Related literature

For the synthesis of compounds containing the pyrazolidinone nucleus and their biological activity, see: Ismail *et al.* (2012); Khodairy (2007); Khloya *et al.* (2013). For biologically active synthetic heterocyclic compounds containing the pyrazol-5(4H)-one core scaffold and displaying some interesting pharmaceutical properties, see: Uramaru *et al.* (2010) for analgesic; Thaker *et al.* (2011) and Chande *et al.* (2007) for antimicrobial; Mariappan *et al.* (2010) and Nishikimi *et al.*

(2012) for anti-inflammatory; Chen *et al.* (2012) for cytotoxicity.



2. Experimental

2.1. Crystal data

C₁₄H₁₅N₃O₄
M_r = 289.29
 Triclinic, *P*1̄
a = 5.4984 (1) Å
b = 7.3585 (2) Å
c = 16.6265 (4) Å
 α = 91.3290 (9)°
 β = 97.325 (1)°
 γ = 99.562 (1)°
V = 657.27 (3) Å³
Z = 2
 Cu *K* α radiation
 μ = 0.91 mm⁻¹
T = 100 K
 0.27 × 0.09 × 0.04 mm

2.2. Data collection

Bruker D8 VENTURE PHOTON
 100 CMOS diffractometer
 Absorption correction: multi-scan
 (*SADABS*; Bruker, 2013)
T_{min} = 0.91, *T_{max}* = 0.96
 5108 measured reflections
 2450 independent reflections
 2171 reflections with *I* > 2σ(*I*)
R_{int} = 0.019

2.3. Refinement

R[*F*² > 2σ(*F*²)] = 0.033
wR(*F*²) = 0.089
S = 1.02
 2450 reflections
 199 parameters
 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max}$ = 0.22 e Å⁻³
 $\Delta\rho_{\min}$ = -0.19 e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> –H··· <i>A</i>	<i>D</i> –H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> –H··· <i>A</i>
N3–H3A···O1	0.914 (18)	2.250 (17)	2.9062 (14)	128.3 (13)
C6–H6···O2	0.95	2.23	2.8833 (16)	126
N1–H1···O1 ⁱ	0.88 (2)	1.89 (2)	2.7573 (14)	170.7 (16)
C2–H2···O1 ⁱ	0.95	2.36	3.2777 (15)	162
C10–H10···O2 ⁱⁱ	0.95	2.28	3.1268 (16)	148
C11–H11A···O2 ⁱⁱ	0.99	2.58	3.1498 (15)	117
C11–H11B···O1 ⁱⁱⁱ	0.99	2.51	3.3386 (15)	142

Symmetry codes: (i) $-x + 2, -y + 2, -z + 1$; (ii) $-x, -y + 1, -z + 1$; (iii) $x - 1, y, z$.

Data collection: *APEX2* (Bruker, 2013); cell refinement: *SAINT* (Bruker, 2013); data reduction: *SAINT*; program(s) used to solve structure: *SHELXT* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg & Putz, 2012); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

Acknowledgements

The support of NSF–MRI grant No. 1228232 for the purchase of the diffractometer is gratefully acknowledged.

Supporting information for this paper is available from the IUCr electronic archives (Reference: SU2759).

References

- Brandenburg, K. & Putz, H. (2012). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2013). *APEX2, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chande, M. S., Barve, P. A. & Suryanarayan, V. (2007). *J. Heterocycl. Chem.* **44**, 49–53.
- Chen, T., Benmohamed, R., Kim, J., Smith, K., Amanta, D., Morimoto, R. I., Kirsch, D. R., Ferrante, R. J. & Silverman, R. B. (2012). *J. Med. Chem.* **55**, 515–527.
- Ismail, M. A. H., Abou El Ella, D. A., Abouzid, K. A. M. & Jaballah, M. (2012). *IJPSR* **3**, 3746–3757.
- Khloya, P., Kumar, P., Mittal, A., Aggarwal, N. K. & Sharma, P. K. (2013). *Org. Med. Chem. Lett.* **3**, 1–7.
- Khodairy, A. (2007). *J. Chin. Chem. Soc.* **54**, 93–102.
- Mariappan, G., Saha, B. P., Satharson, L. & Haldar, A. (2010). *Indian J. Chem. Sect. B*, **49**, 1671–1674.
- Nishikimi, A., Takehito Uruno, T., Duan, X., Cao, Q., Okamura, Y., Saitoh, T., Saito, N., Sakaoka, S., Du, Y., Suenaga, A., Kukimoto-Niino, M., Miyano, K., Gotoh, K., Okabe, T., Sanematsu, F., Tanaka, Y., Sumimoto, H., Honma, T., Yokoyama, S., Nagano, T., Kohda, D., Kanai, M. & Fukui, Y. (2012). *Chem. Biol.* **19**, 488–497.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Thaker, K. M., Ghetiya, R. M., Tala, S. D., Dodiya, B. L., Joshi, K. A., Dubal, K. L. & Joshi, H. S. (2011). *Indian J. Chem. Sect. B*, **50**, 738–744.
- Uramaru, N., Shigematsu, H., Toda, A., Eyanagi, R., Kitamura, S. & Ohta, S. (2010). *J. Med. Chem.* **53**, 8727–8733.

supporting information

Acta Cryst. (2014). E70, o938–o939 [doi:10.1107/S1600536814016766]

Crystal structure of ethyl 2-({[(4*Z*)-3,5-dioxo-1-phenylpyrazolidin-4-ylidene]methyl}amino)acetate

Shaaban K. Mohamed, Mehmet Akkurt, Joel T. Mague, Eman A. Ahmed and Mustafa R. Albayati

S1. Experimental

A mixture of 1 mmol (231 mg) of (4*Z*)-4-[(dimethylamino)methylene]-1-phenylpyrazolidine-3,5-dione and 1 mmol (140 mg) of ethyl aminoacetate hydrochloride and few drops of triethylamine (TEA) as a catalyst in 30 ml 1,4-dioxane was refluxed for 6 h. On cooling the solid product deposited, was filtered off, washed with cold ethanol and dried under vacuum. Crystals of the title compound were obtained as yellow needles by recrystallization of the crude product from dimethyl sulfoxide; *M.p.* 489–491 K.

S2. Refinement

The N-bound H atoms were located in a Fourier difference map and freely refined. The C-bound H atoms were placed in calculated positions and treated as riding atoms: C—H = 0.95 - 0.99 Å with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C-methyl})$ and $= 1.2U_{\text{eq}}(\text{C})$ for other H atoms. The crystal did not diffract well, hence the lower than desirable value for θ_{full} .






Figure 1

The molecular structure of the title molecule, with atom labelling. Displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**

The crystal packing viewed along the *b* axis of the title compound. N—H···O and C—H···O hydrogen bonds are shown, respectively, as purple and black dotted lines (see Table 1 for details).

**Figure 3**

The crystal packing of the title compound, showing the layer structure and the weak C—H···O interlayer hydrogen bonds (black dotted lines; see Table 1 for details).

Ethyl 2-([{(4*Z*)-3,5-dioxo-1-phenylpyrazolidin-4-ylidene]methyl}amino)acetate

Crystal data

$C_{14}H_{15}N_3O_4$

$M_r = 289.29$

Triclinic, $P\bar{1}$

$a = 5.4984$ (1) Å

$b = 7.3585$ (2) Å

$c = 16.6265$ (4) Å

$\alpha = 91.3290$ (9)°

$\beta = 97.325$ (1)°

$\gamma = 99.562$ (1)°

$V = 657.27$ (3) Å³

$Z = 2$

$F(000) = 304$

$D_x = 1.462 \text{ Mg m}^{-3}$
 Cu $K\alpha$ radiation, $\lambda = 1.54178 \text{ \AA}$
 Cell parameters from 3908 reflections
 $\theta = 2.7\text{--}72.2^\circ$

$\mu = 0.91 \text{ mm}^{-1}$
 $T = 100 \text{ K}$
 Needle, yellow
 $0.27 \times 0.09 \times 0.04 \text{ mm}$

Data collection

Bruker D8 VENTURE PHOTON 100 CMOS
 diffractometer
 Radiation source: INCOATEC $I\mu\text{S}$ micro-focus
 source
 Mirror monochromator
 Detector resolution: $10.4167 \text{ pixels mm}^{-1}$
 ω scans
 Absorption correction: multi-scan
 (SADABS; Bruker, 2013)

$T_{\min} = 0.91, T_{\max} = 0.96$
 5108 measured reflections
 2450 independent reflections
 2171 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.019$
 $\theta_{\max} = 72.2^\circ, \theta_{\min} = 5.4^\circ$
 $h = -6 \rightarrow 6$
 $k = -9 \rightarrow 8$
 $l = -20 \rightarrow 20$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.033$
 $wR(F^2) = 0.089$
 $S = 1.02$
 2450 reflections
 199 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: mixed
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0471P)^2 + 0.2143P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.22 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.19 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger. H-atoms attached to carbon were placed in calculated positions (C—H = 0.95 - 0.99 Å) and included as riding contributions with isotropic displacement parameters 1.2 - 1.5 times those of the attached carbon atoms. H-atoms attached to nitrogen were refined independently.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.81993 (17)	0.91924 (12)	0.58190 (5)	0.0208 (2)
O2	0.18424 (17)	0.56447 (13)	0.39606 (5)	0.0232 (2)
O3	0.56220 (17)	0.89351 (13)	0.80229 (5)	0.0238 (2)
O4	0.20347 (17)	0.77726 (12)	0.84951 (5)	0.0214 (2)
N1	0.7714 (2)	0.82957 (15)	0.44586 (6)	0.0193 (2)
N2	0.5788 (2)	0.73223 (15)	0.38878 (6)	0.0182 (2)
N3	0.3705 (2)	0.76426 (15)	0.64837 (6)	0.0185 (2)
C1	0.6132 (2)	0.71090 (16)	0.30668 (7)	0.0171 (3)
C2	0.8342 (2)	0.79219 (17)	0.27959 (7)	0.0187 (3)

H2	0.9640	0.8617	0.3167	0.022*
C3	0.8642 (3)	0.77134 (18)	0.19820 (8)	0.0221 (3)
H3	1.0142	0.8281	0.1799	0.026*
C4	0.6776 (3)	0.66856 (18)	0.14350 (8)	0.0224 (3)
H4	0.6992	0.6543	0.0881	0.027*
C5	0.4589 (3)	0.58691 (18)	0.17083 (8)	0.0224 (3)
H5	0.3308	0.5160	0.1336	0.027*
C6	0.4240 (2)	0.60707 (17)	0.25156 (7)	0.0203 (3)
H6	0.2729	0.5509	0.2693	0.024*
C7	0.3762 (2)	0.66257 (17)	0.42850 (7)	0.0177 (3)
C8	0.4474 (2)	0.72955 (16)	0.51201 (7)	0.0173 (3)
C9	0.6925 (2)	0.83530 (17)	0.51974 (7)	0.0175 (3)
C10	0.3003 (2)	0.69795 (17)	0.57301 (7)	0.0172 (3)
H10	0.1399	0.6246	0.5602	0.021*
C11	0.2115 (2)	0.73107 (18)	0.71157 (7)	0.0187 (3)
H11A	0.1573	0.5967	0.7154	0.022*
H11B	0.0614	0.7881	0.6980	0.022*
C12	0.3509 (2)	0.81183 (17)	0.79164 (7)	0.0188 (3)
C13	0.3112 (3)	0.84585 (19)	0.93050 (7)	0.0239 (3)
H13A	0.3525	0.9822	0.9326	0.029*
H13B	0.4654	0.7958	0.9472	0.029*
C14	0.1209 (3)	0.7838 (2)	0.98568 (8)	0.0326 (3)
H14A	-0.0333	0.8293	0.9670	0.049*
H14B	0.1841	0.8327	1.0411	0.049*
H14C	0.0877	0.6487	0.9850	0.049*
H1	0.896 (4)	0.907 (3)	0.4313 (10)	0.034 (5)*
H3A	0.525 (3)	0.834 (2)	0.6622 (10)	0.027 (4)*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0191 (5)	0.0237 (5)	0.0168 (4)	-0.0039 (4)	0.0017 (3)	-0.0013 (3)
O2	0.0169 (5)	0.0300 (5)	0.0190 (4)	-0.0062 (4)	0.0019 (3)	-0.0010 (4)
O3	0.0199 (5)	0.0270 (5)	0.0218 (4)	-0.0024 (4)	0.0014 (4)	-0.0012 (4)
O4	0.0214 (5)	0.0253 (5)	0.0158 (4)	-0.0016 (4)	0.0035 (3)	-0.0016 (3)
N1	0.0157 (6)	0.0226 (5)	0.0166 (5)	-0.0056 (4)	0.0023 (4)	-0.0017 (4)
N2	0.0154 (5)	0.0210 (5)	0.0158 (5)	-0.0029 (4)	0.0012 (4)	-0.0016 (4)
N3	0.0160 (6)	0.0208 (5)	0.0171 (5)	-0.0018 (4)	0.0028 (4)	0.0003 (4)
C1	0.0190 (6)	0.0162 (6)	0.0163 (6)	0.0036 (5)	0.0020 (5)	0.0012 (5)
C2	0.0182 (6)	0.0179 (6)	0.0193 (6)	0.0010 (5)	0.0023 (5)	0.0005 (5)
C3	0.0220 (7)	0.0228 (6)	0.0222 (6)	0.0034 (5)	0.0062 (5)	0.0028 (5)
C4	0.0275 (7)	0.0245 (7)	0.0161 (6)	0.0062 (6)	0.0037 (5)	0.0009 (5)
C5	0.0230 (7)	0.0236 (7)	0.0189 (6)	0.0022 (5)	-0.0005 (5)	-0.0009 (5)
C6	0.0181 (7)	0.0216 (6)	0.0197 (6)	0.0002 (5)	0.0018 (5)	0.0003 (5)
C7	0.0156 (6)	0.0179 (6)	0.0188 (6)	0.0005 (5)	0.0027 (5)	0.0017 (5)
C8	0.0168 (6)	0.0166 (6)	0.0174 (6)	0.0009 (5)	0.0011 (5)	0.0009 (5)
C9	0.0182 (6)	0.0164 (6)	0.0177 (6)	0.0016 (5)	0.0027 (5)	0.0010 (4)
C10	0.0149 (6)	0.0165 (6)	0.0189 (6)	0.0003 (5)	0.0009 (4)	0.0017 (4)

C11	0.0170 (6)	0.0212 (6)	0.0169 (6)	-0.0005 (5)	0.0036 (5)	0.0006 (5)
C12	0.0193 (7)	0.0177 (6)	0.0189 (6)	0.0022 (5)	0.0022 (5)	0.0015 (5)
C13	0.0270 (7)	0.0268 (7)	0.0157 (6)	0.0006 (6)	0.0008 (5)	-0.0021 (5)
C14	0.0382 (9)	0.0375 (8)	0.0204 (7)	-0.0017 (7)	0.0077 (6)	-0.0002 (6)

Geometric parameters (Å, °)

O1—C9	1.2580 (15)	C4—C5	1.388 (2)
O2—C7	1.2288 (16)	C4—H4	0.9500
O3—C12	1.2052 (17)	C5—C6	1.3879 (18)
O4—C12	1.3378 (16)	C5—H5	0.9500
O4—C13	1.4438 (14)	C6—H6	0.9500
N1—C9	1.3557 (16)	C7—C8	1.4455 (17)
N1—N2	1.4126 (14)	C8—C10	1.3764 (18)
N1—H1	0.88 (2)	C8—C9	1.4289 (18)
N2—C7	1.3975 (17)	C10—H10	0.9500
N2—C1	1.4105 (16)	C11—C12	1.5049 (16)
N3—C10	1.3201 (16)	C11—H11A	0.9900
N3—C11	1.4503 (16)	C11—H11B	0.9900
N3—H3A	0.914 (18)	C13—C14	1.500 (2)
C1—C2	1.3958 (19)	C13—H13A	0.9900
C1—C6	1.4023 (17)	C13—H13B	0.9900
C2—C3	1.3920 (18)	C14—H14A	0.9800
C2—H2	0.9500	C14—H14B	0.9800
C3—C4	1.3873 (18)	C14—H14C	0.9800
C3—H3	0.9500		
C12—O4—C13	115.89 (10)	C10—C8—C9	126.25 (11)
C9—N1—N2	109.72 (10)	C10—C8—C7	125.21 (12)
C9—N1—H1	124.3 (11)	C9—C8—C7	108.54 (11)
N2—N1—H1	122.3 (11)	O1—C9—N1	124.26 (12)
C7—N2—C1	130.04 (10)	O1—C9—C8	128.51 (12)
C7—N2—N1	109.26 (10)	N1—C9—C8	107.22 (11)
C1—N2—N1	120.63 (10)	N3—C10—C8	123.39 (12)
C10—N3—C11	122.52 (11)	N3—C10—H10	118.3
C10—N3—H3A	119.8 (10)	C8—C10—H10	118.3
C11—N3—H3A	117.7 (10)	N3—C11—C12	109.81 (10)
C2—C1—C6	119.51 (11)	N3—C11—H11A	109.7
C2—C1—N2	120.64 (11)	C12—C11—H11A	109.7
C6—C1—N2	119.86 (11)	N3—C11—H11B	109.7
C3—C2—C1	119.92 (12)	C12—C11—H11B	109.7
C3—C2—H2	120.0	H11A—C11—H11B	108.2
C1—C2—H2	120.0	O3—C12—O4	125.26 (12)
C4—C3—C2	120.74 (12)	O3—C12—C11	125.64 (12)
C4—C3—H3	119.6	O4—C12—C11	109.10 (10)
C2—C3—H3	119.6	O4—C13—C14	106.97 (11)
C3—C4—C5	119.13 (12)	O4—C13—H13A	110.3
C3—C4—H4	120.4	C14—C13—H13A	110.3

C5—C4—H4	120.4	O4—C13—H13B	110.3
C4—C5—C6	121.15 (12)	C14—C13—H13B	110.3
C4—C5—H5	119.4	H13A—C13—H13B	108.6
C6—C5—H5	119.4	C13—C14—H14A	109.5
C5—C6—C1	119.55 (12)	C13—C14—H14B	109.5
C5—C6—H6	120.2	H14A—C14—H14B	109.5
C1—C6—H6	120.2	C13—C14—H14C	109.5
O2—C7—N2	124.92 (11)	H14A—C14—H14C	109.5
O2—C7—C8	129.97 (12)	H14B—C14—H14C	109.5
N2—C7—C8	105.10 (10)		
C9—N1—N2—C7	-4.17 (14)	N2—C7—C8—C10	178.32 (11)
C9—N1—N2—C1	178.53 (10)	O2—C7—C8—C9	178.57 (13)
C7—N2—C1—C2	-179.41 (12)	N2—C7—C8—C9	-0.82 (13)
N1—N2—C1—C2	-2.73 (17)	N2—N1—C9—O1	-176.95 (11)
C7—N2—C1—C6	0.56 (19)	N2—N1—C9—C8	3.52 (13)
N1—N2—C1—C6	177.24 (11)	C10—C8—C9—O1	-0.3 (2)
C6—C1—C2—C3	0.66 (18)	C7—C8—C9—O1	178.83 (12)
N2—C1—C2—C3	-179.37 (11)	C10—C8—C9—N1	179.20 (12)
C1—C2—C3—C4	-0.74 (19)	C7—C8—C9—N1	-1.67 (14)
C2—C3—C4—C5	0.28 (19)	C11—N3—C10—C8	179.28 (12)
C3—C4—C5—C6	0.26 (19)	C9—C8—C10—N3	0.2 (2)
C4—C5—C6—C1	-0.33 (19)	C7—C8—C10—N3	-178.81 (11)
C2—C1—C6—C5	-0.13 (19)	C10—N3—C11—C12	175.64 (11)
N2—C1—C6—C5	179.90 (11)	C13—O4—C12—O3	0.30 (18)
C1—N2—C7—O2	0.5 (2)	C13—O4—C12—C11	179.81 (10)
N1—N2—C7—O2	-176.50 (12)	N3—C11—C12—O3	0.61 (18)
C1—N2—C7—C8	179.91 (11)	N3—C11—C12—O4	-178.90 (10)
N1—N2—C7—C8	2.93 (13)	C12—O4—C13—C14	-177.08 (11)
O2—C7—C8—C10	-2.3 (2)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N3—H3A \cdots O1	0.914 (18)	2.250 (17)	2.9062 (14)	128.3 (13)
C6—H6 \cdots O2	0.95	2.23	2.8833 (16)	126
N1—H1 \cdots O1 ⁱ	0.88 (2)	1.89 (2)	2.7573 (14)	170.7 (16)
C2—H2 \cdots O1 ⁱ	0.95	2.36	3.2777 (15)	162
C10—H10 \cdots O2 ⁱⁱ	0.95	2.28	3.1268 (16)	148
C11—H11A \cdots O2 ⁱⁱ	0.99	2.58	3.1498 (15)	117
C11—H11B \cdots O1 ⁱⁱⁱ	0.99	2.51	3.3386 (15)	142

Symmetry codes: (i) $-x+2, -y+2, -z+1$; (ii) $-x, -y+1, -z+1$; (iii) $x-1, y, z$.