

Determinants of Nicotine Dependence in Chronic Obstructive Pulmonary Disease

Yun Su Sim, M.D., Ph.D.¹, Jin Hwa Lee, M.D., Ph.D.², Ki Uk Kim, M.D., Ph.D.³, Seung Won Ra, M.D., Ph.D.⁴, Hye Yun Park, M.D., Ph.D.⁵, Chang-Hoon Lee, M.D., Ph.D.⁶, Deog Kyeom Kim, M.D., Ph.D.⁷, Kyeong-Cheol Shin, M.D., Ph.D.⁸, Sang Haak Lee, M.D., Ph.D.⁹, Hun Gyu Hwang, M.D., Ph.D.¹⁰, Joong Hyun Ahn, M.D., Ph.D.¹¹, Yong Bum Park, M.D.¹², Yu-Il Kim, M.D., Ph.D.¹³, Kwang Ha Yoo, M.D., Ph.D.¹⁴, Ina Jeong, M.D.¹⁵, Yeon-Mok Oh, M.D., Ph.D.¹⁶ and Sang-Do Lee, M.D., Ph.D.¹⁶, on behalf of the KOLD Investigators

*Author affiliations appear at the end of this article.

Background: Smoking cessation is the most powerful intervention to modify progress of chronic obstructive pulmonary disease (COPD), and nicotine dependence is one of the most important determinants of success or failure in smoking cessation. We evaluated nicotine dependence status and investigated factors associated with moderate to high nicotine dependence in patients with COPD.

Methods: We included 53 current smokers with COPD in the Korean Obstructive Lung Disease II cohort enrolled between January 2014 and March 2016. Nicotine dependence was measured by using Fagerstrom test for nicotine dependence (FTND). Cognitive function was assessed by Korean version of Montreal Cognitive Assessment.

Results: The median FTND score was 3, and 32 patients (60%) had moderate to high nicotine dependence. The median smoking amount was 44 pack-years, which was not related to nicotine dependence. Multiple logistic regression analysis revealed that high education status (odds ratio, 1.286; 95% confidence interval, 1.036–1.596; $p=0.023$), age <70 (odds ratio, 6.407; 95% confidence interval, 1.376–29.830; $p=0.018$), and mild to moderate airflow obstruction (odds ratio, 6.969; 95% confidence interval, 1.388–34.998; $p=0.018$) were related to moderate to high nicotine dependence.

Conclusion: Nicotine dependence does not correlate with smoking amount, but with education level, age, and severity of airflow obstruction. Physicians should provide different strategies of smoking cessation intervention for current smokers with COPD according to their education levels, age, and severity of airflow obstruction.

Keywords: Nicotine Dependence; Pulmonary Disease, Chronic Obstructive; Smoking Cessation

Address for correspondence: Jin Hwa Lee, M.D., Ph.D.

Division of Pulmonary and Critical Care Medicine, Department of Internal Medicine, Ewha Womans University School of Medicine, 1071 Anyangcheon-ro, Yangcheon-gu, Seoul 07985, Korea

Phone: 82-2-2650-6007, Fax: 82-2-2655-2076,

E-mail: jinhwalee@ewha.ac.kr

Received: Nov. 30, 2016

Revised: Jan. 31, 2017

Accepted: Mar. 10, 2017

© It is identical to the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>).



Copyright © 2017
The Korean Academy of Tuberculosis and Respiratory Diseases.
All rights reserved.

Introduction

Chronic obstructive pulmonary disease (COPD) is the third leading cause of death worldwide¹ and affects approximately 13.4% of adults aged 40 years and over in Korea². Cigarette smoking is the most important risk factor for development and progress of COPD³. As the amount of cigarette smoke increases, respiratory symptoms become more pronounced and lung function deteriorates³. Therefore, smoking cessation is crucial to prevent the progress of COPD^{1,3,4} and reduces risk of many comorbid conditions such as cardiovascular diseases and lung cancer⁵. Nevertheless, to stop smoking is very difficult, and smokers often fail because of nicotine addiction⁶. Nicotine is a psychoactive agent to regulate stress and/or to

induce relaxed emotions⁷. The intensity of physical addiction to nicotine can be estimated by questionnaire⁸, which is useful for predicting the success rate of smoking cessation and provide a guide for nicotine replacement therapy⁹. It has been known that current smokers with COPD have higher nicotine dependence than current smokers without COPD^{10,11}. However, there were few studies on predictors of moderate to high nicotine dependence in current smokers with COPD^{12,13}.

We evaluated nicotine dependence status and investigated factors associated with moderate to high nicotine dependence in current smokers with COPD in the Korean Obstructive Lung Disease (KOLD) II cohort.

Materials and Methods

1. Patients

All subjects were selected from the KOLD cohort II, which prospectively recruited subjects with obstructive lung disease from the pulmonary clinics of 11 referral hospitals in Korea from January 2014 through March 2016. The inclusion criteria were followings: (1) age ≥ 40 years; (2) post-bronchodilator forced expiratory volume in 1 second (FEV_1)/forced vital capacity (FVC) < 0.7 ; and (3) current smoker, who has smoked greater than 100 cigarettes in his or her lifetime and has smoked in the last 28 days. Patients with severe tuberculosis destroyed lung disease, bronchiectasis or acute lung inflammation were excluded. The study protocol was approved by the institutional review boards of the 11 hospitals.

2. Variables and measures

At the time of inclusion, a standard questionnaire was used to obtain information on smoking history, level of education, and comorbidities. COPD assessment test (CAT) was used for evaluation of COPD impact on health status. The level of dyspnea was assessed by modified Medical Research Council (MMRC) grade.

Spirometry was performed according to the American Thoracic Society/European Respiratory Society guidelines¹⁴ using the Vmax 22 (Sensor Medics, Yorba Linda, CA, USA). Post-bronchodilator spirometry values were measured 15 minutes after inhalation of 400 μg of salbutamol. The spirometry reference values were based on the Korean Choi equation¹⁵.

Severe airflow obstruction is defined as $FEV_1 < 50\%$ predicted.

Patients responded to the Fagerstrom test for nicotine dependence (FTND)⁸. The FTND generates a score based on the sum of the six questionnaire items, which are variably weighted. FTND yields a total score from 0 to 10 points (mild, 0 to 3; moderate, 4 to 6; severe, 7 to 10). Subjects with a score of 4 or higher were defined as having moderate to high nicot-

tine dependence.

Cognitive function was also assessed by Korean version of Montreal Cognitive Assessment (MoCA-K)¹⁶. The MoCA-K is a 30-point test that takes about 10 minutes to administer and consists of 12 individual tasks grouped into cognitive domains including (1) visuospatial/executive functioning, (2) naming, (3) attention, (4) language, (5) abstraction, (6) memory, and (7) orientation. A total score is calculated, and an educational correction is made (one point added for individuals with 12 years of education or less). The cognitive impairment is defined as a total score below 23.

3. Statistical analysis

Descriptive data are expressed as medians with interquartile range, and frequencies are expressed as number (%). A chi-square test was used to compare categorical variables, while continuous variables were compared using the Mann-Whitney U test. A p-value of < 0.05 was regarded as statistically significant. Multiple logistic regression analysis was used to investigate factors associated with moderate to high nicotine dependence including variables, which had p-value of < 0.05 in univariate analysis.

Results

Fifty-three current smokers with COPD were included. Total 268 subjects were selected from the KOLD cohort II and excluded seven non-smokers, 204 ex-smokers, and four non-responders for questionnaire (Figure 1). Clinical characteristics of 53 patients are detailed in Table 1. Fifty-two patients (98%) were male and the median age was 67 years. The median CAT score was 14, and median MMRC dyspnea scale was 1. The median FEV_1/FVC was 48% and the median FEV_1 was 1.69 L (61% predicted). The patients with severe airflow obstruction were 16 (30%). The median smoking amount was 44 pack-years. The median FTND score was 3, and 32 patients (60%) had moderate to high nicotine dependence. The median MoCA-K score for cognitive function was 24, and patients with cognitive dysfunction (MoCA-K < 23) were 24 (45%). Forty-one patients (77%) were using inhalers.

Factors related to moderate to high nicotine dependence were relatively young age (66 years vs. 76 years, $p=0.013$), high education status (12 years vs. 6 years, $p=0.003$), and high cognitive function (24 vs. 22, $p=0.021$). However, there are no differences of MMRC grade, CAT score, FEV_1 , and pack-years of cigarette smoking between two groups according to nicotine dependence (Table 2).

Multiple logistic regression analysis revealed that high education status (odds ratio, 1.286; 95% confidence interval, 1.036–1.596; $p=0.023$), age < 70 (odds ratio, 6.407; 95% confidence interval, 1.376–29.830; $p=0.018$), and mild to moderate

airflow obstruction (odds ratio, 6.969; 95% confidence interval, 1.388–34.998; $p=0.018$) were related to moderate to high nicotine dependence (Table 3).

Discussion

We investigated factors associated with nicotine dependence in current smokers of Korean COPD cohort. High education status was the most significant factor related to nicotine dependence in current smokers with COPD, which is consistent with a previous study¹⁷ in current smokers with diabetes. They reported that nicotine dependence in smokers who had ≥ 8 years of formal education was 2.57 times higher than that in those with less education. On the contrary, three previous studies showed that lower education level was linked with higher nicotine dependence among current smokers^{18–20}. These studies were different in the study subjects, one study was in patients with depressive or anxiety disorders¹⁸, the other study was population-based¹⁹, and another study was in healthy smokers in Korea²⁰. On the other hand, previous studies in current smokers with COPD showed that young age, high anxiety/depression, high smoking amount, low lung function, and emphysematous lung were associated with nicotine dependence^{12,13}. In a study in current smokers with COPD, education level was not related to nicotine dependence¹³. The disparity in the impact of education level on nicotine dependence might be due to differences in the study subjects. Even though the relationship between education levels and nicotine dependence needs to be investigated further, our study indicates that physicians should provide different strategies of smoking cessation intervention for current smokers with COPD according to their education levels. One Korean study about risk factors of high nicotine dependence in male smokers who wanted health checked up showed

that nicotine dependence was related to the score of internal response, the score of external stress, the shorter duration of education, the early age of first smoking²⁰.

In the current study, patients with moderate to high nicotine dependence showed younger age < 70 . It is consistent with results in previous studies: two studies included COPD

Table 1. Clinical characteristics of current smokers with chronic obstructive pulmonary disease

Variable	Value (n=53)
Male sex	52 (98)
Age, yr	67 (63–75)
BMI, kg/m ²	23.1 (20.1–24.9)
Comorbid disease	
Diabetes mellitus	8 (15)
Hypertension	4 (8)
MMRC	1 (1–2)
CAT	14.0 (8.0–19.5)
Spirometry	
FVC, L	3.41 (2.89–4.03)
FVC, % predicted	89 (80–99)
FEV ₁ , L	1.69 (1.32–2.01)
FEV ₁ , % predicted	61 (47–70)
FEV ₁ /FVC, %	48 (42–55)
FEV ₁ <50% predicted	16 (30)
Smoking, pack-yr	44 (36–59)
FTND score	3 (1–5)
FTND ≥ 4	32 (60)
Education, yr	9 (6–12)
MoCA-K score	24 (21–26)
MoCA-K <23	24 (45)
Inhaled treatment	41 (77)
LAMA	37 (69)
LABA	35 (66)
ICS or ICS/LABA	19 (36)
Combined*	36 (67)

Values are presented as number (%) or median (interquartile range).

*LAMA+LABA, LAMA+ICS, or LAMA+ICS/LABA.

BMI: body mass index; MMRC: modified Medical Research Council; CAT: chronic obstructive pulmonary disease assessment test; FVC: forced vital capacity; FEV₁: forced expiratory volume in 1 second; FTND: Fagerstrom test for nicotine dependence; MoCA-K: by Korean version of Montreal Cognitive Assessment; LAMA: long-acting muscarinic antagonists; LABA: long-acting β_2 agonists; ICS: inhaled corticosteroids.




Figure 1. Flow of study: selection of study participants.

Table 2. Factors related to moderate to high nicotine dependence of current smokers with chronic obstructive pulmonary disease

Variable	Low nicotine dependence (n=21)	Moderate to high nicotine dependence* (n=32)	p-value
Male sex	21 (100)	31 (97)	0.413
Age, yr	72 (66–76)	66 (61–70)	0.013
BMI, kg/m ²	23.4 (20.4–25.5)	22.3 (19.6–24.4)	0.244
Comorbid disease			
Heart disease	2 (10)	2 (6)	0.388
Diabetes mellitus	2 (9)	6 (19)	0.359
MMRC	1 (1–2)	1 (1–2)	0.461
CAT	11 (8–18)	15 (7–21)	0.472
Spirometry			
FVC, L	3.34 (2.84–3.88)	3.47 (2.95–4.06)	0.296
FVC, % predicted	86 (75–98)	90 (80–101)	0.439
FEV ₁ , L	1.48 (1.15–2.01)	1.76 (1.49–2.16)	0.127
FEV ₁ , % predicted	52 (44–74)	62 (53–69)	0.288
FEV ₁ /FVC	46 (40–53)	49 (44–56)	0.244
FEV ₁ <50% predicted	10 (48)	6 (19)	0.025
Smoking, pack-yr	50 (33–58)	43 (36–60)	0.749
Education, yr	6 (3–12)	12 (9–12)	0.003
MoCA	22 (18–26)	24 (22–27)	0.021
MoCA <23	13 (62)	11 (34)	0.049
Treatment			
Inhaled treatment	18 (86)	24 (75)	0.347
Compliance	2 (11)	4 (17)	0.611

Values are presented as number (%) or median (interquartile range).

*Moderate to high nicotine dependence is FTND \geq 4.

BMI: body mass index; MMRC: modified Medical Research Council; CAT: chronic obstructive pulmonary disease assessment test; FVC: forced vital capacity; FEV₁: forced expiratory volume in 1 second; MoCA-K: by Korean version of Montreal Cognitive Assessment; FTND: Fagerstrom test for nicotine dependence.

Table 3. Multiple logistic regression analysis to find out factors associated with moderate to high nicotine dependence of current smokers with chronic obstructive pulmonary disease

	Odds ratio	95% CI	p-value
Age <70 yr	6.407	1.376–29.830	0.018
FEV ₁ \geq 50% predicted	6.969	1.388–34.998	0.018
MoCA-K	1.182	0.951–1.470	0.132
Education, yr	1.286	1.036–1.596	0.023

CI: confidence interval; FEV₁: forced expiratory volume in 1 second; MoCA-K: by Korean version of Montreal Cognitive Assessment.

patients and the other included diabetic patients^{12,13,17}. However, another two studies^{18,21} showed higher nicotine dependence in older smokers: one study was population-based and the other study enrolled subjects with depressive or anxiety disorders. Based on the previous conflicting reports, relationship between nicotine dependence and age could not be exactly described. We assume that younger smokers are likely to be healthier than older smokers, which could let them not hesitate smoking cigarettes. Another reason is that working young smokers before retirement might feel more stressed and anxious, which might be linked to high nicotine dependence.

The COPD patients with better lung function showed moderate to higher nicotine dependence in this study. The study of Kim et al.¹² showed similar results that the FEV₁ predicted value of COPD patients with high nicotine dependence was

57.8±14.7% and the FEV₁ predicted value of COPD patients with low nicotine dependence was 53.5±17.2%. The COPD patients with less respiratory symptoms had higher scores on nicotine dependence in the study of Lindberg et al.¹³. These results suggest that the need for smoking is higher when lung function is slightly better and respiratory symptoms are less serious.

The participants had median smoking amount of 44 pack-years and median FTND of 3.0. In a study of Lindberg et al.¹³ of COPD patients, median smoking amount was 23.8 pack-years and mean FTND score was 3.7. They showed lower median smoking amount but higher median FTND compared with those of our study. Factors associated with high FTND in their study were young age, existence of respiratory symptom, anxiety, high smoking amount, and low self efficacy¹³. In the other study on genetics of nicotine dependence in COPD¹², mean smoking amount was 51.7 pack years and mean FTND was 4.7. Both of smoking amount and FTND were higher than those of this study. Mean age of COPD patients with high nicotine dependence was 57.9±7.5 years old and that with low nicotine dependence was 60.0±8.0 years old¹². Younger patients tend to have higher nicotine dependence in our study and previous studies^{12,13}. The relatively low level of median nicotine dependence in our study might be due to the fact that many subjects were older than other studies or different characteristics of subjects including race, sex or social economic status.

Even though nicotine can enhance cognitive functions and emotional processing in some conditions²², chronic smoking is associated with a decline in cognitive functions²³. There has been few study on cognitive function associated to nicotine dependence. In our study, higher cognitive function was observed in subjects with moderate to high nicotine dependence. However, nicotine dependence cannot be a favorable factor for long-term cognitive function. Since COPD patients with moderate to high nicotine dependence in this study showed younger age and higher level of education, these two factors might explain higher cognitive function in this nicotine-dependent group. To understand the relationship between cognitive function and nicotine dependence, follow-up studies on a large number of COPD patients are needed.

Although some studies of COPD smokers showed a positive correlation between smoking amount and nicotine dependence^{12,13}, we found no difference in the amount of smoking between the low and moderate to high nicotine dependence groups. Nicotine dependence reflects current status of nicotine addiction, but not past history of smoking. In other words, it is difficult to estimate the degree of nicotine addiction through the amount of smoking in the past. Therefore, nicotine dependence measures should be made before smoking cessation education or nicotine replacement therapy rather than measuring the amount of cigarette smoking. Because current smokers of COPD exhibit moderate to high nicotine

dependence, nicotine withdrawal symptoms are a major cause of failure to quit smoking, indicating that more aggressive approach for smoking cessation is needed.

The current work has some limitations. The number of participants was small. A large number of current smokers with COPD are needed to identify factors associated with nicotine dependence in COPD. It is difficult to say that our research subjects are representative of the whole COPD because they were recruited from tertiary medical institutions rather than primary medical institutions. Nevertheless, our cohort was well-designed, and surveys were conducted by well-trained nurses. Another limitation is that we did not investigate other factors known to be related to nicotine dependence such as household income, marital status, genetic factors, age at onset of smoking, depression, and drinking status.

This study assessed nicotine dependence in current smokers with COPD. We found that higher education level, mild to moderate airflow obstruction, and age <70 were associated with moderate to high nicotine dependence. In order to provide effective smoking cessation intervention for current smokers with COPD, physicians need to consider the level of individual education, age, and severity of airflow obstruction.

*Author affiliations

¹Division of Pulmonary, Allergy and Critical Care Medicine, Department of Internal Medicine, Hallym University Kangnam Sacred Heart Hospital, Seoul, ²Division of Pulmonary and Critical Care Medicine, Department of Medicine, Ewha Womans University School of Medicine, Seoul, ³Department of Internal Medicine, Pusan National University School of Medicine, Busan, ⁴Division of Pulmonary Medicine, Department of Internal Medicine, Ulsan University Hospital, University of Ulsan College of Medicine, Ulsan, ⁵Division of Pulmonary and Critical Care Medicine, Department of Medicine, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, ⁶Division of Pulmonary and Critical Care Medicine, Department of Internal Medicine, Seoul National University Hospital, Seoul, ⁷Division of Pulmonary and Critical Care Medicine, Department of Internal Medicine, Seoul Metropolitan Government-Seoul National University Boramae Medical Center, Seoul National University College of Medicine, Seoul, ⁸Regional Center for Respiratory Disease, Yeungnam University Medical Center, Yeungnam University College of Medicine, Daegu, ⁹Department of Internal Medicine, St. Paul's Hospital, College of Medicine, The Catholic University of Korea, Seoul, ¹⁰Respiratory Division, Department of Internal Medicine, Soonchunhyang University Gumi Hospital, Soonchunhyang University College of Medicine, Gumi, ¹¹Department of Internal Medicine, Incheon St. Mary's Hospital, College of Medicine, The Catholic University of Korea, Incheon, ¹²Department of Pulmonary and Critical Care Medi-

cine, Hallym University Kangdong Sacred Heart Hospital, Hallym University College of Medicine, Seoul, ¹³Department of Internal Medicine, Chonnam National University Hospital, Gwangju, ¹⁴Division of Pulmonary and Critical Care Medicine, Department of Internal Medicine, Konkuk University Medical Center, Konkuk University School of Medicine, Seoul, ¹⁵Division of Pulmonary and Critical Care Medicine, Department of Internal Medicine, National Medical Center, Seoul, ¹⁶Department of Pulmonary and Critical Care Medicine and Clinical Research Center for Chronic Obstructive Airway Diseases, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Korea

Conflicts of Interest

No potential conflict of interest relevant to this article was reported.

Acknowledgments

This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Science, ICT & Future Planning (2010-0027945).

References

1. Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, Aboyans V, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012;380:2095-128.
2. Yoo KH, Kim YS, Sheen SS, Park JH, Hwang YI, Kim SH, et al. Prevalence of chronic obstructive pulmonary disease in Korea: the fourth Korean National Health and Nutrition Examination Survey, 2008. *Respirology* 2011;16:659-65.
3. Kohansal R, Martinez-Cambor P, Agusti A, Buist AS, Mannino DM, Soriano JB. The natural history of chronic airflow obstruction revisited: an analysis of the Framingham offspring cohort. *Am J Respir Crit Care Med* 2009;180:3-10.
4. Rabe KF, Hurd S, Anzueto A, Barnes PJ, Buist SA, Calverley P, et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med* 2007;176:532-55.
5. Godtfredsen NS, Prescott E. Benefits of smoking cessation with focus on cardiovascular and respiratory comorbidities. *Clin Respir J* 2011;5:187-94.
6. Chandler MA, Rennard SI. Smoking cessation. *Chest* 2010; 137:428-35.
7. Karen LD, Rosecrans JA. Addictive capacity of nicotine. Washington, DC: American Psychiatric Press Inc.; 2000.
8. Fagerstrom KO, Schneider NG. Measuring nicotine dependence: a review of the Fagerstrom Tolerance Questionnaire. *J Behav Med* 1989;12:159-82.
9. Hymowitz N, Cummings KM, Hyland A, Lynn WR, Pechacek TF, Hartwell TD. Predictors of smoking cessation in a cohort of adult smokers followed for five years. *Tob Control* 1997;6 Suppl 2:S57-62.
10. Jimenez-Ruiz CA, Masa F, Miravittles M, Gabriel R, Viejo JL, Villasante C, et al. Smoking characteristics: differences in attitudes and dependence between healthy smokers and smokers with COPD. *Chest* 2001;119:1365-70.
11. Shahab L, Jarvis MJ, Britton J, West R. Prevalence, diagnosis and relation to tobacco dependence of chronic obstructive pulmonary disease in a nationally representative population sample. *Thorax* 2006;61:1043-7.
12. Kim DK, Hersh CP, Washko GR, Hokanson JE, Lynch DA, Newell JD, et al. Epidemiology, radiology, and genetics of nicotine dependence in COPD. *Respir Res* 2011;12:9.
13. Lindberg A, Niska B, Stridsman C, Eklund BM, Eriksson B, Hedman L. Low nicotine dependence and high self-efficacy can predict smoking cessation independent of the presence of chronic obstructive pulmonary disease: a three year follow up of a population-based study. *Tob Induc Dis* 2015;13:27.
14. Miller MR, Hankinson J, Brusasco V, Burgos F, Casaburi R, Coates A, et al. Standardisation of spirometry. *Eur Respir J* 2005;26:319-38.
15. Choi JK, Paek D, Lee JO. Normal predictive values of spirometry in Korean population. *Tuberc Respir Dis* 2005;58:230-42.
16. Lee JY, Lee DW, Cho SJ, Na DL, Jeon HJ, Kim SK, et al. Brief screening for mild cognitive impairment in elderly outpatient clinic: validation of the Korean version of the Montreal Cognitive Assessment. *J Geriatr Psychiatry Neurol* 2008;21:104-10.
17. Osme SF, Ferreira L, Jorge MT, de Souza Andreo J, Jorge M, de Melo Costa Pinto R, et al. Difference between the prevalence of symptoms of depression and anxiety in non-diabetic smokers and in patients with type 2 diabetes with and without nicotine dependence. *Diabetol Metab Syndr* 2012;4:39.
18. Jamal M, Willem Van der Does AJ, Cuijpers P, Penninx BW. Association of smoking and nicotine dependence with severity and course of symptoms in patients with depressive or anxiety disorder. *Drug Alcohol Depend* 2012;126:138-46.
19. Pennanen M, Broms U, Korhonen T, Haukkala A, Partonen T, Tuulio-Henriksson A, et al. Smoking, nicotine dependence and nicotine intake by socio-economic status and marital status. *Addict Behav* 2014;39:1145-51.
20. Pack HJ, Kim JS, Kim KM, Kim SM, Lee DH, Kim SS, et al. Nicotine dependence and internal response to stress in male smokers. *J Korean Acad Fam Med* 2007;28:604-9.
21. Breslau N, Johnson EO, Hiripi E, Kessler R. Nicotine dependence in the United States: prevalence, trends, and smoking persistence. *Arch Gen Psychiatry* 2001;58:810-6.

22. Heishman SJ, Kleykamp BA, Singleton EG. Meta-analysis of the acute effects of nicotine and smoking on human performance. *Psychopharmacology (Berl)* 2010;210:453-69.
23. Richards M, Jarvis MJ, Thompson N, Wadsworth ME. Ciga-

rette smoking and cognitive decline in midlife: evidence from a prospective birth cohort study. *Am J Public Health* 2003;93:994-8.