



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Correspondence to: Shari Lipner, MD, PhD, 1305 York Ave, 9th Floor, New York, New York 10021

E-mail: sbl9032@med.cornell.edu

REFERENCES

1. Zocdoc. How Zocdoc search works. Available at: <https://www.zocdoc.com/about/how-search-works/>. Accessed July 18, 2019.
2. Xiang L, Lipner SR. Analysis of wait times for online dermatology appointments in most and least dermatologist-dense cities. *J Drugs Dermatol*. 2020;19(5):562-565.
3. Glazer AM, Farberg AS, Winkelmann RR, Rigel DS. Analysis of trends in geographic distribution and density of US dermatologists. *JAMA Dermatol*. 2017;153(4):322-325.
4. Bashshur R, Doarn CR, Frenk JM, Kvedar JC, Woolliscroft JO. Telemedicine and the COVID-19 pandemic, lessons for the future. *Telemed J E Health*. 2020;26(5):571-573.
5. Wang RF, Trinidad J, Lawrence J, et al. Improved patient access and outcomes with the integration of an eConsult program (tele dermatology) within a large academic medical center. *J Am Acad Dermatol*. 2019;83(6):1633-1638.

<https://doi.org/10.1016/j.jaad.2020.10.069>

Diagnostic and management considerations for “maskne” in the era of COVID-19



To the Editor: “Maskne,” coined during the coronavirus disease 2019 (COVID-19) pandemic, is a variant of acne mechanica, previously associated with headgear or personal protective equipment. This report focuses on the widespread use of reusable fabric masks and the impact of the skin microenvironment and mechanical factors (ie, textile-skin friction), which to my knowledge has not been previously evaluated.

Our understanding of maskne is largely observational, although it is likely a disorder of follicular occlusion and directly related to mechanical stress¹ (pressure, occlusion, friction) and microbiome dysbiosis² (heat, pH, moisture from biofluids). Both of these are affected by increased duration of mask wear. Tropical climates and outdoor exposure (increased sweating) are risk factors for acne-susceptible populations (active young adults, seborrhea, genetic predisposition).

Clinical criteria proposed for maskne: onset of acne within 6 weeks of start of regular face mask wear or exacerbation of acne over the masked area, distinct pattern, referred to as the O-zone in this report (Fig 1), and exclusion of differential diagnoses, including perioral dermatitis, seborrheic dermatitis, pityrosporum folliculitis, and acne rosacea.

Special consideration for skin care should include antibacterial gentle cleansers and moisturizers formulated as prescription emollient devices, which help maintain a healthy skin barrier/microbiome.





Fig 1. Distinct acne patterns seen in the T zone of physiologic acne, the U zone of adult acne, and the O zone of maskne.

Spot acne treatment with benzoyl peroxide, salicylic acid, sulfur, α -hydroxy acids, and retinoids predispose to irritant contact dermatitis under occlusion. Botanical actives with anti-inflammatory, antioxidant, sebum regulation, and antimicrobial properties are preferred. Hydrogel carrier formulations of retinoid/antibiotic combination topicals can minimize local irritation by ensuring better drug tolerance and efficacy.

Chemical sunscreen sensitivity and comedogenicity increase under occlusion. Ultraviolet protection factor (UPF) 50+ fabric masks should replace the wearing of sunscreen (eliminates constant reapplication) as a practical photoprotective measure for the lower half of the face, to improve compliance to sun protection and incentivize mask wearing. Dermatologists' prescribing patterns for maskne may potentially influence the development of antibiotic resistance worldwide.

Dermatologists can advise on design of face masks, encouraging patient compliance. Natural fibers wick moisture but have increased fluid saturation, heightening discomfort and stickiness sensation (accumulated stickiness magnitude).³ Synthetic biofunctional textiles have a high evaporation/cooling coefficient and are water resistant, preventing biofluid spread. High thread count and tightly woven fabrics have higher UPF and minimize textile-skin friction, relevant to individuals with atopic conditions. Light/reflective colors disperse heat (Fig 2).

Reusable fabric masks should allow for movement while speaking, with minimal displacement over orifices. Fabric masks should omit abrasive metallic parts that also cause nickel sensitization. Allergic contact dermatitis and transcutaneous absorption of



© Dr. Teo Wan Lin 2020

Fig 2. The ideal fabric face mask design to minimize textile-skin friction. *UPF*, Ultraviolet protection factor; *UV*, ultraviolet.

potential carcinogens can occur from textiles used for fabric face masks, which is currently unregulated.

The skin microbiome is influenced by genetic and external factors (ie, environment, pH, and temperature), which are modified with mask wearing. Microbiome dysbiosis is implicated in the pathogenesis of eczema, acne, perioral dermatitis, seborrheic dermatitis, pityrosporum folliculitis, and rosacea. Biofunctional textiles with UPF, antioxidant, and antiaging benefits may incentivize mask wearing. Copper oxide polyesters are associated with reduced facial wrinkles,⁴ elevation of elastin, procollagen 1, and transforming growth factor- β 1 levels.⁵ Textiles that incorporate silver, zinc oxide, and copper oxide have broad-spectrum biocidal properties that are therapeutic for dermatologic conditions, besides reducing antibiotic resistance in acne treatment.

Wan-Lin Teo, MBBS, MRCS (UK), FAMS (Dermatology)

From TWL Specialist Skin & Laser Centre, Singapore.

Funding sources: None.

Conflicts of interest: The author is the chief scientific advisor and owns Dr TWL Dermaceuticals,

a cosmeceutical, biofunctional textile research and development company.

IRB approval status: Not applicable.

Correspondence and reprint requests to: Wan-Lin Teo, MBBS, MRCS (UK), FAMS (Dermatology), 101 Irrawaddy Rd #17-07, Singapore 329565

E-mail: wanlinteo@gmail.com

REFERENCES

1. Boer J, Nazary M, Riis PT. The role of mechanical stress in hidradenitis suppurativa. *Dermatol Clin*. 2016;34(1):37-43.
2. Naik HB, Piquet V. Standardizing hidradenitis suppurativa skin microbiome research: the methods matter. *J Invest Dermatol*. 2020;140(9):1688-1690.
3. Tang KM, Chau KH, Kan CW, Fan JT. Assessing the accumulated stickiness magnitude from fabric-skin friction: effect of wetness level of various fabrics. *R Soc Open Sci*. 2018; 5(8):180860.
4. Baek JH, Yoo MA, Koh JS, Borkow G. Reduction of facial wrinkles depth by sleeping on copper oxide-containing pillowcases: a double blind, placebo controlled, parallel, randomized clinical study. *Cosmet Dermatol*. 2012;11(3): 193-200.
5. Ogen-Shtern N, Chumin K, Cohen G, Borkow G. Increased pro-collagen 1, elastin, and TGF- β 1 expression by copper ions in an ex-vivo human skin model. *J Cosmet Dermatol*. 2020; 19(6):1522-1527.