



Controlling Semi-Invasive Activity of Human Endometrial Stromal Cells by Inhibiting NF- κ B Signaling Pathway Using Aloe-emodin and Aspirin

Nahid Nasiri^{1,2†}, Sara Babaei^{3†}, Ashraf Moini^{4,5,6}, Poopak Eftekhari-Yazdi^{1*}

1- Department of Embryology, Reproductive Biomedicine Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Iran

2- Department of Stem Cells and Developmental Biology, Cell Science Research Center, Royan Institute for Stem Cell Biology and Technology, ACECR, Tehran, Iran

3- Department of Developmental Biology, Faculty of Basic Sciences and Advanced Technologies in Biology, University of Science and Culture, Tehran, Iran

4- Department of Endocrinology and Female Infertility, Reproductive Biomedicine Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Iran

5- Department of Obstetrics and Gynecology, Faculty of Medicine, Tehran University of Medical Sciences, Tehran, Iran

6- Department of Obstetrics and Gynecology, Arash Women's Hospital, Tehran University of Medical Sciences, Tehran, Iran

† The first and the second authors have had equal contribution to this manuscript

Abstract

Background: Inflammation and its master regulator, Nuclear Factor- κ B (NF- κ B), have been implicated in the development of endometriosis. Inhibition of NF- κ B pathway using small molecules ameliorated disease progression and reduced the lesion size; nevertheless, the underlying mechanism is not fully understood. Therefore, this study, is an attempt to assess whether inhibiting NF- κ B signaling by aloe-emodin (AE) or aspirin (Asp), as anti-inflammatory compounds, can suppresses the invasive activity of human endometrial stromal cells at stage IV endometriosis.

Methods: The eutopic and healthy endometrial biopsies from a total of 8 infertile women with confirmed endometriosis and 8 women without endometriosis were digested and the single cells were cultured. Gene and protein markers of proliferation, migration, adhesion, and invasion of eutopic endometrial stromal cells (EuESCs) with and without treatment with AE or Asp, as well as control endometrial stromal cells (CESCs) was analyzed using q-PCR and immunofluorescence staining, respectively. Comparison between groups was performed using one-way ANOVA and the Bonferroni post hoc and $p \leq 0.5$ was considered statistically significant.

Results: There was an association between NF- κ B overexpression and higher proliferation/adhesion capacity in EuESCs. EuESCs (at stage IV endometriosis) displayed no invasive and migratory behaviors. Pre-treatment of EuESCs with AE or Asp significantly attenuated NF- κ B expression and reduced proliferative, adhesive, invasive, and migratory activity of endometrial cells ($p \leq 0.5$).

Conclusion: Eutopic endometrial stromal cells seem to have a semi-invasive activity which is largely suppressed by AE or Asp. It can be suggested that both Asp and AE (as potent NF- κ B inhibitors) can be used as a supplement in conventional endometriosis treatments.

Keywords: Adhesion, Aspirin, Cell proliferation, Endometrial biopsy, Endometriosis.

To cite this article: Nasiri N, Babaei S, Moini As, Eftekhari-Yazdi P. Controlling Semi-Invasive Activity of Human Endometrial Stromal Cells by Inhibiting NF- κ B Signaling Pathway Using Aloe-emodin and Aspirin. *J Reprod Infertil.* 2021;22(4):227-240. <http://dx.doi.org/10.18502/jri.v22i4.7648>.

Introduction

Endometriosis, is a chronic, inflammatory disease affecting 5-10% of women at reproductive age, this gynecological disorder histo-

logically characterized by growth of endometrium-like tissue at abnormal locations and outside the uterus cavity (1, 2). Considering its severe and

long-term health effects including chronic pelvic pain, and infertility, as well as other issues such as difficult diagnosis and lack of definitive treatment, endometriosis has become a great health-care challenge (3).

Concerning the pathogenesis of the endometriosis, it has been indicated that any blockage of the menstrual flow may increase the possibility of endometriosis (4) suggesting the importance of Sampson's theory of reflux menstruation in endometriosis etiology. According to the theory, endometrial debris may adhere to the uterus and proliferate and next be translocated outside the uterus by the retrograde blood flow (4, 5). However, it is worth considering that although in most women, retrograde menstruation occurs, endometriosis develops only in up to 10% of cases. Accordingly, other explanations for endometriosis etiology are existence of extrauterine stem cells such as those brought in the abdominal space by blood or lymphatic vessels, as the main source for development of endometrial lesions (4). It was hypothesized that the altered peritoneal microenvironment as a result of immunological disorders in genetically susceptible patients may lead to the disease development (6, 7).

Progressive endometriosis shares some aspects with cancer. Immortalized endometriotic stromal cells have been extensively used in studies for endometriosis pathophysiology (8). Endometriotic cells can invade adjacent tissues, metastasize to distal organs, recur after treatment, and often, after inducing chronic wounds, the resultant lesions never heal (9, 10). Accordingly, increased cell proliferation and resistance to apoptosis (11, 12), incidence of epithelial–mesenchymal transition (EMT) in endometriotic stromal cells, great potential for migration and invasion (13, 14), and higher ability of attachment and angiogenesis (15), are considered typical characteristics of endometriotic stromal cells. Currently, endometriosis is being managed by available treatments including suppression of inflammation, hormonal inhibition of menstruation, and surgical harvest of ectopic endometrial lesions. However, both medical and surgical interventions are considered sub-optimal due to the high likelihood of disease recurrence and infertility challenge that remain unresolved and necessitate development of novel treatment strategies (16, 17). Considerable evidence introduced inflammation as a key factor playing a significant role in occurrence of the endometriosis (18-20). Endometrial lesions were found to have

elevated levels of various inflammatory mediators including estradiol (E2), prostaglandin E2 (PGE2), TNF- α , interleukin-1 β , *etc.* which form positive feedback and support both the induction and persistence of inflammation followed by fibrosis, pelvic pain, and infertility (21, 22). These key inflammatory mediators were shown to be involved in development of invasive activities of ESCs including proliferation, adhesion, and invasion (23, 24). It was demonstrated that the orchestration of common phenotypes in endometriosis and cancer such as metastasis and angiogenesis, apoptosis suppression, and treatment resistance is carried out by NF-kB signaling pathway (24, 25) that appears to be overexpressed in endometriotic cells and tissues (26, 27).

Literature review shows that most biological or chemical agents blocking NF-kB can effectively modulate endometriosis-associated symptoms and disrupt its progression (28). So, suppression of NF-kB seems to result in promising outcomes in endometriosis management. Therefore, in the current study, the therapeutic effects of aloe-emodin (AE; 1,8-dihydroxy-3-hydroxyl-methylantraquinone) and aspirin (Asp; acetylsalicylate) were investigated on invasive activity of endometriotic cells. AE, the major bioactive phytochemical of rhubarb (*Rheum palmatum*) and aloe vera, which has been widely used in traditional Chinese medicine, is well known for its anti-inflammatory and anti-proliferative effects in different types of cancer (29, 30). Asp, as the next choice for this purpose, is a chemically synthesized compound that serves as one of the most effective globally used medications with the least complications in treatment of fever, inflammation, and pain in patients during the last century. It has been revealed that Asp can halt fibrogenesis, decelerate the epithelial-to-mesenchymal transition phenomenon, reduce cells migration/invasion capacity, and regulate apoptosis (31, 32). In this study, the effect of Asp and AE on NF-kB expression, proliferation, invasion, adhesion, and migration capacity of eutopic endometrial stromal cells (EuESCs) derived from endometrial biopsy of patients with severe endometriosis (Stage IV) was compared with control endometrial stromal cells (CESCs). In addition to blocking the NF-kB signaling, the invasive activity of ESCs following treatment with TNF- α , as a known inducer of the pathway, was evaluated to lay the ground for discussing the role of NF-kB in pathogenesis of endometriosis.

Methods

Study participants and tissue collection: Eutopic endometrium tissues were obtained from 12 patients with primary infertility (23-35 years old) and ovarian endometriosis at stage IV (Proliferative phase; no LH timing; ectopic peritoneal lesions: n=8; and ovarian cysts: n=8) through laparoscopy. The biopsies of control endometrial tissues were obtained from 10 non-endometriosis women with primary infertility (25-35 years old) with laparoscopy indication for ovarian simple cyst or pelvic pain/uterine malformation (proliferative phase). None of the patients had received hormone therapy during 6 months before laparoscopy. All patients gave their written informed consent and the study was approved by the ethical committee of Royan Institute (Code: IR.ACECR.ROYAN.REC.1397.161). Four patients out of the 12 patients included in the eutopic endometriosis group as well as two patients out of 10 patients included in the control group, were excluded from the study due to lack of sufficient cells in the collected biopsies.

Endometrium cell culture and preparation of primary EuESCs and CESC: According to the objectives of the study, five groups were designed. Eutopic endometrial stromal cells were derived from endometrial biopsy of patients with severe endometriosis (Stage IV) and formed EuESCs group. In EuESCs+ASP group, the EuESCs were treated with 2.5 mmol/l of aspirin. In EuESCs+AE group, the EuESCs were treated with 20 μ mol/l of AE. Accordingly, the control endometrial stromal cells (CESCs) were obtained from endometrial biopsy of patients whose normal uterine endometrium was confirmed by laparoscopy. In CESC+TNF, the CESC were treated with 20 ng/ml of TNF- α .

The endometria were washed in PBS solution (Phosphate buffered-tablet, Sigma-Aldrich, USA) and digested using collagenase 0.25% (Thermo Fisher Scientific, USA) for 1 hr at 37 °C for isolation of the ESCs. Removal of undigested tissues was performed using filtration of cell suspension through a 100- μ m filter mesh (Falcon, USA); also, to purify the ESCs, the epithelial cells were removed using supernatant filtration through a 40- μ m sieve (Falcon, USA). The cell suspension was centrifuged at 1000 rpm for 5 min. Isolated cells were then cultured with 5% carbon dioxide at 37 °C in Dulbecco modified Eagle medium F-12 (DMEM/F12; Gibco, Sweden) containing 10% fetal bovine serum (FBS; Gibco, Sweden) and 1%

penicillin-streptomycin (Pen-Strep; 10,000 units of penicillin and 10 mg of streptomycin/ml, Sigma-Aldrich, USA). At passage 3, the purity of ESCs was 95% as determined by flow cytometry evaluation of the cells' immunophenotype. For this aim, the harvested cells were stained with mouse anti-human CD29 (BD, USA), FITC-labelled mouse anti-human CD90 (Daco, USA), and CD45 (BD, USA) antibodies to ensure the mesenchymal origin of the cells, as well as PE-conjugated mouse anti-human vimentin (Santa Cruz, USA) antibody, to identify the stromal phenotype of the isolated cells. Characterized ESCs were used for subsequent experiments.

Cytotoxicity testing and cell viability assay: The MTT assay was performed to check possible deleterious effects of TNF- α (Sigma-Aldrich, USA), Asp (Sigma-Aldrich, USA), and AE (Sigma-Aldrich, USA) on ESCs viability. Briefly, 7×10^3 ESCs were seeded in a 96-well flat-bottomed cell culture plate in a final volume of 100 μ l of DMEM/F12 (Gibco, Sweden) supplemented with 5% FBS, then incubated overnight. After incubation, the medium was removed and the ESCs were incubated for different time periods (24, 48 and 72 hr) with AE, Asp or TNF- α . Then, 50 μ l of MTT solution (Sigma-Aldrich, USA) was added to each well and the ESCs were further incubated for 2 hr. The absorbance of the wells at 570 nm was measured for evaluation of ESCs viability.

Quantitative polymerase chain reaction (q-PCR): Manual extraction of total RNA from ESCs was done using TRIzol (Thermo Fisher Scientific, USA). Concentration of RNA was determined by NanoDrop (Thermo Scientific NanoDrop One/OneC). Human GAPDH reactions were amplified for 40 cycles. Polymerase chain reaction (PCR) was done in Mastercycler Pro S (Eppendorf, Germany) in a 96- μ l reaction system. Next, 1% agarose gel was used for separation of reaction products. Subsequently staining with SYBR Green PCR Master Mix (Applied Biosystems, UK) was done for visualization. Calculation of the gene expression level was performed using the threshold cycle (CT) method ($2^{-\Delta\Delta Ct}$). Table 1 shows the sequences of primers and fragment lengths, used in this study.

Immunofluorescence staining: Semiquantitative evaluation of protein expression was performed using immunofluorescence staining. For this aim, the EuESCs and CESC were fixed using 4% par-

Table 1. Sequences of PCR primers and length of PCR product

Gene	PCR primers (5' → 3')	Length of PCR product (bp)
NF-kB	5'GTGGGGACTACGACCTGAATG3' 5'GGGGCACGATTGTCAAAGATG3'	121 bp
Ki-67	5'GTGCTCAACAACCTTCATTTCCA3' 5'ACTGAAGAACACATTTCTCTCCA3'	231 bp
PCNA	5'TGAACCTCACCAGTATGTCC3' 5'TCATAGTCTGAAACTTTCTCCTG3'	136 bp
C-MYC	5'ACACATCAGCACAACTACG3' 5'CGCCTCTTGACATTCTCC3'	140 bp
V-CAM	5'GAAATGACCTTCATCCCTACC3' 5'ATATTCACAGAACTGCCTTCTC3'	197 bp
I-CAM	5'GTGACCGTGAATGTGCTCTC3' 5'TCTGTATTTCTTGATCTTCCGCT3'	132 bp
CD133	5'GCATCCATCAAGTGAAACGT3' 5'GGTTTGCGGTTG TACTCTGT3'	199 bp
MMP-9	5'GCCACTACTGTGCCTTTGAG3' 5'CAGAGAATCGCCAGTACTTCC3'	121 bp
GAPDH	5'CTCATTTCCTGGTATGACAACGA3' 5'CTTCTGTGCTCTTGCT3'	122 bp

aformaldehyde in PBS (Sigma-Aldrich, USA) for 20 min at room temperature. Then, the cells were washed with PBS and were blocked using 10% goat serum diluted in PBS for 1 hr and incubated with primary antibodies for NF-kB (1:50 dilution; Santa Cruz, USA), C-MYC (1:200 dilution; BD, USA) and I-CAM (1:50 dilution; Santa Cruz, USA) at 4°C overnight. Then, goat anti-mouse IgG conjugated to Alexa Fluor 488 (Invitrogen, USA) for C-MYC and goat anti-mouse IgG conjugated to Alexa Fluor 594 (Invitrogen, USA) were added and kept for 1 hr at room temperature, followed by mounting the cells with DAPI and microscopic imaging (ix71, Olympus, Japan).

ESCs migration and invasion assay: The *in vitro* migration capacity of ESCs was examined by wound healing (wound closure) assay. For this purpose, 24 hr before the assay, the untreated EuESCs and EuESCs treated with AE/Asp, as well as CECs were seeded in DMEM/F12 plus 10% FBS in a six-well plate. At 90% confluence, a low dose of the proliferation inhibitor, 0.1 mg/ml of mitomycin C (Sigma Aldrich, USA) was added (To further avoid the risk of cell proliferation). A wound was created using 200 µl plastic cell scraper on the center of the wells containing confluent cell monolayer. The cells were incubated for 24 hr and ESCs migration was measured 0, 6, 12, 18, 24 and 48 hr after wounding. Wound

closure at each time point was evaluated as follows:

Wound closure %:

$$\frac{[A_{t=0h} - A_{t=\Delta h}] \times 100\%}{A_{t=0h}}$$

A_{t=0h}: is the area of the wound measured immediately after scratching (t=0h),

A_{t=Δh}: is the area of the wound measured hours after the scratch is performed.

Statistical analysis: Statistical analyses were performed using GraphPad Prism v6.0 software (GraphPad Software, USA). TNF-α production was compared using t-test, whereas one-way analysis of variance (one-way ANOVA) and the Bonferroni post hoc were used to compare proliferation, adhesion, invasion, and migration as well as NF-kB expression. The level of significance was set at p≤0.05.

Results

Morphology and immunophenotyping of ESCs: ESCs obtained from digestion of both endometriotic cells and healthy endometrium (EuESCs and CECs, respectively) exhibited a fibroblast-like morphology (flat and spindle-shape). At the third passage, the purity and identity of human ESC cultures were confirmed by flow cytometry evaluation of specific markers for stromal cells with




Figure 1. Immunophenotyping of ESCs isolated from eutopic endometriotic and non-endometriotic endometrial samples, A) immunofluorescence staining of both endometriotic and healthy ESCs, exhibiting the expression of vimentin as a stromal marker, B) flow cytometry analysis of CD90, CD29, CD45 in ESCs. Data are presented as mean±SD

mesenchymal but not hematopoietic origin (Figure 1).

Increased level of TNF- α as an inducer of NF- κ B pathway: The EuESCs supernatant showed different patterns of TNF- α secretion. Using ELISA method, EuESCs at third passage secrete higher concentrations of TNF- α compared with the control (CESCs) (Figure 2). This result indicates that EuESCs express aberrant levels of TNF- α compared with CESCs.

The effect of Asp and AE on ESCs viability: Using MTT assay, the safety of Asp and AE on the ESCs was examined. Up to a concentration of 2.5 mmol/l of Asp and 20 μ M of AE, these compounds did not affect viability of ESCs (Figure 3). In addition, TNF- α , as a known inducer of NF- κ B, was used to improve the experiments.

Restoration of impaired expression of NF- κ B in EuESCs by Asp and AE: EuESCs expressed increased levels of NF- κ B mRNA and protein compared with CESCs, but treatment of CESCs with TNF- α for 24 hr further increased these values (Figure 4A, 4B). Significantly, Asp and AE saved the EuESCs from excess transcripts and proteins of NF- κ B.

Regulation of proliferative phenotype/capacity of EuESCs by Asp and AE: Cell proliferation is a well-known phenotype of endometriotic stromal cells. Three proliferation markers (KI-67, C-MYC, and PCNA) were examined in cells of all




Figure 2. The concentration of inflammatory cytokine, TNF- α protein, produced in cell culture supernatants of EuESCs and CESCs. The EuESCs released higher levels of TNF- α than CESCs evaluated by ELISA. Data are presented as mean± SD (p≤0.01*)

five groups. Based on our results, EuESCs displayed elevated proliferation capacity which was significantly higher than that of TNF- α -treated CESCs at gene (Figure 5A) and protein expression level (Figure 5B). Moreover, our suggested NF- κ B regulators, Asp and AE, decreased the proliferative capacity of endometriotic cells to the level of the control stromal cells.

Reduced response to NF- κ B modulators: Adhesion of EESCs to any ectopic site outside the uterus is likely a landmark in formation and progression of endometrial lesions. Using quantitative assessment of I-CAM and V-CAM gene expression (Figure 6A) and semiquantitative evaluation of I-




Figure 3. The ESCs viability was determined after treatment with 0, 1, 2.5, 5 and 10 mmol/L of Asp (A) 0, 5, 10, 20 and 40 μM of AE (B) and 0, 20, 40 and 60 ng/ml of TNF-α (C) using the MTT (3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide) assay. Results are expressed as the mean measurements from three independent experiments at 24 and 48 hr. Data are presented as mean±SD (p<0.05*)




Figure 4. Comparison of NF-kB expression at gene and protein levels between EuESCs and CESC after treatment with Asp/AE or TNF-α, respectively, A) total extracted RNA of the cells was analyzed by RT-PCR. Quantification was performed using the ratio of the target bands relative to GAPDH. The highest levels of NF-kB were presented in TNF-α treated CECs and then EuESCs. Treatment with Asp or AE reduced the expression levels up to control, B) immunofluorescence analysis of the NF-kB protein expression in endometrial cells at different groups. NF-kB protein was distributed in the cytoplasm of the cells. The nuclei were stained with DAPI (Blue), C) representative chart for NF-kB protein expression. The protein values followed the same pattern of gene expression in different groups. Data are presented as mean±SD.

* Significant difference with the control, # Significant difference with the untreated group (p<0.05), magnification×400




Figure 5. The effect of Asp/AE on proliferation capacity of EuESCs, A) the downregulated levels of proliferative genes; C-MYC, KI67 and PCNA, in untreated EuESCs and those treated with Asp/AE were measured by quantitative polymerase chain reaction (qPCR), B) immunofluorescence assessment of C-MYC protein expression in endometrial cells of all five groups and, C) respective semiquantitative analysis of C-MYC protein levels, indicating reduced proliferative capacity of EuESCs after treatment with Asp or AE. Data are presented as mean±SD.

* Significant difference with the control, # Significant difference with the untreated group ($p \leq 0.01$), magnification×400




Figure 6. Adhesion analysis of CESC and EuESC cells before and after treatment with Asp or AE at gene, A) and protein, B) expression levels, C) the chart for I-CAM protein expression revealed the overexpression and adhesion capacity of EuESCs and CESC treated with TNF- α . As expected, treatment with Asp or AE reduced this activity even lower than the control. Data are presented as mean±SD. * Significant difference with the control, # Significant difference with the untreated group, ($p \leq 0.01$), magnification×400

CAM protein (Figure 6B), adhesive capacity of EuESCs with and without Asp or AE treatment was investigated and compared with CESC. The expression level of adhesion gene markers and

protein was reduced in Asp or AE-treated EuESCs versus the untreated EuESCs.

Expression of invasion markers as the hallmarks of cancer: Endometriosis as a benign disease is not




Figure 7. Effect of Asp/AE on invasive gene marker expression of EuESCs. Compared with CESC, human EuESCs (at stage IV endometriosis) did not express a significant level of invasive gene markers (CD133 and MMP-9). Asp/AE similarly diminished the normal expression of invasive markers in EuESCs. Data are presented as mean±SD.

* Significant difference with the control, # Significant difference with the untreated group ($p \leq 0.01$)

expected to express the invasion markers; however, it shares some features with cancer such as metastasis. Here, the expression pattern of two specified markers (CD-133 and MMP-9) contributing to cancer progression and metastasis was investigated. Both genes had the same response in our studied groups. Figure 7 shows there was no significant difference in expression levels of CD-133 and MMP-9 between EuESCs and CESC. Nevertheless, Asp and AE decreased gene expression to levels even lower than that of the healthy CESC.

Migration potency of EuESCs of patients with stage IV endometriosis: Specifically in endometriosis, the endometrial stromal cells or endometrial lesion cells move and migrate outside the uterus cavity or toward other ectopic sites, leading to formation or progression of the disease. Assuming that the EuESCs have higher migration ability, unexpected results were obtained. Figure 8 displays the migration potential of different groups as assessed by wound healing (scratch) assay. While induction of NF-kB signalling in CESC by TNF- α increased the migration capacity, the EuESCs with higher level of NF-kB expression had lower ability of migration compared to control that was further reduced after treatment with Asp/AE.

Discussion

Considerable evidence demonstrated that NF-kB signaling and its reported ligand, TNF- α , have the most significant effect on endometriosis (33, 34). TNF- α , as an inflammatory cytokine, contributes to inflammation induction through activating sev-

eral transcription factors such as NF-kB (35). In this study, an attempt was made to investigate whether TNF- α could be associated with inflammatory responses in EuESCs. First, it was shown that EuESCs compared with CESC secrete higher levels of TNF- α . Next, higher levels of NF-kB gene expression in EuESCs were displayed compared to CESC. These observations confirmed the hypothesis that NF-kB and TNF- α contribute to formation of an autocrine self-booster cycle and sustain inflammatory reaction including increased capacity of ESCs for adhesion to ectopic sites, neo-vascularization, apoptosis resistance, and proliferation (36). The association between endometriosis and NF-kB signaling was confirmed by the fact that most treatments for endometriosis target NF-kB signaling (37). Therefore, drugs suppressing NF-kB pathway appear to potentially affect endometriosis progression.

In our study, Asp and AE similarly reduced the level of NF-kB gene expression in EuESCs. In contrast, treatment of CESC with TNF- α significantly increased NF-kB expression to levels even higher than those observed in EuESCs, confirming the synergistic role of TNF- α and NF-kB to form the mentioned self-booster cycle. For Asp, it was revealed that its NF-kB-inhibiting role is mainly mediated by preventing the IKK activity. IKK separates the NF-kB from its inhibitor (IKB) and facilitates the NF-kB activation (38). AE also reduces the nuclear translocation of NF-kB and decreases its DNA binding activity (39). In this way, AE stops NF-kB signaling, disrupts the autocrine amplification cycle, and ultimately, reduces NF-kB gene expression.




Figure 8. Migratory behavior in EuESCs of patients with stage IV endometriosis under Asp or AE treatment with *in vitro* wound healing assay, A) microscopic images of migration (Magnification \times 100). ESCs were cultured to 100% confluence and the *in vitro* repair of wounded cells was checked after 0, 6, 12, 18, 24 and 48 hr, B) the statistical curve of calculated migration distance showing reduced capacity of EuESCs for migration compared with CESC. Treatment with Asp or AE further decreased this capacity in EuESCs. Statistical analysis was performed at 24 hr in which the CESC closed the wound

Our results implied that EuESCs and those CESC treated with TNF- α have a high capacity for proliferation, whereas, EuESCs in the presence of Asp/AE, showed much lower proliferation. Studies on the effect of NF- κ B inhibition in a mouse model produced similar results and revealed that both the number and size of the ectopic endometriosis lesions were decreased (40, 41). These results showed that high proliferative capacity of endometriotic cells appears to be controlled by synergistic effect of increased NF- κ B, as a main regulator of cell proliferation, and high concentration of TNF- α in these cells. Treatment with Asp/AE had the same effect on EuESCs pro-

liferation and reduced its level to that of the controls.

Asp, as an efficient medication for treatment of several cancer types (42, 43), is likely to act through adjustment of NF- κ B signaling and simultaneous inhibition of proliferation and regulation of apoptosis (44, 45). AE was also shown to exert an anti-cancer effect in different types of cancer cell lines (46). The therapeutic function of emodin is supposed to be caused by inhibiting EMT and subsequent suppression of cell proliferation and migration (47, 48), all of which are attributed to the NF- κ B inhibition.

Our finding for the higher adhesive capacity of

EuESCs and CESCes treated with TNF- α compared with the untreated CESCes is consistent with previous reports. It was illustrated that endometriosis, as well as TNF- α alone, can significantly increase the expression of cell adhesion genes in ESCs probably through induction of NF- κ B pathway at early stage of inflammation (49, 50). In addition, increased level of I-CAM and V-CAM expression was also shown to be associated with development of invasive activity in cancer cells, including increased cell capacity for migration and proliferation (51).

As expected, the upregulated levels of I-CAM and V-CAM found for EuESCs were significantly suppressed when the EuESCs were treated with Asp and AE. The role of low-dose Asp on preventing the adhesive phenotype of rat platelets was reported previously (52) which confirms our results. Emodin was also revealed to inhibit cancer through reduction of I-CAM, V-CAM, and angiogenesis in human monocytes (53). Furthermore, the expression of I-CAM and V-CAM in EuESCs treated with ASP or AE was even lower than that of the healthy CESCes, although protein expression did not vary significantly. Considering the main physiological role of adhesive molecules which are highly distributed throughout the endometrium during menstruation (54) and implantation (55), there is the possibility of abnormal suppression of these molecules in endometrial cells after treatment with ASP or AE.

Unexpectedly, our findings showed that the invasive capacity of the EuESCs in stage IV endometriosis is not much different from that of the CESCes cells. This result was not consistent with some previous reports indicating higher invasive phenotype of ESCs in endometriosis patients compared with healthy controls (11, 56). Considering the important role of MMPs in metastasis and induction of neo-angiogenesis, they were supposed to have a critical role in pathogenesis of endometriosis (57). However, involvement of MMPs in etiology of endometriosis seems to follow a stage-dependent pattern. In terms of CD133, as a tumor-initiating cell marker, there is little evidence showing the association between certain subsets of endometriosis development and CD133 expression (58). Nevertheless, the results reported by D'Amico et al. confirm our finding, suggesting that CD133 cannot be used as an appropriate cell marker for endometriosis (59).

Our data on cell migration were also somewhat

unexpected, except for those obtained for the TNF- α . It was shown that CESCes, when treated with TNF- α , exhibit higher migration capacity. In contrast, EuESCs displayed lower migration potential compared with the controls, and when treated with ASP or AE, their low ability for migration was further reduced. Nevertheless, contradictory reports showed that the endometriotic cells demonstrate higher migration capacity (60) which is regarded as the main cause for ectopic endometrial growth and wound expansion (61).

In this study, EuESCs at this severe stage have a quite invasive migratory profile. Consistent with our result, Uzan et al. showed that the ovarian endometrioma in women with advanced endometriosis has lower levels of invasive cell markers (MT1-MMP and MMP-2) (62). Accordingly, Liu et al. indicated that the endometriotic cells at different stages of disease display distinct behavior (57). Considering these reports, in the early stages of the disease, cells require higher migration capacity and invasion while following the formation of wounds, the cell is likely to show a higher proliferative capacity. This explains why, despite the intrinsic ability of endometriotic cells to grow and invade, the growth of ovarian endometrioma cysts occurs gradually and over a long period of time.

Here, the effects of AE, as a known member of traditional medicine, as well as Asp, a chemical cure with less complication on human EuESCs (at stage IV endometriosis) activity were investigated and compared with that of healthy CESCes. It was demonstrated that the compounds have a similar effect in terms of inhibiting NF- κ B activation and controlling the invasive behavior of endometriotic cells. This evidence suggests the effectiveness of such anti-inflammatory drugs in controlling endometriosis. Based on the results of this study, AE or Asp at tested doses, reduced the expression levels of adhesion and invasion genes, as well as the immigration capacity of the EuESCs compared to control levels. Due to cyclic pattern of the growth and abscission in the normal ESCs during the menstrual cycle, a threshold level of migration and adhesion is required for their accurate functioning as any changes in these functions in ESCs can be more important than other somatic cells. Therefore, more detailed safety assessments on healthy CESCes are recommended.

Conclusion

In conclusion, considering obvious proliferative

and adhesive phenotype of EuESCs beside their low capacity for migration and invasion, the term "semi-invasive" instead of "invasive" is suggested for description of EuESCs activity. Furthermore, it is hypothesized that human ESCs at stage IV of endometriosis are more involved in developing the old lesions rather than creation of new ones. This may be a cause for the slow progression of endometriosis, as a benign disease, in advanced stages. Our results clearly showed that inhibition of NF- κ B, as a master regulator of inflammation, can be considered an effective treatment strategy to control endometriosis. A chemical-and an herbal-based inhibitor was used in this study and our results may further confirm the wonderful healing effects of Asp and importance of traditional medicine in management of this disease. However, any possible toxic effects of these prodrugs on ESCs should be checked. In addition, further studies on cells at different stages of the disease are needed to first examine the exact behavior of the cells and then the effects of inflammatory inhibitors on the endometriotic cells' activity at each stage.

Acknowledgement

Authors thank all the patients who participated in this study and staff of Arash Women's Hospital and Royan Institute for their help in patient's management and sampling, particularly Dr Reihaneh Hosseini and Dr Mehri Mashayekhi, Mrs Morovati and Ms Ahmadi.

Conflict of Interest

The authors declare no competing financial interest.

Funding: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

- Alimi Y, Iwanaga J, Loukas M, Tubbs RS. The clinical anatomy of endometriosis: a review. *Cureus*. 2018;10(9):e3361.
- Vercellini P, Vigano P, Somigliana E, Fedele L. Endometriosis: pathogenesis and treatment. *Nat Rev Endocrinol*. 2014;10(5):261-75.
- Mosher AA, Tsoulis MW, Lim J, Tan C, Agarwal SK, Leyland NA, et al. Melatonin activity and receptor expression in endometrial tissue and endometriosis. *Hum Reprod*. 2019;34(7):1215-24.
- Burney RO, Giudice LC. Pathogenesis and pathophysiology of endometriosis. *Fertil Steril*. 2012;98(3):511-9.
- Chui MH, Wang TL, Shih IM. Endometriosis: benign, malignant, or something in between? *Oncotarget*. 2017;8(45):78263-4.
- Filippi I, Carrarelli P, Luisi S, Batteux F, Chapron C, Naldini A, et al. Different expression of hypoxic and angiogenic factors in human endometriotic lesions. *Reprod Sci*. 2016;23(4):492-7.
- Liu H, Zhang Z, Xiong W, Zhang L, Xiong Y, Li N, et al. Hypoxia-inducible factor-1 α promotes endometrial stromal cells migration and invasion by upregulating autophagy in endometriosis. *Reproduction*. 2017;153(6):809-20.
- Rekker K, Tasa T, Saare M, Samuel K, Kadastik U, Karro H, et al. Differentially-expressed miRNAs in ectopic stromal cells contribute to endometriosis development: the plausible role of miR-139-5p and miR-375. *Int J Mol Sci*. 2018;19(12):3789.
- Li J, Ma J, Fei X, Zhang T, Zhou J, Lin J. Roles of cell migration and invasion mediated by Twist in endometriosis. *J Obstet Gynaecol Res*. 2019;45(8):1488-96.
- Wang Y, Nicholes K, Shih IM. The origin and pathogenesis of endometriosis. *Annu Rev Pathol*. 2020;15:71-95.
- Banu SK, Lee J, Speights Jr V, Starzinski-Powitz A, Arosh JA. Cyclooxygenase-2 regulates survival, migration, and invasion of human endometriotic cells through multiple mechanisms. *Endocrinology*. 2008;149(3):1180-9.
- Hirakawa T, Nasu K, Abe W, Aoyagi Y, Okamoto M, Kai K, et al. miR-503, a microRNA epigenetically repressed in endometriosis, induces apoptosis and cell-cycle arrest and inhibits cell proliferation, angiogenesis, and contractility of human ovarian endometriotic stromal cells. *Hum Reprod*. 2016;31(11):2587-97.
- Bartley J, Jülicher A, Hotz B, Mechsner S, Hotz H. Epithelial to mesenchymal transition (EMT) seems to be regulated differently in endometriosis and the endometrium. *Arch Gynecol Obstet*. 2014;289(4):871-81.
- Matsuzaki S, Darcha C. Epithelial to mesenchymal transition-like and mesenchymal to epithelial transition-like processes might be involved in the pathogenesis of pelvic endometriosis. *Hum Reprod*. 2012;27(3):712-21.
- Yáñez RA, González MM. [Endometriosis: pathophysiology and investigation trends (first part)]. *Ginecol Obstet Mex*. 2007;75(8):477-83.
- Xu X, Zheng Q, Zhang Z, Zhang X, Liu R, Liu R, et al. Periostin enhances migration, invasion, and adhesion of human endometrial stromal cells through integrin-linked kinase 1/Akt signaling pathway. *Reprod Sci*. 2015;22(9):1098-106.

17. Delbandi AA, Mahmoudi M, Shervin A, Akbari E, Jeddi-Tehrani M, Sankian M, et al. Eutopic and ectopic stromal cells from patients with endometriosis exhibit differential invasive, adhesive, and proliferative behavior. *Fertil Steril.* 2013;100(3):761-9.
18. Wu MH, Wang CA, Lin CC, Chen LC, Chang WC, Tsai SJ. Distinct regulation of cyclooxygenase-2 by interleukin-1 β in normal and endometriotic stromal cells. *J Clin Endocrinol Metab.* 2005;90(1):286-95.
19. Juhasz-Böess I, Fischer C, Latruch C, Skrzypczak M, Malik E, Ortman O, et al. Endometrial expression of estrogen receptor β and its splice variants in patients with and without endometriosis. *Arch Gynecol Obstet.* 2011;284(4):885-91.
20. Noble LS, Takayama K, Zeitoun KM, Putman JM, Johns DA, Hinshelwood MM, et al. Prostaglandin E2 stimulates aromatase expression in endometriosis-derived stromal cells. *J Clin Endocrinol Metab.* 1997;82(2):600-6.
21. Bulun SE, Lin Z, Imir G, Amin S, Demura M, Yilmaz B, et al. Regulation of aromatase expression in estrogen-responsive breast and uterine disease: from bench to treatment. *Pharmacol Rev.* 2005;57(3):359-83.
22. Bulun SE, Monsavais D, Pavone ME, Dyson M, Xue Q, Attar E, et al. Role of estrogen receptor- β in endometriosis. *Semin Reprod Med.* 2012;30(1):39-45.
23. Wu MY, Ho HN. The role of cytokines in endometriosis. *Am J Reprod Immunol.* 2003;49(5):285-96.
24. Taniguchi K, Karin M. NF- κ B, inflammation, immunity and cancer: coming of age. *Nat Rev Immunol.* 2018;18(5):309-24.
25. Jinawath N, Vasoontara C, Jinawath A, Fang X, Zhao K, Yap KL, et al. Oncoproteomic analysis reveals co-upregulation of RELA and STAT5 in carboplatin resistant ovarian carcinoma. *PLoS One.* 2010;5(6):e11198.
26. González-Ramos R, Defrère S, Devoto L. Nuclear factor- κ B: a main regulator of inflammation and cell survival in endometriosis pathophysiology. *Fertil Steril.* 2012;98(3):520-8.
27. González-Ramos R, Van Langendonck A, Defrère S, Lousse JC, Colette S, Devoto L, et al. Involvement of the nuclear factor- κ B pathway in the pathogenesis of endometriosis. *Fertil Steril.* 2010;94(6):1985-94.
28. Celik O, Ersahin A, Acet M, Çelik N, Baykuş Y, Deniz R, et al. Disulfiram, as a candidate NF- κ B and proteasome inhibitor, prevents endometriotic implant growing in a rat model of endometriosis. *Eur Rev Med Pharmacol Sci.* 2016;20(20):4380-9.
29. Tseng HS, Wang YF, Tzeng YM, Chen DR, Liao YF, Chiu HY, et al. Aloe-emodin enhances tamoxifen cytotoxicity by suppressing Ras/ERK and PI3K/mTOR in breast cancer cells. *Am J Chin Med.* 2017;45(2):337-50.
30. Shi GH, Zhou L. Emodin suppresses angiogenesis and metastasis in anaplastic thyroid cancer by affecting TRAF6-mediated pathways in vivo and in vitro. *Mol Med Rep.* 2018;18(6):5191-7.
31. Hsieh CC, Wang CH. Aspirin disrupts the crosstalk of angiogenic and inflammatory cytokines between 4t1 breast cancer cells and macrophages. *Mediators Inflamm.* 2018;2018:6380643.
32. Dai X, Yan J, Fu X, Pan Q, Sun D, Xu Y, et al. Aspirin inhibits cancer metastasis and angiogenesis via targeting heparanase. *Clin Cancer Res.* 2017;23(20):6267-78.
33. Peng Y, Ma J, Lin J. Activation of the CXCL16/CXCR6 axis by TNF- α contributes to ectopic endometrial stromal cells migration and invasion. *Reprod Sci.* 2019;26(3):420-7.
34. Lyu D, Tang N, Wang J, Zhang Y, Chang J, Liu Z, et al. TGR5 agonist INT-777 mitigates inflammatory response in human endometriotic stromal cells: a therapeutic implication for endometriosis. *Int Immunopharmacol.* 2019;71:93-9.
35. Zhang A, Wang G, Jia L, Su T, Zhang L. Exosome-mediated microRNA-138 and vascular endothelial growth factor in endometriosis through inflammation and apoptosis via the nuclear factor- κ B signaling pathway. *Int J Mol Med.* 2019;43(1):358-70.
36. Samimi M, Pourhanifeh MH, Mehdizadehkashi A, Eftekhar T, Asemi Z. The role of inflammation, oxidative stress, angiogenesis, and apoptosis in the pathophysiology of endometriosis: Basic science and new insights based on gene expression. *J Cell Physiol.* 2019;234(11):19384-92.
37. Barra F, Scala C, Mais V, Guerriero S, Ferrero S. Investigational drugs for the treatment of endometriosis, an update on recent developments. *Expert Opin Investig Drugs.* 2018;27(5):445-58.
38. Kim J, Lee KS, Kim JH, Lee DK, Park M, Choi S, et al. Aspirin prevents TNF- α -induced endothelial cell dysfunction by regulating the NF- κ B-dependent miR-155/eNOS pathway: role of a miR-155/eNOS axis in preeclampsia. *Free Radic Biol Med.* 2017;104:185-98.
39. Sanders B, Ray AM, Goldberg S, Clark T, McDaniel HR, Atlas SE, et al. Anti-cancer effects of

- aloe-emodin: a systematic review. *J Clin Transl Res.* 2018;3(3):283-96.
40. Lu Y, Sun Q, Zheng Y, Liu X, Geng JG, Guo SW. The role of nuclear factor-kappa-B p50 subunit in the development of endometriosis. *Front Biosci (Elite Ed).* 2011;3:591-603.
 41. González-Ramos R, Van Langendonck A, Defrère S, Lousse JC, Mettlen M, Guillet A, et al. Agents blocking the nuclear factor- κ B pathway are effective inhibitors of endometriosis in an in vivo experimental model. *Gynecol Obstet Invest.* 2008;65(3):174-86.
 42. Shebl F, Sakoda L, Black A, Koshiol J, Andriole G, Grubb R, et al. Aspirin but not ibuprofen use is associated with reduced risk of prostate cancer: a PLCO study. *Br J Cancer.* 2012;107(1):207-14.
 43. Jeter JM, Han J, Martinez M, Alberts DS, Qureshi A, Feskanich D. Non-steroidal anti-inflammatory drugs, acetaminophen, and risk of skin cancer in the Nurses' Health Study. *Cancer Causes Control.* 2012;23(9):1451-61.
 44. Jin M, Li C, Zhang Q, Xing S, Kan X, Wang J. Effects of aspirin on proliferation, invasion and apoptosis of Hep-2 cells via the PTEN/AKT/NF- κ B/survivin signaling pathway. *Oncol Lett.* 2018;15(6):8454-60.
 45. Zhang X, Feng H, Du J, Sun J, Li D, Hasegawa T, et al. Aspirin promotes apoptosis and inhibits proliferation by blocking G0/G1 into S phase in rheumatoid arthritis fibroblast-like synoviocytes via downregulation of JAK/STAT3 and NF- κ B signaling pathway. *Int J Mol Med.* 2018;42(6):3135-48.
 46. Chen R, Zhang J, Hu Y, Wang S, Chen M, Wang Y. Potential antineoplastic effects of Aloe-emodin: a comprehensive review. *Am J Chin Med.* 2014;42(2):275-88.
 47. Lu J, Xu Y, Zhao Z, Ke X, Wei X, Kang J, et al. Emodin suppresses proliferation, migration and invasion in ovarian cancer cells by down regulating ILK in vitro and in vivo. *Onco Targets Ther.* 2017;10:3579-89.
 48. Li N, Wang C, Zhang P, You S. Emodin inhibits pancreatic cancer EMT and invasion by up-regulating microRNA-1271. *Mol Med Rep.* 2018;18(3):3366-74.
 49. Rahman I, Gilmour PS, Jimenez LA, MacNee W. Oxidative stress and TNF-alpha induce histone acetylation and NF-kappaB/AP-1 activation in alveolar epithelial cells: potential mechanism in gene transcription in lung inflammation. *Mol Cell Biochem.* 2002;234-235(1-2):239-48.
 50. Kuessel L, Wenzl R, Proestling K, Balendran S, Pateisky P, Yerlikaya G, et al. Soluble VCAM-1/soluble ICAM-1 ratio is a promising biomarker for diagnosing endometriosis. *Hum Reprod.* 2017;32(4):770-9.
 51. Jung WC, Jang YJ, Kim JH, Park SS, Park SH, Kim SJ, et al. Expression of intercellular adhesion molecule-1 and e-selectin in gastric cancer and their clinical significance. *J Gastric Cancer.* 2012;12(3):140-8.
 52. Lin HL, Yen HW, Hsieh SL, An LM, Shen KP. Low-dose aspirin ameliorated hyperlipidemia, adhesion molecule, and chemokine production induced by high-fat diet in Sprague-Dawley rats. *Drug Dev Res.* 2014;75(2):97-106.
 53. Heo SK, Yun HJ, Noh EK, Park SD. Emodin and rhein inhibit LIGHT-induced monocytes migration by blocking of ROS production. *Vascul Pharmacol.* 2010;53(1-2):28-37.
 54. Tawia SA, Beaton LA, Rogers PA. Immunolocalization of the cellular adhesion molecules, intercellular adhesion molecule-1 (ICAM-1) and platelet endothelial cell adhesion molecule (PECAM), in human endometrium throughout the menstrual cycle. *Hum Reprod.* 1993;8(2):175-81.
 55. Thomson AJ, Greer MR, Young A, Boswell F, Telfer JF, Cameron IT, et al. Expression of intercellular adhesion molecules ICAM-1 and ICAM-2 in human endometrium: regulation by interferon-gamma. *Mol Hum Reprod.* 1999;5(1):64-70.
 56. Wang D, Liu Y, Han J, Zai D, Ji M, Cheng W, et al. Puerarin suppresses invasion and vascularization of endometriosis tissue stimulated by 17 β -estradiol. *Plos One.* 2011;6(9):e25011.
 57. Liu H, Wang J, Wang H, Tang N, Li Y, Zhang Y, et al. Correlation between matrix metalloproteinase-9 and endometriosis. *Int J Clin Exp Pathol.* 2015;8(10):13399-404.
 58. Liu S, Zhou J, Wen J. Expression and significance of CD133 and ABCG2 in endometriosis. *Clin Exp Obstet Gynecol.* 2015;42(6):771-5.
 59. D'Amico F, Skarmoutsou E, Quaderno G, Malaponte G, La Corte C, Scibilia G, et al. Expression and localisation of osteopontin and prominin-1 (CD133) in patients with endometriosis. *Int J Mol Med.* 2013;31(5):1011-6.
 60. Au HK, Chang JH, Wu YC, Kuo YC, Chen YH, Lee WC, et al. TGF- β I regulates cell migration through pluripotent transcription factor OCT4 in endometriosis. *PloS One.* 2015;10(12):e0145256.
 61. Chang JH, Au HK, Lee WC, Chi CC, Ling TY, Wang LM, et al. Expression of the pluripotent transcription factor OCT4 promotes cell migration

in endometriosis. *Fertil Steril.* 2013;99(5):1332-9.
e5.

62. Uzan C, Cortez A, Dufournet C, Fauvet R, Siffroi JP, Darai E. Eutopic endometrium and peritoneal,

ovarian and bowel endometriotic tissues express a different profile of matrix metalloproteinases-2, -3 and -11, and of tissue inhibitor metalloproteinases-1 and -2. *Virchows Arch.* 2004;445(6):603-9.