

Letter to the Editor

Additional Measurement Approaches for Sleep Disturbances. Comment on “A Transdiagnostic Self-management Web-Based App for Sleep Disturbance in Adolescents and Young Adults: Feasibility and Acceptability Study”

Wan-Tong Tsai; Tzung-Liang Liu

Chung Shan Medical University, Taichung City, Taiwan

Corresponding Author:

Wan-Tong Tsai
Chung Shan Medical University
No 110, Sec 1, Jianguo N Rd
South District
Taichung City, 40201
Taiwan
Phone: 886 968938360
Email: science.tsai@gmail.com

Related Articles:

Comment on: <http://www.jmir.org/2021/11/e25392/>

Comment in: <http://mhealth.jmir.org/2022/6/e39198/>

(*JMIR Form Res* 2022;6(6):e35959) doi: [10.2196/35959](https://doi.org/10.2196/35959)

KEYWORDS

youth; sleep; technology; mHealth; self-management; adolescents; young adults; mobile phone; smartphone; polysomnography

Carmona et al [1] analyzed the effect of DOZE (Delivering Online Zzz's with Empirical Support) on adolescents and young adults. However, this was done only via feedback from the participants. To make measurements more quantitative and less subjective, we propose a few scientific measurement approaches.

Polysomnography [2], a method to measure one's physiological state during sleep, is a common way to assess sleep quality. With the following devices, 7 parts of the physiological state can be detected:

1. Electroencephalogram: Multiple brain electrode patches are placed on the scalp to record the various stages of sleep. Thus, we can obtain a brainwave diagram for analysis.
2. Electromyography: 4 electrode patches are used to monitor muscle tension. Muscle tension will decrease significantly during sleep. We can also detect whether periodic limb movement disorder occurs and then compare the results with other data.
3. Electrocardiography: this can record the activity of the heart, including T waves, P waves, and QRS waves.
4. Electrooculography: this can record the potential difference between the cornea and the retina, so that we know when

the rapid eye movement period occurs, allowing us to judge the stages of sleep.

5. Oxygen saturation: by recording this, we can know if the body remains in good condition during sleep.
6. Thoracic abdominal effort.
7. Nasal and oral airflow.

In addition, a serology examination can also be applied to quantify the survey. For instance, we can detect the melatonin levels of participants, so as to detect fluctuations from baseline to the end point. Thus, we can see if the concentration of melatonin reaches appropriate levels after participants use a web-based self-management app [3].

To keep the self-management app easy to use and convenient, any approach that requires specific tests in a hospital should not be used. Regardless, there are some take-home tests that can be administered easily. Examples include self-adhesive electrodes; heart rate monitoring equipment; or even techniques like the Freestyle Libre Pro for glucose monitoring [4], which can be used to track the physiological state of each participant.

In summary, quantifying the influence of DOZE with physiological states could make it more credible, and monitoring tests can be put into practice feasibly.

Conflicts of Interest

None declared.

References

1. Carmona NE, Usyatynsky A, Kutana S, Corkum P, Henderson J, McShane K, et al. A transdiagnostic self-management web-based app for sleep disturbance in adolescents and young adults: Feasibility and acceptability study. *JMIR Form Res* 2021 Nov 01;5(11):e25392 [FREE Full text] [doi: [10.2196/25392](https://doi.org/10.2196/25392)] [Medline: [34723820](https://pubmed.ncbi.nlm.nih.gov/34723820/)]
2. Jafari B, Mohsenin V. Polysomnography. *Clin Chest Med* 2010 Jun;31(2):287-297. [doi: [10.1016/j.ccm.2010.02.005](https://doi.org/10.1016/j.ccm.2010.02.005)] [Medline: [20488287](https://pubmed.ncbi.nlm.nih.gov/20488287/)]
3. Hsing AW, Meyer TE, Niwa S, Quraishi SM, Chu LW. Measuring serum melatonin in epidemiologic studies. *Cancer Epidemiol Biomarkers Prev* 2010 Apr;19(4):932-937 [FREE Full text] [doi: [10.1158/1055-9965.EPI-10-0004](https://doi.org/10.1158/1055-9965.EPI-10-0004)] [Medline: [20332275](https://pubmed.ncbi.nlm.nih.gov/20332275/)]
4. Blum A. Freestyle Libre glucose monitoring system. *Clin Diabetes* 2018 Apr;36(2):203-204 [FREE Full text] [doi: [10.2337/cd17-0130](https://doi.org/10.2337/cd17-0130)] [Medline: [29686463](https://pubmed.ncbi.nlm.nih.gov/29686463/)]

Abbreviations

DOZE: Delivering Online Zzz's with Empirical Support

Edited by T Leung; this is a non-peer-reviewed article. Submitted 24.12.21; accepted 05.05.22; published 13.06.22.

Please cite as:

Tsai WT, Liu TL

Additional Measurement Approaches for Sleep Disturbances. Comment on "A Transdiagnostic Self-management Web-Based App for Sleep Disturbance in Adolescents and Young Adults: Feasibility and Acceptability Study"

JMIR Form Res 2022;6(6):e35959

URL: <https://formative.jmir.org/2022/6/e35959>

doi: [10.2196/35959](https://doi.org/10.2196/35959)

PMID:

©Wan-Tong Tsai, Tzung-Liang Liu. Originally published in *JMIR Formative Research* (<https://formative.jmir.org>), 13.06.2022. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in *JMIR Formative Research*, is properly cited. The complete bibliographic information, a link to the original publication on <https://formative.jmir.org>, as well as this copyright and license information must be included.