

CORRECTION

Open Access



Correction: External validation of a new predictive model for falls among inpatients using the official Japanese ADL scale, Bedriddenness ranks: a double-centered prospective cohort study

Masaki Tago^{1*}, Naoko E. Katsuki¹, Eiji Nakatani^{2,3}, Midori Tokushima¹, Akiko Dogomori¹, Kazumi Mori¹, Shun Yamashita¹, Yoshimasa Oda⁴ and Shu-ichi Yamashita¹

Correction: *BMC Geriatrics* 22, 331 (2022)

<https://doi.org/10.1186/s12877-022-02871-5>

After publication of this article [1], the authors reported that there was a mistake in the formula of model 2 in Additional file 3. The correct formula of model 2 is as follows:

$$\begin{aligned} & - 5.8563 + 0.0096 \times (\text{Age}) + (\text{Male} = 0.5684) + (\text{Emergency admission} = 0.4418) + (\text{Admitted department; Neurosurgery} = 0.6520) \\ & + (\text{Hypnotics; Using} = 0.3612, \text{Missing data} = 0.2139) + (\text{History of fall} = 0.4362) + (\text{Ability of eating; Independent} = 0.2352, \text{Missing data} = -1.0436) \\ & + (\text{Bedriddenness rank; J} = 1.3758, \text{A} = 1.8317, \text{B} = 1.9186, \text{C} = 1.7205, \text{Not assessable} = -0.1505) \end{aligned}$$

The original article [1] has been corrected.

Author details

¹Department of General Medicine, Saga University Hospital, 5-1-1 Nabeshima, Saga 849-8501, Japan. ²Graduate School of Public Health, Shizuoka Graduate University of Public Health, Shizuoka, Japan. ³Translational Research Center for Medical Innovation, Foundation for Biomedical Research and Innovation at Kobe, Hyogo, Japan. ⁴Department of General Medicine, Yuai-Kai Foundation and Oda Hospital, Saga, Japan.

Published online: 27 July 2022

The original article can be found online at <https://doi.org/10.1186/s12877-022-02871-5>.

*Correspondence: tagomas@cc.saga-u.ac.jp

¹Department of General Medicine, Saga University Hospital, 5-1-1 Nabeshima, Saga 849-8501, Japan

Full list of author information is available at the end of the article

Reference

1. Tago M, et al. External validation of a new predictive model for falls among inpatients using the official Japanese ADL scale, Bedriddenness ranks: a double-centered prospective cohort study. *BMC Geriatrics*. 2022;22:331. <https://doi.org/10.1186/s12877-022-02871-5>.



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.