

Received 29 May 2015
Accepted 29 June 2015

Edited by M. Zeller, Youngstown State University, USA

Keywords: crystal structure; iridium(III); cyclometalated; 4,4':2',2'':4'',4'''-quaterpyridyl ligand

CCDC reference: 1409461

Supporting information: this article has supporting information at journals.iucr.org/e

Crystal structure of [1,1''-bis(pyrimidin-2-yl)-4,4':2',2'':4'',4'''-quaterpyridine-1,1''-diium- $\kappa^2N^{1'},N^{1''}$]bis[2-(pyridin-2-yl)phenyl- κ^2N,C^1]-iridium(III) tris(hexafluoridophosphate) acetonitrile trisolvate

Benjamin J. Coe,^{a*} Martyn K. Peers,^b James Raftery^a and Nigel S. Scrutton^b



^aSchool of Chemistry, University of Manchester, Manchester M13 9PL, England, and ^bManchester Institute of Biotechnology, Faculty of Life Sciences, University of Manchester, 131 Princess Street, Manchester M1 7DN, England.

*Correspondence e-mail: b.coe@manchester.ac.uk

In the title compound, $[Ir(C_{11}H_8N)_2(C_{28}H_{20}N_8)](PF_6)_3 \cdot 3CH_3CN$ or $[Ir^{III}(ppy)_2((2-pym)_2qpy^{2+})](PF_6)_3 \cdot 3CH_3CN$ (ppy = deprotonated 2-phenylpyridine, pym = pyrimidyl and qpy = 4,4':2',2'':4'',4'''-quaterpyridyl), the Ir³⁺ cation is coordinated by two C atoms and four N atoms in a slightly distorted octahedral geometry. The asymmetric unit consists of one complex trication, three hexafluoridophosphate anions and three acetonitrile solvent molecules. The average Ir—C distance is 2.011 (14) Å, the average Ir—N(ppy) distance is 2.05 (6) Å and the average Ir—N(qpy) distance is longer at 2.132 (10) Å. The dihedral angles within the 4,4'-bipyridyl units are 31.5 (6) and 23.8 (7) $^\circ$, while those between the 2-pym and attached pyridyl rings are rather smaller, at 11.7 (9) and 7.1 (9) $^\circ$. The title compound was refined as a two-component inversion twin.

1. Chemical context

Iridium complexes of cyclometalating ligands have been studied widely, mainly due to their interesting photophysical properties (Flamigni *et al.*, 2007; You & Nam, 2012; Ladouceur & Zysman-Colman, 2013). Complexes of the form $[Ir^{III}(ppy)_2(N-N)]^+$ ($N-N$ = 2,2'-bipyridyl or a related α -diimine ligand) are well known, and many examples have been structurally characterized (*e.g.* Ladouceur *et al.*, 2010; Zhao *et al.*, 2010; Constable *et al.*, 2013; Schneider *et al.*, 2014).



However, such compounds containing ligands with pyridinium substituents are scarce, and the only ones reported to our knowledge are the complex salts $[Ir^{III}(C-N)_2(Me_2qpy^{2+})][PF_6]_3$ ($L-L$ = ppy or benzo[*h*]quoline) (Ahmad *et al.*, 2014). We report here a related new compound and what appears to be the first X-ray crystal structure determination of an iridium complex containing a qpy-based ligand.




Figure 1

View of the molecular components of (I) (50% probability displacement ellipsoids)

2. Structural commentary

The molecular structure (Fig. 1) of the complex cation in $[\text{Ir}^{\text{III}}(\text{ppy})_2\{(\text{2-pym})_2\text{qpy}^{2+}\}][\text{PF}_6]_3 \cdot 3\text{CH}_3\text{CN}$ (I) is as indicated by ^1H NMR spectroscopy, with a slightly distorted octahedral coordination geometry. The bite angle of the qpy-based ligand is $76.6(2)^\circ$, while those of the ppy ligands are slightly larger at $80.1(6)$ and $80.8(5)^\circ$. As for other related complexes (Ladouceur *et al.*, 2010; Zhao *et al.*, 2010; Constable *et al.*, 2013; Schneider *et al.*, 2014), the strong *trans* effects of a σ -bonded phenyl ring (Coe & Glenwright, 2000) causes these units to adopt a *cis* orientation, so that the pyridyl rings of the ppy




Figure 2

Crystal packing diagram, viewed approximately along the b axis, showing the alignment of the qpy fragments. The H atoms, PF_6^- anions and acetonitrile solvent molecules have been removed for clarity.

Table 1
Selected geometric parameters (\AA , $^\circ$).

C35–Ir1	2.021 (16)	Ir1–N9	2.095 (13)
C46–Ir1	2.000 (15)	Ir1–N1	2.125 (11)
Ir1–N10	2.011 (14)	Ir1–N5	2.139 (11)
C46–Ir1–N10	80.8 (5)	C35–Ir1–N1	172.7 (6)
C46–Ir1–C35	86.0 (3)	N9–Ir1–N1	93.8 (4)
N10–Ir1–C35	94.2 (6)	C46–Ir1–N5	173.1 (6)
C46–Ir1–N9	93.9 (5)	N10–Ir1–N5	94.4 (5)
N10–Ir1–N9	172.5 (2)	C35–Ir1–N5	99.4 (6)
C35–Ir1–N9	80.1 (6)	N9–Ir1–N5	91.3 (4)
C46–Ir1–N1	98.4 (6)	N1–Ir1–N5	76.6 (2)
N10–Ir1–N1	92.3 (5)		

ligands are oriented *trans*. The structural *trans* effect of the phenyl rings is shown by the *ca* 0.08 \AA lengthening of the Ir–N(qpy) distances [average = $2.132(10) \text{ \AA}$] with respect to the Ir–N(ppy) ones [average = $2.05(6) \text{ \AA}$]. The Ir–C distances (Table 1) are shorter still, with an average value of $2.01(14) \text{ \AA}$. All of the geometric parameters around the Ir^{3+} cation are similar to those reported for related structures.

The dihedral angles within the 4,4'-bipyridyl units in (I) are larger than those [$20.8(6)$ and $21.0(5)^\circ$] in the only other structurally characterized complex of the $(2\text{-pym})_2\text{qpy}^{2+}$ ligand, $[\text{Ru}^{\text{II}}(\text{bpy})_2\{(\text{2-pym})_2\text{qpy}^{2+}\}][\text{PF}_6]_4$ (bpy = 2,2'-bipyridyl) (Coe *et al.*, 2011). On the other hand, the dihedral angles between the 2-pyrimidyl and attached pyridyl rings are closely similar in (I), whereas two quite different such angles are observed in $[\text{Ru}^{\text{II}}(\text{bpy})_2\{(\text{2-pym})_2\text{qpy}^{2+}\}][\text{PF}_6]_4$ [$6.0(9)$ and $20.0(5)^\circ$].




Figure 3

Crystal packing diagram, viewed approximately along the a axis, showing the antiparallel alignment of the molecular dipoles (represented by arrows for the extreme left and right complexes). The H atoms, PF_6^- anions and acetonitrile solvent molecules have been removed for clarity.

3. Supramolecular features

The unit cell contains four complex cations with their qpy units aligned approximately parallel (Fig. 2). There may be a weak π -stacking interaction between a 2-pym ring and one of the rings of the bpy fragment in an adjacent complex, with a centroid-to-centroid distance of 3.854 (8) Å and a dihedral angle of 9.8 (6)°. Ru^{II} complexes of (2-pym)₂qpy²⁺ and related ligands show interesting non-linear optical (NLO) properties (Coe *et al.* 2005). In this context, crystal packing arrangements are of great importance because macroscopic polarity is necessary for the existence of bulk quadratic NLO effects. The space group *Cc* adopted by (I) is non-centrosymmetric, potentially affording a polar material that could display such NLO properties. However, the overall orientation of the dipoles formed by the electron-donating Ir^{III}(ppy)₂ units and the accepting (2-pym)₂qpy²⁺ ligands is antiparallel (Fig. 3). Therefore, significant bulk quadratic NLO behaviour is not expected for this particular crystal form.

4. Synthesis and crystallization

The new compound (I) was synthesised simply by cleaving the commercial chloride-bridged dimer [Ir^{III}(ppy)₂Cl]₂ with the proligand salt [(2-pym)₂qpy²⁺]Cl₂ (Coe *et al.*, 2011) in refluxing 2-methoxyethanol/water.

[Ir^{III}(ppy)₂Cl]₂ (40 mg, 0.037 mmol) and *N,N'*,*N'''*-di(2-pyrimidyl)-4,4':2',2''-4'',4'''-quaterpyridinium chloride·2.3H₂O (47 mg, 0.081 mmol) in argon-sparged 2-methoxyethanol/water (3:1, 10 ml) were heated at reflux for 20 h. After cooling to room temperature, the solvent was removed by rotary evaporation and the residue redissolved in a minimum volume of methanol to which was added an excess of solid NH₄PF₆. Cold water was added and the precipitate was filtered off and washed with water. The product was purified by column chromatography on silica gel, eluting with 0.1 M NH₄PF₆ in acetonitrile, to afford a brown-green solid. Yield: 68 mg (65%). Analysis calculated for C₅₀H₃₆F₁₈IrN₁₀P₃·H₂O: C 42.2, H 2.7, N 9.8%; found: C 42.0, H 2.5, N 9.6%. Spectroscopic analysis: ¹H NMR (400 MHz, CD₃CN, δ , p.p.m.) 10.12 (4H, *dd*, *J* = 7.5, 1.9 Hz), 9.18 (2H, *d*, *J* = 1.4 Hz), 9.13 (4H, *d*, *J* = 4.9 Hz), 8.68 (4H, *dd*, *J* = 7.4, 1.8 Hz), 8.31 (2H, *d*, *J* = 5.6 Hz), 8.13 (2H, *dt*, *J* = 8.1, 0.8 Hz), 8.05 (2H, *dd*, *J* = 5.7, 1.8 Hz), 7.93–7.87 (8H), 7.71 (2H, *ddd*, *J* = 5.9, 1.5, 0.7 Hz), 7.14–6.98 (8H), 6.33 (2H, *dd*, *J* = 7.6, 0.9 Hz). MALDI-MS *m/z* = 1405 ($\{M\}^+$), 1260 ($\{M - PF_6\}^+$), 1115 ($\{M - 2PF_6\}^+$), 970 ($\{M - 3PF_6\}^+$).

Single crystals (amber plates) suitable for X-ray diffraction studies were grown by slow diffusion of diethyl ether vapour into an acetonitrile solution at room temperature.

5. Other Characterization

The complex salt (I) shows a relatively weak, broad visible absorption band at $\lambda_{\text{max}} = 562$ nm ($\varepsilon = 1,800 \text{ M}^{-1} \text{ dm}^3$) in acetonitrile. Based on the results of time-dependent density functional theory (TD-DFT) calculations on the related

Table 2
Experimental details.

Crystal data	
Chemical formula	[Ir(C ₁₁ H ₈ N) ₂ (C ₂₈ H ₂₀ N ₈)](PF ₆) ₃ ·3C ₂ H ₃ N
<i>M</i> _r	1527.16
Crystal system, space group	Monoclinic, <i>Cc</i>
Temperature (K)	100
<i>a</i> , <i>b</i> , <i>c</i> (Å)	22.2647 (16), 14.6139 (11), 18.6288 (14)
β (°)	102.447 (1)
<i>V</i> (Å ³)	5918.9 (8)
<i>Z</i>	4
Radiation type	Mo <i>K</i> α
μ (mm ⁻¹)	2.45
Crystal size (mm)	0.25 × 0.20 × 0.03
Data collection	
Diffractometer	Bruker SMART CCD area detector
Absorption correction	Multiscan (SADABS; Bruker, 2001)
<i>T</i> _{min} , <i>T</i> _{max}	0.697, 0.930
No. of measured, independent and observed [<i>I</i> > 2 <i>σ</i> (<i>I</i>)] reflections	25080, 13146, 11295
<i>R</i> _{int}	0.037
(sin θ/λ) _{max} (Å ⁻¹)	0.669
Refinement	
<i>R</i> [$F^2 > 2\sigma(F^2)$], <i>wR</i> (F^2), <i>S</i>	0.041, 0.093, 1.00
No. of reflections	13146
No. of parameters	824
No. of restraints	434
H-atom treatment	H-atom parameters constrained
$\Delta\rho_{\text{max}}$, $\Delta\rho_{\text{min}}$ (e Å ⁻³)	1.24, -1.00
Absolute structure	Refined as an inversion twin
Absolute structure parameter	0.384 (7)

Computer programs: SMART and SAINT-Plus (Bruker, 2003), SHELXTL (Sheldrick, 2008) and SHELXL2014/7 (Sheldrick, 2015).

complex [Ir^{III}(ppy)₂(Me₂qpy²⁺)]³⁺ (Peers, 2012), this absorption is attributable to d→π* metal-to-ligand charge-transfer (MLCT) transitions directed towards the qpy-based ligand, with significant contributions by the ppy ligands to the donor orbitals introducing also ligand-to-ligand CT character. Below 500 nm, absorption increases steadily into the UV region, with another maximum at 378 nm ($\varepsilon = 16,600 \text{ M}^{-1} \text{ dm}^3$), and a shoulder at ca 410 nm. By way of contrast, the lowest energy band for [Ir^{III}(ppy)₂(Me₂qpy²⁺)][PF₆]₃ appears at $\lambda_{\text{max}} = 531$ nm ($\varepsilon = 1,200 \text{ M}^{-1} \text{ dm}^3$) in acetonitrile (Peers, 2012). The substantial red-shift of this band on moving to (I) is due to the enhanced electron-accepting ability of the *N*-(2-pyrimidyl)pyridinium groups. The higher intensity for (I) is a consequence of extended π-conjugation involving the 2-pym rings.

Cyclic voltammetric studies on (I) reveal an irreversible oxidation process at $E_{\text{pa}} = 1.43$ V vs Ag–AgCl {acetonitrile, 0.1 M [N(*n*-Bu₄)PF₆, 2 mm Pt disc working electrode, 100 mV s⁻¹, ferrocene/ferrocenium standard at 0.44 V ($\Delta E_p = 70$ –90 mV)}. The reductive region shows a reversible wave at $E_{1/2} = -0.29$ V ($\Delta E_p = 80$ mV), followed by an irreversible process with $E_{\text{pc}} = -0.79$ V. Based on the relative peak currents, the reversible wave is assigned as a two-electron process involving reduction of both pyridinium units. The redox behaviour of this complex can be rationalized with the aid of DFT results obtained for [Ir^{III}(ppy)₂(Me₂qpy²⁺)]³⁺

(Peers, 2012). The irreversible oxidation wave corresponds with removing an electron from the HOMO comprising the Ir and ppy ligands. The first and second reductions involve adding electrons to the LUMO based on the $(2\text{-pym})_2\text{qpy}^{2+}$ ligand. The oxidation occurs at the same E_{pa} value for (I) and its methylated analogue, $[\text{Ir}^{\text{III}}(\text{ppy})_2(\text{Me}_2\text{qpy}^{2+})][\text{PF}_6]_3$, but the first two reductions appear as overlapping reversible waves at $E_{1/2} = -0.62 \text{ V}$ ($\Delta E_p = 70 \text{ mV}$) and $E_{1/2} = -0.73 \text{ V}$ ($\Delta E_p = 60 \text{ mV}$) in the latter compound. These waves can be resolved by using differential pulse voltammetry (potential increment = 2 mV, amplitude = 50 mV, pulse width = 0.01 s). The anodic shift in the reduction waves is consistent with the qpy-based ligand being more electron-deficient, and therefore easier to reduce, in (I). The lack of splitting of these waves in (I) indicates that electronic communication between the pyridyl radicals is diminished with respect to its methyl analogue. Interestingly, for the related compound $[\text{Ru}^{\text{II}}(\text{bpy})_2\{(2\text{-pym})_2\text{qpy}^{2+}\}][\text{PF}_6]_4$, the first two reductions are irreversible under the same conditions using a glassy carbon working electrode (Coe *et al.*, 2011).

6. Refinement

The structure was solved by direct methods. The two rings of one of the ppy ligands are indistinguishable by bond lengths, and the presented structure gives the lowest R factors. Crystal twinning is present. There is a pseudo-twofold axis that manifests itself as high correlation between parameters during refinement. The non-hydrogen atoms were refined anisotropically, but a rigid bond restraint (RIGU in *SHELX*) was applied for atoms with pseudo-symmetry-related counterparts. H atoms were included in calculated positions with C–H bond lengths of 0.95 (CH), 0.99 (CH₂) and 0.98 (CH₃) Å; $U_{\text{iso}}(\text{H})$ values were fixed at $1.2U_{\text{eq}}(\text{C})$ except for CH₃ where it was $1.5U_{\text{eq}}(\text{C})$. Crystal data, data collection and structure refinement details are given in Table 2.

Acknowledgements

We thank the BBSRC for support in the form of a PhD studentship (MKP). NSS is supported by an EPSRC Established Career Fellowship (grant EP/J020192/1) and was funded by a Royal Society Wolfson Merit Award.

References

- Ahmad, H., Wragg, A., Cullen, W., Wombwell, C., Meijer, A. J. H. M. & Thomas, J. A. (2014). *Chem. Eur. J.* **20**, 3089–3096.
- Bruker (2001). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2003). *SMART* and *SAINT-Plus*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Coe, B. J. & Glenwright, S. J. (2000). *Coord. Chem. Rev.* **203**, 5–80.
- Coe, B. J., Harper, E. C., Helliwell, M. & Ta, Y. T. (2011). *Polyhedron*, **30**, 1830–1841.
- Coe, B. J., Harris, J. A., Jones, L. A., Brunschwig, B. S., Song, K., Clays, K., Garín, J., Orduna, J., Coles, S. J. & Hursthouse, M. B. (2005). *J. Am. Chem. Soc.* **127**, 4845–4859.
- Constable, E. C., Housecroft, C. E., Kopecky, P., Martin, C. J., Wright, I. A., Zampese, J. A., Bolink, H. J. & Pertegás, A. (2013). *Dalton Trans.* **42**, 8086–8103.
- Flamigni, L., Barbieri, A., Sabatini, C., Ventura, B. & Barigelli, F. (2007). *Top. Curr. Chem.* **281**, 143–203.
- Ladouceur, S., Fortin, D. & Zysman-Colman, E. (2010). *Inorg. Chem.* **49**, 5625–5641.
- Ladouceur, S. & Zysman-Colman, E. (2013). *Eur. J. Inorg. Chem.* pp. 2985–3007.
- Peers, M. K. (2012). PhD thesis, The University of Manchester, England.
- Schneider, G. E., Pertegás, A., Constable, E. C., Housecroft, C. E., Hostettler, N., Morris, C. D., Zampese, J. A., Bolink, H. J., Junquera-Hernández, J. M., Ortí, E. & Sessolo, M. (2014). *J. Mater. Chem. C*, **2**, 7047–7055.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst. C* **71**, 3–8.
- You, Y.-M. & Nam, W.-W. (2012). *Chem. Soc. Rev.* **41**, 7061–7084.
- Zhao, N., Wu, Y.-H., Shi, L.-X., Lin, Q.-P. & Chen, Z.-N. (2010). *Dalton Trans.* **39**, 8288–8295.

supporting information

Acta Cryst. (2015). E71, 879-882 [doi:10.1107/S2056989015012463]

Crystal structure of [1,1''-bis(pyrimidin-2-yl)-4,4':2',2'':4'',4'''-quaterpyridine-1,1''-diium- κ^2N^1,N^1'']bis[2-(pyridin-2-yl)phenyl- κ^2N,C^1]iridium(III) tris(hexafluoridophosphate) acetonitrile trisolvate

Benjamin J. Coe, Martyn K. Peers, James Raftery and Nigel S. Scrutton

Computing details

Data collection: SMART (Bruker, 2003); cell refinement: SAINT-Plus (Bruker, 2003); data reduction: SAINT-Plus (Bruker, 2003); program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXL2014/7 (Sheldrick, 2015); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL (Sheldrick, 2008).

[1,1''-Bis(pyrimidin-2-yl)-4,4':2',2'':4'',4'''-quaterpyridine-1,1''-diium- κ^2N^1,N^1'']bis[2-(pyridin-2-yl)phenyl- κ^2N,C^1]iridium(III) tris(hexafluoridophosphate) acetonitrile trisolvate

Crystal data

$[Ir(C_{11}H_8N)_2(C_{28}H_{20}N_8)](PF_6)_3 \cdot 3C_2H_3N$
 $M_r = 1527.16$
Monoclinic, Cc
 $a = 22.2647 (16)$ Å
 $b = 14.6139 (11)$ Å
 $c = 18.6288 (14)$ Å
 $\beta = 102.447 (1)$ °
 $V = 5918.9 (8)$ Å³
 $Z = 4$

$F(000) = 3024$
 $D_x = 1.714$ Mg m⁻³
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 5333 reflections
 $\theta = 2.2\text{--}23.6$ °
 $\mu = 2.44$ mm⁻¹
 $T = 100$ K
Plate, brown
0.25 × 0.20 × 0.03 mm

Data collection

Bruker SMART CCD area-detector
diffractometer
 φ and ω scans
Absorption correction: multi-scan
(SADABS; Bruker, 2001)
 $T_{\min} = 0.697$, $T_{\max} = 0.930$
25080 measured reflections

13146 independent reflections
11295 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.037$
 $\theta_{\max} = 28.4$ °, $\theta_{\min} = 1.7$ °
 $h = -29 \rightarrow 29$
 $k = -19 \rightarrow 19$
 $l = -24 \rightarrow 24$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.093$
 $S = 1.00$
13146 reflections
824 parameters
434 restraints

Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0493P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 1.24$ e Å⁻³
 $\Delta\rho_{\min} = -1.00$ e Å⁻³

Absolute structure: Refined as an inversion twin.

Absolute structure parameter: 0.384 (7)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refined as a 2-component inversion twin.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.2259 (6)	0.2672 (9)	0.2388 (7)	0.016 (2)
C2	0.1948 (6)	0.3457 (9)	0.2574 (7)	0.022 (3)
H2	0.2117	0.4048	0.2540	0.027*
C3	0.1400 (6)	0.3367 (9)	0.2803 (7)	0.022 (3)
C4	0.1148 (6)	0.2499 (9)	0.2826 (7)	0.021 (3)
H4	0.0773	0.2415	0.2982	0.025*
C5	0.1461 (6)	0.1756 (9)	0.2613 (8)	0.022 (3)
H5	0.1283	0.1165	0.2612	0.027*
C6	0.1124 (8)	0.4180 (9)	0.3043 (10)	0.025 (3)
C7	0.1177 (6)	0.5038 (10)	0.2731 (7)	0.029 (3)
H7	0.1422	0.5094	0.2374	0.035*
C8	0.0887 (6)	0.5808 (9)	0.2923 (8)	0.028 (3)
H8	0.0926	0.6382	0.2699	0.033*
C9	0.0490 (6)	0.4917 (9)	0.3774 (8)	0.029 (3)
H9	0.0257	0.4879	0.4144	0.034*
C10	0.0776 (6)	0.4132 (9)	0.3577 (8)	0.022 (2)
H10	0.0732	0.3564	0.3809	0.027*
C11	0.0220 (6)	0.6533 (9)	0.3638 (7)	0.024 (3)
C12	-0.0487 (6)	0.7102 (9)	0.4174 (8)	0.045 (4)
H12	-0.0782	0.7039	0.4471	0.055*
C13	-0.0393 (7)	0.7960 (10)	0.3933 (8)	0.039 (3)
H13	-0.0618	0.8481	0.4030	0.047*
C14	0.0060 (7)	0.7995 (10)	0.3538 (9)	0.045 (4)
H14	0.0163	0.8577	0.3371	0.054*
C15	0.2866 (7)	0.2711 (10)	0.2175 (9)	0.029 (3)
C16	0.3129 (6)	0.3483 (9)	0.2000 (7)	0.023 (3)
H16	0.2934	0.4058	0.2024	0.028*
C17	0.3691 (6)	0.3434 (10)	0.1782 (8)	0.025 (3)
C18	0.3968 (6)	0.2579 (10)	0.1791 (8)	0.029 (3)
H18	0.4362	0.2525	0.1675	0.035*
C19	0.3678 (6)	0.1829 (10)	0.1964 (7)	0.027 (3)
H19	0.3869	0.1249	0.1962	0.032*
C20	0.4021 (8)	0.4263 (8)	0.1574 (11)	0.027 (4)
C21	0.3913 (7)	0.5118 (11)	0.1809 (10)	0.049 (5)
H21	0.3612	0.5207	0.2094	0.058*

C22	0.4228 (9)	0.5822 (11)	0.1639 (12)	0.054 (6)
H22	0.4137	0.6414	0.1799	0.065*
C23	0.4759 (6)	0.4923 (9)	0.0972 (8)	0.033 (3)
H23	0.5045	0.4867	0.0663	0.039*
C24	0.4456 (8)	0.4188 (10)	0.1122 (10)	0.043 (4)
H24	0.4531	0.3610	0.0925	0.051*
C25	0.5026 (5)	0.6526 (9)	0.1111 (7)	0.023 (2)
C26	0.5170 (7)	0.8075 (10)	0.1180 (8)	0.039 (3)
H26	0.5070	0.8670	0.1324	0.047*
C27	0.5625 (7)	0.7970 (10)	0.0824 (9)	0.041 (3)
H27	0.5830	0.8492	0.0688	0.049*
C28	0.5797 (5)	0.7123 (9)	0.0654 (8)	0.039 (3)
H28	0.6146	0.7043	0.0444	0.047*
C29	0.3416 (6)	0.0041 (8)	0.3583 (7)	0.021 (2)
C30	0.3697 (6)	-0.0094 (10)	0.4323 (7)	0.029 (3)
H30	0.4010	-0.0544	0.4455	0.035*
C31	0.3525 (7)	0.0409 (11)	0.4848 (8)	0.029 (3)
H31	0.3710	0.0319	0.5352	0.035*
C32	0.3089 (7)	0.1038 (12)	0.4643 (8)	0.032 (3)
H32	0.2968	0.1414	0.5002	0.038*
C33	0.2812 (5)	0.1147 (11)	0.3905 (7)	0.025 (3)
H33	0.2495	0.1588	0.3764	0.030*
C34	0.3566 (7)	-0.0476 (10)	0.2978 (8)	0.025 (3)
C35	0.3202 (7)	-0.0283 (11)	0.2284 (9)	0.024 (3)
C36	0.3307 (8)	-0.0773 (8)	0.1672 (10)	0.025 (4)
H36	0.3059	-0.0649	0.1200	0.030*
C37	0.3771 (6)	-0.1442 (9)	0.1744 (8)	0.025 (3)
H37	0.3836	-0.1781	0.1332	0.030*
C38	0.4124 (6)	-0.1580 (9)	0.2432 (7)	0.026 (3)
H38	0.4454	-0.2006	0.2492	0.032*
C39	0.4022 (6)	-0.1135 (10)	0.3030 (8)	0.027 (3)
H39	0.4269	-0.1277	0.3499	0.032*
C40	0.1739 (6)	-0.0114 (9)	0.1020 (8)	0.025 (3)
C41	0.1474 (6)	-0.0268 (10)	0.0296 (7)	0.031 (3)
H41	0.1184	-0.0750	0.0171	0.037*
C42	0.1623 (7)	0.0272 (11)	-0.0272 (9)	0.036 (3)
H42	0.1429	0.0162	-0.0771	0.044*
C43	0.2074 (7)	0.0991 (11)	-0.0083 (8)	0.026 (3)
H43	0.2188	0.1363	-0.0450	0.031*
C44	0.2327 (6)	0.1113 (11)	0.0641 (7)	0.029 (3)
H44	0.2619	0.1591	0.0770	0.034*
C45	0.1616 (7)	-0.0586 (8)	0.1663 (8)	0.021 (3)
C46	0.1983 (7)	-0.0308 (10)	0.2348 (8)	0.020 (3)
C47	0.1897 (7)	-0.0784 (9)	0.2961 (9)	0.026 (4)
H47	0.2142	-0.0643	0.3432	0.031*
C48	0.1464 (7)	-0.1454 (10)	0.2891 (8)	0.033 (4)
H48	0.1405	-0.1745	0.3326	0.040*
C49	0.1101 (6)	-0.1740 (9)	0.2227 (7)	0.032 (3)

H49	0.0807	-0.2216	0.2205	0.038*
C50	0.1188 (6)	-0.1296 (10)	0.1591 (7)	0.033 (3)
H50	0.0960	-0.1474	0.1120	0.039*
C51	0.2934 (8)	0.7144 (9)	0.3228 (9)	0.083 (5)
H51A	0.2700	0.7210	0.3614	0.125*
H51B	0.2848	0.7665	0.2890	0.125*
H51C	0.3375	0.7127	0.3452	0.125*
C52	0.2755 (5)	0.6290 (7)	0.2821 (7)	0.050 (3)
C53	0.2087 (8)	0.6642 (11)	0.0818 (8)	0.084 (5)
H53A	0.2270	0.7128	0.1157	0.126*
H53B	0.2356	0.6104	0.0892	0.126*
H53C	0.1684	0.6477	0.0911	0.126*
C54	0.2017 (6)	0.6948 (9)	0.0098 (7)	0.059 (3)
C55	0.3156 (8)	0.4400 (11)	0.4141 (8)	0.079 (4)
H55A	0.3164	0.4092	0.3676	0.118*
H55B	0.3409	0.4955	0.4184	0.118*
H55C	0.2732	0.4565	0.4152	0.118*
C56	0.3386 (7)	0.3822 (11)	0.4716 (8)	0.057 (4)
F1	0.5218 (5)	0.5503 (8)	0.3177 (7)	0.107 (4)
F2	0.4542 (7)	0.6128 (10)	0.4395 (6)	0.140 (5)
F3	0.4492 (7)	0.6520 (8)	0.3254 (6)	0.158 (6)
F4	0.5169 (5)	0.5071 (8)	0.4329 (5)	0.095 (4)
F5	0.5426 (7)	0.6515 (8)	0.4094 (6)	0.129 (5)
F6	0.4350 (4)	0.5105 (8)	0.3415 (5)	0.101 (3)
F7	0.2000 (3)	0.6676 (4)	0.5720 (3)	0.0510 (16)
F8	0.2549 (3)	0.5503 (5)	0.5384 (4)	0.062 (2)
F9	0.2081 (4)	0.5770 (5)	0.4199 (4)	0.068 (2)
F10	0.1541 (4)	0.6900 (7)	0.4542 (5)	0.063 (2)
F11	0.2579 (3)	0.6905 (5)	0.4913 (4)	0.066 (2)
F12	0.1508 (3)	0.5525 (5)	0.5020 (5)	0.068 (2)
F13	-0.0032 (5)	0.6054 (9)	0.1681 (5)	0.109 (4)
F14	0.0776 (3)	0.5921 (8)	0.1137 (5)	0.079 (3)
F15	0.0179 (4)	0.6271 (6)	0.0040 (5)	0.072 (2)
F16	-0.0633 (3)	0.6303 (5)	0.0573 (4)	0.062 (2)
F17	0.0169 (3)	0.7190 (5)	0.0984 (4)	0.065 (2)
F18	-0.0041 (4)	0.5076 (5)	0.0676 (7)	0.102 (4)
Ir1	0.25717 (3)	0.07100 (2)	0.22916 (3)	0.01852 (8)
N1	0.1991 (5)	0.1828 (8)	0.2413 (6)	0.018 (2)
N2	0.0540 (7)	0.5722 (7)	0.3447 (8)	0.028 (3)
N3	-0.0202 (5)	0.6343 (8)	0.4028 (6)	0.041 (3)
N4	0.0357 (6)	0.7290 (9)	0.3376 (8)	0.041 (3)
N5	0.3110 (5)	0.1880 (8)	0.2146 (7)	0.024 (3)
N6	0.4664 (6)	0.5746 (6)	0.1255 (8)	0.025 (3)
N7	0.4839 (6)	0.7316 (8)	0.1344 (7)	0.036 (3)
N8	0.5457 (5)	0.6371 (7)	0.0792 (6)	0.032 (2)
N9	0.2980 (6)	0.0656 (6)	0.3415 (7)	0.020 (3)
N10	0.2188 (6)	0.0582 (7)	0.1215 (7)	0.023 (3)
N11	0.2633 (5)	0.5614 (6)	0.2520 (5)	0.047 (3)

N12	0.1991 (8)	0.7186 (11)	-0.0486 (7)	0.109 (6)
N13	0.3612 (8)	0.3294 (13)	0.5157 (10)	0.093 (6)
P1	0.4866 (2)	0.5830 (3)	0.3762 (3)	0.0604 (11)
P2	0.20527 (14)	0.6201 (2)	0.49735 (18)	0.0384 (7)
P3	0.00657 (16)	0.6120 (3)	0.0852 (2)	0.0445 (8)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.019 (4)	0.016 (5)	0.009 (5)	-0.002 (4)	0.000 (3)	0.002 (4)
C2	0.027 (5)	0.019 (5)	0.021 (6)	-0.001 (4)	0.004 (4)	-0.001 (4)
C3	0.025 (5)	0.023 (5)	0.018 (6)	-0.001 (4)	0.006 (4)	-0.001 (4)
C4	0.017 (5)	0.024 (5)	0.020 (6)	-0.003 (4)	0.003 (4)	-0.001 (4)
C5	0.023 (5)	0.015 (5)	0.032 (8)	-0.005 (4)	0.012 (5)	-0.005 (5)
C6	0.026 (7)	0.022 (5)	0.026 (7)	-0.004 (4)	0.006 (6)	-0.001 (4)
C7	0.031 (6)	0.033 (5)	0.027 (6)	0.001 (4)	0.013 (5)	0.003 (4)
C8	0.025 (6)	0.031 (6)	0.027 (5)	0.002 (4)	0.007 (4)	0.003 (4)
C9	0.025 (5)	0.031 (5)	0.032 (5)	-0.002 (4)	0.010 (4)	-0.002 (4)
C10	0.024 (5)	0.019 (5)	0.024 (5)	0.003 (4)	0.005 (4)	0.002 (4)
C11	0.018 (5)	0.029 (5)	0.025 (6)	-0.001 (3)	0.003 (4)	-0.004 (4)
C12	0.035 (7)	0.031 (5)	0.082 (10)	0.005 (4)	0.038 (7)	0.003 (5)
C13	0.043 (6)	0.028 (5)	0.050 (7)	0.000 (4)	0.020 (6)	-0.006 (5)
C14	0.042 (7)	0.021 (5)	0.080 (10)	0.009 (4)	0.031 (7)	0.015 (5)
C15	0.029 (5)	0.028 (5)	0.034 (8)	0.000 (4)	0.014 (5)	-0.001 (5)
C16	0.016 (5)	0.027 (6)	0.030 (7)	0.001 (4)	0.013 (4)	-0.001 (5)
C17	0.019 (5)	0.026 (5)	0.032 (8)	0.000 (4)	0.010 (5)	-0.001 (4)
C18	0.019 (6)	0.028 (6)	0.043 (9)	0.001 (4)	0.015 (5)	-0.001 (5)
C19	0.022 (5)	0.035 (6)	0.024 (7)	-0.003 (4)	0.005 (5)	-0.001 (5)
C20	0.022 (7)	0.026 (6)	0.034 (8)	-0.005 (4)	0.008 (6)	0.000 (4)
C21	0.044 (8)	0.029 (6)	0.085 (11)	-0.007 (5)	0.043 (8)	-0.011 (6)
C22	0.060 (10)	0.029 (7)	0.093 (13)	-0.012 (6)	0.059 (10)	-0.017 (6)
C23	0.028 (6)	0.021 (5)	0.054 (8)	0.005 (4)	0.022 (6)	0.002 (4)
C24	0.041 (8)	0.024 (6)	0.072 (10)	0.004 (5)	0.032 (7)	0.006 (5)
C25	0.018 (5)	0.027 (4)	0.024 (6)	-0.001 (3)	0.001 (4)	0.002 (4)
C26	0.047 (6)	0.020 (5)	0.052 (7)	-0.015 (4)	0.017 (5)	-0.003 (5)
C27	0.045 (7)	0.025 (5)	0.056 (8)	-0.011 (4)	0.022 (6)	0.002 (5)
C28	0.021 (5)	0.042 (5)	0.057 (8)	-0.007 (4)	0.012 (5)	0.008 (5)
C29	0.024 (5)	0.015 (5)	0.021 (4)	-0.004 (4)	0.002 (3)	0.000 (3)
C30	0.032 (6)	0.033 (6)	0.020 (5)	0.002 (4)	-0.001 (4)	-0.002 (4)
C31	0.034 (7)	0.036 (6)	0.016 (5)	0.000 (5)	0.002 (4)	0.003 (4)
C32	0.036 (7)	0.040 (7)	0.017 (6)	0.002 (6)	0.001 (5)	-0.003 (5)
C33	0.020 (6)	0.033 (7)	0.020 (5)	-0.004 (5)	0.005 (4)	0.003 (4)
C34	0.024 (6)	0.027 (5)	0.021 (5)	-0.001 (5)	0.000 (4)	-0.001 (4)
C35	0.021 (6)	0.023 (7)	0.027 (5)	-0.002 (5)	0.003 (4)	-0.003 (5)
C36	0.030 (7)	0.020 (7)	0.025 (6)	0.005 (5)	0.003 (5)	0.003 (4)
C37	0.026 (5)	0.021 (6)	0.032 (6)	0.003 (4)	0.014 (4)	0.003 (5)
C38	0.025 (5)	0.020 (6)	0.032 (5)	0.006 (4)	0.003 (4)	-0.001 (4)
C39	0.024 (5)	0.029 (5)	0.027 (5)	0.001 (4)	0.007 (4)	-0.001 (4)

C40	0.025 (5)	0.019 (5)	0.033 (5)	-0.003 (4)	0.010 (4)	0.001 (4)
C41	0.031 (6)	0.032 (6)	0.029 (5)	-0.009 (5)	0.007 (4)	0.005 (4)
C42	0.042 (7)	0.044 (7)	0.024 (6)	-0.008 (6)	0.008 (5)	-0.001 (5)
C43	0.027 (6)	0.028 (6)	0.025 (6)	0.006 (5)	0.010 (5)	0.003 (5)
C44	0.039 (8)	0.022 (6)	0.027 (6)	-0.004 (6)	0.010 (5)	0.006 (5)
C45	0.026 (6)	0.020 (6)	0.021 (6)	-0.001 (4)	0.009 (5)	0.005 (4)
C46	0.022 (7)	0.020 (7)	0.021 (6)	0.003 (5)	0.011 (5)	-0.002 (5)
C47	0.020 (6)	0.036 (9)	0.023 (7)	-0.004 (5)	0.005 (5)	0.007 (5)
C48	0.036 (7)	0.042 (9)	0.021 (6)	-0.003 (6)	0.006 (5)	0.006 (6)
C49	0.040 (7)	0.017 (6)	0.041 (7)	-0.009 (5)	0.011 (5)	0.004 (5)
C50	0.042 (7)	0.030 (7)	0.024 (6)	-0.018 (5)	0.003 (5)	-0.008 (5)
C51	0.105 (12)	0.045 (8)	0.122 (13)	-0.019 (8)	0.076 (11)	-0.025 (8)
C52	0.058 (7)	0.033 (6)	0.073 (8)	0.000 (5)	0.044 (6)	-0.001 (6)
C53	0.110 (12)	0.086 (11)	0.066 (7)	0.013 (9)	0.041 (7)	0.010 (7)
C54	0.064 (8)	0.058 (8)	0.061 (6)	-0.007 (6)	0.023 (6)	0.010 (6)
C55	0.079 (11)	0.088 (10)	0.076 (9)	0.018 (8)	0.032 (8)	0.003 (7)
C56	0.071 (10)	0.060 (8)	0.041 (7)	0.001 (8)	0.016 (7)	-0.019 (5)
F1	0.093 (7)	0.144 (10)	0.097 (9)	-0.042 (7)	0.048 (7)	-0.016 (7)
F2	0.207 (14)	0.163 (11)	0.066 (7)	0.032 (12)	0.062 (8)	0.009 (7)
F3	0.286 (17)	0.090 (8)	0.071 (7)	0.108 (10)	-0.018 (8)	0.007 (6)
F4	0.101 (7)	0.099 (8)	0.069 (6)	-0.028 (6)	-0.018 (5)	0.019 (5)
F5	0.200 (13)	0.091 (8)	0.085 (7)	-0.057 (9)	0.009 (9)	-0.017 (6)
F6	0.073 (6)	0.112 (8)	0.098 (7)	-0.013 (6)	-0.024 (5)	0.024 (6)
F7	0.038 (3)	0.067 (4)	0.046 (3)	0.011 (3)	0.005 (3)	-0.020 (3)
F8	0.058 (4)	0.057 (4)	0.060 (4)	0.024 (3)	-0.010 (3)	-0.011 (3)
F9	0.082 (5)	0.077 (5)	0.046 (4)	0.021 (4)	0.014 (4)	-0.023 (3)
F10	0.056 (5)	0.064 (5)	0.065 (5)	0.025 (4)	0.003 (4)	0.000 (4)
F11	0.043 (4)	0.068 (5)	0.096 (6)	-0.005 (3)	0.031 (4)	-0.009 (4)
F12	0.055 (5)	0.070 (5)	0.076 (5)	-0.022 (4)	0.009 (4)	-0.011 (4)
F13	0.083 (6)	0.194 (11)	0.063 (6)	0.087 (8)	0.045 (5)	0.054 (7)
F14	0.030 (4)	0.139 (9)	0.067 (5)	0.016 (4)	0.010 (4)	-0.012 (5)
F15	0.090 (6)	0.071 (5)	0.062 (5)	0.023 (5)	0.033 (4)	-0.004 (4)
F16	0.054 (4)	0.065 (5)	0.068 (5)	-0.011 (4)	0.014 (4)	-0.039 (4)
F17	0.053 (4)	0.058 (4)	0.094 (5)	-0.021 (3)	0.036 (4)	-0.041 (4)
F18	0.080 (6)	0.024 (4)	0.202 (13)	0.002 (4)	0.031 (8)	0.002 (6)
Ir1	0.01925 (12)	0.01816 (12)	0.01878 (12)	0.0000 (3)	0.00550 (8)	0.0008 (3)
N1	0.019 (4)	0.017 (5)	0.015 (5)	-0.005 (3)	-0.002 (4)	0.003 (4)
N2	0.027 (5)	0.030 (5)	0.030 (5)	0.008 (4)	0.013 (4)	0.004 (3)
N3	0.049 (6)	0.029 (5)	0.060 (7)	0.001 (4)	0.041 (5)	0.001 (4)
N4	0.030 (5)	0.034 (5)	0.062 (7)	0.002 (4)	0.021 (5)	0.004 (4)
N5	0.029 (5)	0.020 (5)	0.027 (6)	-0.005 (4)	0.013 (5)	0.003 (4)
N6	0.021 (5)	0.021 (4)	0.034 (6)	0.003 (3)	0.010 (4)	-0.002 (3)
N7	0.043 (6)	0.021 (4)	0.053 (6)	-0.006 (4)	0.028 (5)	-0.005 (4)
N8	0.029 (4)	0.030 (5)	0.042 (6)	-0.001 (3)	0.019 (4)	0.005 (4)
N9	0.018 (5)	0.020 (5)	0.020 (5)	-0.007 (3)	0.002 (4)	0.002 (4)
N10	0.027 (6)	0.019 (5)	0.024 (5)	-0.001 (4)	0.007 (4)	0.005 (4)
N11	0.054 (6)	0.036 (5)	0.056 (7)	0.004 (4)	0.026 (6)	-0.006 (4)
N12	0.136 (13)	0.121 (12)	0.063 (7)	-0.068 (11)	0.008 (7)	0.025 (7)

N13	0.078 (11)	0.098 (11)	0.095 (11)	0.003 (8)	0.000 (9)	0.015 (9)
P1	0.059 (3)	0.055 (3)	0.065 (2)	0.0109 (19)	0.010 (2)	0.0109 (19)
P2	0.0353 (17)	0.0401 (17)	0.0383 (16)	0.0025 (14)	0.0047 (13)	-0.0089 (13)
P3	0.0349 (19)	0.051 (2)	0.0489 (19)	-0.0018 (15)	0.0122 (15)	-0.0149 (17)

Geometric parameters (\AA , $\text{^{\circ}}$)

C1—N1	1.375 (16)	C33—H33	0.9500
C1—C2	1.421 (18)	C34—C35	1.40 (2)
C1—C15	1.489 (9)	C34—C39	1.39 (2)
C2—C3	1.382 (17)	C35—C36	1.41 (2)
C2—H2	0.9500	C35—Ir1	2.021 (16)
C3—C4	1.392 (18)	C36—C37	1.408 (19)
C3—C6	1.452 (19)	C36—H36	0.9500
C4—C5	1.394 (17)	C37—C38	1.367 (19)
C4—H4	0.9500	C37—H37	0.9500
C5—N1	1.317 (15)	C38—C39	1.352 (18)
C5—H5	0.9500	C38—H38	0.9500
C6—C10	1.39 (2)	C39—H39	0.9500
C6—C7	1.397 (18)	C40—C41	1.370 (18)
C7—C8	1.383 (19)	C40—N10	1.418 (16)
C7—H7	0.9500	C40—C45	1.459 (17)
C8—N2	1.37 (2)	C41—C42	1.414 (19)
C8—H8	0.9500	C41—H41	0.9500
C9—N2	1.341 (16)	C42—C43	1.44 (2)
C9—C10	1.400 (17)	C42—H42	0.9500
C9—H9	0.9500	C43—C44	1.357 (19)
C10—H10	0.9500	C43—H43	0.9500
C11—N4	1.273 (17)	C44—N10	1.409 (18)
C11—N3	1.334 (14)	C44—H44	0.9500
C11—N2	1.466 (16)	C45—C50	1.396 (18)
C12—N3	1.335 (15)	C45—C46	1.42 (2)
C12—C13	1.362 (18)	C46—C47	1.39 (2)
C12—H12	0.9500	C46—Ir1	2.000 (15)
C13—C14	1.372 (18)	C47—C48	1.361 (19)
C13—H13	0.9500	C47—H47	0.9500
C14—N4	1.294 (18)	C48—C49	1.388 (19)
C14—H14	0.9500	C48—H48	0.9500
C15—N5	1.336 (18)	C49—C50	1.401 (17)
C15—C16	1.343 (19)	C49—H49	0.9500
C16—C17	1.400 (16)	C50—H50	0.9500
C16—H16	0.9500	C51—C52	1.470 (17)
C17—C18	1.392 (19)	C51—H51A	0.9800
C17—C20	1.509 (19)	C51—H51B	0.9800
C18—C19	1.347 (19)	C51—H51C	0.9800
C18—H18	0.9500	C52—N11	1.140 (13)
C19—N5	1.379 (16)	C53—C54	1.390 (18)
C19—H19	0.9500	C53—H53A	0.9800

C20—C21	1.362 (19)	C53—H53B	0.9800
C20—C24	1.42 (2)	C53—H53C	0.9800
C21—C22	1.32 (2)	C54—N12	1.132 (16)
C21—H21	0.9500	C55—C56	1.37 (2)
C22—N6	1.33 (2)	C55—H55A	0.9800
C22—H22	0.9500	C55—H55B	0.9800
C23—C24	1.330 (19)	C55—H55C	0.9800
C23—N6	1.349 (15)	C56—N13	1.16 (2)
C23—H23	0.9500	F1—P1	1.549 (12)
C24—H24	0.9500	F2—P1	1.571 (12)
C25—N8	1.256 (14)	F3—P1	1.505 (10)
C25—N7	1.331 (16)	F4—P1	1.579 (11)
C25—N6	1.453 (16)	F5—P1	1.614 (12)
C26—C27	1.336 (17)	F6—P1	1.594 (10)
C26—N7	1.402 (16)	F7—P2	1.581 (6)
C26—H26	0.9500	F8—P2	1.577 (7)
C27—C28	1.352 (19)	F9—P2	1.589 (7)
C27—H27	0.9500	F10—P2	1.611 (9)
C28—N8	1.388 (15)	F11—P2	1.582 (8)
C28—H28	0.9500	F12—P2	1.582 (8)
C29—N9	1.311 (16)	F13—P3	1.608 (9)
C29—C30	1.400 (16)	F14—P3	1.582 (8)
C29—C34	1.453 (19)	F15—P3	1.602 (9)
C30—C31	1.343 (18)	F16—P3	1.553 (8)
C30—H30	0.9500	F17—P3	1.592 (8)
C31—C32	1.33 (2)	F18—P3	1.568 (9)
C31—H31	0.9500	Ir1—N10	2.011 (14)
C32—C33	1.388 (18)	Ir1—N9	2.095 (13)
C32—H32	0.9500	Ir1—N1	2.125 (11)
C33—N9	1.280 (18)	Ir1—N5	2.139 (11)
N1—C1—C2	118.6 (12)	C44—C43—H43	121.5
N1—C1—C15	117.9 (14)	C42—C43—H43	121.5
C2—C1—C15	123.5 (15)	C43—C44—N10	124.9 (14)
C3—C2—C1	120.5 (12)	C43—C44—H44	117.6
C3—C2—H2	119.7	N10—C44—H44	117.6
C1—C2—H2	119.7	C50—C45—C46	123.3 (12)
C4—C3—C2	119.0 (13)	C50—C45—C40	121.2 (12)
C4—C3—C6	122.4 (12)	C46—C45—C40	115.4 (12)
C2—C3—C6	118.5 (13)	C47—C46—C45	116.3 (14)
C3—C4—C5	118.1 (12)	C47—C46—Ir1	128.7 (12)
C3—C4—H4	120.9	C45—C46—Ir1	115.0 (10)
C5—C4—H4	120.9	C46—C47—C48	120.2 (15)
N1—C5—C4	123.5 (12)	C46—C47—H47	119.9
N1—C5—H5	118.3	C48—C47—H47	119.9
C4—C5—H5	118.3	C47—C48—C49	124.6 (15)
C10—C6—C7	117.0 (14)	C47—C48—H48	117.7
C10—C6—C3	121.2 (12)	C49—C48—H48	117.7

C7—C6—C3	121.8 (16)	C48—C49—C50	117.0 (12)
C8—C7—C6	122.3 (14)	C48—C49—H49	121.5
C8—C7—H7	118.8	C50—C49—H49	121.5
C6—C7—H7	118.8	C49—C50—C45	118.6 (12)
N2—C8—C7	118.4 (13)	C49—C50—H50	120.7
N2—C8—H8	120.8	C45—C50—H50	120.7
C7—C8—H8	120.8	C52—C51—H51A	109.5
N2—C9—C10	120.8 (13)	C52—C51—H51B	109.5
N2—C9—H9	119.6	H51A—C51—H51B	109.5
C10—C9—H9	119.6	C52—C51—H51C	109.5
C6—C10—C9	120.1 (13)	H51A—C51—H51C	109.5
C6—C10—H10	119.9	H51B—C51—H51C	109.5
C9—C10—H10	119.9	N11—C52—C51	177.7 (15)
N4—C11—N3	130.1 (13)	C54—C53—H53A	109.5
N4—C11—N2	116.1 (12)	C54—C53—H53B	109.5
N3—C11—N2	113.6 (12)	H53A—C53—H53B	109.5
N3—C12—C13	125.7 (13)	C54—C53—H53C	109.5
N3—C12—H12	117.2	H53A—C53—H53C	109.5
C13—C12—H12	117.2	H53B—C53—H53C	109.5
C12—C13—C14	113.3 (14)	N12—C54—C53	176.4 (18)
C12—C13—H13	123.4	C56—C55—H55A	109.5
C14—C13—H13	123.4	C56—C55—H55B	109.5
N4—C14—C13	124.4 (14)	H55A—C55—H55B	109.5
N4—C14—H14	117.8	C56—C55—H55C	109.5
C13—C14—H14	117.8	H55A—C55—H55C	109.5
N5—C15—C16	123.5 (14)	H55B—C55—H55C	109.5
N5—C15—C1	112.1 (15)	N13—C56—C55	173.4 (18)
C16—C15—C1	124.4 (16)	C46—Ir1—N10	80.8 (5)
C15—C16—C17	119.4 (13)	C46—Ir1—C35	86.0 (3)
C15—C16—H16	120.3	N10—Ir1—C35	94.2 (6)
C17—C16—H16	120.3	C46—Ir1—N9	93.9 (5)
C16—C17—C18	117.7 (13)	N10—Ir1—N9	172.5 (2)
C16—C17—C20	123.3 (13)	C35—Ir1—N9	80.1 (6)
C18—C17—C20	119.0 (12)	C46—Ir1—N1	98.4 (6)
C19—C18—C17	119.9 (13)	N10—Ir1—N1	92.3 (5)
C19—C18—H18	120.0	C35—Ir1—N1	172.7 (6)
C17—C18—H18	120.0	N9—Ir1—N1	93.8 (4)
C18—C19—N5	121.9 (14)	C46—Ir1—N5	173.1 (6)
C18—C19—H19	119.1	N10—Ir1—N5	94.4 (5)
N5—C19—H19	119.1	C35—Ir1—N5	99.4 (6)
C21—C20—C24	116.8 (14)	N9—Ir1—N5	91.3 (4)
C21—C20—C17	121.7 (16)	N1—Ir1—N5	76.6 (2)
C24—C20—C17	121.5 (13)	C5—N1—C1	120.2 (12)
C22—C21—C20	119.9 (16)	C5—N1—Ir1	124.8 (9)
C22—C21—H21	120.0	C1—N1—Ir1	114.1 (8)
C20—C21—H21	120.0	C9—N2—C8	121.2 (12)
C21—C22—N6	123.4 (15)	C9—N2—C11	120.4 (13)
C21—C22—H22	118.3	C8—N2—C11	118.5 (11)

N6—C22—H22	118.3	C12—N3—C11	111.1 (11)
C24—C23—N6	120.5 (14)	C11—N4—C14	115.3 (13)
C24—C23—H23	119.7	C15—N5—C19	117.5 (12)
N6—C23—H23	119.7	C15—N5—Ir1	118.5 (10)
C23—C24—C20	120.4 (14)	C19—N5—Ir1	123.8 (10)
C23—C24—H24	119.8	C22—N6—C23	118.7 (12)
C20—C24—H24	119.8	C22—N6—C25	122.1 (11)
N8—C25—N7	129.5 (12)	C23—N6—C25	119.2 (13)
N8—C25—N6	117.2 (12)	C25—N7—C26	113.6 (12)
N7—C25—N6	113.3 (11)	C25—N8—C28	116.6 (12)
C27—C26—N7	120.5 (14)	C33—N9—C29	122.0 (13)
C27—C26—H26	119.7	C33—N9—Ir1	124.1 (10)
N7—C26—H26	119.7	C29—N9—Ir1	113.8 (10)
C26—C27—C28	120.3 (14)	C44—N10—C40	117.2 (12)
C26—C27—H27	119.8	C44—N10—Ir1	126.2 (10)
C28—C27—H27	119.8	C40—N10—Ir1	116.6 (9)
C27—C28—N8	119.3 (12)	F3—P1—F1	93.4 (8)
C27—C28—H28	120.4	F3—P1—F2	90.3 (8)
N8—C28—H28	120.4	F1—P1—F2	176.2 (8)
N9—C29—C30	118.8 (13)	F3—P1—F4	171.8 (9)
N9—C29—C34	116.9 (12)	F1—P1—F4	92.6 (7)
C30—C29—C34	124.2 (12)	F2—P1—F4	83.8 (7)
C31—C30—C29	120.2 (13)	F3—P1—F6	86.4 (7)
C31—C30—H30	119.9	F1—P1—F6	86.5 (6)
C29—C30—H30	119.9	F2—P1—F6	94.7 (8)
C32—C31—C30	118.3 (13)	F4—P1—F6	88.4 (5)
C32—C31—H31	120.8	F3—P1—F5	95.9 (8)
C30—C31—H31	120.8	F1—P1—F5	89.7 (7)
C31—C32—C33	120.3 (15)	F2—P1—F5	88.9 (8)
C31—C32—H32	119.9	F4—P1—F5	89.7 (6)
C33—C32—H32	119.9	F6—P1—F5	175.7 (7)
N9—C33—C32	120.4 (14)	F8—P2—F12	91.9 (4)
N9—C33—H33	119.8	F8—P2—F7	92.0 (4)
C32—C33—H33	119.8	F12—P2—F7	91.3 (4)
C35—C34—C39	118.3 (14)	F8—P2—F11	90.2 (4)
C35—C34—C29	115.3 (13)	F12—P2—F11	177.9 (5)
C39—C34—C29	126.4 (13)	F7—P2—F11	88.5 (4)
C34—C35—C36	118.8 (15)	F8—P2—F9	91.3 (4)
C34—C35—Ir1	113.8 (12)	F12—P2—F9	89.0 (5)
C36—C35—Ir1	127.4 (12)	F7—P2—F9	176.7 (4)
C37—C36—C35	121.5 (16)	F11—P2—F9	91.0 (4)
C37—C36—H36	119.2	F8—P2—F10	178.9 (5)
C35—C36—H36	119.2	F12—P2—F10	87.5 (5)
C38—C37—C36	117.0 (14)	F7—P2—F10	89.0 (4)
C38—C37—H37	121.5	F11—P2—F10	90.4 (5)
C36—C37—H37	121.5	F9—P2—F10	87.7 (5)
C39—C38—C37	122.5 (13)	F16—P3—F18	90.1 (5)
C39—C38—H38	118.7	F16—P3—F14	179.3 (6)

C37—C38—H38	118.7	F18—P3—F14	89.2 (6)
C38—C39—C34	121.8 (13)	F16—P3—F17	89.0 (4)
C38—C39—H39	119.1	F18—P3—F17	176.9 (6)
C34—C39—H39	119.1	F14—P3—F17	91.7 (5)
C41—C40—N10	120.1 (12)	F16—P3—F15	90.6 (5)
C41—C40—C45	127.9 (12)	F18—P3—F15	89.1 (6)
N10—C40—C45	112.0 (12)	F14—P3—F15	89.6 (5)
C40—C41—C42	121.6 (13)	F17—P3—F15	87.9 (5)
C40—C41—H41	119.2	F16—P3—F13	89.9 (5)
C42—C41—H41	119.2	F18—P3—F13	95.6 (7)
C41—C42—C43	119.1 (13)	F14—P3—F13	90.0 (5)
C41—C42—H42	120.4	F17—P3—F13	87.4 (6)
C43—C42—H42	120.4	F15—P3—F13	175.3 (7)
C44—C43—C42	117.1 (14)		
N1—C1—C2—C3	2.6 (18)	C41—C40—C45—C50	1 (2)
C15—C1—C2—C3	-176.6 (10)	N10—C40—C45—C50	-180.0 (13)
C1—C2—C3—C4	-2.1 (19)	C41—C40—C45—C46	177.1 (14)
C1—C2—C3—C6	175.2 (13)	N10—C40—C45—C46	-3.4 (18)
C2—C3—C4—C5	-0.2 (19)	C50—C45—C46—C47	0 (2)
C6—C3—C4—C5	-177.3 (14)	C40—C45—C46—C47	-176.5 (13)
C3—C4—C5—N1	2 (2)	C50—C45—C46—Ir1	178.8 (12)
C4—C3—C6—C10	28 (2)	C40—C45—C46—Ir1	2.3 (17)
C2—C3—C6—C10	-149.1 (15)	C45—C46—C47—C48	-2 (2)
C4—C3—C6—C7	-149.6 (14)	Ir1—C46—C47—C48	179.1 (12)
C2—C3—C6—C7	33 (2)	C46—C47—C48—C49	3 (2)
C10—C6—C7—C8	-2 (2)	C47—C48—C49—C50	-1 (2)
C3—C6—C7—C8	176.1 (14)	C48—C49—C50—C45	-2 (2)
C6—C7—C8—N2	1 (2)	C46—C45—C50—C49	2 (2)
C7—C6—C10—C9	1 (2)	C40—C45—C50—C49	178.3 (13)
C3—C6—C10—C9	-176.9 (14)	C4—C5—N1—C1	-2 (2)
N2—C9—C10—C6	1 (2)	C4—C5—N1—Ir1	167.1 (10)
N3—C12—C13—C14	-3 (2)	C2—C1—N1—C5	-0.7 (18)
C12—C13—C14—N4	3 (3)	C15—C1—N1—C5	178.5 (10)
N1—C1—C15—N5	-9.7 (9)	C2—C1—N1—Ir1	-170.6 (9)
C2—C1—C15—N5	169.5 (15)	C15—C1—N1—Ir1	8.7 (10)
N1—C1—C15—C16	167.5 (16)	C10—C9—N2—C8	-2 (2)
C2—C1—C15—C16	-13.3 (11)	C10—C9—N2—C11	177.7 (13)
N5—C15—C16—C17	-1 (2)	C7—C8—N2—C9	1 (2)
C1—C15—C16—C17	-177.7 (10)	C7—C8—N2—C11	-178.5 (12)
C15—C16—C17—C18	-3 (2)	N4—C11—N2—C9	171.0 (14)
C15—C16—C17—C20	180.0 (15)	N3—C11—N2—C9	-12.9 (19)
C16—C17—C18—C19	4 (2)	N4—C11—N2—C8	-10 (2)
C20—C17—C18—C19	-179.1 (14)	N3—C11—N2—C8	166.6 (13)
C17—C18—C19—N5	-1 (2)	C13—C12—N3—C11	3 (2)
C16—C17—C20—C21	22 (3)	N4—C11—N3—C12	-3 (2)
C18—C17—C20—C21	-154.8 (16)	N2—C11—N3—C12	-178.8 (12)
C16—C17—C20—C24	-157.4 (17)	N3—C11—N4—C14	3 (2)

C18—C17—C20—C24	26 (2)	N2—C11—N4—C14	178.6 (14)
C24—C20—C21—C22	-3 (3)	C13—C14—N4—C11	-3 (3)
C17—C20—C21—C22	177.5 (18)	C16—C15—N5—C19	4 (2)
C20—C21—C22—N6	-1 (3)	C1—C15—N5—C19	-179.1 (9)
N6—C23—C24—C20	1 (3)	C16—C15—N5—Ir1	-171.3 (12)
C21—C20—C24—C23	3 (3)	C1—C15—N5—Ir1	6.0 (13)
C17—C20—C24—C23	-177.2 (15)	C18—C19—N5—C15	-3 (2)
N7—C26—C27—C28	-4 (2)	C18—C19—N5—Ir1	171.8 (11)
C26—C27—C28—N8	6 (2)	C21—C22—N6—C23	6 (3)
N9—C29—C30—C31	-1 (2)	C21—C22—N6—C25	-176.5 (18)
C34—C29—C30—C31	-179.4 (14)	C24—C23—N6—C22	-5 (2)
C29—C30—C31—C32	-1 (2)	C24—C23—N6—C25	176.9 (14)
C30—C31—C32—C33	2 (3)	N8—C25—N6—C22	174.8 (17)
C31—C32—C33—N9	-1 (2)	N7—C25—N6—C22	-5 (2)
N9—C29—C34—C35	-3 (2)	N8—C25—N6—C23	-7.3 (19)
C30—C29—C34—C35	175.5 (13)	N7—C25—N6—C23	172.4 (13)
N9—C29—C34—C39	177.9 (14)	N8—C25—N7—C26	2 (2)
C30—C29—C34—C39	-4 (2)	N6—C25—N7—C26	-177.5 (12)
C39—C34—C35—C36	1 (2)	C27—C26—N7—C25	0 (2)
C29—C34—C35—C36	-178.2 (13)	N7—C25—N8—C28	0 (2)
C39—C34—C35—Ir1	-178.7 (12)	N6—C25—N8—C28	179.7 (12)
C29—C34—C35—Ir1	2.1 (18)	C27—C28—N8—C25	-4.2 (19)
C34—C35—C36—C37	-1 (2)	C32—C33—N9—C29	0 (2)
Ir1—C35—C36—C37	179.0 (11)	C32—C33—N9—Ir1	177.0 (11)
C35—C36—C37—C38	-1 (2)	C30—C29—N9—C33	1 (2)
C36—C37—C38—C39	3 (2)	C34—C29—N9—C33	179.8 (13)
C37—C38—C39—C34	-3 (2)	C30—C29—N9—Ir1	-176.2 (10)
C35—C34—C39—C38	1 (2)	C34—C29—N9—Ir1	2.4 (16)
C29—C34—C39—C38	179.8 (13)	C43—C44—N10—C40	-2 (2)
N10—C40—C41—C42	-2 (2)	C43—C44—N10—Ir1	177.9 (12)
C45—C40—C41—C42	177.4 (15)	C41—C40—N10—C44	2 (2)
C40—C41—C42—C43	1 (2)	C45—C40—N10—C44	-177.3 (12)
C41—C42—C43—C44	-1 (2)	C41—C40—N10—Ir1	-177.4 (10)
C42—C43—C44—N10	1 (2)	C45—C40—N10—Ir1	3.0 (15)