

miR-223-3p regulates cell growth and apoptosis via FBXW7 suggesting an oncogenic role in human testicular germ cell tumors

JIKAI LIU^{1,2}, HAO SHI¹, XIDAN LI³, GANG CHEN², CATHARINA LARSSON¹ and WENG-ONN LUI¹

¹Department of Oncology-Pathology, Karolinska Institutet, Cancer Center Karolinska, Karolinska University Hospital, SE-171 76 Stockholm, Sweden; ²Department of Urology, Jinshan Hospital, Fudan University, Shanghai 201508, P.R. China; ³Department of Medicine-Huddinge, Karolinska University Hospital-Huddinge, SE-141 86 Stockholm, Sweden

Received September 11, 2016; Accepted November 25, 2016

DOI: 10.3892/ijo.2016.3807

Abstract. *miR-223-3p* is deregulated in several tumor types and plays an important role in tumorigenesis and progression. However, its role in the pathogenesis of testicular germ cell tumor (TGCT) remains uncharacterized. We previously demonstrated that *miR-223-3p* expression was increased in TGCTs compared with normal testes (NT), suggesting that *miR-223-3p* may have an oncogenic role in TGCT. Using published dataset and The Cancer Genome Atlas database, we validated higher *miR-223-3p* expression in TGCTs than NT, and found a negative correlation between *miR-223-3p* and *FBXW7* mRNA expression levels. Using both gain- and loss-of-function experiments, we show that *miR-223-3p* regulates *FBXW7* protein expression, cell growth and apoptosis in TGCT cell lines. Additionally, we demonstrate that ectopic expression of the full-length coding sequence of *FBXW7* could rescue the cell growth and apoptotic effects mediated by *miR-223-3p*. Our findings suggest an oncogenic role for *miR-223-3p* in TGCT, which promotes cell growth and inhibits apoptosis through repression of *FBXW7*.

Introduction

Testicular germ cell tumor (TGCT) is the most frequent solid malignancy occurring in males between the ages of 15 and 34 years (1), with a steadily rising incidence for the past few decades in the United States and Europe (2). Histopathologically, ~55% of all TGCTs are classified as seminomas, and the remaining cases as non-seminomas (3). The vast majority of TGCTs have an excellent cure rate with cisplatin-based treatment. Nevertheless, a subset of patients develops cisplatin resistance resulting in tumor progression

and reduced survival (4). Therefore, a better understanding of the molecular mechanisms of TGCT tumorigenesis is needed for identification of new therapeutic targets and treatment development.

MicroRNAs (miRNAs) are small non-coding RNAs of ~20-24 nucleotides in length, which play important roles in a broad range of cellular processes, including tumor development and drug response (5). Genome-wide miRNA profiling studies have provided evidence of miRNA deregulations in TGCT. For example, the *miR-371-373* cluster is frequently overexpressed in malignant TGCTs of all histopathological subtypes (6,7). Other miRNAs, such as the *miR-302* cluster and *miR-301*, are differentially expressed based on the cellular differentiation of the tumor (7,8). To date, very few miRNAs have been functionally characterized in TGCT. *miR-372* and *miR-373* have been shown to play oncogenic roles in TGCT by targeting the tumor suppressor *LATS2* (9). However, the functional roles of other differentially expressed miRNAs in TGCT have yet to be characterized.

We previously identified a subset of miRNAs that were differentially expressed between TGCTs and normal testes (NT) using a deep sequencing approach (10). Among these, *miR-223-3p* expression was higher in TGCTs as compared to NT. *miR-223-3p* is known to be deregulated in a broad range of hematological malignancies and solid tumors (11,12). However, its role in TGCT remains uncharacterized. *miR-223-3p* has been shown to regulate multiple targets in different cancer types. Among them, F-box/WD repeat-containing protein (FBXW7) is the most common target, which has been reported in acute T-cell lymphoblastic leukemia, esophageal squamous cell carcinoma and gastric cancer (13-15). FBXW7 is the substrate-recognition component of the SCF-(SKP1, CUL1, F-box protein)-ubiquitin-ligase complex, which has been demonstrated to function as a tumor suppressor by promoting the degradation of several oncoprotein substrates, including c-Myc, cyclin E, MCL-1, c-JUN, NFκB2 and Notch1 (16,17). Therefore, suppression of FBXW7 by *miR-223-3p* can promote tumor development and progression.

In this study, we investigated the expression and function of *miR-223-3p* and FBXW7 in TGCT clinical samples and cell lines. Our data show that *miR-223-3p* plays an oncogenic role in TGCT by promoting cell proliferation and inhibiting apoptosis via FBXW7.

Correspondence to: Jikai Liu or Dr Weng-Onn Lui, Department of Oncology-Pathology, Karolinska Institutet, Karolinska University Hospital-Solna, SE-171 76 Stockholm, Sweden
E-mail: 14111270004@fudan.edu.cn
E-mail: weng-onn.lui@ki.se

Key words: *miR-223-3p*, cell growth, apoptosis, FBXW7, TGCT

Materials and methods

Clinical samples and cell lines. Fifteen frozen TGCTs and five NT were provided by the Cooperative Human Tissue Network, which is funded by the National Cancer Institute, USA. All samples were included in our previous small RNA-sequencing study (10). The study was approved by the Stanford Human Subjects Review Committee.

Two established TGCT cell lines were included in this study: the TCam-2 seminoma cell line and the 2102Ep non-seminoma cell line (18,19). TCam-2 was kindly provided by Dr Leendert H.J. Looijenga (Department of Pathology, Erasmus MC-University Medical Center Rotterdam, The Netherlands) and 2102Ep by Dr Peter Andrews (Department of Biomedical Science, University of Sheffield, UK). TCam-2 cells were grown in RPMI-1640 and 2102Ep cells were cultured in DMEM medium, supplemented with 10% fetal bovine serum. All cells were cultured at 37°C with 5% CO₂. Authentication of the cell lines was verified by short tandem repeat profiling in our recent study (10).

Data extraction and analysis from published data and The Cancer Genome Atlas database. For comparison of *miR-223-3p* expression between TGCTs and NT, we extracted global TaqMan miRNA profiling data from the study of Gillis *et al.* (7), which analyzed 61 germ cell tumors, three NT and five embryonal carcinoma cell lines. We excluded the 10 dysgerminomas (ovarian germ cell tumors), one ovarian embryonal carcinoma, one ovarian yolk sac carcinoma and five cell lines, and re-analyzed the *miR-223-3p* expression by normalization to *miR-16* in the 49 TGCTs and three NT.

For *FBXW7* mRNA, we extracted the microarray gene expression data of 101 TCGTs and five NT from Gene Expression Omnibus (GEO accession no. GSE3218; <http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE3218>).

For analysis of correlation between *miR-223-3p* expression and *FBXW7* mRNA levels, we extracted *miR-223-3p* and *FBXW7* mRNA data from The Cancer Genome Atlas (TCGA) testicular cancer database using the UCSC Xena browser (<http://xena.ucsc.edu/>). These *miR-223-3p* and *FBXW7* expression data had been generated by miRNA expression Illumina HiSeq and exon expression RNAseq, respectively.

RNA extraction. Total RNA was extracted using the mirVana miRNA isolation kit (AM1560; Ambion/Thermo Fisher Scientific, Waltham, MA, USA) and RNA concentration was measured with a NanoDrop ND-1000 spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA). All RNA samples were stored at -80°C until further use.

TaqMan reverse transcription quantitative polymerase chain reaction (RT-qPCR). RT-qPCR was performed to evaluate the transfection efficiency of *miR-223-3p* overexpression or inhibition using the StepOnePlus Real-Time PCR system (Applied Biosystems/Thermo Fisher Scientific). cDNA was synthesized from 20 ng of total RNA and used to quantify *miR-223-3p* (ID 002295) and *RNU48* (ID 001093). All reactions were performed in triplicate. The relative expression of *miR-223-3p* was normalized to *RNU48*, and the fold change of *miR-223-3p* in cells transfected with *miR-223-3p*

mimic/inhibitor relative to their respective control was reported as 2^{-ΔΔC_t}.

Transfection. For *miR-223* overexpression and inhibition, 2x10⁵ cells were transfected with 30 nM of miRNA inhibitor (anti-miR-223, AM12301 or anti-miR negative control no. 1, AM17010; Ambion) or 10 nM of miRNA mimic (pre-miR-223, PM12301 or pre-miR negative control no. 1, AM17110; Ambion) using siPORT NeoFX transfection agent (AM4511; Ambion).

For co-transfection of *miR-223* mimic and *FBXW7*-expressing plasmid, 1.5x10⁵ cells were co-transfected with 500 ng of pCMV-Myc *FBXW7* and 10 nM of pre-miR-223 or pre-miR-NC using Lipofectamine 2000 (no. 11668-019; Invitrogen/Thermo Fisher Scientific). Cells co-transfected with an empty vector and pre-miR-NC was used as a control. Cells were collected 48 h after transfection for subsequent analysis. The pCMV-Myc *FBXW7* plasmid was obtained from Addgene (no. 16652; Cambridge, MA, USA; <https://www.addgene.org/>). The empty vector was prepared by cleavage of pCMV-Myc *FBXW7* with *Bgl*III and *Not*I to remove the full-length coding sequence of *FBXW7*.

Annexin V cell apoptosis and EdU (5-ethynyl-2'-deoxyuridine) cell proliferation assays. Cell apoptosis and proliferation were evaluated in TCam-2 and 2102Ep cells 72 h after transfection using Annexin V FITC Apoptosis kit (PHN1018; Invitrogen) and Click-iT EdU Alexa Fluor 488 flow cytometry assay (C10425; Invitrogen), respectively. All experimental conditions were according to the manufacturer's instructions and analyzed by NovoCyte flow cytometer (ACEA Biosciences, San Diego, CA, USA). At least three independent experiments were performed in each cell line.

Trypan blue exclusion assay. Trypan blue exclusion assay was performed in TCam-2 and 2102Ep cells 48 or 72 h after transfection. Cells were stained with 0.4% trypan blue solution and counted by TC10™ Automated Cell Counter (Bio-Rad, Hercules, CA, USA).

WST-1 assay. Cell growth was measured by WST-1 colorimetric assay (no. 11644807001; Roche Diagnostics, Indianapolis, IN, USA) in TCam-2 and 2102Ep cells 72 h after transfection. Cells were plated into a 96-well plate at a concentration of 5x10³/well in 100 μl culture medium. At different time intervals (0, 24, 48 or 72 h after transfection), 10 μl of WST-1 reagent was added to each well and incubated for 3 h at 37°C. After incubation, absorbance values were detected at the wavelengths 450 nm (measurement) and 650 nm (reference) using the VERSAmax ELISA Microplate Reader (Molecular Devices, Sunnyvale, CA, USA). Each experimental group was performed in six replicates for each time-point and all experiments were repeated three times independently.

Western blotting. Total protein lysates were extracted using NP-40 cell lysis buffer (FNN0021; Invitrogen), supplemented with 1 mM of phenylmethanesulfonyl fluoride (P7626; Sigma-Aldrich, St. Louis, MO, USA) and protease inhibitor (P8340; Sigma-Aldrich). Protein concentrations were determined using the Pierce™ BCA Protein assay kit (no. 23227;




Figure 1. Expression analysis of *miR-223-3p* and *FBXW7* mRNA and protein in testicular germ cell tumors (TGCTs) and normal testes (NT). (A and B) The graphs depict *miR-223-3p* expression between NT and TGCTs in Özata *et al* (10) (A) and Gillis *et al* (7) (B). In both cohorts, *miR-223-3p* expression was quantified by RT-qPCR and *miR-16* was used for normalization. (C) The graph shows the normalized expression data of *FBXW7* mRNA in NT (n=5) and TGCTs (n=101), which were extracted from the microarray gene expression of GEO database accession no. GSE3218. (D) *miR-223-3p* expression and *FBXW7* mRNA data were obtained from the TCGA database. Correlation was assessed using the Pearson's correlation analysis. (E) Western blot analysis of *FBXW7* protein in NT (n=3) and TGCTs (n=14). GAPDH was used as a loading control. Data represent mean \pm SD. All comparisons were evaluated using Mann-Whitney U test. *P<0.05; ***P<0.001.

Pierce Biotechnology, Thermo Fisher Scientific). Thirty micrograms of protein lysates were separated in NuPAGE Novex 4-12% Bis-Tris gels (NP0321BOX; Invitrogen) and transferred to 0.2 μ m nitrocellulose membranes (no. 88024; Invitrogen). After blocking with 5% skim milk powder (no. 70166; Sigma-Aldrich) in Tris-buffered saline/0.05% Tween-20, membranes were incubated with anti-*FBXW7* (NBP1-59631; Novus Biologicals, Littleton, CO, USA; 1:1,000 dilution), anti-cleaved PARP (ab32064; Abcam, Cambridge, UK; 1:1,000 dilution) or anti-Myc-Tag (no. 2276; Cell Signaling Technologies, Danvers, MA, USA; 1:500 dilution) overnight at 4°C. Anti-rabbit IgG-HRP (no. 170-6515; Bio-Rad Laboratories; 1:3,000 dilution) or anti-mouse IgG-HRP (sc-2005; Santa Cruz Biotechnology, Dallas, TX, USA; 1:10,000 dilution) was used as secondary antibodies. For normalization purpose, the membrane was incubated with anti-GAPDH (sc-47724; Santa Cruz Biotechnology; 1:1,000 dilution). Signals were detected using the Novex ECL HRP chemiluminescent substrate reagent (WP20005; Invitrogen) and LAS-1000 image analyzer (Fujifilm, Tokyo, Japan).

Statistical analyses. All statistical analyses were performed using MS Office Excel 2007 or SPSS 22.0 (IBM Corp., Armonk, NY, USA). Comparisons between TGCT and NT were performed by Mann-Whitney U test, and the transfection experiments were assessed by Student's paired t-test. Correlation between *miR-223-3p* and *FBXW7* mRNA expression levels was evaluated using Pearson's correlation analysis. All statistical tests were two-sided and P-values <0.05 were considered as statistically significant.

Results

Expression of *miR-223-3p* and *FBXW7* in TGCTs and NT. To validate our previous observation of *miR-223-3p* overexpression in TGCTs, we re-analyzed *miR-223-3p* expression from the miRNA profiling data of Gillis *et al* (7), with inclusion of 49 TGCTs and 3 NT. In agreement with our previous finding (10) (Fig. 1A), *miR-223-3p* was overexpressed in TGCTs compared to NT (P=0.011; Fig. 1B).

To determine whether *FBXW7* could be a candidate target of *miR-223-3p* in TGCT, we analyzed *FBXW7* expression from the microarray gene expression data of 101 TGCTs and 5 NT in the Gene Expression Omnibus (GEO) database (accession no. GSE3218). Indeed, we found that *FBXW7* mRNA expression was decreased in TGCTs as compared to NT (P<0.001; Fig. 1C). We further assessed the correlation between *miR-223-3p* and *FBXW7* expression levels using miRNA and gene expression profiles from the TCGA testicular cancer datasets. The analysis revealed an inverse correlation ($r=-0.309$; P<0.001; Fig. 1D), supporting the miRNA-target relationship.

Additionally, we also quantified *FBXW7* protein expression in 3 NT and 14 TGCT samples by western blot analysis. As shown in Fig. 1E, the expression of *FBXW7* was low or undetectable in 10/14 TGCTs (71.4%), and moderate or high in the remaining four tumors. By contrast, all three NT showed moderate to high expression of *FBXW7*. Consistent with the mRNA expression pattern, the *FBXW7* protein level in TGCTs was lower than in NT (P=0.023).




Figure 2. Effect of *miR-223-3p* regulation on FBXW7 protein expression in TGCT cells. (A) TCam-2 and 2102Ep cells were transfected with anti-*miR-223* or pre-*miR-NC* and their respective negative controls. After 72 h of transfection, *miR-223-3p* expression was quantified by RT-qPCR and normalized to *RNU48*. (B) FBXW7 protein expression was measured in both TGCT cell lines transfected with anti-*miR-223* or pre-*miR-223* and controls using western blotting. GAPDH was used as a loading control. Data represent mean \pm SD from at least three independent experiments. P-values were calculated by paired t-test. * $P < 0.05$; ** $P < 0.01$.

Effect of miR-223-3p modulation on FBXW7 in TGCT cell lines. To further determine whether *miR-223-3p* could regulate FBXW7 in TGCT, we performed *miR-223-3p* overexpression and inhibition in two TGCT cell lines and evaluated the effect on FBXW7 protein expression using western blot analysis. As shown in Fig. 2A, cells transfected with anti-*miR-223* showed significantly lower *miR-223-3p* expression than the anti-*miR-NC*-treated cells in both cell lines ($P=0.018$ for both). Similarly, *miR-223-3p* expression was significantly increased in cells transfected with pre-*miR-223* relative to its negative control (TCam-2: $P=0.014$ and 2102Ep: $P=0.003$). The data support the efficiency of transfection.

Furthermore, inhibition of *miR-223-3p* led to a significant increase of FBXW7 expression in TCam-2 (1.9-fold; $P=0.003$) and 2102Ep (1.3-fold, $P=0.015$) cells. Similarly, overexpression of *miR-223-3p* significantly reduced FBXW7 expression in both cell lines (0.8-fold and $P < 0.01$ for both) (Fig. 2B). The findings indicate that *miR-223-3p* suppresses FBXW7 expression in human TGCT cells.

Functional consequences of miR-223-3p regulation in TGCT cells. To explore the functional role of *miR-223-3p* on apoptosis, we investigated the effect using flow cytometric detection of Annexin V-stained cells as well as by western blot analysis of cleaved PARP (cPARP, 25 kDa), which is an apoptosis marker. For the Annexin V assay, we observed that inhibition of *miR-223-3p* in TCam-2 cells significantly increased apoptotic cells by 80% ($P=0.018$), while overexpression of *miR-223-3p* reduced apoptotic cells by 38% ($P=0.038$), relative to their respective negative controls (Fig. 3A). Similar effects were also observed in 2102Ep cells, however, the effect was less pronounced compared to TCam-2 cells (27% increase in the *miR-223-3p* inhibition, $P=0.009$; 34% decrease in the *miR-223-3p* overexpression, $P=0.016$; Fig. 3A).

For the cPARP detection, silencing of *miR-223-3p* led to a significant increase of cPARP expression in both TCam-2 (1.6-fold, $P=0.003$) and 2102Ep (1.3-fold, $P=0.007$) cells, while overexpressing *miR-223-3p* resulted in a significant decrease of cPARP expression (TCam-2: 0.7-fold, $P=0.016$;




Figure 3. Modulation of cell apoptosis by *miR-223-3p* in TGCT cells. (A) Representative flow cytometric images of cells co-stained with Annexin V-FITC and propidium iodide (PI) upon silencing or overexpression of *miR-223-3p*. The early and late apoptotic cells are presented by Annexin V⁺/PI⁻ (lower right) and Annexin V⁺/PI⁺ (upper right) cells, respectively. Quantification of total apoptotic cells (Annexin V⁺) is shown below. The percentage of apoptotic cells represents both early and late apoptotic cells. (B) Western blot analysis of the 25-kDa cleaved PARP (cPARP, an apoptosis marker) in both cell lines with *miR-223-3p* inhibition or overexpression. The relative expression of cPARP was normalized to GAPDH. Data represent mean \pm SD from at least three independent experiments. P-values were calculated by paired t-test. *P<0.05; **P<0.01.

2102Ep: 0.8-fold, P=0.002) (Fig. 3B). These results indicate that *miR-223-3p* inhibits apoptosis in TGCT cells.

For cell proliferation, we applied three different assays: Click-iT EdU, WST-1 and trypan blue exclusion. Using the EdU assay, we observed reduction of EdU-positive cells upon silencing of *miR-223-3p* in both TCam-2 (50.3 vs. 30.8%, P=0.026) and 2102Ep cells (46.5 vs. 30.0%, P=0.045), and increase of EdU-positive cells upon overexpression of *miR-223-3p* (TCam-2: 48.1 vs. 61.6%, P=0.003; 2102Ep: 47.4 vs. 57.7%, P=0.027) (Fig. 4A). Similarly, the trypan blue exclusion assay revealed reduction of cell count upon silencing of *miR-223-3p* (TCam-2: 0.8-fold, P=0.004; 2102Ep: 0.8-fold, P=0.016) and increase of cell number upon overexpression of *miR-223-3p* (TCam-2: 1.3-fold, P=0.006; 2102Ep: 1.2-fold,

P=0.011) (Fig. 4B). The WST-1 assay also showed that silencing of *miR-223-3p* reduced cell growth at 72-h post-transfection in both TCam-2 (P=0.043) and 2102Ep (P=0.041) cells (Fig. 4C). Taken together, the results support that *miR-223-3p* promotes cell proliferation in TGCT cell lines.

miR-223-3p mediates regulation of cell growth and apoptosis through *FBXW7* in TGCT. Given that *FBXW7* expression is a well-characterized target of *miR-223-3p*, we tested whether ectopically expressed *FBXW7* could rescue the *miR-223-3p*-mediated apoptotic and proliferative effects. We co-transfected TCam-2 cells with pre-miR-223 together with a plasmid expressing the entire open reading frame of *FBXW7* without the *miR-223-3p* binding site (pCMV-Myc *FBXW7*) or




Figure 4. Effect of *miR-223-3p* regulation on cell growth in TGCT cells. (A) Representative flow cytometric images of proliferating cells with EdU incorporation in anti-*miR-223* or pre-*miR-223* cells and their respective negative controls at 72 h post-transfection (upper). The graph shows the changes of EdU-positive cells in both cell lines upon silencing or overexpression of *miR-223-3p* from three independent experiments (lower). (B) Total live cells were counted using trypan blue dye exclusion assay (n=4). (C) Relative cell growth was examined at different time-points in both cell lines transfected with *miR-223-3p* inhibitor or negative control using WST-1 assay (n=3 for each time-point). Data represent mean \pm SD. P-values were calculated by paired t-test. *P<0.05; **P<0.01.

a vector control. The effects on cell apoptosis and proliferation were determined using western blot analysis of cPARP and trypan blue exclusion assay, respectively. As shown in Fig. 5A, the endogenous FBXW7 was reduced in both cells co-transfected with pre-*miR-223* and Myc-FBXW7 or vector control as compared with the negative control-transfected cells, indicating the suppression of endogenous FBXW7 by *miR-223-3p* overexpression.

Ectopic expression of *miR-223-3p* significantly reduced the abundance of cPARP (P=0.043) and increased the number of live cells (P=0.006) as compared to their respective controls; the effects were abolished by the ectopically expressed

FBXW7 (Fig. 5). Together, our data indicate that *miR-223-3p* regulates cell growth and apoptosis in TGCT cells through FBXW7.

Discussion

miR-223-3p expression was found higher in TGCTs than NTs in our previous study (10), and here, we validated the findings in independent cohorts using previously published dataset (7). Deregulation of *miR-223-3p* has been observed in a variety of tumor types. Overexpression was found in T-cell acute lymphoblastic leukemia (11), oral (12), esophageal (14), gastric




Figure 5. Rescue of *miR-223-3p* mediated apoptosis and cell growth effects by ectopically expressed FBXW7. TCam-2 cells were co-transfected with *miR-223-3p* and Myc-FBXW7 or an empty vector, and compared to cells co-transfected with pre-miR-NC and empty vector. After 48 h of transfection, cells were harvested for apoptosis assay using western blot analysis of cPARP (A) and cell growth using trypan blue exclusion assays (B). (A) Exogenous Myc-FBXW7 and endogenous FBXW7 were detected by immunoblotting with anti-Myc-tag and anti-FBXW7, respectively. The relative cPARP expression was normalized to GAPDH. (B) Relative cell growth was determined by counting trypan blue negative cells. Data represent mean \pm SD from four independent experiments. P-values were calculated by paired t-test. *P<0.05; **P<0.01.

(20,21), bladder (22), and pancreatic (23) cancers, while its reduced expression has been reported in osteosarcoma (24), chronic lymphocytic leukemia (25), and intrahepatic cholangiocarcinoma (26). These findings indicate that *miR-223-3p* plays vital roles in a variety of tumor types, either as an oncogene or tumor suppressor depending on the cellular contexts. Consistent with its dual role, *miR-223-3p* has been shown to function as an oncogene in T-cell acute lymphoblastic leukemia, gastric and lung cancers (13,15,21,27,28), and as a tumor suppressor in cutaneous T-cell lymphoma and prostate cancer (29,30). Given its diverse function in different cancer types, we characterized the functional role of *miR-223-3p* in TGCT cells. *miR-223-3p* was shown to promote cell proliferation in TGCT cell lines in all three methods applied and which are based on different principles: the Click-iT EdU assay allows the detection of the thymidine analog EdU incorporated into cellular DNA during replication; the WST-1 assay is based on the metabolic activity of cells for conversion of the tetrazolium salt WST-1 into a colored dye, and the trypan blue exclusion assay provides direct counting of the number of live cells. Our findings support its oncogenic role in TGCT by promoting cell growth and inhibiting apoptosis in TGCT cell lines.

Additionally, *miR-223-3p* has been shown to modulate drug response in several cancer types (31-35). Importantly, *miR-223-3p* regulates cisplatin sensitivity in gastric and esophageal cancers (31,32). Given that most TGCTs are responsive to cisplatin treatment, it is intriguing to speculate that *miR-223-3p* may play an important role in cisplatin sensitivity in TGCT. Further investigations are warranted to evaluate the role of *miR-223-3p* in cisplatin response in TGCT.

FBXW7 has been demonstrated as a direct target of *miR-223-3p* using luciferase reporter assays (13,15). Here, we

show that FBXW7 expression is lower in TGCTs than NT and inversely correlated with *miR-223-3p*, and *miR-223-3p* regulates FBXW7 protein expression using both gain- and loss-of-function studies. Most importantly, ectopic expression of the *FBXW7* open reading frame can rescue the cell growth and apoptosis effects mediated by *miR-223-3p*. Together, our findings suggest that *miR-223-3p* regulates FBXW7 in TGCT and this regulatory pathway plays an important role in TGCT pathogenesis.

As afore-mentioned, FBXW7 is an E3 ubiquitin ligase that degrades several proto-oncogenes involved in cell growth, apoptosis, cell cycle regulation and differentiation (16). Therefore, the functional phenotypes observed in this study could be due to the loss of FBXW7-mediated degradation of its substrates. Furthermore, numerous cancer-associated mutations in *FBXW7* have been found in many cancer types (36), and loss of FBXW7 function can lead to chromosomal instability and tumorigenesis (37,38). These findings support the tumor suppressor function of FBXW7 in human cancers. Although nothing is known about its role in TGCT, FBXW7 is expressed specifically in the undifferentiated spermatogonia and suppresses cell proliferation of spermatogonial stem cell in mice (39). It is tempting to speculate that loss of FBXW7 expression could lead to uncontrolled cell growth in TGCT.

In conclusion, we report deregulation of *miR-223-3p* and FBXW7 in human TGCT. Our findings also reveal an oncogenic role of *miR-223-3p* through repression of the FBXW7 tumor suppressor, suggesting that this regulation is important for cell proliferation and apoptosis in TGCT. This study provides additional evidence of miRNA function in testicular germ cell tumorigenesis.

Acknowledgements

We thank Dr Leendert H.J. Looijenga and Dr Peter Andrews for TGCT cell lines; the Cooperative Human Tissue Network for frozen tissue samples; and the members of the sRNA Group for valuable discussions and suggestions. This study was supported by the Swedish Research Council (523-2009-3517 and 521-2010-3518), the Swedish Cancer Society, the Cancer Research Funds of Radiumhemmet, Karolinska Institutet and Stockholm County Council. J. Liu and H. Shi are the recipients of China Scholarship Council training grants.

References

- Mannuel HD, Mitikiri N, Khan M and Hussain A: Testicular germ cell tumors: Biology and clinical update. *Curr Opin Oncol* 24: 266-271, 2012.
- Nigam M, Aschebrook-Kilfoy B, Shikanov S and Eggener S: Increasing incidence of testicular cancer in the United States and Europe between 1992 and 2009. *World J Urol* 33: 623-631, 2015.
- McGlynn KA and Cook MB: Etiologic factors in testicular germ-cell tumors. *Future Oncol* 5: 1389-1402, 2009.
- Koychev D, Oechsle K, Bokemeyer C and Honecker F: Treatment of patients with relapsed and/or cisplatin-refractory metastatic germ cell tumours: An update. *Int J Androl* 34: e266-e273, 2011.
- Calin GA and Croce CM: MicroRNA signatures in human cancers. *Nat Rev Cancer* 6: 857-866, 2006.
- Palmer RD, Murray MJ, Saini HK, van Dongen S, Abreu-Goodger C, Muralidhar B, Pett MR, Thornton CM, Nicholson JC, Enright AJ, *et al*: Children's Cancer and Leukaemia Group: Malignant germ cell tumors display common microRNA profiles resulting in global changes in expression of messenger RNA targets. *Cancer Res* 70: 2911-2923, 2010.
- Gillis AJ, Stoop HJ, Hersmus R, Oosterhuis JW, Sun Y, Chen C, Guenther S, Sherlock J, Veltman I, Baeten J, *et al*: High-throughput microRNAome analysis in human germ cell tumours. *J Pathol* 213: 319-328, 2007.
- Suh MR, Lee Y, Kim JY, Kim SK, Moon SH, Lee JY, Cha KY, Chung HM, Yoon HS, Moon SY, *et al*: Human embryonic stem cells express a unique set of microRNAs. *Dev Biol* 270: 488-498, 2004.
- Voorhoeve PM, le Sage C, Schrier M, Gillis AJ, Stoop H, Nagel R, Liu YP, van Duijse J, Drost J, Griekspoor A, *et al*: A genetic screen implicates miRNA-372 and miRNA-373 as oncogenes in testicular germ cell tumors. *Cell* 124: 1169-1181, 2006.
- Özata DM, Li X, Lee L, Liu J, Warsito D, Hajeri P, Hultman I, Fotouhi O, Marklund S, Åhrlund-Richter L, *et al*: Loss of miR-514a-3p regulation of PEG3 activates the NF-kappa B pathway in human testicular germ cell tumors. *Cell Death Dis* (In press).
- Chiaretti S, Messina M, Tavoraro S, Zardo G, Elia L, Vitale A, Fatica A, Gorello P, Picciocchi A, Scappucci G, *et al*: Gene expression profiling identifies a subset of adult T-cell acute lymphoblastic leukemia with myeloid-like gene features and over-expression of miR-223. *Haematologica* 95: 1114-1121, 2010.
- Manikandan M, Deva Magendhra Rao AK, Arunkumar G, Manickavasagam M, Rajkumar KS, Rajaraman R and Munirajan AK: Oral squamous cell carcinoma: microRNA expression profiling and integrative analyses for elucidation of tumorigenesis mechanism. *Mol Cancer* 15: 28, 2016.
- Mavrakis KJ, Van Der Meulen J, Wolfe AL, Liu X, Mets E, Taghon T, Khan AA, Setty M, Rondou P, Vandenberghe P, *et al*: A cooperative microRNA-tumor suppressor gene network in acute T-cell lymphoblastic leukemia (T-ALL). *Nat Genet* 43: 673-678, 2011.
- Kurashige J, Watanabe M, Iwatsuki M, Kinoshita K, Saito S, Hiyoshi Y, Kamohara H, Baba Y, Mimori K and Baba H: Overexpression of microRNA-223 regulates the ubiquitin ligase FBXW7 in oesophageal squamous cell carcinoma. *Br J Cancer* 106: 182-188, 2012.
- Li J, Guo Y, Liang X, Sun M, Wang G, De W and Wu W: MicroRNA-223 functions as an oncogene in human gastric cancer by targeting FBXW7/hCdc4. *J Cancer Res Clin Oncol* 138: 763-774, 2012.
- Welcker M and Clurman BE: FBW7 ubiquitin ligase: A tumour suppressor at the crossroads of cell division, growth and differentiation. *Nat Rev Cancer* 8: 83-93, 2008.
- Wang L, Ye X, Liu Y, Wei W and Wang Z: Aberrant regulation of FBW7 in cancer. *Oncotarget* 5: 2000-2015, 2014.
- Mizuno Y, Gotoh A, Kamidono S and Kitazawa S: Establishment and characterization of a new human testicular germ cell tumor cell line (TCam-2). *Nippon Hinyokika Gakkai Zasshi* 84: 1211-1218, 1993 (In Japanese).
- Andrews PW, Goodfellow PN, Shevinsky LH, Bronson DL and Knowles BB: Cell-surface antigens of a clonal human embryonal carcinoma cell line: Morphological and antigenic differentiation in culture. *Int J Cancer* 29: 523-531, 1982.
- Petrocca F, Visone R, Onelli MR, Shah MH, Nicoloso MS, de Martino I, Iliopoulos D, Pilozzi E, Liu CG, Negrini M, *et al*: E2F1-regulated microRNAs impair TGFbeta-dependent cell-cycle arrest and apoptosis in gastric cancer. *Cancer Cell* 13: 272-286, 2008.
- Li X, Zhang Y, Zhang H, Liu X, Gong T, Li M, Sun L, Ji G, Shi Y, Han Z, *et al*: miRNA-223 promotes gastric cancer invasion and metastasis by targeting tumor suppressor EPB41L3. *Mol Cancer Res* 9: 824-833, 2011.
- Gottardo F, Liu CG, Ferracin M, Calin GA, Fassan M, Bassi P, Sevignani C, Byrne D, Negrini M, Pagano F, *et al*: Micro-RNA profiling in kidney and bladder cancers. *Urol Oncol* 25: 387-392, 2007.
- Bloomston M, Frankel WL, Petrocca F, Volinia S, Alder H, Hagan JP, Liu CG, Bhatt D, Taccioli C and Croce CM: MicroRNA expression patterns to differentiate pancreatic adenocarcinoma from normal pancreas and chronic pancreatitis. *JAMA* 297: 1901-1908, 2007.
- Xu J, Yao Q, Hou Y, Xu M, Liu S, Yang L, Zhang L and Xu H: MiR-223/Ect2/p21 signaling regulates osteosarcoma cell cycle progression and proliferation. *Biomed Pharmacother* 67: 381-386, 2013.
- Fazi F, Racanicchi S, Zardo G, Starnes LM, Mancini M, Travaglini L, Diverio D, Ammatuna E, Cimino G, Lo-Coco F, *et al*: Epigenetic silencing of the myelopoiesis regulator microRNA-223 by the AML1/ETO oncoprotein. *Cancer Cell* 12: 457-466, 2007.
- Karakatsanis A, Papaconstantinou I, Gazouli M, Lyberopoulou A, Polymeneas G and Voros D: Expression of microRNAs, miR-21, miR-31, miR-122, miR-145, miR-146a, miR-200c, miR-221, miR-222, and miR-223 in patients with hepatocellular carcinoma or intrahepatic cholangiocarcinoma and its prognostic significance. *Mol Carcinog* 52: 297-303, 2013.
- Mansour MR, Sanda T, Lawton LN, Li X, Kreslavsky T, Novina CD, Brand M, Gutierrez A, Kelliher MA, Jamieson CH, *et al*: The TAL1 complex targets the FBXW7 tumor suppressor by activating miR-223 in human T cell acute lymphoblastic leukemia. *J Exp Med* 210: 1545-1557, 2013.
- Liang H, Yan X, Pan Y, Wang Y, Wang N, Li L, Liu Y, Chen X, Zhang CY, Gu H, *et al*: MicroRNA-223 delivered by platelet-derived microvesicles promotes lung cancer cell invasion via targeting tumor suppressor EPB41L3. *Mol Cancer* 14: 58, 2015.
- McGirt LY, Adams CM, Baerenwald DA, Zwerner JP, Zic JA and Eischen CM: miR-223 regulates cell growth and targets proto-oncogenes in mycosis fungoides/cutaneous T-cell lymphoma. *J Invest Dermatol* 134: 1101-1107, 2014.
- Kurozumi A, Goto Y, Matsushita R, Fukumoto I, Kato M, Nishikawa R, Sakamoto S, Enokida H, Nakagawa M, Ichikawa T, *et al*: Tumor-suppressive microRNA-223 inhibits cancer cell migration and invasion by targeting ITGA3/ITGB1 signaling in prostate cancer. *Cancer Sci* 107: 84-94, 2016.
- Streppel MM, Pai S, Campbell NR, Hu C, Yabuuchi S, Canto MI, Wang JS, Montgomery EA and Maitra A: MicroRNA 223 is upregulated in the multistep progression of Barrett's esophagus and modulates sensitivity to chemotherapy by targeting PARP1. *Clin Cancer Res* 19: 4067-4078, 2013.
- Zhou X, Jin W, Jia H, Yan J and Zhang G: MiR-223 promotes the cisplatin resistance of human gastric cancer cells via regulating cell cycle by targeting FBXW7. *J Exp Clin Cancer Res* 34: 28, 2015.
- Eto K, Iwatsuki M, Watanabe M, Ishimoto T, Ida S, Imamura Y, Iwagami S, Baba Y, Sakamoto Y, Miyamoto Y, *et al*: The sensitivity of gastric cancer to trastuzumab is regulated by the miR-223/FBXW7 pathway. *Int J Cancer* 136: 1537-1545, 2015.
- Kumar V, Palermo R, Talora C, Campese AF, Checquolo S, Bellavia D, Tottone L, Testa G, Miele E, Indraco S, *et al*: Notch and NF-kB signaling pathways regulate miR-223/FBXW7 axis in T-cell acute lymphoblastic leukemia. *Leukemia* 28: 2324-2335, 2014.

35. Li R, Wu S, Chen X, Xu H, Teng P and Li W: miR-223/FBW7 axis regulates doxorubicin sensitivity through epithelial mesenchymal transition in non-small cell lung cancer. *Am J Transl Res* 8: 2512-2524, 2016.
36. Akhoondi S, Sun D, von der Lehr N, Apostolidou S, Klotz K, Maljukova A, Cepeda D, Fiegl H, Dafou D, Marth C, *et al*: FBXW7/hCDC4 is a general tumor suppressor in human cancer. *Cancer Res* 67: 9006-9012, 2007.
37. Rajagopalan H, Jallepalli PV, Rago C, Velculescu VE, Kinzler KW, Vogelstein B and Lengauer C: Inactivation of hCDC4 can cause chromosomal instability. *Nature* 428: 77-81, 2004.
38. Mao JH, Perez-Losada J, Wu D, Delrosario R, Tsunematsu R, Nakayama KI, Brown K, Bryson S and Balmain A: Fbxw7/Cdc4 is a p53-dependent, haploinsufficient tumour suppressor gene. *Nature* 432: 775-779, 2004.
39. Kanatsu-Shinohara M, Onoyama I, Nakayama KI and Shinohara T: Skp1-Cullin-F-box (SCF)-type ubiquitin ligase FBXW7 negatively regulates spermatogonial stem cell self-renewal. *Proc Natl Acad Sci USA* 111: 8826-8831, 2014.