

# The acute vs. chronic effect of exercise on insulin sensitivity: nothing lasts forever

Fred J. DiMenna and Avigdor D. Arad

Regular exercise causes chronic adaptations in anatomy/physiology that provide first-line defense for disease prevention/treatment ('exercise is medicine'). However, transient changes in function that occur following each exercise bout (acute effect) are also important to consider. For example, in contrast to chronic adaptations, the effect of exercise on insulin sensitivity is predominantly rooted in a prolonged acute effect (PAE) that can last up to 72 h. Untrained individuals and individuals with lower insulin sensitivity benefit more from this effect and even trained individuals with high insulin sensitivity restore most of a detraining-induced loss following one session of resumed training. Consequently, exercise to combat insulin resistance that begins the pathological journey to cardiometabolic diseases including type 2 diabetes (T2D) should be prescribed with precision to elicit a PAE on insulin sensitivity to serve as a first-line defense prior to

pharmaceutical intervention or, when such intervention is necessary, a potential adjunct to it. *Cardiovasc Endocrinol Metab* 10: 149–161 Copyright © 2020 The Author(s). Published by Wolters Kluwer Health, Inc.

Cardiovascular Endocrinology & Metabolism 2021, 10:149–161

Video Abstract: <http://links.lww.com/CAEN/A27>

Keywords: acute exercise effect, exercise, exercise is medicine, exercise prescription, exercise intensity, high-intensity interval training, insulin sensitivity, insulin resistance, prediabetes, type 2 diabetes

Division of Endocrinology, Diabetes and Bone, Department of Medicine, Icahn School of Medicine at Mount Sinai, New York, USA

Correspondence to Fred DiMenna, PhD, Division of Endocrinology, Diabetes and Bone, Department of Medicine, Icahn School of Medicine at Mount Sinai, One Gustave L. Levy Place Box 1055, New York 10029, NY, USA  
E-mail: [fjdimenna2020@gmail.com](mailto:fjdimenna2020@gmail.com)

Received 22 May 2020 Accepted 22 September 2020

## Introduction

The lingering effects of a single exercise bout (i.e. acute effect) can provide misleading information regarding a person's health status in some circumstances. For example, if a habitually-sedentary person performed a bout of exercise the day before a physician's visit, the prolonged acute effect (PAE) of exercise on insulin sensitivity could mask an elevated reading on an oral glucose tolerance test (OGTT). Conversely, for those who exercise regularly and, therefore, benefit from the 'continuous' presence of such an effect, the representation would be accurate. Importantly, this is also the case in the research setting where it is necessary to control for PAE when assessing previously-sedentary individuals; for example, before and after a training intervention if the objective is to determine the chronic effect that accumulates from repeat training (i.e. 'training effect'). However, the need to discount the lingering effect of an exercise bout in these circumstances contrasts an acute effect's potential role as a therapeutic agent for individuals who exercise frequently enough such that it is present most/all of the time. Moreover, in reality, the highly-touted chronic effect is also a bit

of a misrepresentation because it, too, must be maintained. Granted, 'detraining' has a longer time course compared to dissipation of the acute effect; however, chronic adaptations do indeed wane if regular workouts are not continued. Finally, referring to either as a singular entity is an oversimplification because each likely includes multiple alterations in anatomy/physiology which might have different time courses for dissipation. For example, some adaptations comprising the acute effect might be longer lasting while some chronic changes are shorter lived. However, one thing is certain: when it comes to the effect of exercise on insulin sensitivity, nothing lasts forever.

## Type 2 diabetes and the obesity epidemic

Estimates indicate that ~34 million US citizens have diabetes [1] with 90–95% suffering from T2D [1]. Furthermore, it is estimated that ~88 million have prediabetes, which means blood-glucose concentration is higher than normal, but not yet high enough for T2D diagnosis (e.g. fasting level of 100–125 mg/dL compared to <100 and >125 for normal and T2D, respectively). Predictions suggest up to 70% of individuals with prediabetes will develop T2D [2] such that total prevalence in 2050 could increase to as much as 33% of the population (compared to, for example, 14% in 2010) [3]. Intervention at the prediabetic stage is, therefore, critical. Excess body-fat deposition is a risk factor for T2D with 10-year risk increasing by 4.6- and 10.0-fold for females and 3.5- and 11.2-fold

Supplemental Digital Content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's website, [www.cardiovascularendocrinology.com](http://www.cardiovascularendocrinology.com).

This is an open access article distributed under the Creative Commons Attribution License 4.0 (CCBY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

for males classified as overweight (BMI, 25.0–29.9 kg/m<sup>2</sup>) or obese (BMI, 30.0–34.9 kg/m<sup>2</sup>), respectively [4]. However, ‘metabolically-healthy obese’ and ‘metabolically-obese normal-weight’ (MONW) phenotypes indicate greater complexity [4] at least partially rooted in genetic predisposition [5].

### Exercise is medicine

‘Exercise is medicine’ is an initiative launched by the American Medical Association and American College of Sports Medicine in 2007 [6]. The objective was to make physical-activity assessment/promotion standard in clinical practice so that healthcare can be augmented by evidence-based physical-activity resources [6]. Given the prevalence of insulin resistance and its association with obesity as a disorder of fuel storage/usage, exercise is medicine might be particularly appropriate for addressing the T2D epidemic [7]. Insulin-stimulated glucose uptake is primarily directed to skeletal muscle [8] and, in addition to energy use during exercise, post-exercise recovery and the additional fat-free mass gained by sedentary individuals who begin exercising each serve to favorably alter energy balance. This could prove beneficial for prevention/treatment of these diseases. However, to isolate the influence of chronic exercise per se on insulin sensitivity, in addition to differentiating chronic from acute effect, it is important to consider the independent effect of body-fat loss that often occurs when sedentary individuals begin exercising. When body fat is reduced, insulin resistance improves even when reduction occurs via surgical intervention [9,10]. Conversely, some evidence suggests that chronic exercise neither augments the improvement in insulin sensitivity induced by weight loss due to dietary restriction without exercise [11,12; *c.f.* 13] nor improves insulin sensitivity when weight loss is prevented [14–17; *c.f.* 18]. This suggests that the chronic effect of exercise per se might be insufficient as medicine for insulin resistance. Furthermore, considering the growing appreciation for location in addition to quantity of fat storage [19,20], the ability for chronic exercise to redistribute fat is also important to consider. Finally, some of the structural changes that result in improved insulin sensitivity consequent to chronic training might be unattainable for individuals most in need of an intervention to combat insulin resistance. Collectively, these complexities associated with the chronic effect coupled with the relatively ‘simple’ (not mechanistically but from a practical standpoint) influence an individual session can evoke implies that the acute effect might be more relevant to consider when prescribing exercise to combat insulin resistance.

### Contraction-mediated insulin-independent glucose uptake

The acute effect of exercise as a stimulant for glucose uptake by skeletal muscle begins during exercise when a number of factors work in concert to facilitate

an increase [21]. Exercise-induced increased perfusion to and capillary recruitment within contracting muscle allows for greater glucose delivery/exchange [22] while contraction-mediated insulin-independent activation of intracellular signaling pathways results in glucose extraction which can exceed that which is present with maximal insulin stimulation depending on muscle fibre type [22]. Glucose transport occurs via facilitated diffusion utilising glucose transporter isoform 4 (GLUT-4), a protein found in muscle and adipose tissue [23]. Similar to insulin stimulation, contractions increase GLUT-4 translocation to the sarcolemma of muscle fibers; however, in this case, there is evidence which indicates that GLUT-4 might emanate from a different intracellular pool [24]. This distinction is important because it suggests that the effects of insulin and contractions are additive [25]. Exercise-induced activation of GLUT-4 is predominantly driven by AMP-activated protein kinase (AMPK; an enzyme that plays an important role in regulating cellular energy status) [26] and Rac1, a member of the guanosine triphosphatase (GTP) family of regulatory proteins [27]. However, other pathways (e.g. related to calcium, reactive oxygen species and nitric oxide signaling) appear to provide protective redundancy [28]. During exercise, increased AMPK results in deactivation of *tre-2/USP6*, *BUB2*, *cdc16* domain family member 1 (*TBC1D1*), a member of the Rab GTPase-activating protein family [29]. Deactivation of *TBC1D1* removes its inhibitory influence thereby promoting GLUT-4 translocation [29,30]. This ‘early phase’ comprising insulin-independent glucose uptake during exercise persists after the bout for 30 (but not 120) min [31]; however, the acute effect of exercise on insulin sensitivity also includes long-lasting enhancement that can persist for up to 72 h [32]. As previously mentioned, this PAE of exercise on insulin sensitivity might be most relevant when prescribing exercise as medicine for insulin resistance.

### The prolonged acute exercise effect on insulin sensitivity: mechanistic bases

Four hours following exercise, insulin-stimulated GLUT4 translocation and glucose uptake by rat skeletal muscle is increased compared to the pre-exercise condition in association with increased phosphorylation of Akt substrate of 160 kDa (*AS160*; also known as *TBC1D4*), a Rab GTPase-activating protein paralogue [33] that appears to be ‘primed’ by AMPK [30]. However, unlike the insulin-independent route, *TBC1D1* is not directly involved in this process [33]. For example, *Kjøbsted et al.* recently confirmed that 3 h after contraction or 6 h after stimulation of 5-Aminoimidazole-4-carboxamide ribonucleotide (the first direct AMPK activator), the increase in insulin-stimulated glucose uptake driven by enhanced insulin sensitivity in muscle of wild-type mice is absent in mice lacking *TBC1D4* even though the phosphorylation pattern of *TBC1D1* in these animals was unaltered [30]. Moreover,

this prolonged post-exercise effect on insulin sensitivity occurs with unaltered insulin binding or insulin receptor substrate (IRS) activation (i.e. proximal insulin signaling) [30]; however, a simple cause-effect relationship between events at/distal to AS160 and the exercise-induced increase in insulin-stimulated glucose uptake has not been clarified and likely involves greater complexity [29]. To differentiate the PAE of exercise on insulin sensitivity from the chronic effect, Prior *et al.* had previously-sedentary older individuals perform aerobic training for 6 months followed by 2 weeks of detraining to ‘wash out’ PAE and found that insulin sensitivity (as evaluated during euglycemic hyperinsulinemic clamp; EHC) increased by ~25% after training with an ~18% increase still present after detraining [34]. Reduction of the training-induced increase from ~25 to ~18% after 2 weeks of detraining indicates that ~25% of the improvement that was achieved due to 6 months of regular training was attributable to short-term effects [34]. Mechanistically, the loss of these short-term effects occurred in conjunction with return of GLUT4 expression, AMPK $\alpha$ 1 expression and insulin-mediated activation of glycogen synthase to baseline levels whereas the increase that remained was associated with an increase in capillary density that training elicited [34]. Importance of microvascular structural changes as a chronic adaptation to exercise training for improved glycemic control agreed with earlier observations by Williamson *et al.* who found that the increase in width of the capillary basement membrane that occurs with sedentary aging is normalised after 9 months of endurance training regardless of baseline glucose tolerance [35]. However, being that body-fat modulates insulin resistance, especially when it is located in the abdominal region [36], the most important factor that positively influences insulin sensitivity as a chronic adaptation to exercise training appears to be the change in body composition that typically accompanies it [11,12,14–17].

#### The effect of exercise on insulin sensitivity: early findings

In a landmark study, Heath *et al.* found that when chronically-trained individuals with high insulin sensitivity performed no exercise for 10 days, an OGTT revealed a reduction in insulin sensitivity compared to the trained state despite no change in fitness level (as indicated by maximal rate of oxygen uptake;  $\dot{V}O_{2max}$ ) or body weight/fat [37]. However, a single bout following detraining resulted in an insulin response that was not different at any post-exercise time point (30, 60, 120 and 180 min) compared to the trained state [37]. Collectively, the marked reduction in insulin sensitivity following a short period of detraining and almost complete restoration of the ‘trained insulin sensitivity’ after the initial bout of resumption indicates the important role that the PAE of exercise plays in maintaining insulin sensitivity even for trained individuals [37].

The relevance of the PAE in relation to the overall influence of exercise on insulin sensitivity reported by

Heath *et al.* provided information in humans that complemented early research in animals. For example, 1 year prior, Richter *et al.* observed greater glucose utilisation in the isolated perfused hindlimb muscle for 4 h following 45 min of treadmill running by untrained rats when insulin was added to the perfusate [38]. Importantly, glucose utilisation was unchanged compared to the no-exercise control condition in the absence of added insulin which confirmed that acute exercise influenced the insulin-dependent pathway [38]. In a follow-up study, Ivy *et al.* compared glucose uptake at various insulin concentrations in the isolated perfused hindlimb muscle of rats that were trained for 14–16 weeks and found that the rate of glucose uptake the day after the final training session was ~50% greater than that which was present in sedentary controls [39]. However, 40–46 h later, there was no difference regardless of the presence of insulin at physiological or maximally-effective levels [39]. Furthermore, a swimming bout to exhaustion and muscle contractions induced by electrical stimulation both resulted in an increase in glucose uptake the following day that was of similar magnitude regardless of whether the rat was trained or untrained [39]. The authors concluded that the increase in insulin-stimulated glucose uptake demonstrated by chronically-trained rats is a function of the PAE of exercise on insulin sensitivity as opposed to a long-term adaptive response [39]. In 1989, Cartee *et al.* provided further insight when they found that the PAE of exercise on insulin sensitivity could persist for up to 48 h in rats subject to carbohydrate deprivation whereas high-carbohydrate feeding during the post-exercise period resulted in more rapid dissipation [40]. This was consistent with earlier findings which indicated that insulin-stimulated carbohydrate storage rate is increased after glycogen-depleting exercise without, but not with re-feeding, even though post-exercise carbohydrate oxidation rate is decreased in both cases [41].

More information regarding the PAE of exercise on insulin sensitivity in humans was provided by Rogers *et al.* in 1990 [42]. These researchers replicated the 10-day detraining methodology employed by Heath *et al.* (see above); however, in this case, their subjects were 14 masters athletes whose life-long training allowed them to avoid the development of insulin resistance that typically accompanies aging. Interestingly, for 10 of these athletes, responses for glucose and insulin after 10 days of detraining were not different than those of young, lean controls [42]. This meant that for these individuals, in addition to the PAE of exercise on insulin sensitivity, exercise training conveyed a chronic effect that protected them from aging-related insulin resistance. The authors speculated that the positive effect of chronic exercise on body fatness was responsible [42]. However, despite similar leanness, the other four athletes experienced a reduction in glucose tolerance due to detraining which confirmed that for these individuals, the PAE of exercise was responsible for their ‘youthful’ insulin sensitivity [42]. The authors

speculated that this subgroup had a genetic predisposition to develop insulin resistance that was offset by the PAE of exercise [42]. This supports the contention that for individuals most at need for an intervention to maintain insulin sensitivity/prevent insulin resistance, it is the PAE as opposed to chronic effect of exercise on insulin sensitivity that is the more relevant to consider. However, in a previous study, these same researchers found that the improvement in OGTT demonstrated by individuals with mild T2D 18h after the final bout of a 7-day aerobic training intervention was not yet present following the initial session [43]. This suggests greater complexity that might be rooted in the counterbalancing effect of reduced insulin concentration or the intensity of the exercise being performed [43]. Nevertheless, the fact that insulin sensitivity was improved after 7 days (i.e. a training period that was too short to result in changes in body composition or  $\dot{V}O_{2\max}$ ) indicates that the PAE of exercise on insulin sensitivity (albeit apparently derived from the accumulation of multiple bouts) was capable of providing therapeutic benefit for patients with T2D [43].

Definitive proof that the improved insulin action demonstrated by endurance-trained individuals was due to increased insulin sensitivity was provided by King *et al.* in 1987 [44]. These researchers used a two-stage (i.e. submaximal and maximal stimulation) EHC to tease out the difference between changes in insulin sensitivity per se as opposed to changes in responsiveness to maximal insulin stimulation [44]. In a follow-up study, the same group also confirmed that the superior insulin sensitivity demonstrated by trained individuals is predominantly transient as the same rate of whole-body glucose disposal required an ~67% higher plasma insulin response after 14 days of inactivity compared to 16h post-exercise [45]. Finally, in addition to these longitudinal studies, a cross-sectional analysis by Helmrich *et al.* confirmed that leisure-time physical activity (quantified as energy expended per week in walking, stair climbing, and sports) was inversely related to the development of T2D for 5990 male alumni of the University of Pennsylvania during 98524 man-years of follow-up from 1962 to 1976 [46]. Importantly, this protective effect of physical activity was strongest in persons with high BMI, a history of hypertension or a parental history of diabetes, which confirmed that those at greatest risk for developing T2D stood to benefit most from exercise training per se as a preventive intervention [46].

### The prolonged acute exercise effect on insulin sensitivity: recent findings

Recent research provides more insight regarding the effect of exercise on insulin sensitivity and, specifically, the importance of the PAE in this regard. In 2019, Steenberg *et al.* had healthy men perform one-legged knee-extensor exercise at the same relative intensity before and after 12 weeks of bilateral cycle training and

observed enhanced insulin sensitivity (EHC) 4–6h after exercise in the trained compared to untrained state [47]. However, in the trained state, when insulin sensitivity was compared between tested (PAE plus chronic effect) and untested (chronic effect only) leg, the difference between the two was reduced by ~50% compared to the difference that was present pre-training (i.e. when tested and untested leg were experiencing only the PAE and no effect, respectively) [47]. This implies that the chronic effect occurred at the ‘expense’ of the PAE that was present in the untrained state [47]. One possible explanation is that in the trained state, exercise-induced AMPK-mediated phosphorylation of TBC1D4 was decreased despite the fact that pre- and post-training bouts were performed at the same ‘relative’ intensity [47]. Regardless of mechanistic basis; however, assuming that the exercised muscle’s rate of glucose uptake in the trained state was not maximal (i.e. there was ‘room’ for a further increase), reciprocity between acute and chronic enhancements implies that the positive influence of exercise per se on insulin sensitivity is constrained by an upper limit that cannot be surpassed [47]. A similar conclusion can be drawn from the work of Nelson and Horowitz who found that the superior insulin sensitivity (i.e. compared to sedentary individuals with similar body composition; however, still inferior compared to healthy lean individuals) displayed by chronically-trained overweight/obese individuals after 3 days of detraining (i.e. when only the chronic effect should still be present) was not further enhanced by the PAE following one bout of resumption of training whereas the PAE following the single bout increased insulin sensitivity to the same level for their sedentary counterparts [48]. Collectively, these findings support the contention that while chronic effects from accumulated training should not be discounted, apparently, as Cartee asserts, ‘once is enough’ to benefit insulin sensitivity [49], particularly for individuals most in need of a preventive/treatment intervention.

### Eliciting the prolonged acute exercise effect on insulin sensitivity

‘Physical activity’ refers to bodily movement by skeletal muscle which results in energy expenditure [50]. ‘Exercise’ has been defined as a subset of physical activity because it must be planned, structured and repetitive, and have a final or intermediate objective of improving or maintaining physical fitness [50]. While more specific, this classification is still broad and the term, therefore, describes many types of physical activities that individuals perform. While such variety likely improves program adherence [51], if exercise is to represent a true alternative to pharmaceutical intervention [6], it must be ‘prescribed’ with greater rigor [7]. For example, the objective in the research setting should be to define exercise for eliciting the PAE of exercise on insulin sensitivity with evidence-based precision by better identifying the most effective mode (e.g. endurance or resistance training and



in the case of the former, lower-body exclusive like the oft-performed leg cycling or full body), duration (the time for which exercise must be maintained during each individual session; e.g. a minimal and optimal dose), frequency (based on how long the effect lasts, elucidation of which requires investigations with measurements of insulin sensitivity at multiple time points post-exercise) and intensity (e.g. low sustained for long duration, moderate/high sustainable for lesser duration or very high unsustainable intermittent) with participant characteristics and post-exercise feeding taken into account. Table 1 provides a chronologically-ordered list of studies that have contributed to our current understanding of the PAE of exercise on insulin sensitivity which is important to consider when making these determinations and planning future research.

### Mode

Studies investigating the PAE of exercise on insulin sensitivity have typically involved endurance ('aerobic') exercise; specifically, leg-cycle ergometry, treadmill exercise or a combination of the two (see Table 1). This stands to reason being that cycling and walking/jogging are the most accessible modalities for the average person. Furthermore, exercise's acute insulin-sensitising effect is mediated by local (as opposed to systemic) contraction-induced factors [87,88]; hence, modalities like these which require sustained involvement of a greater quantity of muscle and larger muscles would likely be superior. Resistance ('strength') training can involve larger muscles; however, it can also be performed for upper-body musculature resulting in activation of more total muscle mass. Given that resistance training induces muscular hypertrophy and skeletal muscle is the predominant site for glucose disposal [89], it is not surprising that chronic resistance training improves insulin sensitivity [90]. However, unlike endurance exercise, the acute (and, therefore, not hypertrophy-/body-composition related) effect of resistance training has not been investigated frequently and findings are equivocal. With respect to the ability to improve glycemic control, van Dijk *et al.* used subcutaneous glucose monitoring following a single 45-min bout of endurance (cycling at 50% maximum work rate) or resistance (75% one-repetition maximum) exercise performed by overweight/obese males with impaired glucose tolerance or treated T2D and found that hyperglycemic glucose concentrations were similarly lowered in all three groups for 24h after both types of exercise [91]. As for insulin action, Fluckey *et al.* found that unlike older healthy individuals, older individuals with T2D and young control participants each demonstrated increased insulin clearance as evidenced by a reduced total insulin response (area under curve; AUC) with no change in C-peptide response (an indirect index of insulin secretion) during OGTT 18h after resistance training [59]. Fenicchia *et al.* assessed women with T2D and healthy controls and found that for the former, but

not the latter, full-body resistance training reduced glucose AUC and peak glucose concentration (OGTT) 12–24h post-exercise with no change in insulin concentration [65]. Finally, Black *et al.* found that individuals with impaired fasting glucose demonstrated increased insulin sensitivity (homeostatic model assessment; HOMA) 24h following two different full-body resistance training bouts [73]. Conversely, Chapman *et al.* observed no change compared to the no-exercise control condition for insulin sensitivity or AUC for glucose or insulin during an intravenous glucose tolerance test (IVGTT) for sedentary postmenopausal women with normal glucose tolerance 15h after resistance training [64]. In addition to metabolic health, these discrepant findings might be attributable to participant age, gender [64] or the method used to assess insulin sensitivity [66]. Specifically, with respect to the latter, instead of OGTT or IVGTT, Koopman *et al.* used an intravenous insulin tolerance test (ITT) and found that healthy male participants demonstrated increased insulin sensitivity 24h post-resistance training that mainly consisted of leg exercises [66]. In addition to testing method, resistance training intensity and control of pre-test diet and physical activity levels might be responsible for this finding [67]. However, large intersubject variability and absence of response in some participants suggest greater complexity [67]. Finally, Howlett *et al.* found that compared to the no-exercise condition, insulin sensitivity (EHC) was unchanged 24h following the final of three resistance training bouts performed over the course of a week by young recreationally-active men [69]. Interestingly, the authors also observed decreased insulin sensitivity immediately following the initial session (i.e. reversal of the immediate acute-exercise effect after endurance exercise; see above) in association with a decrease in AS160 phosphorylation [69]. These findings and others [92] indicate that even if resistance training provides a viable alternative to endurance exercise for inducing the PAE of exercise on insulin sensitivity, the post-exercise response by signaling pathways that affect glucose metabolism is different. In this regard, using an electrical stimulation resistance training model in rats, Kido *et al.* recently found that the increase in insulin-stimulated glucose uptake 6h after a resistance-training bout is greater when mechanistic target of rapamycin complex 1 (mTORC1) and associated downstream phosphorylation of IRS-1 serine residues (IRS-1 Ser) is inhibited [93]. This has resonance because IRS-1 Ser phosphorylation is associated with insulin resistance. Importantly, compared to endurance exercise, mTORC1 activation, which stimulates skeletal-muscle hypertrophy, is higher and has a longer duration after resistance training. Collectively, this implies that the mTORC1-induced phosphorylation of IRS-1 Ser and consequent elicitation of insulin resistance that is present following resistance training counters the AMPK-driven increase in insulin sensitivity that occurs following exercise per se resulting in reduction/elimination of the PAE of exercise

Table 1 A chronologically-ordered listing of studies that have contributed to our current understanding of the acute effect of exercise on insulin sensitivity in different human populations

Study	Participants	Insulin sensitivity assessment				Exercise effect	
		Mode	Volume	Intensity	Time post-exercise		
Devlin and Horton (1985) [52]	Obese (IGT or hyperinsulinemia untrained males (n = 6) vs. lean healthy untrained males (n = 6))	Cycle	2 min on/3 min off to limit of tolerance (obese, 33 ± 2 min; lean, 66 ± 3 min)	85% $\dot{V}O_{2max}$	12–16 h	EHC	↑ Insulin sensitivity post bout for obese, but not for lean
Devlin et al. (1987) [53]	T2D untrained males (n = 5)	Cycle	2 min on/3 min off to limit of tolerance (41 ± 10 min)	85% $\dot{V}O_{2max}$	12–16 h	EHC	↑ Insulin sensitivity post bout
Mikines et al. (1988) [54]	Healthy untrained young males (n = 7)	Cycle	60 min	150 W	0 h, 48 h, 120 h	EHC	↑ Insulin sensitivity at 48 h post bout; ↔ insulin sensitivity at 120 h post bout
Richter et al. (1989) [55]	Healthy young RA males (n = 6)	One-legged knee-extensor exercise	60 min	75% $W_{max}$	4 h	EHC	↑ Insulin sensitivity post bout for exercised, but not for rested leg
Young et al. (1989) [56]	Healthy trained males post ≥40 h detraining (n = 7) vs. healthy RA males (n = 7)	Cycle	40 min	~40% $\dot{V}O_{2max}$	14 h	OGTT	↓ I-AUC/↔ G-AUC post bout for untrained so that they were no longer different vs. trained who displayed superior I-AUC pre bout that was unchanged post bout; no difference between conditions
Burnstein et al. (1990) [57]	Obese T2D non-AA males (n = 3) and females (n = 3) vs. obese NG non-AA males (n = 2) and females (n = 5) vs. lean healthy non-AA males (n = 6)	Cycle	40 min	~80% $\dot{V}O_{2max}$	14 h	(I-AUC/G-AUC)	↑ Insulin sensitivity post bout for obese T2D and NG; ↔ insulin sensitivity post bout for lean
Kjaer et al. (1990) [58]	Healthy sedentary males (n = 7) vs. T2D sedentary males (n = 7)	Treadmill	60 min	HR = 150–160 bpm	60 min	EHC	↑ Insulin sensitivity post bout for T2D, but not for healthy
Fluckey et al. (1994) [59]	Young healthy untrained males (n = 3) and females (n = 4) vs. older T2D untrained males (n = 3) and females (n = 4)	Machine full-body resistance exercise	3 sets/exercise x 10 reps or to failure if it occurred sooner	calculated 10-RM	18 h	OGTT	↓ I-AUC/↔ G-AUC post bout for both groups
Ben Ezra et al. (1995) [60]	Healthy untrained females (n = 24)	Treadmill	~87 min	~40% $\dot{V}O_{2max}$	15–17 h	OGTT	↔ I-AUC/↔ G-AUC post bout for MIE; ↓ I-AUC/↔ G-AUC post bout for HIE
Braun et al. (1995) [61]	Obese T2D untrained females (n = 8)	Treadmill	143 ± 23 min (3 bouts over 2 days)	~70% $\dot{V}O_{2max}$	~18 h	IST	↑ Insulin sensitivity post bout for both conditions; no difference between conditions
Perseghin et al. (1996) [62]	Young lean non-HPA non-excessively sedentary male (n = 6) and female (n = 2) offspring of T2D vs. young lean non-HPA non-excessively sedentary males (n = 3) and females (n = 5) with no T2D family history	Stair climbing	90 ± 8 min (3 bouts over 2 days)	74 ± 3% $\dot{V}O_{2max}$	48 h	HHC	↑ Insulin sensitivity post bout for both groups
Wojtaszewski et al. (2000) [63]	Healthy young males (n = 7)	One-legged knee-extensor exercise	60 min	75–95% $W_{max}$	4 h	EHC	↑ Insulin sensitivity post bout for exercised, but not rested leg
Chapman et al. (2002) [64]	Sedentary PM females (n = 10)	Machine full-body resistance exercise	3 sets x 10 reps or failure if sooner	50, 75 and 100% calculated 10-RM	14 h	IVGTT (minimal model)	↔ Insulin sensitivity post bout
Fenicchia et al. (2004) [65]	Healthy untrained females (n = 8) vs. T2D untrained females (n = 7)	Free weight and machine full-body resistance exercise	3 sets x 8–12 reps to failure	80% 3-RM	12–24 h	OGTT	↔ I-AUC/↓ G-AUC post bout for T2D; ↔ I-AUC/↔ G-AUC post bout for healthy
Koopman et al. (2005) [66]	Healthy untrained males (n = 12)	Machine leg resistance exercise	8 sets x 10 reps	75% 1-RM	24 h	ITT ( $\Delta BG_{t=4}$ and HOMA)	↑ Insulin sensitivity post bout
O'Gorman et al. (2006) [67]	Obese T2D males (n = 6) and females (n = 2) vs. obese NGT males (n = 6) and females (n = 1)	Cycle	60 min	~75% $\dot{V}O_{2peak}$	16 h	EHC	↔ Insulin sensitivity post single bout (acute effect); ↑ insulin sensitivity post final bout of 7 performed over 7 days (chronic effect)

(Continued)

Table 1 (Continued)

Study	Insulin sensitivity assessment			Exercise			Participants	Mode	Volume	Intensity	Time post-exercise	Method	Exercise effect
	Study	Time	Method	Exercise	Volume	Intensity							
Zhang <i>et al.</i> (2006) [68]	MS sedentary males (n = 10)	14–20h	Post fat-meal HOMA	Treadmill	60 min 60 min 60 min	~40% $\dot{V}O_{2max}$ ~60% $\dot{V}O_{2max}$ ~70% $\dot{V}O_{2max}$	14–20h	Post fat-meal HOMA	↔ Insulin sensitivity post bout/fat meal for MIE; ↑ insulin sensitivity post bout/fat meal for both HIE				
Howlett <i>et al.</i> (2007) [69]	RA males (n = 8)	0 h	EHC	Machine leg resistance exercise	3 sets x 10–12 reps	80% 1-RM	0 h	EHC	↓ Insulin sensitivity post single bout (acute effect); ↔ insulin sensitivity post final bout of 4 performed over 7 days (chronic effect)				
Schenk and Horowitz (2007) [70]	Healthy inactive females (n = 8)	15–18h	IVGTT (minimal model)	Treadmill + cycle	45 min + 45 min	66 ± 1% $\dot{V}O_{2peak}$	15–18h	IVGTT (minimal model)	↑ Insulin sensitivity (i.e. removal of lipid-induced impairment + enhancement vs. baseline) in response to overnight lipid infusion post bout				
Magkos <i>et al.</i> (2008) [71]	Healthy RA males (n = 30)	12h	HOMA	Cycle or treadmill	30, 60, 90 or 120 min	61–63 ± 2% $\dot{V}O_{2peak}$	12h	HOMA	↑ Insulin sensitivity post bout for <900 kcal expenditure				
Brestoff <i>et al.</i> (2009) [72]	Healthy RA young males (n = 8) and females (n = 5)	12–16h	OGTT (Matsuda) and HOMA	Cycle	3 x 15 min 5 x 30 s	~75% $\dot{V}O_{2peak}$ All-out	12–16h	OGTT (Matsuda) and HOMA	↑ Insulin sensitivity post bout for 75% $\dot{V}O_{2peak}$ ; ↔ insulin sensitivity post all-out				
Black <i>et al.</i> (2010) [73]	IFG sedentary males (n = 12) and females (n = 5)	24h	HOMA	Free weight and machine full-body resistance exercise	1 set or 4 sets x 12–15 reps/ 1 set or 4 sets x 6–8 reps/exercise	65% 1-RM 85% 1-RM	24h	HOMA	↑ Insulin sensitivity post bout for all protocols; 85% 1-RM > 65% 1-RM and 4 sets > 1 set				
Hasson <i>et al.</i> (2010) [74]	Healthy sedentary black males (n = 4) and females (n = 7) vs. healthy sedentary white males (n = 4) and females (n = 6)	12–15h	EHC	Treadmill	4 x 15 min	75% HR <sub>max</sub>	12–15h	EHC	↔ Insulin sensitivity post bout for both groups				
Manders <i>et al.</i> (2010) [75]	T2D sedentary males (n = 10)	24h	OGTT (OGIS) and HOMA	Cycle	60 min	35% W <sub>max</sub>	24h	OGTT (OGIS) and HOMA	↓ 24-h average glucose concentration post 35% W <sub>max</sub> ; ↔ 24-h average glucose concentration post 70% W <sub>max</sub>				
Newsom <i>et al.</i> (2010) [76]	Healthy sedentary males (n = 9)	24h	IVGTT (minimal model)	Treadmill + Cycle	45 min + 45 min	70% W <sub>max</sub> 60–65% $\dot{V}O_{2peak}$	24h	IVGTT (minimal model)	↔ Insulin sensitivity post bout when energy and carbohydrate balance were maintained and when carbohydrate balance was maintained with energy deficit; ↑ insulin sensitivity post bout when energy balance was maintained with carbohydrate deficit				
Richards <i>et al.</i> (2010) [77]	Healthy sedentary/RA young males (n = 2) and females (n = 7)	72h	EHC	Cycle	4 x 30 s	All-out	72h	EHC	↔ Insulin sensitivity post single bout (acute effect); ↑ insulin sensitivity post final bout of 6 performed over 14 days (chronic effect)				
Short <i>et al.</i> (2012) [78]	Healthy young RA/non-RA (n = 4/7) males (n = 4) and females (n = 7)	1 h, 17 h	IVGTT (Minimal Model) and MMTT (Matsuda) EHC	Treadmill + cycle + video-game boxing	15 min + 15 min	75% HR <sub>max</sub>	1 h, 17 h	IVGTT (Minimal Model) and MMTT (Matsuda) EHC	↑ Insulin sensitivity in response to meal post bout at 1 h, but not 17 h				
Newsom <i>et al.</i> (2013) [79]	Obese sedentary males (n = 3) and females (n = 8)	19h	OGTT (Matsuda) and HOMA	Treadmill + cycle	70 ± 3 min 54 ± 2 min	51% $\dot{V}O_{2peak}$ 66% $\dot{V}O_{2peak}$	19h	OGTT (Matsuda) and HOMA	↑ Insulin sensitivity post bout for MIE; ↔ insulin sensitivity post bout for HIE; no difference between conditions; Δ insulin sensitivity correlated with Δ fatty-acid disappearance from plasma				
Whyte <i>et al.</i> (2013) [80]	Overweight/obese sedentary males (n = 10)	18–22h	OGTT (Matsuda) and HOMA	Cycle	4 x 30 s 1 x 198 ± 10 s	All out All out	18–22h	OGTT (Matsuda) and HOMA	↑ Insulin sensitivity post bout when work was performed in one continuous bout; ↔ insulin sensitivity post bout when work was performed in 4 bouts separated by 270 s				
De Matos <i>et al.</i> (2014) [81]	Lean healthy untrained males (n = 3) and females (n = 6) vs. obese-insulin sensitive healthy untrained males (n = 3) and females (n = 6) vs. obese-insulin resistant untrained males (n = 3) and females (n = 6)	30 min	Biopsy (western blot for measurement of protein expression of cellular stressors related to insulin resistance; no direct measurement of insulin sensitivity)	Cycle	3 x 20 min	50% W <sub>peak</sub>	30 min	Biopsy (western blot for measurement of protein expression of cellular stressors related to insulin resistance; no direct measurement of insulin sensitivity)	↓ JNK post bout for obese-insulin resistant; ↔ JNK post bout for healthy lean and obese-insulin sensitive; ↓ p-IRS-1 <sup>ser612</sup> post-bout for obese-insulin resistant and obese-insulin sensitive; ↔ p-IRS-1 <sup>ser612</sup> for healthy lean; ↑HSP70 post-bout main effect				

(Continued)

Table 1 (Continued)

Study	Participants	Exercise			Insulin sensitivity assessment			Exercise effect
		Mode	Volume	Intensity	Time post-exercise	Method		
Nelson and Horowitz (2014) [48]	Overweight/obese male (n = 5) and female (n = 7) exercisers post 72h detraining vs. overweight/obese male (n = 5) and female (n = 7) non-exercisers	Treadmill	60 min	70% APMHR	24h	OGTT (Matsuda)	↑ Insulin sensitivity post bout for non-exercisers so that they were no longer different vs. exercisers who displayed superior insulin sensitivity pre-bout that was unchanged post bout	
Rynders et al. (2014) [82]	Pre-T2D sedentary males (n = 9) and females (n = 9)	Cycle	40 ± 9 min 24 ± 5 min 4 × 30 s 60 min	-50% $\dot{V}O_{2peak}$ -83% $\dot{V}O_{2peak}$ All out	60 min	OGTT (Minimal Model)	↑ Insulin sensitivity post bout that was not different between conditions	
Ortega et al. (2015) [83]	Healthy, young RA males (n = 10)	Cycle	35 ± 2 min 42 ± 2 min 24 ± 1 min	46 ± 3% $\dot{V}O_{2peak}$ 77 ± 5% $\dot{V}O_{2peak}$ 68 ± 1% $\dot{V}O_{2peak}$ 90 ± 1% $\dot{V}O_{2peak}$	30 min, 24h, 48h	IVGTT	↑ Insulin sensitivity 30 min, 24h and 48h post bout for all conditions; all out ↑ > continuous ↑ at 30 min	
Malin et al. (2016) [84]	Pre-T2D middle-aged sedentary males (n = 7) and females (n = 8)	Treadmill	24 ± 1 min	68 ± 1% $\dot{V}O_{2peak}$ 90 ± 1% $\dot{V}O_{2peak}$	60 min	OGTT (Minimal Model)	↓ Insulin resistance-SM post bout that was not different between conditions; ↔ insulin resistance-HEP and insulin resistance-AT post bout for MIE; ↑ insulin resistance-HEP and insulin resistance-AT post bout for both conditions	
Metcalf et al. (2016) [85]	Healthy sedentary/RA young males (n = 8) and females (n = 6)	Cycle	3 × 15 min 5 × 30 s	76 ± 4% $\dot{V}O_{2max}$ All out	14–16 h	OGTT (Cederholm)	↔ Insulin sensitivity post bout for both conditions	
Castleberry et al. (2019) [86]	Healthy inactive males (n = 10)	Treadmill	1 × 60 min 3 × 60 min (consecutive days) 3 × 60 min (alternate days)	78 ± 6% $\dot{V}O_{2max}$ 78 ± 6% $\dot{V}O_{2max}$ 77 ± 5% $\dot{V}O_{2max}$	12–14h	OGTT (Matsuda)	↔ Insulin sensitivity post bout for all conditions	

AA, athletically active; APMHR, age-predicted maximum heart rate; bpm, beats per minute; Cederholm index; Δ, change in variable;  $\Delta BG_{6-4}$ , change in blood-glucose concentration from 4 to 16 min; EHC, euglycemic hyperinsulinemic clamp; G-AUC, glucose area under the curve; HHC, hyperglycemic hyperinsulinemic clamp; HIE, high-intensity exercise; HOMA, homeostatic model assessment; HPA, participates in heavy physical activity; HR, heart rate;  $HR_{max}$ , maximum heart rate; HSP70, heat shock protein 70; I-AUC, insulin area under the curve; IFG, impaired fasting glucose; IGT, impaired glucose tolerance; Insulin resistance-AT, adipose tissue insulin resistance; Insulin resistance-HEP, hepatic insulin resistance; Insulin resistance-SM, skeletal muscle insulin resistance; IT, insulin suppression test; ITT, insulin tolerance test; IVGTT, intravenous glucose tolerance test; JNK, C-jun N-terminal kinase phosphorylation; Matsuda, Matsuda index; MIE, moderate-intensity exercise; MMTT, mixed meal tolerance test; MS, with metabolic syndrome; NG, normoglycemic; NGT, normal glucose tolerance; OGIS, oral glucose insulin sensitivity index; OGTT, oral glucose tolerance test; p-IRS-1<sup>ser612</sup>, insulin receptor substrate 1 serine 612 phosphorylation; PM, postmenopausal; Pre-T2D, with prediabetes; RA, recreationally active; RM, repetition maximum; T2D, with type 2 diabetes mellitus;  $\dot{V}O_{2peak}$ , peak rate of oxygen uptake;  $\dot{V}O_{2max}$ , maximum rate of oxygen uptake;  $W_{max}$ , maximum work rate.



on insulin sensitivity following a bout of this type of exercise [93]. More research is required to further clarify the differences between resistance training and conventional endurance exercise for eliciting the PAE of exercise on insulin sensitivity and, with respect to the latter, whether specific modalities are more effective (e.g. more muscles involved vs. leg cycling, body weight supported vs. not supported, etc.).

### Frequency

To determine the frequency at which bouts must be 'strung together' to elicit a 'continuous' PAE of exercise on insulin sensitivity, the time course for dissipation of the effect must be considered. Studies that have assessed the PAE of exercise on insulin sensitivity at multiple time points provide insight. Oshida *et al.* observed elevated insulin sensitivity (EHC) in young trained athletes compared to sedentary controls at 14, but not 38 h after cessation of training [94]. King *et al.* had middle-aged moderately-trained individuals perform five consecutive days of exercise (45 min per day at  $\sim 75\% \dot{V}O_{2peak}$ ) and observed increased insulin sensitivity (OGTT) at 24 and 72, but not 90 h following the final bout [32]. In addition to age, another possible explanation for these contrasting findings is that compared to OGTT, the higher insulin stimulus during EHC might have resulted in complete suppression of hepatic glucose production thereby rendering the test incapable of detecting an increase in hepatic insulin sensitivity that might have a longer time course compared to skeletal muscle [32]. However, 72-h duration cannot be confirmed or refuted by most studies that have used EHC (the 'gold standard' means of assessment) to determine the PAE of exercise on insulin sensitivity because they have only involved a single measurement; for example, at  $\leq 24$  h [57,79] or 48 h [54,62]. One exception is Richards *et al.* who found unchanged insulin sensitivity in sedentary/recreationally-active individuals 72 h after the first bout of a 14-day protocol involving sprint interval training (SIT), a form of high-intensity interval training (HIIT; see below) [77]. This agrees with findings by Whyte *et al.* that increased insulin sensitivity (OGTT) following a 2-week SIT intervention was no longer present 72 h following the final bout [95]. The reason that these reports contrast those by King *et al.* might be related to volume of exercise (4–6 30-s all-out cycling sprints compared to five 45-min bouts over multiple days) [32,77,95]. In this regard, Magkos *et al.* observed a curvilinear relationship between energy expenditure during evening exercise (30–120 min on a treadmill or cycle ergometer at  $60\% \dot{V}O_{2max}$ ) and increased insulin sensitivity (HOMA) the following morning indicating that outlay  $\geq 900$  kcal was required to elicit the effect for healthy, nonobese untrained men [71]. However, this blanket statement should be interpreted with caution because in addition to energy use per se, energy use per unit time (i.e. intensity) is also important to consider (see below) [60,83; *c.f.* 61]. Finally, post bout nutrition influences the

duration of the PAE of exercise on insulin sensitivity as carbohydrate ingestion has the potential to decrease the time course from 48 to 18 h in rats [40] and eliminate it completely in humans [76]. More research involving measurement of the PAE of exercise on insulin sensitivity at multiple time points is required to clarify how long the effect lasts (and, by extension, how frequently bouts should be performed) in different types of participants (e.g. healthy, but at risk for a loss of insulin sensitivity with aging vs. at risk of developing or already possessing insulin resistance vs. already possessing T2D) with different nutritional intake pre- and immediately post-bout.

### Duration

Pharmacological evaluation includes determining a drug's minimum effective dose (MED) to inform prescription in an effective/well tolerated manner. For exercise to be considered medicine [7,8], similar determinations are required. With respect to prescribing exercise to elicit the PAE of exercise on insulin sensitivity, in addition to frequency, duration of the exercise bout is important to consider. However, like frequency, duration likely depends on intensity so establishing a single optimal frequency/duration combination is attractive in theory, but flawed in practice. Indeed, unlike prescribing pharmaceuticals, manipulating the 'exercise dose' is far more complicated because there are more variables to consider. Moreover, not all 'side effects' associated with exceeding MED in the exercise setting are deleterious. For example, while excessive volume (i.e. frequency  $\times$  duration) might increase the likelihood of overuse injury and lack of program adherence, the MED required to achieve/maintain the PAE of exercise on insulin sensitivity might be less than what is necessary to establish an energy deficit to lose body fat at a sufficient rate for those who are also exercising for this purpose. This is important given the association between insulin resistance and the overweight/obese condition. However, for individuals in the MONW subgroup [96], exceeding the MED for the PAE of exercise on insulin sensitivity for this purpose would be unnecessary.

As previously mentioned, Magkos *et al.* suggest that the PAE of exercise on insulin sensitivity increased proportionally with exercise-bout energy expenditure as long as a 900-kcal threshold was surpassed, which meant that  $\geq 60$ –90 min of sustained exercise was required for healthy nonobese untrained men [71]. However, even though participants were normoglycemic, the PAE of exercise on insulin sensitivity was inversely related to baseline insulin sensitivity, which raises the possibility that less might be needed for individuals with insulin resistance. Moreover, bouts were performed at a moderate intensity ( $60\% \dot{V}O_{2peak}$ ) in that study [71]. In this regard, Ben Ezra *et al.* found that despite equal energy expenditure ( $\sim 400$  kcal), treadmill walking at  $70\% \dot{V}O_{2max}$  for 50 min resulted in a decreased insulin response (OGTT) 15–17 h post-exercise for untrained normoglycemic

women whereas walking at 40% for 87 min did not [60]. Conversely, Ortega *et al.* found that young healthy non-obese untrained men experienced improved insulin sensitivity 48 h following continuous leg cycling that required ~570 kcal regardless of whether it was performed at ~46%  $\dot{V}O_{2\text{peak}}$  for 60 min or ~77%  $\dot{V}O_{2\text{peak}}$  for ~35 min [83]. Moreover, despite requiring ~25% of the energy outlay of the two continuous bouts, similar improvement was derived from a bout comprising four 30-s all-out sprints (SIT) [83]. While such time efficiency of HIIT/SIT has been highly touted [97], it is important to recognise that these sprints were separated by 270 s of unloaded cycling; hence, the entire bout excluding warm-up and cool-down still required >15 min to perform [83]. Interestingly, Whyte *et al.* found that a similar SIT protocol resulted in no improvement in insulin sensitivity (OGTT) for inactive overweight/obese young men 18–22 h post-exercise whereas the same amount of work performed during a single extended sprint (i.e. ‘all-out’ cycling for ~200 s) resulted in marked improvement [80]. The authors advance this as evidence that including warm-up and cool-down, as little as 10–12 min of exercise is required to derive the PAE of exercise on insulin sensitivity [80]. More research involving measurement following exercise bouts with different duration/intensity combinations is required to clarify what is required to elicit the PAE of exercise on insulin sensitivity for different types of participants (e.g. healthy, at risk for disease or already with disease).

### Intensity

As indicated by mention in previous sections, in all likelihood, intensity is the variable that provides the foundation upon which prescription of the PAE of exercise on insulin sensitivity should be built. With respect to continuous exercise at a sustainable work rate, Braun *et al.* found that energy-matched treadmill exercise at ~50 and ~75%  $\dot{V}O_{2\text{max}}$  increased insulin sensitivity (insulin suppression test; IST) to a similar extent for obese women with T2D [61]. Rynders *et al.* also report similar enhancement of insulin sensitivity (OGTT) 3 h following energy-matched cycling at ~50 and ~83%  $\dot{V}O_{2\text{peak}}$  for pre-diabetic individuals [82]. These findings, which suggest that energy expenditure per se is the key determinant, contrast the observation that normoglycemic untrained women achieved the PAE of exercise on insulin sensitivity (OGTT) from exercise at ~70%, but not energy-matched exercise at ~40%  $\dot{V}O_{2\text{max}}$  (see above) [60]. Furthermore, Newsom *et al.* observed ~35% improvement in insulin sensitivity (EHC) for sedentary obese adults ~19 h after participants expended 350 kcal at ~50%  $\dot{V}O_{2\text{max}}$  whereas similar expenditure at ~65% elicited a nonsignificant ~20% increase [79]. On the surface, equal if not greater enhancement at energy-matched lower-intensity exercise is counterintuitive being that, presumably, the PAE of exercise on insulin sensitivity is at least in part determined by muscle-glycogen depletion [41].

However, Newsom *et al.* also report a correlation between the PAE of exercise on insulin sensitivity and the change in fatty acid rate of disappearance from plasma, which was reduced following the ~50%, but not ~65% protocol [79]. Given the link between skeletal-muscle fatty acid uptake and the obesity-related decline in insulin action [98], this raises the possibility that exercise of lower intensity can also play a role in establishing the PAE of exercise on insulin sensitivity via its effect on lipid metabolism even though it does not substantially influence muscle-glycogen stores [79].

More information regarding intensity and the PAE of exercise on insulin sensitivity comes from research that has ‘stretched’ the intensity extreme by having subjects perform intermittent exercise involving unsustainable work rates (HIIT) or exercise with all-out effort (SIT) [99]. As previously mentioned, Ortega *et al.* found similar enhancement of insulin sensitivity (IVGTT) for young healthy nonobese untrained men 48 h following exercise regardless of whether it involved four 30-second cycling sprints or continuous cycling at moderate or high intensity that required four times the energy expenditure of the sprint bouts [83]. While this finding suggests intensity is important to consider, the fact that PAE of exercise on insulin sensitivity induced by the energy-matched continuous bouts in that study did not differ indicates that energy expenditure per se is a key determinant at least during sustainable exercise. Conversely, Brestoff *et al.* found that SIT did not elicit a similar effect as continuous cycling in recreationally-active individuals who experienced no change in insulin sensitivity (OGTT) 12–16 h after completing five 30-s sprints separated by 240–300 s whereas a 70–100% increase was present following 45 min of cycling at ~75%  $\dot{V}O_{2\text{max}}$  [72]. Importantly, energy expenditure was not matched for the two conditions; hence, the lesser outlay during SIT might have played a role [72; *cf.*, 83]. Using the same methodology, Metcalfe *et al.* found that ‘reduced-exertion’ SIT (two 20-s sprints separated by 200 s) also did not induce PAE on insulin sensitivity; however, in that study, the 45-min bout also elicited no effect [85]. Baseline insulin sensitivity and post-exercise feeding are two factors that might explain this unexpected finding [85]. Finally, while Richards *et al.* observed increased insulin sensitivity following 14 days of SIT for sedentary/recreationally-active individuals, there was no acute effect after the initial bout of the protocol (four 30-s sprints separated by 240 s) [77]. However, as previously mentioned, in that study, the PAE of exercise on insulin sensitivity was only assessed (EHC) 72 h after completion of the initial bout [77], which means that the degree to which the effect might have been present earlier cannot be determined. However, in this regard, Whyte *et al.* found no increase in insulin sensitivity (OGTT) 24 h following SIT (four 30-s sprints separated by 270 s) although as previously mentioned, insulin sensitivity

was increased when recovery intervals were removed from an all-out work-matched bout [80]. More research is required to determine whether it is energy use per se or energy use per unit time (i.e. intensity) which is the primary factor that drives the PAE of exercise on insulin sensitivity and whether low-intensity sustained exercise that maximizes fat use, high-intensity sustained exercise that maximizes energy use (e.g. at the 'critical power/velocity') or very-high-intensity unsustainable intermittent exercise that provides for maximal mitochondrial stimulation and glycogen depletion provides for the optimal PAE of exercise on insulin sensitivity.

### Conclusion

Exercise is medicine implies that exercise can serve as a first-line defense for prevention/treatment of disease with pharmaceuticals representing a surrogate for exercise only when necessary. Regular exercise results in chronic adaptations that help serve this purpose; however, while recognized for many years, there is growing appreciation for beneficial residual effects of each individual exercise bout on myriad aspects of function [100,101]. With respect to insulin sensitivity, the PAE of exercise appears to be the predominant effect induced by training even for well-trained individuals with high insulin sensitivity [37]. Furthermore, untrained individuals [47] and individuals with lower insulin sensitivity [57,71] benefit more from the PAE of exercise whereas the chronic-exercise effect on insulin sensitivity for individuals with 'metabolic syndrome' might depend on weight loss that accompanies exercise as opposed to exercise per se [102,103]. This means that for at-risk individuals for whom it is not necessary or feasible to lose weight (MONW, for example), the PAE of exercise on insulin sensitivity might be the only option available. Collectively, these observations suggest that an intervention designed to combat the demise in insulin sensitivity that begins the pathological journey to cardiometabolic diseases including T2D should be prescribed with precision to elicit a PAE of exercise on insulin sensitivity that can serve as a surrogate for pharmaceutical intervention or, in cases where such intervention is necessary, an adjunct to it [104]. While prescribing exercise is far more complicated than prescribing pharmaceuticals due to the many variables that can be manipulated (e.g. mode, volume, frequency, duration and, perhaps most importantly, intensity), beginning with the seminal work of the Holloszy group in the 1980s, there have been a number of studies that have provided information regarding the PAE of exercise on insulin sensitivity that are important to consider for determining the specifics of prescribing exercise to elicit/maintain this effect. Table 1 provides a chronically-ordered listing of some of the major ones involving human subjects. However, more standardised research with gold-standard testing (e.g. EHC, as opposed to assessments used in many of the studies included in this review; e.g. OGTT, ITT, IST and HOMA) would be

helpful to further determine dose-response specifics (e.g. session frequency and duration, importance of energy use per se vs. intensity of effort and degree to which lower- and higher-intensity efforts elicit the PAE of exercise on insulin sensitivity via different mechanisms) for individuals with varying degrees of insulin sensitivity/insulin resistance or T2D. Differentiation between the minimal effective and optimal dose is also important to clarify because many of the individuals that can benefit most from the PAE of exercise on insulin sensitivity might be unable to achieve/sustain the latter. Finally, the influence of post-exercise feeding requires further investigation because the PAE of exercise on insulin sensitivity can be eliminated via carbohydrate replenishment despite negative energy balance or maintained despite energy replenishment with carbohydrate restriction [76].

### Acknowledgements

#### Conflicts of interest

There are no conflicts of interest.

### References

- Centers for Disease Control and Prevention. *National Diabetes Statistics Report*. Atlanta, GA: Centers for Disease Control and Prevention, US Department of Health and Human Services; 2020.
- Centers for Disease Control and Prevention. National Diabetes Fact Sheet, 2011. 2011. [http://www.cdc.gov/diabetes/pubs/pdf/ndfs\\_2011.pdf](http://www.cdc.gov/diabetes/pubs/pdf/ndfs_2011.pdf). [Accessed October 2013]. p. 1.
- Boyle JP, Thompson TJ, Gregg EW, Barker LE, Williamson DF. Projection of the year 2050 burden of diabetes in the US adult population: dynamic modeling of incidence, mortality, and prediabetes prevalence. *Popul Health Metr* 2010; **8**:29.
- Field AE, Coakley EH, Must A, Spadano JL, Laird N, Dietz WH, *et al*. Impact of overweight on the risk of developing common chronic diseases during a 10-year period. *Arch Intern Med* 2001; **161**:1581–1586.
- Huang LO, Loos RJF, Kilpeläinen TO. Evidence of genetic predisposition for metabolically healthy obesity and metabolically obese normal weight. *Physiol Genomics* 2018; **50**:169–178.
- Berryman JW. Exercise is medicine: a historical perspective. *Curr Sports Med Rep* 2010; **9**:195–201.
- DiMenna FJ, Arad AD. Exercise as 'precision medicine' for insulin resistance and its progression to type 2 diabetes: a research review. *BMC Sports Sci Med Rehabil* 2018; **10**:21.
- DeFronzo RA, Jacot E, Jequier E, Maeder E, Wahren J, Felber JP. The effect of insulin on the disposal of intravenous glucose. Results from indirect calorimetry and hepatic and femoral venous catheterization. *Diabetes* 1981; **30**:1000–1007.
- Giugliano G, Nicoletti G, Grella E, Giugliano F, Esposito K, Scuderi N, D'Andrea F. Effect of liposuction on insulin resistance and vascular inflammatory markers in obese women. *Br J Plast Surg* 2004; **57**:190–194.
- D'Andrea F, Grella R, Rizzo MR, Grella E, Grella R, Nicoletti G, *et al*. Changing the metabolic profile by large-volume liposuction: a clinical study conducted with 123 obese women. *Aesthetic Plast Surg* 2005; **29**:472–478; discussion 479.
- Brochu M, Malita MF, Messier V, Doucet E, Strychar I, Lavoie JM, *et al*. Resistance training does not contribute to improving the metabolic profile after a 6-month weight loss program in overweight and obese postmenopausal women. *J Clin Endocrinol Metab* 2009; **94**:3226–3233.
- Schenk S, Harber MP, Shrivastava CR, Burant CF, Horowitz JF. Improved insulin sensitivity after weight loss and exercise training is mediated by a reduction in plasma fatty acid mobilization, not enhanced oxidative capacity. *J Physiol* 2009; **587**:4949–4961.
- Bouchonville M, Armamento-Villareal R, Shah K, Napoli N, Sinacore DR, Qualls C, Villareal DT. Weight loss, exercise or both and cardiometabolic risk factors in obese older adults: results of a randomized controlled trial. *Int J Obes (Lond)* 2014; **38**:423–431.
- Ross R. Does exercise without weight loss improve insulin sensitivity? *Diabetes Care* 2003; **26**:944–945.



- 15 Arad AD, DiMenna FJ, Thomas N, Tamis-Holland J, Weil R, Geliebter A, Albu JB. High-intensity interval training without weight loss improves exercise but not basal or insulin-induced metabolism in overweight/obese African American women. *J Appl Physiol* (1985) 2015; **119**:352–362.
- 16 Segal KR, Edano A, Abalos A, Albu J, Blando L, Tomas MB, Pi-Sunyer FX. Effect of exercise training on insulin sensitivity and glucose metabolism in lean, obese, and diabetic men. *J Appl Physiol* (1985) 1991; **71**:2402–2411.
- 17 Dengel DR, Pratley RE, Hagberg JM, Rogus EM, Goldberg AP. Distinct effects of aerobic exercise training and weight loss on glucose homeostasis in obese sedentary men. *J Appl Physiol* (1985) 1996; **81**:318–325.
- 18 Ross R, Dagnone D, Jones PJ, Smith H, Paddags A, Hudson R, Janssen I. Reduction in obesity and related comorbid conditions after diet-induced weight loss or exercise-induced weight loss in men. A randomized, controlled trial. *Ann Intern Med* 2000; **133**:92–103.
- 19 Lettner A, Roden M. Ectopic fat and insulin resistance. *Curr Diab Rep* 2008; **8**:185–191.
- 20 Shimabukuro M, Kozuka C, Taira S, Yabiku K, Dagvasumberel M, Ishida M, et al. Ectopic fat deposition and global cardiometabolic risk: new paradigm in cardiovascular medicine. *J Med Invest* 2013; **60**:1–14.
- 21 Richter EA, Hargreaves M. Exercise, GLUT4, and skeletal muscle glucose uptake. *Physiol Rev* 2013; **93**:993–1017.
- 22 James DE, Kraegen EV, Chisholm DJ. Muscle glucose metabolism in exercising rats: comparison with insulin stimulation. *Am J Physiol* 1985; **248**:E575–E580.
- 23 James DE, Brown R, Navarro J, Pilch PF. Insulin-regulatable tissues express a unique insulin-sensitive glucose transport protein. *Nature* 1988; **333**:183–185.
- 24 Douen AG, Ramlal T, Rastogi S, Bilan PJ, Cartee GD, Vranic M, et al. Exercise induces recruitment of the “insulin-responsive glucose transporter”. Evidence for distinct intracellular insulin- and exercise-recruitable transporter pools in skeletal muscle. *J Biol Chem* 1990; **265**:13427–13430.
- 25 Neshler R, Karl IE, Kipnis DM. Dissociation of effects of insulin and contraction on glucose transport in rat epitrochlearis muscle. *Am J Physiol* 1985; **249**:C226–C232.
- 26 O’Neill HM. AMPK and exercise: glucose uptake and insulin sensitivity. *Diabetes Metab J* 2013; **37**:1–21.
- 27 Yue Y, Zhang C, Zhang X, Zhang S, Liu Q, Hu F, et al. An AMPK/Axin1-Rac1 signaling pathway mediates contraction-regulated glucose uptake in skeletal muscle cells. *Am J Physiol Endocrinol Metab* 2020; **318**:E330–E342.
- 28 Merry TL, McConell GK. Skeletal muscle glucose uptake during exercise: a focus on reactive oxygen species and nitric oxide signaling. *IUBMB Life* 2009; **61**:479–484.
- 29 Cartee GD. Roles of TBC1D1 and TBC1D4 in insulin- and exercise-stimulated glucose transport of skeletal muscle. *Diabetologia* 2015; **58**:19–30.
- 30 Kjøbsted R, Chadt A, Jørgensen NO, Kido K, Larsen JK, de Wendt C, et al. TBC1D4 is necessary for enhancing muscle insulin sensitivity in response to AICAR and contraction. *Diabetes* 2019; **68**:1756–1766.
- 31 Goodyear LJ, Hirshman MF, King PA, Horton ED, Thompson CM, Horton ES. Skeletal muscle plasma membrane glucose transport and glucose transporters after exercise. *J Appl Physiol* (1985) 1990; **68**:193–198.
- 32 King DS, Baldus PJ, Sharp RL, Kesl LD, Feltmeyer TL, Riddle MS. Time course for exercise-induced alterations in insulin action and glucose tolerance in middle-aged people. *J Appl Physiol* (1985) 1995; **78**:17–22.
- 33 Funai K, Schweitzer GG, Sharma N, Kanzaki M, Cartee GD. Increased AS160 phosphorylation, but not TBC1D1 phosphorylation, with increased postexercise insulin sensitivity in rat skeletal muscle. *Am J Physiol Endocrinol Metab* 2009; **297**:E242–E251.
- 34 Prior SJ, Goldberg AP, Ortmeyer HK, Chin ER, Chen D, Blumenthal JB, Ryan AS. Increased skeletal muscle capillarization independently enhances insulin sensitivity in older adults after exercise training and detraining. *Diabetes* 2015; **64**:3386–3395.
- 35 Williamson JR, Hoffmann PL, Kohrt WM, Spina RJ, Coggan AR, Holloszy O. Endurance exercise training decreases capillary basement membrane width in older nondiabetic and diabetic adults. *J Appl Physiol* (1985) 1996; **80**:747–753.
- 36 Kohrt WM, Kirwan JP, Staten MA, Bourey RE, King DS, Holloszy JO. Insulin resistance in aging is related to abdominal obesity. *Diabetes* 1993; **42**:273–281.
- 37 Heath GW, Gavin JR 3rd, Hinderliter JM, Hagberg JM, Bloomfield SA, Holloszy JO. Effects of exercise and lack of exercise on glucose tolerance and insulin sensitivity. *J Appl Physiol Respir Environ Exerc Physiol* 1983; **55**:512–517.
- 38 Richter EA, Garetto LP, Goodman MN, Ruderman NB. Muscle glucose metabolism following exercise in the rat: increased sensitivity to insulin. *J Clin Invest* 1982; **69**:785–793.
- 39 Ivy JL, Young JC, McLane JA, Fell RD, Holloszy JO. Exercise training and glucose uptake by skeletal muscle in rats. *J Appl Physiol Respir Environ Exerc Physiol* 1983; **55**:1393–1396.
- 40 Cartee GD, Young DA, Sleeper MD, Zierath J, Wallberg-Henriksson H, Holloszy JO. Prolonged increase in insulin-stimulated glucose transport in muscle after exercise. *Am J Physiol* 1989; **256**:E494–E499.
- 41 Bogardus C, Thuillez P, Ravussin E, Vasquez B, Narimiga M, Azhar S. Effect of muscle glycogen depletion on *in vivo* insulin action in man. *J Clin Invest* 1983; **72**:1605–1610.
- 42 Rogers MA, King DS, Hagberg JM, Ehsani AA, Holloszy JO. Effect of 10 days of physical inactivity on glucose tolerance in master athletes. *J Appl Physiol* (1985) 1990; **68**:1833–1837.
- 43 Rogers MA, Yamamoto C, King DS, Hagberg JM, Ehsani AA, Holloszy JO. Improvement in glucose tolerance after 1 wk of exercise in patients with mild NIDDM. *Diabetes Care* 1988; **11**:613–618.
- 44 King DS, Dalsky GP, Staten MA, Clutter WE, Van Houten DR, Holloszy JO. Insulin action and secretion in endurance-trained and untrained humans. *J Appl Physiol* (1985) 1987; **63**:2247–2252.
- 45 King DS, Dalsky GP, Clutter WE, Young DA, Staten MA, Cryer PE, Holloszy JO. Effects of lack of exercise on insulin secretion and action in trained subjects. *Am J Physiol* 1988; **254**:E537–E542.
- 46 Helmrich SP, Ragland DR, Leung RW, Paffenbarger RS Jr. Physical activity and reduced occurrence of non-insulin-dependent diabetes mellitus. *N Engl J Med* 1991; **325**:147–152.
- 47 Steenberg DE, Jørgensen NB, Birk JB, Sjøberg KA, Kiens B, Richter EA, Wojtaszewski JFP. Exercise training reduces the insulin-sensitizing effect of a single bout of exercise in human skeletal muscle. *J Physiol* 2019; **597**:89–103.
- 48 Nelson RK, Horowitz JF. Acute exercise ameliorates differences in insulin resistance between physically active and sedentary overweight adults. *Appl Physiol Nutr Metab* 2014; **39**:811–818.
- 49 Cartee GD. Once is enough for acute exercise benefits on insulin sensitivity. *J Physiol* 2019; **597**:7–8.
- 50 Caspersen CJ, Powell KE, Christenson GM. Physical activity, exercise, and physical fitness: definitions and distinctions for health-related research. *Public Health Rep* 1985; **100**:126–131.
- 51 Sylvester BD, Standage M, Ark TK, Sweet SN, Crocker PR, Zumbo BD, Beauchamp MR. Is variety a spice of (an active) life?: perceived variety, exercise behavior, and the mediating role of autonomous motivation. *J Sport Exerc Psychol* 2014; **36**:516–527.
- 52 Devlin JT, Horton ES. Effects of prior high-intensity exercise on glucose metabolism in normal and insulin-resistant men. *Diabetes* 1985; **34**:973–979.
- 53 Devlin JT, Hirshman M, Horton ED, Horton ES. Enhanced peripheral and splanchnic insulin sensitivity in NIDDM men after single bout of exercise. *Diabetes* 1987; **36**:434–439.
- 54 Mikines KJ, Sonne B, Farrell PA, Tronier B, Galbo H. Effect of physical exercise on sensitivity and responsiveness to insulin in humans. *Am J Physiol* 1988; **254**:E248–E259.
- 55 Richter EA, Mikines KJ, Galbo H, Kiens B. Effect of exercise on insulin action in human skeletal muscle. *J Appl Physiol* (1985) 1989; **66**:876–885.
- 56 Young JC, Enslin J, Kuca B. Exercise intensity and glucose tolerance in trained and nontrained subjects. *J Appl Physiol* (1985) 1989; **67**:39–43.
- 57 Burstein R, Epstein Y, Shapiro Y, Charuzi I, Karnieli E. Effect of an acute bout of exercise on glucose disposal in human obesity. *J Appl Physiol* (1985) 1990; **69**:299–304.
- 58 Kjaer M, Hollenbeck CB, Frey-Hewitt B, Galbo H, Haskell W, Reaven GM. Glucoregulation and hormonal responses to maximal exercise in non-insulin-dependent diabetes. *J Appl Physiol* (1985) 1990; **68**:2067–2074.
- 59 Fluckey JD, Hickey MS, Brambrink JK, Hart KK, Alexander K, Craig BW. Effects of resistance exercise on glucose tolerance in normal and glucose-intolerant subjects. *J Appl Physiol* (1985) 1994; **77**:1087–1092.
- 60 Ben-Ezra V, Jankowski C, Kendrick K, Nichols D. Effect of intensity and energy expenditure on postexercise insulin responses in women. *J Appl Physiol* (1985) 1995; **79**:2029–2034.
- 61 Braun B, Zimmermann MB, Kretschmer N. Effects of exercise intensity on insulin sensitivity in women with non-insulin-dependent diabetes mellitus. *J Appl Physiol* (1985) 1995; **78**:300–306.
- 62 Perseghin G, Price TB, Petersen KF, Roden M, Cline GW, Gerow K, et al. Increased glucose transport-phosphorylation and muscle glycogen synthesis after exercise training in insulin-resistant subjects. *N Engl J Med* 1996; **335**:1357–1362.
- 63 Wojtaszewski JF, Hansen BF, Gade, Kiens B, Markuns JF, Goodyear LJ, Richter EA. Insulin signaling and insulin sensitivity after exercise in human skeletal muscle. *Diabetes* 2000; **49**:325–331.
- 64 Chapman J, Garvin AW, Ward A, Cartee GD. Unaltered insulin sensitivity after resistance exercise bout by postmenopausal women. *Med Sci Sports Exerc* 2002; **34**:936–941.

- 65 Fenicchia LM, Kanaley JA, Azevedo JL Jr, Miller CS, Weinstock RS, Carhart RL, Ploutz-Snyder LL. Influence of resistance exercise training on glucose control in women with type 2 diabetes. *Metabolism* 2004; **53**:284–289.
- 66 Koopman R, Manders RJ, Zorenc AH, Hul GB, Kuipers H, Keizer HA, van Loon LJ. A single session of resistance exercise enhances insulin sensitivity for at least 24 h in healthy men. *Eur J Appl Physiol* 2005; **94**:180–187.
- 67 O'Gorman DJ, Karlsson HK, McQuaid S, Yousif O, Rahman Y, Gasparro D, et al. Exercise training increases insulin-stimulated glucose disposal and GLUT4 (SLC2A4) protein content in patients with type 2 diabetes. *Diabetologia* 2006; **49**:2983–2992.
- 68 Zhang JQ, Ji LL, Fretwell VS, Nunez G. Effect of exercise on postprandial lipemia in men with hypertriglyceridemia. *Eur J Appl Physiol* 2006; **98**:575–582.
- 69 Howlett KF, Sakamoto K, Garnham A, Cameron-Smith D, Hargreaves M. Resistance exercise and insulin regulate AS160 and interaction with 14-3-3 in human skeletal muscle. *Diabetes* 2007; **56**:1608–1614.
- 70 Schenk S, Horowitz JF. Acute exercise increases triglyceride synthesis in skeletal muscle and prevents fatty acid-induced insulin resistance. *J Clin Invest* 2007; **117**:1690–1698.
- 71 Magkos F, Tsekouras Y, Kavouras SA, Mittendorfer B, Sidossis LS. Improved insulin sensitivity after a single bout of exercise is curvilinearly related to exercise energy expenditure. *Clin Sci (Lond)* 2008; **114**:59–64.
- 72 Brestoff JR, Clippinger B, Spinella T, von Duvillard SP, Nindl B, Arciero PJ. An acute bout of endurance exercise but not sprint interval exercise enhances insulin sensitivity. *Appl Physiol Nutr Metab* 2009; **34**:25–32.
- 73 Black LE, Swan PD, Alvar BA. Effects of intensity and volume on insulin sensitivity during acute bouts of resistance training. *J Strength Cond Res* 2010; **24**:1109–1116.
- 74 Hasson RE, Granados K, Chipkin S, Freedson PS, Braun B. Effects of a single exercise bout on insulin sensitivity in black and white individuals. *J Clin Endocrinol Metab* 2010; **95**:E219–E223.
- 75 Manders RJ, Van Dijk JW, van Loon LJ. Low-intensity exercise reduces the prevalence of hyperglycemia in type 2 diabetes. *Med Sci Sports Exerc* 2010; **42**:219–225.
- 76 Newsom SA, Schenk S, Thomas KM, Harber MP, Knuth ND, Goldenberg N, Horowitz JF. Energy deficit after exercise augments lipid mobilization but does not contribute to the exercise-induced increase in insulin sensitivity. *J Appl Physiol (1985)* 2010; **108**:554–560.
- 77 Richards JC, Johnson TK, Kuzma JN, Lonac MC, Schweder MM, Voyles WF, Bell C. Short-term sprint interval training increases insulin sensitivity in healthy adults but does not affect the thermogenic response to beta-adrenergic stimulation. *J Physiol* 2010; **588**:2961–2972.
- 78 Short KR, Pratt LV, Teague AM. The acute and residual effect of a single exercise session on meal glucose tolerance in sedentary young adults. *J Nutr Metab* 2012; **2012**:278678.
- 79 Newsom SA, Everett AC, Hinko A, Horowitz JF. A single session of low-intensity exercise is sufficient to enhance insulin sensitivity into the next day in obese adults. *Diabetes Care* 2013; **36**:2516–2522.
- 80 Whyte LJ, Ferguson C, Wilson J, Scott RA, Gill JM. Effects of single bout of very high-intensity exercise on metabolic health biomarkers in overweight/obese sedentary men. *Metabolism* 2013; **62**:212–219.
- 81 de Matos MA, Ottono Vde O, Duarte TC, Sampaio PF, Costa KB, Fonseca CA, et al. Exercise reduces cellular stress related to skeletal muscle insulin resistance. *Cell Stress Chaperones* 2014; **19**:263–270.
- 82 Rynders CA, Weltman JY, Jiang B, Breton M, Patrie J, Barrett EJ, Weltman A. Effects of exercise intensity on postprandial improvement in glucose disposal and insulin sensitivity in prediabetic adults. *J Clin Endocrinol Metab* 2014; **99**:220–228.
- 83 Ortega JF, Fernández-Eliás VE, Hamouti N, Pallarés JG, Mora-Rodríguez R. Higher insulin-sensitizing response after sprint interval compared to continuous exercise. *Int J Sports Med* 2015; **36**:209–214.
- 84 Malin SK, Rynders CA, Weltman JY, Barrett EJ, Weltman A. Exercise intensity modulates glucose-stimulated insulin secretion when adjusted for adipose, liver and skeletal muscle insulin resistance. *PLoS One* 2016; **11**:e0154063.
- 85 Metcalfe R, Fawcner S, Vollaard N. No acute effect of reduced-exertion high-intensity interval training (REHIT) on insulin sensitivity. *Int J Sports Med* 2016; **37**:354–358.
- 86 Castleberry T, Irvine C, Deemer SE, Brisebois MF, Gordon R, Oldham MD, et al. Consecutive days of exercise decrease insulin response more than a single exercise session in healthy, inactive men. *Eur J Appl Physiol* 2019; **119**:1591–1598.
- 87 Richter EA, Garetto LP, Goodman MN, Ruderman NB. Enhanced muscle glucose metabolism after exercise: modulation by local factors. *Am J Physiol* 1984; **246**:E476–E482.
- 88 Annuzzi G, Riccardi G, Capaldo B, Kaijser L. Increased insulin-stimulated glucose uptake by exercised human muscles one day after prolonged physical exercise. *Eur J Clin Invest* 1991; **21**:6–12.
- 89 DeFronzo RA, Ferrannini E, Sato Y, Felig P, Wahren J. Synergistic interaction between exercise and insulin on peripheral glucose uptake. *J Clin Invest* 1981; **68**:1468–1474.
- 90 Pesta DH, Goncalves RLS, Madiraju AK, Strasser B, Sparks LM. Resistance training to improve type 2 diabetes: working toward a prescription for the future. *Nutr Metab (Lond)* 2017; **14**:24.
- 91 van Dijk JW, Manders RJ, Tummers K, Bonomi AG, Stehouwer CD, Hartgens F, van Loon LJ. Both resistance- and endurance-type exercise reduce the prevalence of hyperglycaemia in individuals with impaired glucose tolerance and in insulin-treated and non-insulin-treated type 2 diabetic patients. *Diabetologia* 2012; **55**:1273–1282.
- 92 Camera DM, Edge J, Short MJ, Hawley JA, Coffey VG. Early time course of Akt phosphorylation after endurance and resistance exercise. *Med Sci Sports Exerc* 2010; **42**:1843–1852.
- 93 Kido K, Sase K, Yokokawa T, Fujita S. Enhanced skeletal muscle insulin sensitivity after acute resistance-type exercise is upregulated by rapamycin-sensitive mTOR complex 1 inhibition. *Sci Rep* 2020; **10**:8509.
- 94 Oshida Y, Yamanouchi K, Hayamizu S, Nagasawa J, Ohsawa I, Sato Y. Effects of training and training cessation on insulin action. *Int J Sports Med* 1991; **12**:484–486.
- 95 Whyte LJ, Gill JM, Cathcart AJ. Effect of 2 weeks of sprint interval training on health-related outcomes in sedentary overweight/obese men. *Metabolism* 2010; **59**:1421–1428.
- 96 Choi JY, Ha HS, Kwon HS, Lee SH, Cho HH, Yim HW, et al. Characteristics of metabolically obese, normal-weight women differ by menopause status: the Fourth Korea National Health and Nutrition Examination Survey. *Menopause* 2013; **20**:85–93.
- 97 Gibala MJ. High-intensity interval training: a time-efficient strategy for health promotion? *Curr Sports Med Rep* 2007; **6**:211–213.
- 98 Turcotte LP, Swenberger JR, Zavitz Tucker M, Yee AJ. Increased fatty acid uptake and altered fatty acid metabolism in insulin-resistant muscle of obese Zucker rats. *Diabetes* 2001; **50**:1389–1396.
- 99 Bird SR, Hawley JA. Update on the effects of physical activity on insulin sensitivity in humans. *BMJ Open Sport Exerc Med* 2016; **2**:e000143.
- 100 da Nobrega AC. The subacute effects of exercise: concept, characteristics, and clinical implications. *Exerc Sport Sci Rev* 2005; **33**:84–87.
- 101 Contrepois K, Wu S, Moneghetti KJ, Hornburg D, Ahadi S, Tsai MS, et al. Molecular choreography of acute exercise. *Cell* 2020; **181**:1112–1130. e16.
- 102 Dumortier M, Brandou F, Perez-Martin A, Fedou C, Mercier J, Brun JF. Low intensity endurance exercise targeted for lipid oxidation improves body composition and insulin sensitivity in patients with the metabolic syndrome. *Diabetes Metab* 2003; **29**:509–518.
- 103 Mora-Rodríguez R, Ortega JF, Ramirez-Jimenez M, Moreno-Cabañas A, Morales-Palomo F. Insulin sensitivity improvement with exercise training is mediated by body weight loss in subjects with metabolic syndrome. *Diabetes Metab* 2020; **46**:210–218.
- 104 Ortega JF, Hamouti N, Fernández-Eliás VE, de Prada MV, Martínez-Vizcaino V, Mora-Rodríguez R. Metformin does not attenuate the acute insulin-sensitizing effect of a single bout of exercise in individuals with insulin resistance. *Acta Diabetol* 2014; **51**:749–755.