

POSTER PRESENTATION

Open Access

Yeast as a simple eukaryotic model to study human diseases linked to membrane trafficking

J De Craene^{1*}, S Bär¹, J Laporte², V Marion³, H Dollfus³, S Friant¹

From Cilia 2014 - Second International Conference
Paris, France. 18-21 November 2014

Many human diseases are linked to mutations in genes encoding membrane trafficking proteins, among them the X-linked centronuclear myopathy (XLCNM) and ciliopathies. We use yeast *Saccharomyces cerevisiae* as a model system to study these human disease trafficking genes. It is a good model because yeast and human cells present similar intracellular organization and its membrane trafficking and metabolism are well characterized. We first studied the MTM1 gene, responsible for XLCNM, which codes for a myotubularin, a lipid phosphatase required for endosomal sorting. The heterologous expression of wild-type MTM1 or XLCNM patient mutants in yeast allowed comparing their *in vivo* phosphatase activity and their cellular function in trafficking. The yeast results showed that the phosphatase activity is not defective in patient mutants, further confirmed in the mouse *Mtm1* KO model for myopathy [1].

Ciliopathies are complex genetic disorders that result from defects in the formation and/or function of the primary cilium. They include disorders such as Bardet-Biedl syndrome (BBS), Joubert syndrome (JBTS) and Alström syndrome (ALS) [2]. Cilia are involved in development and in signaling cascades. Despite the identification of a large number of genes involved in ciliopathies (<http://www.syscilia.org/goldstandard.shtml>), the molecular and cellular mechanisms by which they cause the disease are largely unknown. In collaboration with the laboratory of Hélène Dollfus, we aim at understanding the cellular roles of membrane trafficking genes involved in ciliopathies by using the yeast model system. Indeed, organization of the primary cilium depends on membrane trafficking since many genes linked to ciliopathies are involved in membrane trafficking [3].

Authors' details

¹UMR7156 Trafic Intracellulaire et Signalisation Lipidique, Université de Strasbourg, Strasbourg, France. ²Laboratoire de Physiopathologie des Maladies Neuromusculaires, IGBMC, Illkirch, France. ³Unité INSERM U1112, Université de Strasbourg, Strasbourg, France.

Published: 13 July 2015

References

1. Amoasii L, Bertazzi DL, Tronchere H, Hnia K, Chicanne G, Rinaldi B, et al: Phosphatase-dead myotubularin ameliorates X-linked centronuclear myopathy phenotypes in mice. *PLoS Genet* 2012, **8**:e1002965.
2. Baker K, Beales PL: Making sense of cilia in disease: the human ciliopathies. *Am J Med Genet C Semin Med Genet* 2009, **151C**:281-295.
3. Leroux MR: Taking vesicular transport to the cilium. *Cell* 2007, **129**:1041-1043.

doi:10.1186/2046-2530-4-S1-P31

Cite this article as: Craene et al.: Yeast as a simple eukaryotic model to study human diseases linked to membrane trafficking. *Cilia* 2015 **4**(Suppl 1):P31.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹UMR7156 Trafic Intracellulaire et Signalisation Lipidique, Université de Strasbourg, Strasbourg, France
Full list of author information is available at the end of the article