



Atopic Dermatitis Is Associated with Preeclampsia and Endometriosis

JID Innovations (2022);2:100123 doi:10.1016/j.jid.2022.100123

TO THE EDITOR

NK cell function has been shown to be associated with atopic dermatitis (AD), with patients with AD usually having decreased circulating numbers of NK cells but increasing lesional skin numbers of NK cells (Kabashima and Weidinger, 2020; Mack et al., 2020; Möbus et al., 2021; Sun et al., 2021). An important regulator of NK cell function are KIR genes, which can have either inhibitory or activating effects on NK cells (Pende et al., 2019). In addition to having a potential role in the pathogenesis of AD, NK cells have also been implicated in the development of preeclampsia and endometriosis. Interactions between maternal uterine NK cells and the extravillous trophoblast, potentially mediated by KIR and HLA-C ligands interactions, can influence the development of preeclampsia and pregnancy outcomes (Hiby et al., 2004). KIR and HLA-C ligands have also been associated with the development of endometriosis (Chou et al., 2020). A recent study revealed that KIR gene variation is associated with AD (Margolis et al., 2021). Because NK cell function and KIR genes appear to be involved in the pathogenesis of AD, preeclampsia, and endometriosis, it is possible that there may be an association between these conditions.

To evaluate whether there is an association between AD with preeclampsia and endometriosis, a retrospective cohort study was performed using the Optum Clininformatics Data Mart (Optum) between January 2016 and July 2020, as well as using The Health Improvement Network (THIN) through February 2015. Patients

were identified in Optum through the presence of diagnostic codes for AD (L20x) and either asthma (J45x) or allergic rhinitis (J30x), and in THIN using a previously validated algorithm for this dataset (Abuabara et al., 2017).

To evaluate for a potential association with preeclampsia, in Optum, patients with AD were matched with up to five patients without AD, based on age and index date of pregnancy delivery. In THIN, patients with AD were matched with up to five patients without AD patients based on age, practice, and an encounter within ± 6 months of the latter of practice registration and diagnosis dates for the patient with AD. Logistic regression was used to examine for differences in the frequency of preeclampsia (O14x) during the first recorded pregnancy. In Optum, preeclampsia was defined as any encounter with a diagnosis code for preeclampsia within 180 days before or 90 days after the delivery date. In THIN, preeclampsia was defined by any diagnosis of preeclampsia within 180 days before or after the first pregnancy code.

To evaluate for a potential association with endometriosis, in Optum, patients with AD were matched 1:1 with patients who were diagnosed with a nevus or seborrheic keratosis, based on age and index date of diagnosis. In THIN, the same matching approach was used as described above. Logistic regression was used to examine for differences in the frequency of endometriosis (N80x). All analyses were restricted to female patients between the ages of 15 and 45 years at the time of the index date. Analyses in Optum

were additionally adjusted for follow-up time before and after the index date, and all individuals were required to have at least 1 year of continuous enrollment before and after the index date.

Analyses performed in Optum and THIN were synthesized using random effects meta-analysis. Statistical analyses were performed in Stata 15 (StataCorp, College Station, TX). This study was reported in adherence with the Strengthening the Reporting of Observational Studies in Epidemiology guidelines (von Elm et al., 2007).

Preeclampsia was more common in women with AD than controls in both the Optum (1.59% vs. 1.10%; adjusted odds ratio [aOR] = 1.27, 95% confidence interval [CI] 0.83–1.94) and THIN cohorts (0.74% vs. 0.67%; OR = 1.11, 95% CI = 1.03–1.19), although this association did not reach statistical significance in the Optum cohort (Table 1). In the meta-analysis, there was an association between AD and preeclampsia (OR = 1.11, 95% CI = 1.03–1.19). Endometriosis was more common in women with AD than controls in both the Optum (4.29% vs. 3.00%; aOR = 1.48, 95% CI = 1.24–1.78) and THIN cohorts (1.17% vs. 0.95%; OR = 1.24, 95% CI = 1.20–1.27) (Table 2). In the meta-analysis, there was an association between AD and endometriosis (OR = 1.32, 95% CI = 1.09–1.55).

AD is a common persistent dermatologic illness that often begins in early childhood. Classically, other allergic illnesses have been associated with AD such as asthma, food allergies, and seasonal allergies. In this cohort study, using two administrative databases, we identify an association of AD with preeclampsia and endometriosis. The effect estimates were small but similar in both datasets, revealing the generalizability of this finding. The findings of our work and others support that nonallergic illnesses are also associated

Abbreviations: AD, atopic dermatitis; CI, confidence interval; Optum, Optum Clininformatics Data Mart; THIN, The Health Improvement Network

Accepted manuscript published online XXX; corrected proof published online XXX

Cite this article as: JID Innovations 2022;2:100123

© 2022 The Authors. Published by Elsevier, Inc. on behalf of the Society for Investigative Dermatology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Table 1. Association of Atopic Dermatitis with Preeclampsia

	Atopic Dermatitis	Without Atopic Dermatitis
Optum		
N	1,821	7,946
Continuous enrollment prior to index, d, mean (SD)	1,240 (1,082)	1,097 (985)
Continuous enrollment after index, d, mean (SD)	630 (304)	553 (288)
Preeclampsia, n (%)	29 (1.59)	88 (1.10)
OR preeclampsia (95% CI), adj ¹	1.27 (0.83–1.94)	P = 0.276
THIN		
N	142,773	515,542
Preeclampsia, n (%)	1,067 (0.74)	3,482 (0.67)
OR preeclampsia (95% CI)	1.10 (1.03–1.19)	P = 0.004

Abbreviations: adj, adjusted; CI, confidence interval; d, days; Optum, Optum Clininformatics Data Mart; THIN, The Health Improvement Network

¹Adjusted for follow-up time

Table 2. Association of Atopic Dermatitis with Endometriosis

	Atopic Dermatitis	Without Atopic Dermatitis
Optum		
N	18,342	18,342
Continuous enrollment prior to index, d, mean (SD)	1,639 (1,317)	1,170 (793)
Continuous enrollment after index, d, mean (SD)	709 (253)	747 (271)
Endometriosis, n (%)	304 (4.3)	213 (3.0)
OR endometriosis preeclampsia (95% CI), adj ¹	1.48 (1.24–1.78)	P < 0.001
THIN		
N	567,557	2,295,873
Endometriosis, n (%)	6,725 (1.17)	21,995 (0.95)
OR endometriosis (95% CI)	1.23 (1.08–1.40)	P = 0.002

Abbreviations: adj, adjusted; CI, confidence interval; d, days; Optum, Optum Clininformatics Data Mart; THIN, The Health Improvement Network

¹Adjusted for follow-up time

with AD (Sinaii et al., 2002; Stokholm et al., 2017). Furthermore, the associations with preeclampsia and endometriosis are potentially related to genetic variation in *KIR* genes and NK cell function, revealing additional immunologic pathways that may be associated with AD (Margolis et al., 2021; Nakimuli et al., 2015).

This study has several strengths, including replicating the findings in two independent datasets from two different countries (United States and United Kingdom). This study is limited in that the Optum cohort was relatively small, limiting power to detect differences in our analysis of preeclampsia. Although we have attempted to mitigate this risk using matched cohorts, there is the potential for unmeasured confounding

related to degree of interaction with the healthcare system, race and ethnicity, obesity, and smoking; these variables are all poorly measured in our cohorts, making adjustment unreliable. In addition, because we do not have associated genetic data, we are unable to assess whether *KIR* gene function may be a key factor in these observed associations. Future studies are needed to replicate our findings further and to examine for potential biologic mediations of these observed associations, including potential interactions with ancestry.

Data availability statement

Data are not available from the authors because this dataset is restricted.

ORCIDs

- John S. Barbieri: <http://orcid.org/0000-0002-5467-4102>
 Daniel B. Shin: <http://orcid.org/0000-0002-4974-2561>
 David J. Margolis: <http://orcid.org/0000-0002-0506-8085>

AUTHOR CONTRIBUTIONS

Conceptualization: DJM; Data Curation: JSB, DBS; Formal Analysis: JSB, DBS; Funding Acquisition: DJM; Investigation: JSB, DBS, DJM; Methodology: JSB, DBS, DJM; Project Administration: DJM; Resources: DJM; Software: JSB, DBS; Supervision: DJM; Validation: JSB, DBS; Visualization: JSB, DBS, DJM; Writing - Original Draft Preparation: JSB; Writing - Review and Editing: JSB, DBS, DJM

CONFLICT OF INTEREST

The authors state no conflict of interest.

ACKNOWLEDGMENTS

Support for this work was provided by the Penn Skin Biology and Diseases Resource-based Center, funded by National Institutes of Health/National Institute of Arthritis and Musculoskeletal and Skin Diseases grant P30-AR069589 and the University of Pennsylvania Perelman School of Medicine.

John S. Barbieri^{1,*}, Daniel B. Shin² and David J. Margolis^{3,4}

¹Department of Dermatology, Brigham and Women's Hospital, Boston, Massachusetts, USA; ²Department of Dermatology, University of Pennsylvania, Philadelphia, Pennsylvania, USA; ³Department of Dermatology, University of Pennsylvania, Philadelphia, Pennsylvania, USA; and ⁴Department of Biostatistics, Epidemiology and Informatics, University of Pennsylvania, Philadelphia, Pennsylvania, USA

*Corresponding author
 e-mail: jbarbieri@bwh.harvard.edu

REFERENCES

- Abuabara K, Magyari AM, Hoffstad O, Jabbar-Lopez ZK, Smeeth L, Williams HC, et al. Development and validation of an algorithm to accurately identify atopic eczema patients in primary care electronic health records from the UK. *J Invest Dermatol* 2017;137:1655–62.
 Chou YC, Chen CH, Chen MJ, Chang CW, Chen PH, Yu MH, et al. Killer cell immunoglobulin-like receptors (KIR) and human leukocyte antigen-C (HLA-C) allor-ecognition patterns in women with endometriosis. *Sci Rep* 2020;10:4897.
 Hiby SE, Walker JJ, O'Shaughnessy KM, Redman CWG, Carrington M, Trowsdale J, et al. Combinations of maternal KIR and fetal HLA-C genes influence the risk of preeclampsia and reproductive success. *J Exp Med* 2004;200:957–65.
 Kabashima K, Weidinger S. NK cells as a possible new player in atopic dermatitis. *J Allergy Clin Immunol* 2020;146:276–7.
 Mack MR, Brestoff JR, Berrien-Elliott MM, Trier AM, Yang TLB, McCullen M, et al. Blood natural killer cell deficiency reveals an

immunotherapy strategy for atopic dermatitis. *Sci Transl Med* 2020;12:eaay1005.

Margolis DJ, Mitra N, Hoffstad OJ, Kim BS, Monos DS, Phillips EJ. Association of KIR genes and MHC class I ligands with atopic dermatitis. *J Immunol* 2021;207:1522–9.

Möbus L, Rodriguez E, Harder I, Schwarz A, Wehkamp U, Stölzl D, et al. Elevated NK-cell transcriptional signature and dysbalance of resting and activated NK cells in atopic dermatitis. *J Allergy Clin Immunol* 2021;147:1959–65.e2.

Nakimuli A, Chazara O, Hiby SE, Farrell L, Tukwasibwe S, Jayaraman J, et al. A KIR B centromeric region present in Africans but not Europeans protects pregnant women from pre-eclampsia. *Proc Natl Acad Sci USA* 2015;112:845–50.

Pende D, Falco M, Vitale M, Cantoni C, Vitale C, Munari E, et al. Killer Ig-like receptors (KIRs): their role in NK cell modulation and developments leading to their clinical exploitation. *Front Immunol* 2019;10:1179.

Sinaii N, Cleary SD, Ballweg ML, Nieman LK, Stratton P. High rates of autoimmune and endocrine disorders, fibromyalgia, chronic fatigue syndrome and atopic diseases among women with endometriosis: a survey analysis. *Hum Reprod* 2002;17:2715–24.

Stokholm J, Sevelsted A, Anderson UD, Bisgaard H. Preeclampsia associates with asthma, allergy, and eczema in childhood. *Am J Respir Crit Care Med* 2017;195:614–21.

Sun Z, Kim JH, Kim SH, Kim HR, Zhang K, Pan Y, et al. Skin-resident natural killer T cells participate in cutaneous allergic inflammation in

atopic dermatitis. *J Allergy Clin Immunol* 2021;147:1764–77.

von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandebroucke JP, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet* 2007;370:1453–7.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>