



OPEN

Nitrogen and phosphorus losses from paddy fields and the yield of rice with different water and nitrogen management practices

Dongliang Qi, Qixia Wu & Jianqiang Zhu[✉]

The assessment and control of losses of nitrogen (N) and phosphorus (P) from paddy fields is critical to improve the quality of water and atmosphere on earth. A field experiment was conducted to investigate the effect of three N managements (local common N fertilization practice, urea mixed with controlled-release N fertilizer, and optimized and reduced N fertilizer, designated CN, U + CRF and ON, respectively) on N and P losses through runoff and leaching from a paddy field, and yield of rice under shallow-irrigation and deep-slucice (SIDS) and continuous flooding irrigation (FI) in the Jianhan Plain of China in 2016. The results showed that, compared with FI, SIDS significantly reduced the frequency of irrigation and amount of irrigation water, resulting in an increase of 16.2% in rainfall use efficiency, and therefore, a reduction in the amount of surface runoff and water that had leached. This was responsible for the decreased total N (TN) and total P (TP) losses through runoff leaching under SIDS. The U + CRF and ON treatments resulted in a significant reduction in losses of TN through runoff and leaching and the loss of TP through leaching compared to CN. SIDS resulted in comparable or greater soil TN and TP contents in the 0–40 cm soil depths after rice harvest; N and P accumulation at the jointing, filling and maturity stages; and yield of grain compared to FI. Moreover, the U + CRF and ON improved or maintained accumulation of N and P and yield of rice compared to CN. Compared with FI coupled with CN, SIDS coupled with the U + CRF or ON treatments significantly reduced losses of N and P from paddy fields and enhanced or maintained the accumulation of N and P and yield of rice grains. In conclusion, SIDS coupled with the new N management could be an effective approach to reduce losses of N and P from paddy fields and would be a positive improvement for high yield of middle-season rice grains in the Jianhan Plain of central China and other regions with similar environments.

Rice (*Oryza sativa* L.) is one of the primary crops in the world and the foremost staple food in Asia, supplying 35–60% of the dietary calories consumed by more than three billion people¹. In China, the average annual area of rice planted and total production had reached 30.1 million ha and 18.6 billion t in 2011, respectively². Both the planting area and total production ranked first in the world³. However, the growth of rice requires a substantial amount of fresh water, so that the rice planting system accounted for 45–50% of total water consumption in China⁴. Moreover, a shortage of water resources is a serious problem in China, and its spatial and temporal distribution is extremely uneven⁵. The water scarcity is further strengthened by climate change, a limited supply of water, and the increasing water consumption by cities, industries and other sectors of the economy^{6,7}. This has encouraged more researchers to develop novel irrigation strategies to improve crop water use efficiency (WUE), so that the sustainability of rice production could be assured^{8,9}.

Various water-efficient irrigation management modes are currently practiced in different paddy fields in China, including alternate wetting and drying, shallow-irrigation and deep-slucice (SIDS), intermittent irrigation, controlled irrigation, flooding-midseason drainage-frequent water logging with intermittent irrigation, and semi-dry cultivation among others^{10–13}. Among these irrigation methods, SIDS is considered to be an efficient irrigation method to maintain the yield of rice, while reducing no-point pollution and the amount of irrigation compared to continuous flood irrigation in the Jiangnan Plain of China^{13–15}. In SIDS, the precipitation is sluciced

Engineering Research Center of Ecology and Agriculture Use of Wetland, Ministry of Education, Yangtze University, Jingzhou, Hubei, 434025, China. ✉e-mail: 200572@yangtzeu.edu.cn

to some extent, and the field remains non-flooded unless cracks appear on the soil surface; thus, alternate wetting and drying cycles occur in paddy fields during the whole rice growing season¹⁴. Because of the high economic return and its ease of application, SIDS has been widely practiced in several provinces in China, including Hubei, Hunan, Jiangsu, and Anhui.

Nitrogen (N) fertilizer is another important input for intensive rice production¹⁶. The average yield of rice per unit area in China is $6.18 \times 10^3 \text{ kg ha}^{-1}$, which was 65% higher than that of the average yield in world; the amount of N fertilizer applied for rice production in China accounts for 37% of the N fertilizer used for rice in the world¹⁷. Thus, the recovery efficiency of fertilizer N is only approximately 30%, which is approximately 40–50% lower than the world average¹⁸. Over and/or improper fertilization is a serious issue in intensive agricultural production areas in China, contributing to soil degradation, lake eutrophication, groundwater pollution, and the emission of ammonia and greenhouse gases^{19–22}. The loss of nutrients from agricultural fields is one of the main contributors to these types of pollution^{20,21}, particularly in the central and southern regions of China²³. Additionally, N fertilizer is becoming expensive because of increasing demand for fossil fuels worldwide. Therefore, exploring effective measures to reduce nutrient losses from farmland, while maintaining crop yield has become a priority for the sustainable development of agriculture, particularly for rice production.

To not only favor the profitability of farmers but also reduce the negative environmental impacts, several researchers have tested different N management practices to improve N use efficiency (NUE)^{7,18,23}. Some N saving methods (site-specific N management, balanced N fertilization, integrated N management, the use of nitrification/urease inhibitors and slow/controlled-release fertilizers) are already showing promise^{2,23,24}. Recently, the use of controlled release N fertilizer has been found to be an effective way to increase the NUE and reduce N losses from paddy fields²³. Moreover, a reduction in the level of N fertilizer supplied and amount of irrigation is the most effective measure to control nitrate leaching in cropland farming^{25,26}. In addition, optimized fertilization could increase the NUE of rice²⁷ and reduce the N and phosphorous (P) losses from source¹⁹. However, the effects of N-saving methods vary depending on different environmental conditions.

The key growth period of rice usually coincides with the plum-rain (a weather phenomenon characterized by continuous overcast and rainy weather) season in the Jiangnan Plain of China, leading to more frequent surface and underground drainage occurring from paddy fields, and an indirect drought often happens after the plum-rain²⁸. Moreover, earlier research has illustrated that soil nutrients and water availability are closely linked and mutually influence one another²⁹. Fertilization and water management are two important factors that influence the migration and use efficiency of N and P in paddy fields³⁰. Thus, it is necessary to explore an individual appropriate N application mode under water-saving irrigation based on specific weather conditions in the rice planting areas, such as the Jiangnan Plain of China. In addition, the manner in which controlled release N fertilizer and optimized N management influence the losses of N and P from paddy fields and yield of rice has yet to be addressed under SIDS.

This study was designed to quantify the losses of N and P through leaching and runoff from a paddy field, N and P accumulation in rice, and yield of rice with controlled release N fertilizer and optimized N management under SIDS. Moreover, the contributions of various forms of N and P compounds to the total N and total P losses through leaching and runoff were quantified. The results should provide effective measures to reduce the losses of N and P in paddy fields and improve rice yield in the Jiangnan Plain of central China and other regions with similar environments.

Materials and Methods

Experimental site and materials. A field study was conducted in 2016 at the agricultural experimental station of Yangtze University in Jingzhou, Hubei Province, central China (latitude $30^\circ 21' \text{N}$, longitude $112^\circ 09' \text{E}$, altitude 28 m). The site is in a typical subtropical monsoon climate zone with a mean annual precipitation of 1,095 mm. The mean annual sunshine duration is more than 1,718 h, and the mean annual temperature is 16.5°C . The accumulated temperature of $>10^\circ \text{C}$ is 5,094.9–5,294.3 $^\circ \text{C}$. The soil pH was 7.4, total N 2.04 g kg^{-1} , total P 0.48 g kg^{-1} , available N 79.5 mg kg^{-1} , available P 38.5 mg kg^{-1} and available potassium (K) 108.7 mg kg^{-1} in the plough layer (0–40 cm soil depth).

The experimental rice used was mid-maturation two-line hybrid rice (*Oryza sativa* L.) variety “Huifeng8”. The controlled-release N fertilizer used was CRF (N:P₂O₅:K₂O = 28:5:9) (Kingenta Ecological Engineering Co., Ltd., Shandong, China) with 70 days of N released period. The two conventional N fertilizers used were urea (46% N) and common compound fertilizer (N:P₂O₅:K₂O = 18:8:15). The P and K fertilizers used were superphosphate (12% P₂O₅) and potassium chloride (60% K₂O), respectively.

Experimental design. The experimental factors were comprised of the irrigation method and N fertilizer management. The irrigation treatments included conventional flooding irrigation (FI) and shallow-irrigation and deep-sluice (SIDS). The N treatments included local common N fertilization practice, 30% of fertilizer N (urea) mixed with the other 70% of controlled release N fertilizer (CRF), and optimized and reduced N fertilization, designated CN, U + CRF and ON, respectively. This experimental plan yielded six treatments. There were three replicates for each treatment.

At 10–14 days after the transplantation of rice, a standing water depth of 10–40 mm was maintained for all the treatments to facilitate recovery of seedlings and their ability to turn green. FI and SIDS were then managed differently. In FI, a standing water depth of 10–80 mm in the field was maintained until the terminal drainage at approximately 10 days before the harvest of rice. In SIDS, the field was allowed to be intermittently submerged and was not irrigated unless the standing water depth dropped to approximately 100 mm below the topsoil. The field was re-flooded to a standing water depth of 40–60 mm in each irrigation event. The cycles were repeated until the necessary drainage before harvest¹⁴. However, there was one week of exception to the drainage to maintain a standing water depth of 30–50 mm at the flowering stage. Wu *et al.*²⁸ reported that the sluice water above

Treatment	Basal application					Topdressing one	Topdressing two
	CRF (kg ha ⁻¹)	CCF (kg ha ⁻¹)	U (kg ha ⁻¹)	S (kg ha ⁻¹)	PC (kg ha ⁻¹)	U (kg ha ⁻¹)	U (kg ha ⁻¹)
CN	/	700	/	158	/	117	/
U + CRF	450	/	117	438	108	/	/
ON	/	417	/	347	70.8	114	49

Table 1. Fertilization scheme for rice under the different nitrogen fertilizer treatments. Note: / represents that there was no fertilization. S, superphosphate; U, urea; CRF, controlled-release nitrogen fertilizer; CCF, common compound fertilizer; PC, potassium chloride.

soil surface of paddy field could be maintained up to 50, 100, and 150 mm at re-greening, tillering to jointing, and booting to maturity stages of rice, respectively. In the CN treatment, 70% N (common compound fertilizer) was applied at the basal, and 30% N (urea) was applied as topdressing at the tillering stage. In the U + CRF treatment, basal applications of 70% N used CRF, and the other 30% N used urea as N source were utilized. In the ON treatment, common compound fertilizer was applied at the basal (50% N), and the urea was top-dressed at tillering (35% N) and heading (15% N) stages, respectively. The corresponding dates were June 4, July 6, and August 1, 2016, respectively. The application of N fertilizer rate for CN, U + CRF and ON treatments was 180, 150 and 150 kg N ha⁻¹, respectively. A total of 75 kg P₂O₅ ha⁻¹ and 105 kg K₂O ha⁻¹ was utilized as a basal application for each N treatment. Details of the fertilization scheme, including fertilizer sources and the corresponding application rates used for each N treatment, are shown in Table 1.

Experimental management. The experimental field was planted with rice year round and served as a representative paddy field. Each plot was 30 m² (6 m × 5 m), and the circumference of each plot was isolated by a PVC sheet that was 60 cm high (30 cm below the soil surface and 30 cm above the soil surface) to form a barrier. Each plot was separately irrigated and drained. The 4-week-old seedlings were artificially transplanted at 25 cm × 30 cm with 3 seedlings per plant on June 5, 2016. The plots were regularly hand-weeded to prevent weed damage during the rice grown season. Diseases and insects were intensively controlled by chemicals to avoid a loss of yield. The whole growth period was 136 d in all treatments. The rice was harvested on September 18, 2016. All of rice straw was returned to paddy field after the crop harvest. A water meter was installed at the discharging end of water inlet to measure amount of water applied, and a container was installed at the terminal of drainpipe to collect runoff.

Measurements

Precipitation. The amount of precipitation during the rice grown season was recorded by an automatic weather station (ICT, Australia) within the experimental base (approximately 100 m away from the plots).

Runoff and leaching water. Before the experiment, a homemade iron leaching bucket (30 cm in diameter, 100 cm long) was installed vertically at a depth of 60 cm to collect the daily percolation water of each plot. The pipe orifice was 40 cm above the soil surface, and the upper part was covered to prevent rain, dust or insects from entering pipe. A water level indicator was used to measure water level at an interval of 2–3 days. The daily volume of percolation water in paddy fields was calculated according to the water level difference proposed by Yang *et al.*²³. Ceramic suction cups were installed at 3, 9 and 12 m of the water inlet and vertically at a depth of 30 cm to measure leaching water after irrigation or precipitation (≥ 30 mm) during the growth stages of rice. After rainfall, the excessive water was drained on the basis of upper limit of sluice water at the different growth stages in SIDS. The runoff water was collected by an overflow bucket and sampled from the outlet of each plot. The amount of drainage was calculated from the difference between the water depth of the paddy field before and after drainage as described by Yang *et al.*²³.

N and P of water and soil. Soil samples were collected after the rice harvest using a stainless-steel auger. The S-shaped 5-point collection method was used for sampling. The samples were collected from 0–20 and 20–40 cm soil depths in each plot. Root debris and soil samples were sieved and stored as described by Si *et al.*³¹. In short, the plant root remnants and rocks were removed, and the samples were passed through a 0.85 mm sieve and stored at 4 °C to determine relevant indices.

The water quality parameters determined included the following: TP, TN, dissolved phosphorus (DP), ammonia-nitrogen (NH₄⁺-N), nitrate-nitrogen (NO₃⁻-N) and particulate phosphorus (PP). Soil nutrient parameters that were monitored included TN, TP, NH₄[±]-N, NO₃⁻-N and available P. All the analyses were conducted as recommended by the State Environmental Protection Administration of China (1997). Dissolved organic N (DON) was calculated by the differences between TN and sum of NH₄⁺-N and NO₃⁻-N.

Nutrient uptake. Five representative rice plants in each plot were cut at the re-greening, tillering, joint-booting, filling and maturity stages. The plants were divided into stems and leaves before heading stage, and stems, leaves and panicles after the heading stage. All the plant samples were oven-dried to a constant biomass at 70 °C and weighed. The samples were passed through a 0.15 mm sieve, and subsamples were collected for total N and total P determination as described by Ye *et al.*². The total N was analyzed using the semi-micro Kjeldahl method, and the total P was analyzed using the vanadium molybdate yellow colorimetric method³². Tissue N or P concentrations were multiplied by the yield of dry matter to calculate total N or P uptake.




Figure 1. The volume of precipitation, irrigation and runoff water during rice grown season under continuously flooding (FI) irrigation and shallow-irrigation and deep-sluice (SIDS).

Rice yield. Yield of grain was measured from a 5 m² area in the center of each plot at maturity stage. The rice was harvested manually, and the grains were air dried.

Data analysis. According to Zhu *et al.*³³, the deep underground leaching was ignored owing to high water table (0.8–1.5 m below the surface) in the paddy fields at Jiangnan Plain and the negligible plant interception (approximately 0.2% of precipitation amount). Precipitation use efficiency (Y) was calculated as follows:

$$Y = \frac{RT - R_t}{R_T} \times 100\% \quad (1)$$

where R_T is the total amount of precipitation (mm) during a certain period of time, and R_t is the amount of runoff water (mm) during a certain period of time.

The amount of N or P loss was calculated as follows:

$$P = \sum_{i=1}^n (C_i \times V_i) \quad (2)$$

where C_i is the concentration of N or P (mg L⁻¹) in the runoff or leaching water at the i time, and V_i is the volume of runoff or leaching water at the i time (L).

An analysis of variance (ANOVA) was performed using the general linear model-univariate procedure from SPSS 12.0 software (USA). ANOVAs were conducted with irrigation method and fertilizer management as the main effects and included their interactions. The mean values were compared for any significant differences among different treatments using the Duncan's multiple range tests at a significance level of $P \leq 0.05$

Results

In total, the leaching water was measured 20 times for SIDS and FI; the runoff water was measured four times for SIDS and seven times for FI during the rice growing season. The average of TN concentration from leaching water was 3.78 and 3.89 mg L⁻¹ for FI and SIDS, respectively, which was comparable between the two irrigation regimes. However, the average of TN concentration from leaching water was 4.21 mg L⁻¹ for the CN treatment, which was 18.7% and 20.3% significantly greater than that for the U + CRF and ON treatments, respectively. The average of TN concentration from runoff water and average of TP concentration from runoff and leaching water showed a highly similar amount of variation compared with the average of TN concentration from leaching water.

Precipitation, runoff, irrigation and leaching water. Within 110 days after the transplantation (DAT) of rice, the accumulated precipitation was 550.3 mm. The maximum daily precipitation of 70.3 mm was recorded on July 2, 2016 (Fig. 1). For the FI treatments, the field was irrigated seven times, and the amount of total irrigation water, total water consumption (sum of the accumulated precipitation and the total irrigation) and runoff water was 440.5, 990.8 and 194.8 mm, respectively. The runoff water of 157.3 mm was measured at tillering




Figure 2. Total nitrogen (TN) and total phosphorus (TP) losses through runoff during the whole grown season of rice as affected by different water and nitrogen management strategies. Note: Values (mean \pm standard error, $n = 3$) by different letters are significantly different at $P < 0.05$.




Figure 3. Total nitrogen (TN) and total phosphorus (TP) losses through leaching during the whole growth season of rice as affected by different water and nitrogen management strategies. Note: Values (mean \pm standard error, $n = 3$) by different letters are significantly different at $P < 0.05$.

stage and accounted for 80.8% of total runoff water. The rest of total runoff water (37.5 mm) was measured at jointing-booting stage of rice. For the SIDS treatments, the field was irrigated four times, and the amount of total irrigation water, water consumption and runoff water was 257.0, 807.3 and 105.6 mm, respectively. The runoff water was only measurable at the tillering stage of rice. Compared with the FI treatment, the SIDS treatment significantly decreased the amount of irrigation water, total water consumption and runoff water by 41.7%, 18.5% and 45.8%, respectively, and significantly increased the precipitation use efficiency by 16.2% (Fig. 1). The amount of total leachate water from paddy field was 268 mm for SIDS and 343 mm for FI.

Losses of N and P during the whole grown season. The FI treatments had the following runoff losses during the whole growth period of rice: $\text{NH}_4^+\text{-N}$ 1.99–2.69 kg ha^{-1} ; $\text{NO}_3^-\text{-N}$ 0.77–1.16 kg ha^{-1} ; DON 2.69–4.87 kg ha^{-1} ; TN 4.30–6.07 kg ha^{-1} ; DP 0.14–0.16 kg ha^{-1} ; PP 0.17–0.19 kg ha^{-1} ; and TP 0.32–0.34 kg ha^{-1} . Compared with the FI treatments, the runoff losses of $\text{NH}_4^+\text{-N}$, $\text{NO}_3^-\text{-N}$, DON, TN, DP, PP and TP were 28.5–35.7%, 22.4–54.5%, 26.1–48.9%, 32.6–35.9%, 35.7–60.0%, 36.8–47.1% and 36.4–53.1% significantly smaller in the SIDS treatments, respectively. Both the U + CRF and ON treatments significantly decreased the loss of TN through runoff compared with that of CN treatment under the two irrigation treatments. FI coupled with CN resulted in the greatest runoff loss of TN, while SIDS coupled with U + CRF or CN resulted in the smallest one (Fig. 2). However, the runoff loss of TP was comparable among the different N treatments under FI and SIDS (Fig. 2). $\text{NH}_4^+\text{-N}$ and DON were the major components of TN lost through runoff and accounted for 41.1–41.3% and 45.1–46.1% of the TN loss, respectively. PP was the major component of TP lost through runoff and accounted for 52.9–60.0% of the TP loss.

The FI treatments had the following amount of leaching losses during the whole growth period of rice: $\text{NH}_4^+\text{-N}$ 8.98–13.83 kg ha^{-1} ; $\text{NO}_3^-\text{-N}$ 1.40–2.83 kg ha^{-1} ; DON 4.38–13.57 kg ha^{-1} ; TN 14.26–19.38 kg ha^{-1} ; DP 0.20–0.32 kg ha^{-1} ; PP 0.17–0.17 kg ha^{-1} ; and TP 0.37–0.49 kg ha^{-1} . Compared with the FI treatments, the total amount of $\text{NH}_4^+\text{-N}$, $\text{NO}_3^-\text{-N}$, DON, TN, DP, PP and TP lost through leaching was 23.5–28.1%, 12.9–37.5%, 16.8–28.8%, 22.8–32.0%, 5.0–36.4%, 23.5–29.4% and 16.2–33.3% significantly smaller in the SIDS treatments, respectively. Both the U + CRF and ON treatments significantly decreased leaching loss of TN compared with that of CN under the two irrigation treatments (Fig. 3). FI coupled with CN resulted in the greatest loss of TN by leaching, while SIDS coupled with U + CRF or CN resulted in the smallest loss of TN (Fig. 3). The loss of TP by leaching was very similar to that observed from the loss of TN by leaching (Fig. 3). $\text{NH}_4^+\text{-N}$ was the major

Growth stage	Index	FI			SIDS		
		CN	U + CRF	ON	CN	U + CRF	ON
Tillering	NH ₄ ⁺ -N	2.27 a	1.53 b	1.54 b	1.73 b	1.43 c	1.28 c
	NO ₃ ⁻ -N	1.07 a	0.90 b	0.73 b	0.90 b	0.66 b	0.35 c
	DON	1.42 a	0.77 c	0.73 c	1.26 a	1.11 b	1.27 b
	DP	0.11 a	0.12 a	0.12 a	0.09 b	0.08b	0.06 b
	PP	0.16a	0.15a	0.14a	0.12a	0.11a	0.09b
Jointing-booting	NH ₄ ⁺ -N	0.43 a	0.47 a	0.45 a	/	/	/
	NO ₃ ⁻ -N	0.09 a	0.06 b	0.04 b	/	/	/
	DON	0.79 a	1.14 a	1.02 a	/	/	/
	DP	0.03 a	0.03 a	0.03 a	/	/	/
	PP	0.03a	0.04a	0.03a			

Table 2. Amount of various nitrogen (N) and phosphorus (P) form losses through runoff (kg ha⁻¹) as affected by different water and nitrogen management strategies. Note: DON, dissolved organic nitrogen; DP, dissolved phosphorus; PP, particulate phosphorus. Means within a row followed by different letters are significantly different at $P < 0.05$.

Growth stage	Index	FI			SIDS		
		CN	U + CRF	ON	CN	U + CRF	ON
Regreening-booting	NH ₄ ⁺ -N	11.32 a	8.64 b	7.06 b	8.58 b	6.87 c	5.96 c
	NO ₃ ⁻ -N	1.92 a	1.26 a	0.85 a	1.48 a	1.00 ab	0.90 b
	DON	1.84b	1.38b	2.44a	0.95c	1.32b	1.72b
	DP	0.23 a	0.15 a	0.14 a	0.21 a	0.16 a	0.13 a
	PP	0.10 a	0.13 a	0.10 a	0.09 a	0.08 a	0.09 a
Heading-maturity	NH ₄ ⁺ -N	2.51 a	1.88 b	1.92 b	1.36b	1.16c	0.91 c
	NO ₃ ⁻ -N	0.91 a	0.28b	0.55 ab	0.32 a	0.26 a	0.29 a
	DON	0.88c	1.05b	1.45a	0.52d	0.58d	0.61d
	DP	0.09 a	0.05 a	0.08 a	0.04 a	0.03 a	0.04 a
	PP	0.07 a	0.04 a	0.07 a	0.04 a	0.04 a	0.03 a

Table 3. Amount of various nitrogen (N) and phosphorus (P) form losses through leaching (kg ha⁻¹) as affected by different water and nitrogen management strategies. Note: DON, dissolved organic nitrogen; DP, dissolved phosphorus; PP, particulate phosphorus. Means within a row followed by different letters are significantly different at $P < 0.05$.

component of TN lost through leaching and accounted for 65.9–75.4% of the TN loss. DP was the major component of TP lost through leaching and accounted for 53.8–65.8% of the TP loss.

Runoff losses of various forms of N and P. For FI, approximately 70% of the various N and P forms lost through runoff took place at tillering stage, and the other 30% took place at jointing-booting stage of rice. For SIDS, all of the various N and P forms lost through runoff took place at tillering stage (Table 2). Compared with FI, SIDS decreased the amount of NH₄⁺-N, NO₃⁻-N, DP and PP losses. However, SIDS increased the amount of DON loss at U + CRF and ON (Table 2). Compared with the CN treatment, the U + CRF and ON treatments decreased amount of NH₄⁺-N and NO₃⁻-N at tillering stage. The amount of DP and PP losses was comparable among the three N treatments under FI and SIDS (Table 2).

Leachate losses of various forms of N and P. The amount of various N and P forms lost through leaching during the re-greening to booting stages was approximately 75% of the whole rice growing stage. The other 25% was lost during heading to maturity. Compared with FI, SIDS decreased the amount of losses of NH₄⁺-N, NO₃⁻-N and DON in each N treatment (Table 3). Compared with CN, the U + CRF and ON treatments decreased the amount of NH₄⁺-N and NO₃⁻-N under FI and SIDS. However, the losses of DP and PP were comparable among different treatments (Table 3).

N and P uptake by rice. As shown in Table 4, the N and P uptake by rice at re-greening and tillering stages was comparable between FI and SIDS treatments in each N treatment with the exception of uptake of P in the ON treatment. Compared with FI, SIDS increased the uptake of N by rice at the jointing, filling and maturity stages in each N treatment. Similarly, SIDS treatments increased the P uptake by rice at jointing, filling and maturity stages in each N treatment. For the two irrigation regimes, compared with CN treatment, the U + CRF treatment increased the N and P uptake at all measured stages with the exception of uptake of P at the filling stage. The ON treatments increased uptake of N and P in general at the jointing, filling and maturity stages of rice. SIDS coupled with U + CRF achieved the greatest uptake of N and P by rice at maturity stage.

Irrigation method	Fertilizer management	Re-greening		Tillering		Jointing		Filling		Maturity	
		NU	PU	NU	PU	NU	PU	NU	PU	NU	PU
FI	CN	1.12a	0.13ab	41.53a	2.91b	61.19b	12.31b	124.51c	30.40a	153.64c	48.86b
	U + CRF	1.40a	0.23a	49.40a	5.49a	73.14a	13.56b	157.55a	25.13b	171.87b	55.96b
	ON	1.05a	0.13b	26.21b	3.38b	70.59a	13.13b	138.82b	20.68a	154.79c	53.82b
SIDS	CN	1.13ab	0.16ab	41.66a	3.39b	71.13a	14.85b	137.51b	34.82a	157.37c	52.27b
	U + CRF	1.21a	0.19a	56.99a	6.39a	82.95a	16.78a	158.11a	30.38a	179.02a	63.29a
	ON	0.92ab	0.10b	30.23b	5.37a	76.06a	15.90a	139.00b	30.93a	167.72b	57.68b

Table 4. Nitrogen (N) and phosphorus (P) uptake by rice (kg ha^{-1}) as affected by different water and nitrogen management strategies. Note: Means within a column followed by different letters are significantly different at $P < 0.05$. NU, nitrogen uptake; PU, phosphorus uptake.

Residual soil N and P after the rice harvest. As shown in Fig. 4, the amounts of soil $\text{NH}_4^+\text{-N}$, $\text{NO}_3^-\text{-N}$, TN, available P and TP content in the 0–40 cm soil depths were comparable between the FI and SIDS treatments for the three N treatments. For the two irrigation regimes, the content of $\text{NH}_4^+\text{-N}$ in the 0–40 cm soil depths was significantly greater in the U + CRF and ON treatments than that in the CN (Fig. 4A,B). The content of $\text{NO}_3^-\text{-N}$ in the 0–40 cm soil depths was greater in U + CRF than in the other N fertilizer treatments (Fig. 4C,D). The contents of TN and TP in the 0–40 cm soil depths were comparable among three N fertilizer treatments with the exception of TP content in the 0–20 cm soil depth under SIDS (Fig. 4E–H). The U + CRF and ON treatments increased soil available P content compared to FFP under FI and SIDS (Fig. 4I,J).

Yield of rice. As shown in Fig. 5, the SIDS treatments increased yield of rice grains compared with the FI treatments, although a significant difference was only observed from the U + CRF treatment. The U + CRF resulted in a greater yield of grain compared with the other N treatments under SIDS. SIDS coupled with U + CRF resulted in the greatest yield of grain among the different treatments.

Discussion

N and P losses from paddy fields. Alternate wetting and drying irrigation could slightly increase the concentration of N compared with conventional flooding irrigation²¹. Consistently, SIDS resulted in an average of 2.9% higher TN concentration from leachate water compared with FI. This could be related to the smaller water depth of field and irrigation amount under water-saving irrigation, which led to an increase in substrate concentration³⁴.

Earlier studies have shown that the amount of N and P lost through runoff and leaching are closely related to the water management measures^{23,35}. Greater amounts of irrigation lead to more substantial nutrient losses through runoff and leaching from a paddy field²³. In this study, the SIDS treatments significantly reduced the irrigation frequency by 42.3% and amount of irrigation by 41.7%, leading to the lower runoff and leaching water (reduced by 45.8% and 21.9%, respectively). This resulted in the significantly reduced amount of TN and TP lost through runoff and leaching under SIDS (Figs. 2 and 3), which is consistent with the findings of Liang *et al.*³⁶ derived from alternate wetting and drying irrigation. This confirms the substantial water-saving and environmental protection effects of SIDS^{13,14}. The possible reasons include the fact that field water depth is an important factor that contributes to occurrence of surface runoff and deep leaching from paddy fields^{2,20,21}. A high field water depth results in more surface runoff when faced with concentrated or heavy precipitation². SIDS significantly decreased the field water depth compared with FI (data not shown) and thereby decreased the occurrence and amount of surface runoff (Fig. 1), resulting in the increased use of precipitation and reduction in losses of N and P. Moreover, the low water level and large air exchange capacity at water-soil interface under SIDS helped to increase the P fixation by soil. Fe^{2+} is oxidized to Fe^{3+} , and then Fe^{3+} easily combines with PO_4^{3-} to form insoluble $\text{Fe}(\text{PO}_4)$ ³⁷. In addition, Liang *et al.*³⁶ suggested that alternate wetting and drying irrigation reduced the irrigation water, runoff water and TN loss through runoff by 13.4–27.5%, 30.2–36.7% and 23.3–30.4%, respectively, in comparison with FI, which were lower than the corresponding values obtained from this study. This could be related to the ability of SIDS to more effectively use precipitation compared with alternate wetting and drying irrigation¹³, thereby enhancing the WUE and reducing losses of nutrients. In addition, the reduced frequency of irrigation results in a more economical use of the labor force³⁸.

An earlier study has shown that there was a substantial difference in the loss of TN through runoff during rice growing season, which ranged from 0.5 to 54.3 kg ha^{-1} ³⁹. The variation was associated with differences in precipitation, soil types, crop growth conditions and management of irrigation water and mineral fertilizers^{23,36}. In this study, the loss of TN through runoff during the rice grown season was 2.90–6.07 kg ha^{-1} among different treatments, which are relatively low levels³⁹. Nevertheless, our measured losses are within the range of those in other studies that quantified losses of N from paddy fields^{21,23}. This was primarily attributed to the low total runoff volume (105.6–194.8 mm). Moreover, the time of runoff events was long behind N fertilization (>8 d), resulting in the low N concentration in the surface water of paddy fields. It has been shown that the amount of N lost through runoff can be significantly reduced when the runoff occurred one week after the N fertilization³⁴.

In this study, the U + CRF and ON treatments reduced loss of TN through runoff by 19.7–29.2% and 25.4–51.7%, respectively, and loss of TN through leaching by 15.1–25.2% and 20.9–26.4%, respectively, in comparison with the CN treatment (Fig. 2). These results are consistent with the findings of Yang *et al.*²³ and Ji *et al.*⁴⁰ Because controlled-release N fertilizer has the characteristics of “peak cutting and valley filling,” namely its N release




Figure 4. Soil $\text{NH}_4^+\text{-N}$, $\text{NO}_3^-\text{-N}$, total nitrogen, available phosphorus and total phosphorus content in the 0–40 cm soil depths after rice harvest as affected by different water and nitrogen management strategies. Note: Values (mean \pm standard error, $n = 3$) within the same soil depth and item followed by different letters are significantly different at $P < 0.05$.

amount is not too much at the early growth stages and not too small at the later growth stages of rice^{30,41}, this effect was conducive to improving the metabolism of N in rice⁴⁰, which corresponded to the increased N uptake (Table 4) and decreased TN concentration from leaching and runoff water. Consequently, this reduced the loss of N. In terms of the ON, numerous studies have shown that the optimal application of fertilizer can reduce concentration of nutrients from source^{42,43}, which is consistent with our results. Moreover, an earlier study has shown that TN leaching increases in a significantly linear fashion in parallel with the increase in rate of application of N




Figure 5. Grain yield of rice as affected by different water and nitrogen management strategies. Note: Values (mean \pm standard error, $n = 3$) by different letters are significantly different at $P < 0.05$.

in a typical open field for vegetable planting²⁶. A 40% reduction in traditional N rate of synthetic fertilizer could reduce amount of NO_3^- -N leaching by 39.6%⁴⁴. Thus, these results strongly suggest that appropriately reducing fertilizer inputs can be an efficient approach to reduce loss of N from paddy fields. In addition, both the U + CRF and ON treatments could partially avoid adverse effects of plum-rain season owing to relatively low concentration of N in paddy water, resulting in a reduction in loss of N. Most dramatically, SIDS coupled with the U + CRF or ON treatments achieved the smallest N and P losses through runoff and leaching, suggesting that SIDS coupled with U + CRF or ON can help reduce the N and P losses through runoff and leaching from paddy fields.

The U + CRF and ON treatments reduced loss of TP through leaching by 18.4–24.5% and 20.4–31.6%, respectively, compared with that of the CN treatment (Fig. 3). This was attributed to the lower concentration of P in the leachate water. It has been shown that the concentration of P in leachate water increased with increase in rate of N fertilization²¹. Since the N fertilization could occupy the soil colloid or iron and aluminum oxide surface adsorption and decrease the ability of soil to absorb P, this type of fertilization could result in the enhancement of dissolution of P and its release into water⁴⁵. In the U + CRF treatment, the controlled release N fertilizer enabled the process of release of N to become slower and longer compared to that of CN⁴⁶. In the ON treatment, the rate of N fertilizer input was reduced by approximately 17%. As a result, both the U + CRF and ON treatments resulted in a lower concentration of N in leachate water. In addition, Peng *et al.*³⁰ found that controlled-release N fertilizer reduced the loss of TN through leaching by 53.6%, which was far greater than the corresponding data in this study, which was 18.4–24.5%. This was attributed to the N rate of controlled release N fertilizer treatment being far smaller than that of standard fertilization by the farmer (180 vs 403 kg N ha⁻¹) in Peng's study³⁰, but the same N rate was used for the U + CRF and CN treatments (180 vs 180 kg N ha⁻¹).

In this study, NH_4^+ -N was the major component of loss of TN through leaching (Table 3). This is consistent with the findings of Peng *et al.*³⁰ and Ji *et al.*⁴⁷. Owing to long-term flooding and the high groundwater table in paddy fields at Jiangnan Plain (less than 150 cm below the surface), the paddy soil is maintained in a reduced state²³. The anaerobic environment inhibits the activity of autotrophic nitrifying bacteria, resulting in limited soil nitrification. Moreover, ammonification, denitrification, and biological nitrogen fixation are the three primary forms of N transformation in flooded soil, with the result that most inorganic N exists in the form of NH_4^+ -N^{21,48}. Moreover, PP was the major component of TP loss through runoff (Table 2). This is consistent with the findings of Liang *et al.*⁴⁹ and Ye *et al.*²⁰. Because precipitation or irrigation would impact the soil surface, which causes a substantial amount of PP in the soil to move to paddy water and consequently be lost through runoff. However, DP was the major component of loss of TP through leaching (Table 3). This was attributed to the fertility of paddy soil in Jiangnan Plain and the generally high contents of clay and organic matter in soil²⁷, leading to the strong adsorption and filtration function of PP in percolating water.

N and P uptake by rice and rice yield. The optimization of water management can realize the purpose of promoting effect of fertilizer with water, which is of substantial importance on the efficiency of improvement of water and fertilizer use to achieve a stable and high yield of rice⁵⁰. In this study, the SIDS treatments resulted in a higher uptake of N and P at maturity and yield of rice grain compared with those of FI treatments (Table 4, Fig. 5). This can be explained as follows: SIDS enhanced the air exchange between soil and the atmosphere owing to the alternate wetting and drying cycles^{11,51}. Therefore, the root system was surrounded by relatively sufficient oxygen to accelerate mineralization of soil organic matter and inhibit soil N immobilization, resulting in the increased soil available nutrients for rice growth^{52,53}. It has been shown that water-saving irrigation resulted in higher activities of glutamine synthetase, glutamate synthase, and glutamate dehydrogenase (the main enzymes involved in plant N metabolism) compared to conventional flooded irrigation⁷. Therefore, the root growth, N metabolism, and photosynthetic rate in leaves of rice were improved under SIDS, leading to a high yield of grain^{54,55}.

Optimal fertilization measures help to improve the uptake of nutrients, yield of grain, conserve soil nutrients and reduce excessive nutrient loss to prevent water eutrophication¹⁹. In this study, the U + CRF and ON treatments resulted in comparable TN and TP contents in the 0–40 cm soil depths after rice harvest (Fig. 4) and resulted in a better or comparable uptake of N and P at the latter growth stages (Table 4). These results indicate that the U + CRF and ON treatments improved uptake of N and P by rice and did not lead to excessive accumulation of TN and TP in soil. Similar results were also reported from other studies^{21,35}. The increased uptake of N and P can be explained as follows: in the U + CRF treatment, the use of controlled-release N fertilizer prolonged

the amount of soil available N at latter growth stages of rice, which is conducive to promote effective panicles per m² and spikelets per panicle, and thereby enhance the uptake of N². In the ON treatment, postponing and decreasing the application of N fertilizer reduces the transfer rate of N in leaves and delays leaf senescence, resulting in a high accumulation of N in rice⁵⁶. In addition, approximately reducing the rate of fertilizer usage could enhance uptake of N and P by rice⁵⁷. The relatively lower TN and TP losses through leaching under ON (Fig. 3) supported this conclusion. Thus, it is no surprise that the U + CRF treatment significantly increased yield of grain, and the ON treatment maintained yield of grain compared with that of CN treatment (Fig. 5). Higher grain yield obtained under controlled release N fertilizer than under urea with an equivalent rate of N had been reported previously^{2,58}. Most dramatically, compared to FI coupled with CN, SIDS coupled with U + CRF or ON resulted in a better or comparable uptake of N and P (Table 4) and yield of rice grains (Fig. 5). These results suggest that SIDS coupled with U + CRF or ON was effective at improving or maintaining the N and P uptake and grain yield of rice.

This study strongly illustrates that water-saving irrigation can reduce the losses of N and P from runoff and leaching from paddy fields owing to decreasing amounts of runoff and water. Controlled release N fertilizer and optimized and reduced N fertilization reduce the losses of N and P by lowering concentrations of N and P in the runoff and leaching water. Thus, the combination of SIDS and U + CRF or ON obviously reduced N and P losses during rice grown season. In the future, the mechanism of reducing N and P losses from SIDS coupled with new N management should be considered from perspective of soil N and P cycles.

Conclusions

This study demonstrated that SIDS enabled the paddy field to receive fewer irrigation frequencies (a decrease of 42.3%) and less irrigation water (savings of 41.7%) while using more water from precipitation (an increase of 16.2%), resulting in the reduced total amount of surface runoff and leaching water by 45.8% and 21.9%, respectively. Consequently, SIDS significantly reduced the losses of TN and TP through runoff and leaching. The U + CRF and ON treatments generated a lower loss of TN through runoff and leaching and loss of TP through leaching compared with that of CN treatment. The combined SIDS and U + CRF or ON reduced the N and P loss through runoff and leaching and enhanced or maintained N and P uptake at the later growth stages of rice, subsequently improving the yield of rice grains.

Received: 8 January 2020; Accepted: 27 May 2020;

Published online: 16 June 2020

References

1. Fageria, N. K. Yield physiology of rice. *J. Plant Nutri.* **30**, 843–879 (2007).
2. Ye, Y. *et al.* Alternate wetting and drying irrigation and controlled-release nitrogen fertilizer in late-season rice. Effects on dry matter accumulation, yield, water and nitrogen use. *Field Crops Res.* **144**, 212–244 (2013).
3. Nie, L. & Peng, S. Rice Production in China. In: Chauhan B., Jabran K., Mahajan G. (eds) *Rice Production Worldwide*. *Cham: Springer* **14**, 33–52 (2017).
4. Cai, H. & Chen, Q. Rice research in China in the early 21st century. *Chin. Rice Res. News.* **8**, 14–16 (2000).
5. Dou, X. China's inter-basin water management in the context of regional water shortage. *Sustain. Water Res. Manage.* **4**, 519–526 (2018).
6. Li, H. & Li, M. Sub-group formation and the adoption of the alternate wetting and drying irrigation method for rice in China. *Agric. Water Manage.* **97**, 700–706 (2010).
7. Sun, Y. *et al.* The effects of different water and nitrogen managements on yield and nitrogen use efficiency in hybrid rice of China. *Field Crops Res.* **127**, 85–98 (2012).
8. Morison, J. L., Baker, N. R., Mullineaux, P. M. & Davies, W. J. Improving water use in crop production. *Philos. T. Roy. Soc. B.* **363**, 639–658 (2008).
9. Yao, F. *et al.* Agronomic performance of high-yielding rice variety grown under alternate wetting and drying irrigation. *Field Crops Res.* **126**, 16–22 (2012).
10. Belder, P. *et al.* Effect of water-saving irrigation on rice yield and water use in typical lowland conditions in Asia. *Agric. Water Manage.* **65**, 193–210 (2004).
11. Mao, Z. *et al.* Irrigation and drainage technology of high efficient saving water and continue high yield for rice production. *Water Res. Hydro.* **33**, 65–67 (2002).
12. Zhang, Z., Zhang, S., Yang, J. & Zhang, J. Yield, grain quality and water use efficiency of rice under non-flooded mulching cultivation. *Field Crops Res.* **108**, 71–81 (2008).
13. Shao, D., Li, Y., Yang, P. & Gao, D. Analysis on nitrogen utilization and environmental effects under water-saving irrigation in paddy field. *J. Hydraulic Eng.* **46**(2), 146–152 (2015).
14. Shao, D., Qiao, X., Liu, H., Chai, M. & Wang, J. Study of moving and transformation law of water and fertilizer under different treatments of irrigation and drainage. *Eng. J. Wuhan Uni.* **43**(4), 409–418 (2010).
15. Qiao, X., Shao, D., Liu, H. & Yuan, J. Study on the moving and transforming law of N and P under water-saving irrigation and controlled drainage. *J. Hydraulic Eng.* **42**(7), 862–867 (2011).
16. Spiertz, J. H. J. Nitrogen, sustainable agriculture and food security. A review. *Agronomy Sustain. Dev.* **30**, 43–55 (2010).
17. Peng, S. *et al.* Research strategy in improving fertilizer-nitrogen use efficiency of irrigated rice in China. *Sci. Agric. Sin.* **35**(9), 1095–1103 (2002).
18. Peng, S. *et al.* Strategies for overcoming low agronomic nitrogen use efficiency in irrigated rice systems in China. *Field Crops Res.* **96**, 37–47 (2006).
19. Ju, X. *et al.* Reducing environmental risk by improving N management in intensive Chinese agricultural systems. *PANS.* **106**, 3041–3046 (2009).
20. Ye, Y. *et al.* Effects of different water and nitrogen managements on phosphorus loss via runoff and leaching from paddy fields in Taihu Lake basin. *Acta Sci. Circ.* **35**(4), 1125–1135 (2015).
21. Ye, Y. *et al.* Effects of water-saving irrigation and controlled-release fertilizer application on nitrogen leaching loss of paddy soil in Taihu Lake region. *Acta Sci. Circ.* **35**(1), 270–279 (2015).
22. Li, S. An approach to accelerating innovative development of the lake science in China. *Bull. Chin. Acad. Sci.* **21**(5), 399–405 (2006).
23. Yang, S., Peng, S., Xu, J., He, Y. & Wang, Y. Effects of water saving irrigation and controlled release nitrogen fertilizer managements on nitrogen losses from paddy fields. *Paddy Water Environ.* **13**, 71–78 (2013).

24. Pathak, H. Mitigating greenhouse gas and nitrogen loss with improved fertilizer management in rice: quantification and economic assessment. *Nutri. Cyc. Agro.* **87**(3), 443–454 (2010).
25. Zhang, B. *et al.* Reducing nitrogen leaching in a subtropical vegetable system. *Agric. Ecosys. Environ.* **241**, 133–141 (2017).
26. Wang, D. *et al.* Effects of nitrogen fertilizer and water management practices on nitrogen leaching from a typical open field used for vegetable planting in northern China. *Agri. Water Manage.* **213**, 913–921 (2019).
27. Han, B. *et al.* Effects of fertilization technology of nitrogen (N) on grain yield, grain quality and N-use efficiency of midseason rice in Hubei Province. *Sci. Agric. Sin.* **44**(4), 842–850 (2011).
28. Wu, Q., Zhu, J., Yan, J. & Huang, C. Morphology of middle-season hybrid rice in Hubei Province and its yield under different waterlogging stresses. *Chinese J. Agr. Meteor.* **37**(2), 188–198 (2016).
29. Hu, T., Kang, S., Li, F. & Zhang, J. Effects of partial root-zone irrigation on the nitrogen absorption and utilization of maize. *Agri. Water Manage.* **96**, 208–214 (2009).
30. Peng, S., Yang, S., Xu, J., Luo, Y. & Hou, H. Nitrogen and phosphorus leaching losses from paddy fields with different water and nitrogen managements. *Paddy Water Environ.* **9**, 333–342 (2011).
31. Si, G. *et al.* Effects of an integrated rice-crayfish farming system on soil organic carbon, enzyme activity, and microbial diversity in waterlogged paddy soil. *Acta Eco. Sin.* **38**, 29–35 (2018).
32. Bao, S. Soil and agricultural chemistry analysis. Edition III. Beijing: Chinese Agriculture Press (2000).
33. Zhu, S., Sun, A., Zhang, Z., Wang, Z. & Du, P. Study on rainfall utilization rate of rice under different water-saving irrigation methods. *J. Irrigation Drain.* **32**(5), 1–13 (2013).
34. Zhao *et al.* Nitrogen runoff dominates water nitrogen pollution from rice-wheat rotation in the Taihu Lake region of China. *Agri. Ecology Environ.* **156**, 1–11 (2012).
35. Ye, Y. *et al.* Dynamic variation and runoff loss of nitrogen in surface water of paddy field as affected by water-saving irrigation and controlled-release fertilizer application. *J. Soil Water Conservation.* **28**(5), 105–112 (2014).
36. Liang, X. *et al.* Mitigation of nutrient losses via surface runoff from rice cropping systems with alternate wetting and drying irrigation and site-specific nutrient management practices. *Env. Sci. Pollution Res.* **20**, 6980–6991 (2013).
37. Zhang, Z. *et al.* Improving water management practices to reduce nutrient export from rice paddy fields. *Environ. Tech.* **32**(2), 197–209 (2011).
38. Rejesus, R. M., Palis, F. G., Rodriguez, D. G. P., Lampayan, R. M. & Bouman, B. A. M. Impact of the alternate wetting and drying (AWD) water-saving irrigation technique: evidence from rice producers in the Philippines. *Food Policy.* **36**, 280–288 (2011).
39. Zhu, L. *et al.* Effects of different tillage and straw return on nitrogen and phosphorus runoff loss from paddy fields. *J. Soil Water Cons.* **26**(6), 6–10 (2012).
40. Ji, X., Zhang, S., Lu, Y. & Liao, Y. Dynamics of floodwater nitrogen and its runoff loss, urea and controlled release nitrogen fertilizer application regulation in rice. *Sci. Agri. Sin.* **39**(12), 2521–2530 (2006).
41. Chien, S. H., Prochnow, L. I. & Cantarella, H. Recent developments of fertilizer production and use to improve nutrient efficiency and minimize environmental impacts. *Adv. Agronomy* **102**, 267–322 (2009).
42. Malhi, S., Zentner, R. & Herier, K. Effectiveness of alfalfa in reducing fertilizer N input for optimum forage yield, protein concentration, returns and energy performance of brome-grass-alfalfa mixtures. *Nutri. Cyc. Agri.* **62**, 219–227 (2002).
43. Wang, E., Bell, M., Luo, Z., Moody, P. & Probert, M. Modelling crop response to phosphorus inputs and phosphorus use efficiency in a crop rotation. *Field Crop Res.* **155**, 120–132 (2014).
44. Min, J., Zhang, H. & Shi, W. Optimizing nitrogen input to reduce nitrate leaching loss in greenhouse vegetable production. *Agri. Water Manage.* **111**, 53–59 (2012).
45. Wu, J. *et al.* Study on rice and dynamics of nitrogen and phosphorus in surface water of paddy field under different fertilizations in Xiaoxi river Basin. *Soil.* **45**(2), 207–213 (2013).
46. Jat, R. A. *et al.* Recent approaches in nitrogen management for sustainable agricultural production and eco-safety. *Arch. Agronomy Soil Sci.* **58**, 1033–1060 (2012).
47. Ji, X., Zheng, S., Shi, L. & Liu, Z. Systematic studies of nitrogen loss from paddy soils through leaching in the Dongting Lake area of China. *Pedosphere* **21**(6), 753–762 (2011).
48. Tian, Y. *et al.* Nitrogen runoff and leaching losses during rice-wheat rotations in Taihu Lake region, China. *Pedosphere* **17**(4), 445–456 (2007).
49. Liang, X., Tian, G., Li, H., Chen, Y. & Zhu, S. Study on characteristic of nitrogen and phosphorus loss from rice field by natural rainfall runoff. *J. Soil Water Cons.* **19**(1), 59–63 (2005).
50. Chu, G. *et al.* Effects of interaction between irrigation regimes and nitrogen rates on rice yield and water and nitrogen use efficiencies. *Chin. J. Rice Sci.* **31**(5), 513–523 (2017).
51. Tan, X. *et al.* Effects of alternate wetting and drying irrigation on percolation and nitrogen leaching in paddy fields. *Paddy Water Environ.* **11**, 1–15 (2013).
52. Bouman, B.A.M., Lampayan, R.M. & Tuong, T.P. Water management in irrigated rice: Coping with water scarcity. *International Rice Research Institute, Los Banos, Philippines* pp. 19–46 (2007).
53. Dong, N. M. *et al.* Effects of alternating wetting and drying versus continuous flooding on fertilizer nitrogen fate in rice fields in the Mekong Delta, Vietnam. *Soil Biol. Biochemistry* **47**, 166–174 (2012).
54. Zhang, H., Xue, Y., Wang, Z., Yang, J. & Zhang, J. An alternate wetting and moderate soil drying regime improves root and shoot growth in rice. *Crop Sci.* **49**, 2246–2260 (2009).
55. Qi, D., Hu, T., Song, X. & Zhang, M. Effect of nitrogen supply method on root growth and grain yield of maize under alternate partial root-zone irrigation. *Sci. Rep.* **9**, 8191 (2019).
56. Ma, W. *et al.* Effects of postponing and decreasing nitrogen application on nitrogen use efficiency and yield of Super rice 'Jijing 88'. *J. Northwest Agri. Sci.* **41**(1), 23–27 (2016).
57. Qi, D., Yan, J. & Zhu, J. Effect of a reduced fertilizer rate on the water quality of paddy fields and rice yields under fishpond effluent irrigation. *Agri. Water Manage.* **231**, 105999 (2020).
58. Kiran, J. K., Khanif, Y. M., Amminuddin, H. & Anuar, A. R. Effects of controlled release urea on the yield and nitrogen nutrition of flooded rice. *Comm. Soil Sci. Plant Anal.* **41**, 811–819 (2010).

Acknowledgements

We are grateful to the National Key Research and Development Project (2016YFD0800503), the Innovation System Found Project of Ecological and Circular Agriculture in Hubei Province (2018skjcx01) and the Engineering Research Center of Ecology and Agricultural Use of Wetland, Ministry of Education, China (KFT201904) for providing funding in the form of research grants.

Author contributions

D.L.Q. and Q.X.W. were involved in acquisition and analysis of data for the work. J.Q.Z. was involved in the design of the work. D.L.Q. wrote this paper. J.Q.Z. revised it critically for important intellectual content. All authors approved the submitted and final versions.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to J.Z.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2020