

## Original Article

# The evaluation of antibiotic resistance genes of *Arcobacter butzleri* isolated from animal products, and chicken slaughterhouse sewage in Mazandaran province, Northern Iran

Pourabbasgholi, Z.<sup>1</sup>; Kaboosi, H.<sup>2\*</sup>; Ghane, M.<sup>3\*\*</sup>; Khoshbakht, R.<sup>4</sup> and Ghiamirad, M.<sup>5</sup>

<sup>1</sup>Ph.D. Student in Microbiology, Department of Microbiology, Faculty of Basic Sciences, Ayatollah Amoli Branch, Islamic Azad University, Amol, Iran; <sup>2</sup>Department of Microbiology, Faculty of Basic Sciences, Ayatollah Amoli Branch, Islamic Azad University, Amol, Iran; <sup>3</sup>Department of Microbiology, Faculty of Basic Sciences, Tonekabon Branch, Islamic Azad University, Tonekabon, Iran; <sup>4</sup>Department of Pathobiology, Faculty of Veterinary Medicine, Amol University of Special Modern Technologies, Amol, Iran; <sup>5</sup>Department of Microbiology, Faculty of Basic Sciences, Ahar Branch, Islamic Azad University, Ahar, Iran

\*Correspondence: H. Kaboosi, Department of Microbiology, Faculty of Basic Sciences, Ayatollah Amoli Branch, Islamic Azad University, Amol, Iran. E-mail: h.kaboosi@iauamol.ac.ir

\*\*Co-correspondence: M. Ghane, Department of Microbiology, Faculty of Basic Sciences, Tonekabon Branch, Islamic Azad University, Tonekabon, Iran. E-mail: masood.ghane@iau.ac.ir



10.22099/ijvr.2024.48879.7150

(Received 19 Nov 2023; revised version 2 Aug 2024; accepted 18 Sept 2024)

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

## Abstract

**Background:** *Arcobacter butzleri*, the most common genus of the Campylobacter family, is considered an emerging zoonotic pathogen. **Aims:** This study aimed to evaluate *A. butzleri* from diverse sources, in order to determine the antibiotic resistance pattern of isolates and the frequency of some genes responsible for their antibiotic resistance. **Methods:** In this study, 425 samples were collected from different sources (chicken slaughterhouse sewage, poultry meat, beef, sheep meat, dairy products) during different seasons of 2020-2021. Suspicious colonies were confirmed using biochemical tests. Furthermore, the polymerase chain reaction technique was used to confirm the phenotypic results using the *16S rRNA* gene. The antibiotic resistance pattern of the isolates to 16 antibiotics were determined using the disk diffusion method. Also, the minimum inhibitory concentration (MIC) of their growth was detected using the tube dilution method in the presence of tetracycline, erythromycin, and gentamicin. **Results:** A total of 53 isolates of *A. butzleri* (12.5%) were isolated from (chicken slaughterhouse sewage=36, poultry meat=8, beef=4, sheep meat=5), which contain all three antibiotic resistance genes of *abu\_0814* (90.57%), *OXA\_464* (100%), and *gyrA* (83.02%). The findings of the present investigation showed the presence of *A. butzleri* in different sources and the high prevalence of antimicrobial resistance in the isolates. Nineteen isolates (36%) have extensive drug resistance and 34 isolates (64%) showed multi-drug resistance to the used antibiotics. **Conclusion:** The elevated level of antibiotic resistance observed in *A. butzleri* isolates originating from various samples suggests a significant use of antibiotics and a prevalent environmental contamination.

**Key words:** Animal products, Antimicrobial resistance, *Arcobacter butzleri*, Extensive drug resistance, Multi-drug resistance

## Introduction

The genus *Arcobacter* was included in the family Campylobacteraceae in 1991 (Vandamme *et al.*, 1991). *Arcobacter* has undergone a restructuring, resulting in its division into six distinct genera: *Arcobacter*, *Aliarcobacter*, *Halarcobacter*, *Malacibacter*, *Poseidonibacter*, and *Pseudarcobacter* (Pérez-Cataluña *et al.*, 2019a). Among the twenty-nine officially recognized *Arcobacter* species, they are now spread across these newly defined genera, with the species *Arcobacter butzleri* being reclassified as *Aliarcobacter butzleri* within the genus *Aliarcobacter* (now *Aliarcobacter butzleri*) belonging to the genus

*Aliarcobacter* (Pérez-Cataluña *et al.*, 2019b; Oren and Garrity, 2020). *A. butzleri* typically results in a self-limiting acute enteritis characterized by watery diarrhea, nausea, abdominal pain, and occasionally fever in individuals (Collado and Figueras, 2011; Ramees *et al.*, 2017). The diarrhea is a result of the disruption of epithelial barrier function caused by alterations in tight-junction proteins and the stimulation of epithelial apoptosis (Bucker *et al.*, 2009). An extensive research conducted over an extended period revealed that *A. butzleri* ranks as the fourth most prevalent Campylobacter-like microorganism identified in human fecal samples (Vandenberg *et al.*, 2004). In exceptional instances, *A. butzleri* might lead to bacteremia. In the

case of animals, *A. butzleri* has been linked to the occurrence of enteritis and/or diarrhea in pigs, cattle, and horses; nevertheless, it has also been observed in the fecal matter of healthy animals (Van Driessche *et al.*, 2003; Atabay *et al.*, 2008). *A. butzleri* stands out as the most predominant *Aliarcobacter* spp. identified in food (Lehner *et al.*, 2005). These bacteria have been recovered from animal-derived items like poultry meat, pork, and beef, as well as water sources, dairy products, milk, shellfish, and vegetables (Arias *et al.*, 2011; Yesilmen *et al.*, 2014; Fanelli *et al.*, 2019). The ingestion of tainted food or water represents the most likely pathway of transmission to both humans and animals (Shah *et al.*, 2012; Ferreira *et al.*, 2016). Contact with domestic animals can serve as a potential route of transmission to humans, given that *A. butzleri* has been detected in the oral cavity of felines (Fera *et al.*, 2009; Ramees *et al.*, 2017). Consequently, *A. butzleri* is not solely an emerging pathogen responsible for foodborne illnesses and zoonoses, but it is also deemed a significant threat to human well-being (ICMSF, 2018; Chieffi *et al.*, 2020). Nevertheless, the modes of horizontal and vertical transmission in animals have been documented (Ho *et al.*, 2006). Previous research has predominantly focused on investigating the antibiotic susceptibility of *A. butzleri* phenotypically; however, in the last few years, the underlying antimicrobial resistance (AMR) genes have also been studied more closely (Parisi *et al.*, 2019; Isidro *et al.*, 2020). The excessive use of antimicrobial drugs in animal husbandry and the transfer of contamination through the release of wastewater have led to the evolution and spread of drug-resistant isolates. They have also caused the transfer of resistance genes to other medically important Gram-negative bacteria (Isidro *et al.*, 2020). The increase in multidrug-resistant (MDR) strains, which are resistant to three or more categories of antimicrobial agents, occurs as a consequence of the incorrect use of antibiotics for either preventive purposes or the treatment of infectious diseases. However, there are limited studies on frequency, antibacterial susceptibility, and antibiotic-resistance genes in *Arcobacter* strains (Atabay and Corry, 1997). The present study aimed to isolate and identify *A. butzleri* from different sources, determining the antibiotic resistance pattern of isolates of this bacterium to different antibiotics and the frequency of some genes responsible for antibiotic resistance.

## Materials and Methods

### Samples of collection

In this study, 425 samples were collected from different sources, including chicken slaughterhouse sewage (297 samples), poultry meat (20 samples), beef (20 samples), sheep meat (20 samples), and dairy products (68 samples) during different seasons of 2020-2021. For sampling, sterile swabs (sterilized at 121°C for 15 min) were used. All the collected samples were placed in a container filled with ice. Finally, samples were transferred at 4°C to the laboratory of microbiology

in Islamic Azad University, Tonekabon Branch, Iran.

### Isolation of bacteria from collected samples

*Arcobacter* spp. was isolated using the standard culture technique (Atabay and Corry, 1997; Fanelli *et al.*, 2019; Kim *et al.*, 2019). To isolate bacteria, different samples were enriched in Preston broth containing Preston supplement (Biomark, India). Incubation was carried out at a temperature of 25°C for 24 h. Subsequently, the cultures were passed through filters with 0.45 µm pore size in completely sterile conditions. After 48 h, the filtered samples were cultured on the surface of CAMPY agar (Merck, Germany) containing defibrinated sheep blood and supplemental antibiotics such as Vancomycin and Polymyxin. Then, the cultures were incubated at 25°C for 48-72 h (Fanelli *et al.*, 2019). The isolates were identified using Gram staining, motility test, growth at different temperatures and under aerobic and microaerophilic conditions, growth in Macconkey agar (Merck, Germany), and common biochemical tests such as catalase, oxidase, and urease tests. After the final confirmation, the isolated bacteria were kept at -20°C.

### Evaluation of antibiotic susceptibility of isolates

To determine the pattern of antibiotic resistance of *A. butzleri* isolated from different samples, the disk diffusion test (Hudzicki, 2009) was used based on the standards of the Clinical and Laboratory Standards Institute (CLSI) (Humphries *et al.*, 2021). Briefly, isolates were cultured on Preston broth at 25°C for 48 h. Then, a microbial suspension with a concentration of half McFarland was made using physiological serum. The prepared suspension was densely cultured using an L-shaped rod on the surface of Mueller Hinton agar (Merck, Germany). After placing the antibiotic discs on the surface of the culture medium, the plates were incubated at 25°C for 24-48 h. Finally, the diameter of the bacterial growth inhibition zone around each disc was measured using a caliper and the results were interpreted using CLSI standard tables.

Sixteen antibiotic disks of Penicillin (PEN, 10 u), Oxacillin (OXA, 10 µg), Amikacin (AMK, 30 µg), Ampicillin (AMP, 10 µg), Amoxicillin (AMX, 25 µg), Amoxicillin/Clavulanic acid (AMC, 20/10 µg), Cephalothin (KF, 30 µg), Chloramphenicol (CHL, 30 µg), Erythromycin (ERY, 15 µg), Gentamicin (GEN, 10 µg), Nalidixic acid (NAL, 30 µg), Nitrofurantoin (NIT, 300 µg), Tetracycline (TCY, 30 µg), Ciprofloxacin (CIP, 5 µg), Trimethoprim/Sulfamethoxazole (SXT, 1.25/23.75 µg), and Azithromycin (AZM, 15 µg) were selected to perform this study. All antibiotics were purchased from Padtan Teb Company (Iran). After the selection and screening of *A. butzleri* strains, the minimum inhibitory concentration against Tetracycline, Erythromycin, and Gentamicin antibiotics was determined by broth macrodilution tube method based on CLSI standards.

### Design of primer

Three primers of *OXA\_464*, *abu\_0814*, and *gyrA*

were designed using the Gene Runner 6.5 software package. The primers were validated through cross-referencing with the GenBank database to confirm their lack of significant similarity to genetic sequences from non-*Arcobacter* species. The pairs of *16S rRNA* primers were used by Scullion *et al.* (2006) (Table 1).

### Molecular identification of *A. butzleri* strains and antibiotic resistance genes

All isolates were characterized through the employment of polymerase chain reaction (PCR). Bacterial DNA was isolated using a DNA extraction kit for this specific procedure. To do so, bacterial DNA was extracted using a DNA extraction kit (Kiagen Teb Sadra Company, Iran). The *16S rRNA* gene and resistance genes of *OXA\_464*, *abu\_0814*, and *gyrA* were amplified using the specific primers (Table 1).

In addition, each solution of PCR reaction with a final volume of 20  $\mu$ L included 10  $\mu$ L of master mix (Hot Start TaqPolymerase, dNTP, buffer, MgCl), 1  $\mu$ L of forward primer, 1  $\mu$ L of reverse primer, 7  $\mu$ L of distilled water and 1  $\mu$ L of template DNA. For polymerization of the *16S rRNA* gene, initial denaturation of the DNA samples was performed at 94°C for 4 min. Then, 30 cycles of PCR reaction including 94°C for 1 min (denaturation), 58.5°C for 1 min (annealing), and 72°C for 90 s (extension) were performed. The final extension was performed at 72°C for 10 min. The PCR products were taken on 1.5% agarose gel for final analysis. Thus, 5  $\mu$ L of the amplified DNA was transferred into each of the wells embedded in the agarose gel. The voltage was set to 75 V for 40 min. Finally, the gel was transferred into the UV transilluminator (UV Doc-England) to observe the bands of PCR products. Therefore, to ensure the proper amplification of *16S rRNA* and *OXA\_464*, *abu\_0814*, *gyrA* gene in the PCR experiment, the obtained product was sent to Macrogen, South Korea for

sequencing. All sequences data were subjected to BLAST analysis (<http://www.ncbi.nlm.nih.gov/BLAST/>) to definitively identify each respective *16S rRNA* *OXA\_464*, *abu\_0814*, *gyrA* gene amplicon.

### Multiplex PCR conditions and gel electrophoresis

PCR reactions were conducted in a 25  $\mu$ L reaction mixture containing 3  $\mu$ L template DNA, 10  $\mu$ L of the Master mix (Hot Start TaqPolymerase, dNTP, buffer, MgCl), 1  $\mu$ L of each of the primers, and 6  $\mu$ L of distilled water. The PCR reactions were performed in a thermal cycler lab-cycler 48 (lab-cycler, Germany) with the following amplification conditions: a denaturation step at 94°C for 4 min; 35 amplification cycles: denaturation at 94°C for 60 s, annealing at 55°C for 60 s, and extension at 72°C for 90 s; and the final extension step at 72°C for 10 min. The PCR products were analyzed on 1.5% (w/v) TBE-agarose gel using a 100 bp ladder (BIO BASIC, Canada). The band patterns were analyzed in the gel documentation system (UV Doc-England). DNA from reference strains (*A. butzleri*, ATCC 49616) and sterile deionized water was used as positive and negative controls, respectively (Çelik and Otlu, 2020).

## Results

### Isolation and identification of *A. butzleri* from different sources

From 425 samples collected during different seasons of 2021-2020, 53 isolates of *A. butzleri* (12.5%) were detected in different sources (Table 2).

The highest rate of isolation was related to poultry meat (40%). The highest isolation rate of the bacterium from different sources was 45.28% in the fall while the lowest isolation rate (7.55%) was detected in the winter (Table 3).

**Table 1:** The sequence of primers used for the evaluation and identification of *A. butzleri* from different resources



Gene	5' → 3'	Product size (bp)	Reference
<i>Arco</i>	F: AGAGATTAGCCTGTATTGTATC R: TAGCATCCCCGCTTCGAATGA	1200	Scullion <i>et al.</i> (2006)
<i>gyrA</i>	F: TGCTAAAATTGCAGATGTACCA R: AATTCCTTTTTCAGAACTGTACG	212	This study
<i>OXA_464</i>	F: ATTCGCAAATGATGTGGAAC R: TCCCATAATTAAGCTCTTTTAG	362	This study
<i>Abu_0814</i>	F: AGTCGTTAGTTGCAATATCT R: ATTTGAACCTTTGTATCTGG	1026	This study

**Table 2:** Number of *A. butzleri* isolates based on sample type

Sample type	Positive	Negative	Total	Chi-square (X <sup>2</sup> )	P-value
Chicken slaughterhouse sewage	36 (12.1%)	261 (87.9%)	297 (100%)	17.258	0.001
Poultry meat	8 (40.0%)	12 (60.0%)	20 (100.0%)	2.234	0.027
Beef	4 (20.0%)	16 (80.0%)	20 (100.0%)	14.312	0.001
Sheep meat	5 (25.0%)	15 (75.0%)	20 (100.0%)	3.808	0.001
Dairy products	0 (0.0%)	68 (100.0%)	68 (100.0%)	-	-
Total	53 (12.5%)	372 (87.5%)	425 (100.0%)	13.980	0.001

**Table 3:** The frequency of *A. butzleri* isolated in different seasons of the year



Season	Frequency	Chi-square ( $X^2$ )	P-value
Spring	18 (33.96%)	8.554	0.001
Summer	7 (13.21%)		
Fall	24 (45.28%)		
Winter	4 (7.55%)		
Total	53 (100.0%)		

**Fig. 1:** Electrophoresis pattern of PCR for detection of *Arco* gene in *Arcobacter butzleri* isolates. M: Marker 100 bp, C+: Positive control, C-: Negative control, and Lanes 1, 2, 3, and 4: Amplification of *Arco* gene at 1200 bp**Fig. 2:** Electrophoresis pattern of PCR for detection of *OXA\_464* gene in *Arcobacter butzleri* isolates. M: Marker 100 bp, C+: Positive control, C-: Negative control, and Lanes 1, 2, 3, 4 and 5: Amplification of *OXA\_464* gene at 362 bp

### Authentication of *A. butzleri*


The isolates were confirmed as *A. butzleri* using genus-based PCR and species-specific multiplex PCR. The results of PCR and sequencing of *16S rRNA* genes confirmed the phenotypic results obtained in 53 isolates

(Fig. 1).

**Fig. 3:** Electrophoresis pattern of PCR for detection of *abu\_0814* gene in *Arcobacter butzleri* isolates. M: Marker 100 bp, C+: Positive control, C-: Negative control, and Lanes 1, 2, 3 and 4: Amplification of *abu\_0814* gene at 1029 bp**Fig. 4:** Electrophoresis pattern of PCR for detection of *gyrA* gene in *Arcobacter butzleri* isolates. M: Marker 100 bp, C+: Positive control, C-: Negative control, and Lanes 1, 2 and 3: Amplification of *gyrA* gene at 212 bp

### Evaluation of the presence of antibiotic-resistance genes in the isolates

The presence of *OXA\_464*, *abu\_0814*, and *gyrA* genes in *A. butzleri* was investigated using the PCR technique. According to the study results, the presence of the *OXA\_464* gene was observed in all 53 isolates (100%) (Fig. 2). Also, 48 isolates (90.57%) were positive for the *abu\_0814* gene (Fig. 3), and *gyrA* was detected in 44 isolates (83.02%) (Fig. 4). Finally, after examining all antibiotic resistance genes using the single-PCR method, the frequency of *abu\_0814*, *OXA\_464*, and *gyrA* in all isolates were studied by multiplex-PCR technique (Fig. 5).



**Fig. 5:** Electrophoresis pattern of multiplex PCR for detection of *OXA\_464*, *abu\_0814*, and *gyrA* genes in *Arcobacter butzleri* isolates. M: Marker 100 bp, C+: Positive control, C-: Negative control, and Lanes 1, 2 and 3: Amplification of *gyrA*, *OXA\_464*, and *abu\_0814* genes at 212, 362, and 1029 bp, respectively

### Sequencing analysis

The findings derived *16S rRNA*, *OXA\_464*, *abu\_0814*, and *gyrA* gene sequencing of PCR products were clear. There was 100% alignment agreement for all the samples using the BLAST alignment program. As shown in Table 4, BLAST analysis of the *16S rRNA*, *OXA\_464*, *abu\_0814*, *gyrA* gene sequence data for sample 1-4 indicated a 100% homology with the *A. butzleri* strain ED-1. Samples 5 indicated a 99% homology with the *A. butzleri* strain ED-1. Sample 6 was identified using the BLAST analysis of the sequencing data as *A. butzleri* strain ATCC 49616 (98% homology). Sample 7 was identified using the BLAST analysis of the sequencing data as *A. butzleri* strain 7h1h (100% homology). Sample 8 was identified using the BLAST analysis of the sequencing data as *A. butzleri* strain NCTC12481 (98% homology). Sample 9 was identified using the BLAST analysis of the sequencing data as *A. butzleri* strain JCR7715 (98% homology). Sample 10 was identified using the BLAST analysis of the sequencing data as *A. butzleri* strain P1100 (98%

homology). The data concerning the comparison between phenotyping and genotyping for identifying the isolates revealed that the majority of genotyping results supported our phenotyping identification.

The frequency of *abu\_0814*, *OXA\_464*, and *gyrA* genes in the isolates was 90.57%, 100%, and 83.02%, respectively (Table 5).

**Table 5:** Frequency of antibiotic-resistance genes in *A. butzleri* isolates

Gene	Frequency	Chi-square (X <sup>2</sup> )	P-value
<i>OXA_464</i>	53 (100.0%)		
<i>abu_0814</i>	48 (90.57%)	2.013	0.187
<i>gyrA</i>	44 (83.02%)		

### Determining antimicrobial susceptibility pattern

All isolates showed resistance to Penicillin (100%), Ampicillin (100%), and Oxacillin (100%). Also, the levels of resistance to Trimethoprim/Sulfamethoxazole (68%), Ciprofloxacin (78%), Nalidixic acid (74%), Azithromycin (68%), and Amoxicillin (68%) were detected. Out of the 53 isolates tested, all isolates showed susceptibility to Gentamicin (100%) (Table 6). The MIC and MBC of 72% of the isolates to Tetracycline were  $\geq 128$   $\mu\text{g/ml}$  and  $\geq 256$   $\mu\text{g/ml}$ , respectively. All of the isolates (100%) had MIC  $\geq 8$   $\mu\text{g/ml}$  and MBC  $\geq 32$   $\mu\text{g/ml}$  to Gentamicin. Totally, 75% of the isolates had MIC  $\geq 64$   $\mu\text{g/ml}$  and MBC  $\geq 128$   $\mu\text{g/ml}$  to Erythromycin. Also, 19 isolates were detected as extensively drug-resistant (36%) and 34 isolates had multi-drug resistant (64%).

### Discussion

*Arcobacter* species are globally recognized for their role in the etiology of acute gastroenteritis, worldwide (Zhang *et al.*, 2019; Brückner *et al.*, 2020; Chieffi *et al.*, 2020). The primary mode of transmission of *Arcobacter* spp. to humans is through the ingestion of water and food that have been contaminated (Collado and Figueras, 2011; Zambri *et al.*, 2019). Advancements in research regarding the frequency and virulence of *A. butzleri* and *A. cryaerophilus* have resulted in their classification as significant threats to human health by the International Commission on Microbiological Specifications for Foods (ICMSF, 2018).

**Table 4:** Comparison of phenotyping and genotyping identification

Number of samples	Phenotyping detection	Genotyping detection
1	<i>A. butzleri</i>	100% homology with the <i>A. butzleri</i> strain ED-1
2	<i>A. butzleri</i>	100% homology with the <i>A. butzleri</i> strain ED-1
3	<i>A. butzleri</i>	100% homology with the <i>A. butzleri</i> strain ED-1
4	<i>A. butzleri</i>	100% homology with the <i>A. butzleri</i> strain ED-1
5	<i>A. butzleri</i>	99% homology with the <i>A. butzleri</i> strain ED-1
6	<i>A. butzleri</i>	98% homology with the <i>A. butzleri</i> strain ATCC 49616
7	<i>A. butzleri</i>	100% homology with the <i>A. butzleri</i> strain 7h1h
8	<i>A. butzleri</i>	98% homology with the <i>A. butzleri</i> strain NCTC12481
9	<i>A. butzleri</i>	98% homology with the <i>A. butzleri</i> strain JCR7715
10	<i>A. butzleri</i>	98% homology with the <i>A. butzleri</i> strain P1100

**Table 6:** Antibiotic resistance pattern of *A. butzleri*

Antibiotic agent	Dick content	Interpretive categories and zone diameter breakpoint whole mm			R%	COD
		S	I	R		
Ampicillin	10 mg	17≥	14-16	13≤	100 R	AMP
Amoxicillin	25 mg	17≥	14-16	13≤	68 I	AMX
Amoxicillin-clavulanic acid	20/10 mg	17≥	14-17	13≤	75 I	AMC
Gentamycin	10 mg	15≥	13-14	13≤	100 S	GEN
Azithromycin	15 mg	13≥	14-16	12≤	68 R	AZM
Erythromycin	15 mg	23≥	14-22	13≤	75 S	ERY
Chloramphenicol	30 mg	18≥	13-17	15≤	73.5 I	CHL
Tetracycline	30 mg	15≥	12-14	12≤	68 I	TCY
Trimethoprim/Sulfamethoxazole	1.25/23.75 mg	16≥	11-15	10≤	68 R	SXT
Amikacin	30 mg	17≥	14-16	13≤	89 S	AMK
cephalothin	30 mg	23≥	20-22	19≤	57 I	CEP
ciprofloxacin	5 mg	12≥	16-20	15≤	78 R	CIP
nitrofurantoin	300 mg	17≥	15-16	12≤	67 S	NIT
Nalidixic acid	30 mg	19≥	14-18	14≤	74 R	NAL
Oxacillin	1 mg	18≥	14-18	12≤	100 R	OXA
penicillin	10 u	18≥	14-16	12≤	100 R	PEN

R: Resistant, I: Intermediate, and S: Sensitive

This study offers recent data concerning the prevalence and antibacterial resistance patterns of *A. butzleri*. Specimens were collected from diverse origins in Tonekabon, Iran.

Our findings indicated that the frequency of *A. butzleri* was observed, ranging from high to low, in poultry meat (40.0%), sheep meat (25.0%), beef (20.0%), chicken slaughterhouse sewage (12.1%), and dairy products (0.0%). Several studies investigated the presence of *Arcobacter* spp. in chicken meat. Similar to the results of the present study, Amare *et al.* (2011) identified *Arcobacter* spp. in 39% of fresh chicken meat. Di Noto *et al.* (2018) found *Arcobacter* spp. in 53.3% of the wing and carcasses of chickens. In another study conducted in Mexico by Scullion *et al.* (2006), the prevalence of *Arcobacter* spp. in chicken, beef, and raw milk samples were about 35%, 34%, and 46%, respectively. *A. butzleri* was the dominant species isolated from samples (Aski *et al.*, 2016). The difference in the isolation rate of *Arcobacter* spp. may be due to different conditions in the research process including the geographical location, season of study, analysis methods, design of the study, types of used antimicrobial agents and their concentrations in culture mediums.

In the present study, the resistance rate of *Arcobacter* isolates to Ciprofloxacin, Nalidixic acid, and Azithromycin was 78%, 74%, and 68%. The increase in the rate of antibiotic resistance in Iran may be related to the excessive and arbitrary use of antibiotics. Also, the high resistance of pathogenic bacteria to Penicillins is usually associated with the presence of beta-lactamases. In the present study, almost all strains of *Arcobacter* isolates have  $\beta$ -lactam antibiotic resistance genes. It has been speculated that *A. butzleri* may be resistant to  $\beta$ -lactamase. Several studies reported that  $\beta$ -lactam resistance may be caused by the presence of three putative  $\beta$ -lactamases (AB0578, AB1306, and AB1486) identified in the RM4018 genome, which are enhanced by the occurrence of the *lrgAB* operon (ab0179 and ab0180) and may

regulate tolerance to penicillin in *Staphylococcus* (Bayles, 2000; Groicher *et al.*, 2000; Miller *et al.*, 2007).

In the present study, the frequency of *abu\_0814* gene in the isolates was reported as 90.57%. The presence of this gene corresponds to high resistance to beta-lactams. The presence of *abu\_0814* can lead to resistance against beta-lactams in *Arcobacter* spp. and there was an agreement between the presence of this gene and the results obtained from Antibigram and Whonet software.

The increase in the rate of resistance to Penicillin, Ampicillin, Oxacillin, Trimethoprim/Sulfamethoxazole, Nalidixic acid, Ciprofloxacin, and Azithromycin among *A. butzleri* isolates is a cause for concern (Aski *et al.*, 2016). Van den Abeele *et al.* (2016), conducted a study that showed resistance to Ciprofloxacin is attributed to the development of resistance in *Arcobacter* strains due to mutations in the *gyrA* gene. In this study, by examining the frequency of *gyrA* in isolates and its relationship to their resistance to ciprofloxacin, it is possible to understand the relationship between mutations in *gyrA* and increased resistance to ciprofloxacin. Since high resistance levels of isolates to fluoroquinolones were shown in the present study, the results were consistent with the findings of other researchers (Houf *et al.*, 2001). Several studies showed that the base mutation in the *gyrA* gene was associated with a higher level of resistance to ciprofloxacin (Webb *et al.*, 2018; Hodges *et al.*, 2021). In our study, 13 *A. butzleri* and 14 *A. cryaerophilus* isolates with *gyrA* (C254T) mutation were 100% phenotypically resistant to ciprofloxacin. Moreover, the genes *aph* (3')-IIIa and *ant* (6)-Ia were reported to correlate with resistance to kanamycin and streptomycin, respectively (Hormeño *et al.*, 2018; Cho *et al.*, 2020). All isolates showed sensitivity to Amikacin, Gentamicin, Erythromycin, and Nitrofurantoin. Also, 75%, 73.5%, and 68% of the isolates showed sensitivity to Amoxicillin/clavulanic acid, Chloramphenicol, and Tetracycline, respectively. It can be anticipated that the *Arcobacter* spp. is going to gain resistance to tetracycline.

This phenomenon can be attributed to the excessive use of antibiotics in the food chain of poultry due to the selective stress of antibiotics, the unavailability of other antibiotics, and the acquisition of resistance genes in the life cycle of these microorganisms. Resistance to Penicillin and Ampicillin is not surprising because these antibiotics are commonly used in selective media for the isolation of these organisms as well as in the treatment of *Campylobacter* infections (González *et al.*, 2017).

Houf *et al.* (2001) evaluated 47 strains of *A. butzleri* isolated from different sources against several antimicrobial agents and found that most of the tested strains were resistant to Cefoperazone. Thus, specific attention should be paid to the use of Penicillin and Trimethoprim/Sulfamethoxazole, which are commonly used in the treatment of bacterial infections in humans and animals. In the present study, variable sensitivity to amoxicillin, clavulanic acid, chloramphenicol and tetracycline should also be considered when using these antibiotics against *Arcobacter* spp. infections. Although most of the isolates were sensitive to Erythromycin, the existence of resistant isolates should not be ruled out because several isolates showed resistance to this antibiotic. The present study showed that 100% of isolates were sensitive to Gentamicin. The widespread occurrence of antibiotic resistance within *Arcobacter* spp. across various settings could potentially contribute to the dissemination of resistance. It is crucial to collect information on antimicrobial resistance because these drugs are often recommended as the first treatment for infection. Also, data about the antibiotic sensitivity obtained in this study can be used during the design of the study for the isolation of these bacteria. This study showed that the susceptibility of *A. butzleri* to different types of antibiotics is variable.

According to the results of the present study, *A. butzleri* was found in different environmental samples, poultry meat, beef, sheep meat, and the sewage of slaughterhouses. Assessment of antibiotic resistance in the isolates from this study showed that there is relatively high antibiotic resistance and multiple antibiotic resistance in the majority of the isolates. Methods such as microarray expression profiles and proteomic investigations can provide a more comprehensive insight into *Arcobacter* infection.

## Acknowledgement

The authors wish to extend their appreciation to the Islamic Azad University of Tonekabon Branch for the generous supply of the required apparatus.

## Conflict of interest

The authors declare that they have no conflict of interests.

## References

Amare, L; Saleha, A; Zunita, Z; Jalila, A and Hassan, L

- (2011). Prevalence of *Arcobacter* spp. on chicken meat at retail markets and in farm chickens in Selangor, Malaysia. *Food Control*. 22: 732-736.
- Arias, ML; Cid, A and Fernández, H (2011). *Arcobacter butzleri*: first isolation report from chicken carcasses in Costa Rica. *Brazilian J. Microbiol.*, 42: 703-706.
- Aski, HS; Tabatabaei, M; Khoshbakht, R and Raeisi, M (2016). Occurrence and antimicrobial resistance of emergent *Arcobacter* spp. isolated from cattle and sheep in Iran. *Comp. Immunol. Microbiol. Infect. Dis.*, 44: 37-40.
- Atabay, H and Corry, J (1997). The prevalence of *Campylobacters* and *Arcobacters* in broiler chickens. *J. Applied Microbiol.*, 83: 619-626.
- Atabay, HI; Unver, A; Sahin, M; Otlı, S; Elmali, M and Yaman, H (2008). Isolation of various *Arcobacter* species from domestic geese (*Anser anser*). *Vet. Microbiol.*, 128: 400-405.
- Bayles, KW (2000). The bactericidal action of penicillin: New clues to an unsolved mystery. *Trends Microbiol.*, 8: 274-278. doi: 10.1016/S0966-842X(00)01762-5.
- Brückner, V; Fiebiger, U; Ignatius, R; Friesen, J; Eisenblätter, M and Höck, M (2020). Prevalence and antimicrobial susceptibility of *Arcobacter* species in human stool samples derived from out- and inpatients: the prospective German *Arcobacter* prevalence study Arcopath. *Gut Pathog.*, 12: 1-8. doi: 10.1186/s13099-020-00360-x.
- Bucker, R; Troeger, H; Kleer, J; Fromm, M and Schulzke, JD (2009). *Arcobacter butzleri* induces barrier dysfunction in intestinal HT-29/B6 cells. *J. Infect. Dis.*, 200: 756-764.
- Çelik, E and Otlı, S (2020). Isolation of *Arcobacter* spp. and identification of isolates by multiplex PCR from various domestic poultry and wild avian species. *Ann. Microbiol.*, 70: 1-7.
- Chieffi, D; Fanelli, F and Fusco, V (2020). *Arcobacter butzleri*: Up-to-date taxonomy, ecology, and pathogenicity of an emerging pathogen. *Compr. Rev. Food Sci. Food Saf.*, 19: 2071-2109.
- Cho, S; Barrett, JB; Frye, JG and Jackson, CR (2020). Antimicrobial resistance gene detection and plasmid typing among multidrug resistant *Enterococci* isolated from freshwater. *Microorganisms*. 8: 1-15. doi: 10.3390/microorganisms8091338.
- Collado, L and Figueras, MJ (2011). Taxonomy, epidemiology, and clinical relevance of the genus *Arcobacter*. *Clin. Microbiol. Rev.*, 24: 174-192.
- Di Noto, AM; Sciortino, S; Cardamone, C; Ciravolo, C; Napoli, C; Alio, V and Costa, A (2018). Detection of *Arcobacter* spp. in food products collected from Sicilia region: A preliminary study. *Ital. J. Food Saf.*, 7: 2.
- Fanelli, F; Di Pinto, A; Mottola, A; Mule, G; Chieffi, D; Baruzzi, F and Fusco, V (2019). Genomic characterization of *Arcobacter butzleri* isolated from shellfish: novel insight into antibiotic resistance and virulence determinants. *Front. Microbiol.*, 10: 670-672.
- Fera, MT; La camera, E; Carbone, M; Malara, D and Pennisi, MG (2009). Pet cats as carriers of *Arcobacter* spp. in Southern Italy. *J. Appl. Microbiol.*, 106: 1661-1666.
- Ferreira, S; Queiroz, JA; Oleastro, M and Domingues, F (2016). Insights in the pathogenesis and resistance of *Arcobacter*: A review. *Crit. Rev. Microbiol.*, 42: 364-383.
- González, A; Morejón, I and Ferrús, M (2017). Isolation, molecular identification and quinolone-susceptibility testing of *Arcobacter* spp. isolated from fresh vegetables in Spain. *Food Microbiol.*, 65: 279-283.
- Groicher, KH; Firek, BA; Fujimoto, DF and Bayles, K (2000). The *Staphylococcus aureus* *lrgAB* operon



- modulates murein hydrolase activity and penicillin tolerance. *J. Bacteriol.*, 182: 1794-1801. doi: 10.1128/JB.182.7.1794-1801.200.
- Ho, TK; Lipman, LJ; van der Graaf-van Bloois, L; van Bergen, M and Gaastra, W** (2006). Potential routes of acquisition of *Arcobacter* species by piglets. *Vet. Microbiol.*, 114: 123-133.
- Hodges, LM; Taboada, EN; Koziol, A; Mutschall, S; Blais, BW and Inglis, G** (2021). Systematic evaluation of whole-genome sequencing based prediction of antimicrobial resistance in *Campylobacter jejuni* and *C. coli*. *Front. Microbiol.*, 12: 776967. doi: 10.3389/fmicb.2021.776967.
- Hormeño, L; Ugarte-Ruiz, M; Palomo, G; Borge, C; Florez-Cuadrado, D; Vadillo, S and Quesada, A** (2018). ant (6)-I genes encoding aminoglycoside O-nucleotidyltransferases are widely spread among streptomycin resistant strains of *Campylobacter jejuni* and *Campylobacter coli*. *Front. Microbiol.*, 23: 2515-2520.
- Houf, K; Devriese, LA; Zutter, LD; Van Hoof, J and Vandamme, P** (2001). Susceptibility of *Arcobacter butzleri*, *Arcobacter cryaerophilus*, and *Arcobacter skirrowii* to antimicrobial agents used in selective media. *J. Clin. Microbiol.*, 39: 1654-1656.
- Hudzicki, J** (2009). Kirby-Bauer disk diffusion susceptibility test protocol. *ASM.*, 15: 55-63.
- Humphries, R; Bobenchik, AM; Hindler, JA and Schuetz, A** (2021). Overview of changes to the Clinical and Laboratory Standards Institute performance standards for antimicrobial susceptibility testing, M100. *J. Clin. Microbiol.*, 59: 10-1128.
- International Commission on Microbiological Specifications for Foods (ICMSF)** (2018). Microbiological hazards and their control. Microorganisms in foods 7: Microbiological testing in food safety management. PP: 1-30.
- Isidro, J; Ferreira, S; Pinto, M; Domingues, F; Oleastro, M; Gomes, JP and Borges, V** (2020). Virulence and antibiotic resistance plasticity of *Arcobacter butzleri*: Insights on the genomic diversity of an emerging human pathogen. *Infect. Genet. Evol.*, 80: 104213.
- Kim, NH; Park, SM; Kim, HW; Cho, TJ; Kim, SH; Choi, C and Rhee, M** (2019). Prevalence of pathogenic *Arcobacter* species in South Korea: Comparison of two protocols for isolating the bacteria from foods and examination of nine putative virulence genes. *Food Microbiol.*, 78: 18-24.
- Lehner, A; Tasara, T and Stephan, R** (2005). Relevant aspects of *Arcobacter* spp. as potential foodborne pathogen. *Int. J. Food Microbiol.*, 102: 127-135.
- Miller, WG; Parker, CT; Rubenfield, M; Mendz, GL; Wosten, MM and Ussery, DW** (2007). The complete genome sequence and analysis of the epsilonproteobacterium *Arcobacter butzleri*. *PLoS One*. 2: e1358. doi: 10.1371/journal.pone.0001358. doi: 10.3389/fmicb.2018.02515.
- Oren, A and Garrity, G** (2020). List of new names and new combinations previously effectively, but not validly, published. *Int. J. Syst. Evol. Microbiol.*, 70: 2960-2966.
- Parisi, A; Capozzi, L; Bianco, A; Caruso, M; Latorre, L; Costa, A; Giannico, A; Ridolfi, D; Bulzacchelli, C and Santagada, G** (2019). Identification of virulence and antibiotic resistance factors in *Arcobacter butzleri* isolated from bovine milk by whole genome sequencing. *Ital. J. Food Saf.*, 8: 60-64.
- Pérez-Cataluña, A; Salas-Masso, N; Dieguez, AL; Balboa, S; Lema, A; Romalde, JL and Figueras, M** (2018). Revisiting the taxonomy of the genus *Arcobacter*: Getting order from the chaos. *Food Microbiol.*, 4: 2077-2080.
- Pérez-Cataluña, A; Salas-Masso, N; Dieguez, AL; Balboa, S; Lema, A; Romalde, JL and Figueras, M** (2019a). Revisiting the taxonomy of the genus *Arcobacter*: Getting order from the chaos. *Food Microbiol.*, 10: 2253-2261.
- Pérez-Cataluña, A; Salas-Masso, N and Figueras, M** (2019b). *Arcobacter lacus* sp. nov. and *Arcobacter caeni* sp. nov., two novel species isolated from reclaimed water. *Int. J. Syst. Evol. Microbiol.*, 69: 3326-3331.
- Ramees, TP; Dhama, K; Karthik, K; Rathore, RS; Kumar, A; Saminathan, M; Tiwari, R; Malik, YS and Singh, RK** (2017). *Arcobacter*: An emerging food-borne zoonotic pathogen, its public health concerns and advances in diagnosis and control—A comprehensive review. *Vet. Q.*, 37: 136-161.
- Scullion, R; Harrington, CS and Madden, RH** (2006). Prevalence of *Arcobacter* spp. in raw milk and retail raw meats in Northern Ireland. *J. Food Prot.*, 69: 1986-1990.
- Shah, AH; Saleha, AA; Zunita, Z; Cheah, YK; Murugaiyah, M and Korejo, NA** (2012). Genetic characterization of *Arcobacter* isolates from various sources. *Vet. Microbiol.*, 160: 355-361.
- Vandamme, P; Falsen, E; Rossau, R; Hoste, B; Segers, P; Tytgat, R and De Ley, J** (1991). Revision of *Campylobacter*, *Helicobacter*, and *Wolinella* taxonomy: Emendation of generic descriptions and proposal of *Arcobacter* gen. nov. *Int. J. Syst. Bacteriol.*, 41: 88-103.
- Van den Abeele, AM; Vogelaers, D; Vanlaere, E and Houf, K** (2016). Antimicrobial susceptibility testing of *Arcobacter butzleri* and *Arcobacter cryaerophilus* strains isolated from Belgian patients. *J. Antimicrob. Chem.*, 71: 1241-1244.
- Vandenbergh, O; Dediste, A; Houf, K; Ibekwem, S; Souayah, H; Cadranel, S; Douat, N; Zissis, G; Butzler, JP and Vandamme, P** (2004). *Arcobacter* species in humans. *Emerg. Infect. Dis.*, 10: 1863-1867.
- Van Driessche, E; Houf, K; van Hoof, J; De Zutter, L and Vandamme, P** (2003). Isolation of *Arcobacter* species from animal feces. *FEMS Microbiol. Lett.*, 229: 243-248.
- Webb, AL; Selinger, LB; Taboada, EN and Inglis, GD** (2018). Subtype-Specific selection for resistance to fluoroquinolones but not to tetracyclines is evident in *Campylobacter jejuni* isolates from beef cattle in confined feeding operations in Southern Alberta, Canada. *Appl. Environ. Microb.*, 84: 02713-02717. doi: 10.1128/AEM.02713-17.
- Yesilmen, S; Vural, A; Erkan, ME and Yildirim, IH** (2014). Prevalence and antimicrobial susceptibility of *Arcobacter* species in cow milk, water buffalo milk and fresh village cheese. *Int. J. Food Microbiol.*, 188: 11-14.
- Zambri, M; Cloutier, M; Adam, Z; Lapen, DR; Wilkes, G and Sunohara, M** (2019). Novel virulence, antibiotic resistance and toxin gene-specific PCR based assays for rapid pathogenicity assessment of *Arcobacter faecis* and *Arcobacter lanthieri*. *BMC Microbiol.*, 19: 1-15. doi: 10.1186/s12866-018-1357-7.
- Zhang, X; Alter, T and Götz, G** (2019). Characterization of *Arcobacter* spp. isolated from retail seafood in Germany. *Food Microbiol.*, 82: 254-258. doi: 10.1016/j.fm.2019.02.010.