

Notes on two closely related spider species of the *Pholcus phungiformes* species group (Araneae, Pholcidae) from Beijing, China

Xiang Wang^{1*}, Shumaila Shaheen^{1*}, Qiaoqiao He¹, Zhiyuan Yao^{1,2,3}

1 College of Life Science, Shenyang Normal University, Shenyang 110034, Liaoning, China **2** Liaoning Key Laboratory of Evolution and Biodiversity, Shenyang 110034, Liaoning, China **3** Liaoning Key Laboratory for Biological Evolution and Agricultural Ecology, Shenyang 110034, Liaoning, China

Corresponding author: Zhiyuan Yao (yaozy@synu.edu.cn)

Academic editor: Gergin Blagoev | Received 8 July 2020 | Accepted 13 August 2020 | Published 3 September 2020

<http://zoobank.org/E2C55987-CB0C-463C-9178-FAD50CB5CE15>

Citation: Wang X, Shaheen S, He Q, Yao Z (2020) Notes on two closely related spider species of the *Pholcus phungiformes* species group (Araneae, Pholcidae) from Beijing, China. ZooKeys 965: 1–16. <https://doi.org/10.3897/zookeys.965.56199>

Abstract

The *Pholcus phungiformes* species group is highly diverse and currently contains 53 species. In this study, *Pholcus tongyaoi* Wang & Yao, **sp. nov.** (male, female) from Huairou District, Beijing, China is described while similar congener *Pholcus lexuancanhi* Yao, Pham & Li, 2012 from neighboring Haidian District (type locality) is redescribed; the female of *P. lexuancanhi* is described for the first time. In addition, the DNA barcode COI for the two species was obtained to estimate p-distance.

Keywords

daddy-long-leg spider, DNA barcode, morphology, Pholcinae, taxonomy

Introduction

The spider family Pholcidae C.L. Koch, 1850 contains 94 genera and 1768 species (World Spider Catalog 2020). It is among the most species-rich families and has a worldwide distribution (World Spider Catalog 2020). It is composed of five subfami-

* Contributed equally as the first author.

lies: Ninetinae Simon, 1890, Arteminae Simon, 1893, Modisiminae Simon, 1893, Smeringopinae Simon, 1893, and Pholcinae C.L. Koch, 1850 based on recent morphological and molecular phylogenetic analyses (Huber 2011a; Dimitrov et al. 2013; Eberle et al. 2018). Pholcid spiders occupy a wide range of habitats in a variety of ecosystems, e.g., in buildings, under rocks, in crevices, in caves, in leaf litter, and in webs between trunks and twigs of trees (Huber 2005; Yao and Li 2012). *Pholcus* Walckenaer, 1805 is the most diverse genus in Pholcinae and Pholcidae, with 338 described species mainly distributed in the Palearctic, Indo-Malayan, Afrotropical, and Australasian Region (Huber 2011b; Yao and Li 2012; World Spider Catalog 2020). These species belong to 21 species groups, of which the *Pholcus phungiformes* species group is highly diverse, including 53 known species definitively assigned to this species group (Huber 2011b; Peng and Zhang 2013; Kim and Ye 2015; Zhang et al. 2016; Huber et al. 2018; Zhu et al. 2018). The *P. phungiformes* species group is largely restricted to northeastern China and the Korean Peninsula; only *P. phungiformes* Oligier, 1983 occurs in Maritime Territory, Sakhalin Island, and Kurile Islands, Russia, probably as a result of human transport (Huber 2011b; World Spider Catalog 2020). This species group can often be found in caves, at cave entrances or on rock walls (Figs 1, 2), and diagnosed by the following characters: eight eyes, carapace with radiating marks, cylindrical opisthosoma, male chelicerae usually with frontal apophyses, male pedipalpal tibia with prolatero-ventral projection, procurcus usually with dorsal spines, appendix



Figure 1. Distribution records of two *Pholcus* species **1** *P. tongyai* sp. nov. **2** *P. lexuancahi* Yao, Pham & Li, 2012.




Figure 2. *Pholcus tongyaoi* sp. nov., live specimens and habitat **A, B** adult and juvenile males in old house **C, D** adult male and female with egg-sac on rock walls **E, F** habitat. Photographs by T Jiang (IZCAS).

absent, sometimes with ‘pseudo-appendix’, external female genitalia sclerotized, with knob (Huber 2011b; Zhu et al. 2018).

In this study, we describe one new species based on males and females from Huairou District, Beijing, China assigned to the *P. phungiformes* species group and redescribe its similar species *Pholcus lexuancanhi* Yao, Pham & Li, 2012 from a neighboring locality. The female of *P. lexuancanhi* is reported for the first time and the DNA barcode COI for the two species was obtained to estimate p-distance.

Materials and methods

Specimens were examined and measured with a Leica M205 C stereomicroscope. Left male pedipalps were illustrated. External female genitalia were photographed before dissection. Vulvae were previously treated in a 10% warm solution of potassium hydroxide (KOH) to dissolve soft tissues before illustration. Images were captured with a Canon EOS 750D wide zoom digital camera (24.2 megapixels) mounted on the stereomicroscope mentioned above, and assembled using Helicon Focus 3.10.3 image stacking software (Khmelik et al. 2005). All measurements are given in millimeters (mm). Leg measurements are shown as: total length (femur + patella + tibia + metatarsus + tarsus), missing data were coded as ‘–’. Leg podomeres were measured on their dorsal side. The distribution map was generated with Google Earth Pro 7.3.2 (Google Limited Liability Company). The specimens studied are preserved in 75% ethanol and deposited in the College of Life Science, Shenyang Normal University (SYNU) in Liaoning, China and the Institute of Zoology, Chinese Academy of Sciences (IZCAS) in Beijing, China.

Terminology and taxonomic descriptions follow Huber (2011b) and Yao et al. (2015). The following abbreviations are used in the descriptions:

- ALE** anterior lateral eye,
- AME** anterior median eye,
- PME** posterior median eye,
- L/d** length/diameter.

DNA barcode was obtained for estimation of p-distance between *P. tongyaoi* sp. nov. and *P. lexuancanhi*. A partial fragment of the mitochondrial cytochrome oxidase subunit I (COI) gene was amplified and sequenced, using the following primers: forward: LCO1490-oono (5'-CWACAAAYCATARRGATATTTGG-3') and reverse: C1-N-2776 (5'-GGATAATCAGAATANCGNCGAGG-3'). DNA sample is preserved in TE buffer and stored at -20 °C. The sequences are deposited in GenBank. COI p-distance is computed with MEGA 5 (Tamura et al. 2011). For additional information on extraction, amplification, and sequencing procedures, see Yao et al. (2016).

Taxonomic accounts

Family Pholcidae C.L. Koch, 1850

Subfamily Pholcinae C.L. Koch, 1850

Genus *Pholcus* Walckenaer, 1805

Type species. *Aranea phalangoides* Fuesslin, 1775.

Pholcus phungiformes species group

Diagnosis and description. See Huber (2011b).

Remarks. The 'appendix' in the original figures of four species apparently arises from the uncus: *P. papilionis* Peng & Zhang, 2011, *P. chiakensis* Seo, 2014, *P. gajiensis* Seo, 2014, and *P. palgongensis* Seo, 2014. We consider this a divided 'pseudo-appendix' and assign them to the *P. phungiformes* species group. Moreover, although the species *P. xianrendong* Liu & Tong, 2015 does not possess a prolatero-ventral projection on the male pedipalpal tibia, the bulb without appendix, the locality of this species is within the range of the *P. phungiformes* species group. Therefore, we tentatively assigned *P. xianrendong* to this species group. In total, this species group now contains 59 species. Of these, one species is newly described below.

***Pholcus tongyaoi* Wang & Yao, sp. nov.**

<http://zoobank.org/F7249E28-D367-4CA4-9D17-32977C5E5345>

Figs 3, 4

Type material. **Holotype:** male (SYNU-Ar00016), Pool and Valley Natural Park (40°32.600'N, 116°40.687'E, elevation 574 m), Huairou District, **Beijing, China**, 26 April 2019, T Jiang leg. **Paratypes:** 2 males (SYNU-Ar00017, Ar00018, GenBank number in SYNU-Ar00017: MT843113), same data as holotype; 2 females (SYNU-Ar00019, Ar00020), same data as holotype but 23 April 2019.

Etymology. The specific name is a patronym in honor of the collector Tongyao Jiang (IZCAS) and is a noun (name) in genitive case.

Diagnosis. The species resembles *P. lexuancanhi* Yao, Pham & Li, 2012 (Figs 5, 6; Yao et al. 2012: 313, figs 1–15) with similar bulbal apophyses (Fig. 4C) and external female genitalia (Fig. 4A), but can be easily distinguished by procurus with large, semicircular, ventral membranous process (arrowed in Fig. 3A; prolateral membranous lamella in *P. lexuancanhi*, arrowed 1 in Fig. 5C), small, prolateral membranous lamella provided with sawtooth (arrowed 1 in Fig. 3C; large, dorsal membranous lamella in *P. lexuancanhi*, arrowed 2 in Fig. 5C), and small, angular ventral sclerite provided with




Figure 3. *Pholcus tongyaoi* sp. nov., holotype (**A, B, D**) and paratype (**C**) males **A, B** pedipalp (**A** prolateral view, arrow indicates ventral membranous process **B** retrolateral view, arrow indicates ventral sclerite) **C, D** distal part of procursor (**C** prolateral view, arrow 1 indicates prolateral membranous lamella, arrow 2 indicates spine-shaped prolateral apophysis, arrow 3 indicates ventral membranous process **D** dorsal view, arrow indicates dorsal spine). Abbreviations: b = bulb, e = embolus, pr = procursor, u = uncus. Scale bars: 0.20 (**A, B**), 0.10 (**C, D**).




Figure 4. *Pholcus tongyai* sp. nov., holotype male (C–F) and paratype female (A, B, G, H) A external female genitalia, ventral view B vulva, dorsal view C bulbal apophyses, prolatateral view, arrow indicates ‘pseudo-appendix’ D chelicerae, frontal view E–H habitus (E, G dorsal view F lateral view H ventral view). Abbreviations: da = distal apophysis, e = embolus, fa = frontal apophysis, pa = proximo-lateral apophysis, pp = pore plate, u = uncus. Scale bars: 0.20 (A–D), 1.00 (E–H).

curved tip (arrowed in Fig. 3B; large ventral sclerite and its tip not curved in *P. lexuancanhi*, arrowed in Fig. 5B), by male chelicerae with pair of frontal apophyses (arrowed fa in Fig. 4D; absent in *P. lexuancanhi*, Fig. 6D), and by vulva with n-shaped anterior arch without median sclerite (Fig. 4B; slightly curved anterior arch with median sclerite in *P. lexuancanhi*, arrowed in Fig. 6B) and elliptic pore plates (Fig. 4B; oval in *P. lexuancanhi*, Fig. 6B). This species can also be distinguished from *P. lexuancanhi* by COI p-distance 0.106 between them.

Description. Male (holotype, SYNU-Ar00016): Total length 4.75 (4.93 with clypeus), carapace 1.56 long, 1.75 wide, opisthosoma 3.19 long, 1.34 wide. Leg I: – (11.62 + 0.75 + – + – + –), leg II: 30.89 (8.50 + 0.55 + 7.84 + 12.75 + 1.25), leg III: 20.62 (6.40 + 0.60 + 5.12 + 7.60 + 0.90), leg IV: 28.58 (8.40 + 0.62 + 7.12 + 11.12 + 1.32). Distance PME-PME 0.20, diameter PME 0.12, distance PME-ALE 0.05, distance AME-AME 0.02, diameter AME 0.08. Sternum wider than long (1.04/0.96). Habitus as in Fig. 4E, F. Carapace yellowish, with brown radiating marks and marginal brown bands; ocular area yellowish, with median and lateral brown bands; clypeus yellowish; sternum yellowish, with marginal brown marks. Legs yellowish, but dark brown on patellae and whitish on distal parts of femora and tibiae, with darker rings on subdistal parts of femora and proximal and subdistal parts of tibiae. Opisthosoma yellowish, with dorsal and lateral spots. Ocular area elevated, without eye stalks. Thoracic furrow absent. Chelicerae (Fig. 4D) with pair of proximo-lateral apophyses, pair of distal apophyses provided with two teeth each, and pair of frontal apophyses. Pedipalps as in Fig. 3A, B; trochanter with long, retrolaterally strongly bulged ventral apophysis; femur with indistinct ventral protuberance; tibia with prolatero-ventral projection; procurus simple proximally but complex distally, with large, semicircular, ventral membranous process (arrowed in Fig. 3A), small, prolateral membranous lamella with sawtooth (arrowed 1 in Fig. 3C), small, angular ventral sclerite with curved tip (arrowed in Fig. 3B), spine-shaped prolateral apophysis (arrowed 2 in Fig. 3C), and dorsal spine (arrowed in Fig. 3D); bulb with short curved ‘pseudo-appendix’ (arrowed in Fig. 4C); uncus with scaly edge (Fig. 4C); embolus weakly sclerotized, with some transparent distal projections (Fig. 4C). Legs with short vertical setae on tibiae, metatarsi, and tarsi, without spines or curved setae.

Female (paratype, SYNU-Ar00019): Similar to male, habitus as in Fig. 4G, H. Total length 5.31 (5.56 with clypeus), carapace 1.43 long, 1.14 wide, opisthosoma 3.88 long, 2.43 wide; tibia I: 5.90; tibia I L/d: 54. Distance PME-PME 0.18, diameter PME 0.12, distance PME-ALE 0.04, distance AME-AME 0.03, diameter AME 0.08. Sternum wider than long (1.07/0.83). Clypeus brown. External female genitalia (Fig. 4A) curved posteriorly, with short knob. Vulva (Fig. 4B) with sclerotized, n-shaped anterior arch and pair of elliptic pore plates. Retrolateral trichobothrium of tibia I at 4% proximally; tarsus I with 22 distinct pseudosegments.

Variations. Ventral membranous process on procurus nearly crescent-shaped (arrowed 3 in Fig. 3C) in one paratype male (SYNU-Ar00017). Leg I missing in two paratype males (SYNU-Ar00017, Ar00018), total length 5.36 (5.52 with clypeus) in SYNU-Ar00017, total length 4.50 (4.60 with clypeus) in SYNU-Ar00018. Tibia I in another paratype female (SYNU-Ar00020): 5.75.

Distribution. China (Beijing, type locality; Fig. 1).

Natural history. The species was found in an old house and on rock walls.

***Pholcus lexuancanhi* Yao, Pham & Li, 2012**

Figs 5, 6

Pholcus lexuancanhi Yao et al. 2012: 313, figs 1–15. Yao et al. 2015: 15.

Material examined. 2 males (IZCAS-Ar40901, Ar40902, GenBank number in IZCAS-Ar40901: MT843112) and 2 females (IZCAS-Ar40903, Ar40904), Beijing Botanical Garden (40°00'N, 116°12'E; type locality), Haidian District, **Beijing, China**, 30 July 2017, Z Yao leg.

Diagnosis. See diagnosis for *P. tongyaoi* sp. nov.

Redescription. Male (IZCAS-Ar40901): Total length 5.13 (5.38 with clypeus), carapace 1.41 long, 1.75 wide, opisthosoma 3.72 long, 1.44 wide. Leg I: 47.97 (14.47 + 0.80 + 11.41 + 18.53 + 2.76), leg II: 30.06 (8.46 + 0.78 + 7.24 + 11.79 + 1.79), leg III: 21.37 (6.35 + 0.59 + 5.13 + 8.08 + 1.22), leg IV: 28.03 (8.33 + 0.78 + 7.05 + 10.51 + 1.36); tibia I L/d: 76. Distance PME-PME 0.25, diameter PME 0.10, distance PME-ALE 0.05, distance AME-AME 0.05, diameter AME 0.10. Sternum wider than long (1.05/0.88). Habitus as in Fig. 6E, F. Carapace yellowish, with brown radiating marks and marginal brown bands; ocular area yellowish, with median and lateral brown bands; clypeus yellowish; sternum brown. Legs yellowish, but dark brown on patellae and whitish on distal parts of femora and tibiae, with darker rings on subdistal parts of femora and proximal and subdistal parts of tibiae. Opisthosoma yellowish, with dorsal and lateral spots. Ocular area elevated, without eye stalks. Thoracic furrow absent. Chelicerae (Fig. 6D) with pair of proximo-lateral apophyses and pair of distal apophyses provided with two teeth each. Pedipalps as in Fig. 5A, B; trochanter with long, retrolaterally strongly bulged ventral apophysis; femur with indistinct ventral protuberance; tibia with prolatero-ventral projection; procurus simple proximally but complex distally, with large, prolateral membranous lamella (arrowed 1 in Fig. 5C), large, dorsal membranous lamella with sawtooth (arrowed 2 in Fig. 5C), large, angular ventral sclerite (arrowed in Fig. 5B), spine-shaped prolateral apophysis (arrowed 3 in Fig. 5C), and dorsal spine (arrowed in Fig. 5D); bulb with short curved 'pseudo-appendix' (arrowed in Fig. 6C); uncus with scaly edge (Fig. 6C); embolus weakly sclerotized, with some transparent distal projections (Fig. 6C). Retrolateral trichobothrium of tibia I at 5% proximally; legs with short vertical setae on tibiae, metatarsi, and tarsi, without spines or curved setae; tarsus I with 33 distinct pseudosegments.

Female (IZCAS-Ar40903): Similar to male, habitus as in Fig. 6G, H. Total length 5.19 (5.38 with clypeus), carapace 1.36 long, 1.66 wide, opisthosoma 3.83 long, 1.68 wide; tibia I: 8.50; tibia I L/d: 54. Distance PME-PME 0.20, diameter PME 0.10, distance PME-ALE 0.05, distance AME-AME 0.05, diameter AME 0.08. Sternum wider than long (1.08/0.92). Clypeus brown. External female genitalia (Fig. 6A) curved pos-




Figure 5. *Pholcus lexuancanhi* Yao, Pham & Li, 2012, male **A, B** pedipalp (**A** prolateral view **B** retrolateral view, arrow indicates ventral sclerite) **C, D** distal part of procurus (**C** prolateral view, arrows 1 and 2 indicate prolateral and dorsal membranous lamella, respectively, arrow 3 indicates spine-shaped prolateral apophysis **D** dorsal view, arrow indicates dorsal spine). Abbreviations: b = bulb, e = embolus, pr = procurus, u = uncus. Scale bars: 0.20 (**A, B**), 0.05 (**C, D**).




Figure 6. *Pholcus lexuancanhi* Yao, Pham & Li, 2012, male (C–F) and female (A, B, G, H) **A** external female genitalia, ventral view **B** vulva, dorsal view, arrow indicates median sclerite of anterior arch **C** bulbal apophyses, prolateral view, arrow indicates ‘pseudo-appendix’ **D** chelicerae, frontal view **E–H** habitus (**E, G** dorsal view **F** lateral view **H** ventral view). Abbreviations: da = distal apophysis, e = embolus, pa = proximo-lateral apophysis, pp = pore plate, u = uncus. Scale bars: 0.20 (A–D), 1.00 (E–H).

teriorly, with short knob. Vulva (Fig. 6B) with slightly curved, sclerotized anterior arch provided with median sclerite (arrowed in Fig. 6B) and pair of oval pore plates.

Variations. Tibia I in another male (IZCAS-Ar40902): 11.54. Tibia I in another female (IZCAS-Ar40904): 9.12.

Distribution. China (Beijing, type locality; Fig. 1).

Natural history. The species was found on rock walls.

Discussion

The *P. phungiformes* species group is highly diverse and currently contains 59 species including one new species in this study. These species are mainly distributed in three large mountain ranges (see the checklist below): the Mountain Taihang from southern North China (22 spp.), the Mountain Changbai (also called Mountain Paekdu in North Korea) from the border between northeastern China and North Korea (15 spp., of which *P. phungiformes* also occurs further east), and the Mountain Taebaek from Korean Peninsula (22 spp.) (Huber 2011b; Peng and Zhang 2011; Yao and Li 2012; Yao et al. 2012; Peng and Zhang 2013; Seo 2014; Kim and Ye 2015; Liu and Tong 2015; Zhang et al. 2016; Zhu et al. 2018). Nevertheless, the survey of the *P. phungiformes* species group is uneven. The highest diversity (43 spp.) concentrates in the Mountain Taihang and the southern Mountain Taebaek (South Korea). In contrast, only 15 species from the Mountain Changbai and one species from the northern Mountain Taebaek (North Korea) are recorded. Based on the high diversity of this species group from the southern Mountain Taebaek and the Mountain Taihang, as well as the similar landforms and habitats in neighboring northern Mountain Taebaek and Mountain Changbai, we strongly believe that additional species diversity likely remains undiscovered in the neighboring areas (e.g., Li 2020). Further survey in these areas is needed to fully understand the diversity that exists within this group.

A checklist of the *P. phungiformes* species group from three large mountain ranges is provided (for the complete list of references, see World Spider Catalog 2020):

The Mountain Taihang:

1. *Pholcus allectospilus* Zhu & Gong, 1991
2. *Pholcus auricularis* Zhang, Zhang & Liu, 2016
3. *Pholcus babao* Tong & Li, 2010
4. *Pholcus beijingensis* Zhu & Song, 1999
5. *Pholcus brevis* Yao & Li, 2012
6. *Pholcus chicheng* Tong & Li, 2010
7. *Pholcus clavimaculatus* Zhu & Song, 1999
8. *Pholcus curvus* Zhang, Zhang & Liu, 2016
9. *Pholcus datan* Tong & Li, 2010
10. *Pholcus exilis* Tong & Li, 2010

11. *Pholcus jinniu* Tong & Li, 2010
12. *Pholcus lexuancanhi* Yao, Pham & Li, 2012
13. *Pholcus luya* Peng & Zhang, 2013
14. *Pholcus papilionis* Peng & Zhang, 2011
15. *Pholcus papillatus* Zhang, Zhang & Liu, 2016
16. *Pholcus pennatus* Zhang, Zhu & Song, 2005
17. *Pholcus suizhongicus* Zhu & Song, 1999
18. *Pholcus tongyaoi* sp. nov.
19. *Pholcus triangulatus* Zhang & Zhang, 2000
20. *Pholcus wangxidong* Zhang & Zhu, 2009
21. *Pholcus wuling* Tong & Li, 2010
22. *Pholcus zhuolu* Zhang & Zhu, 2009

The Mountain Changbai:

1. *Pholcus decorus* Yao & Li, 2012
2. *Pholcus fengcheng* Zhang & Zhu, 2009
3. *Pholcus foliaceus* Peng & Zhang, 2013
4. *Pholcus gaoi* Song & Ren, 1994
5. *Pholcus hamatus* Tong & Ji, 2010
6. *Pholcus jiuwei* Tong & Ji, 2010
7. *Pholcus lingulatus* Gao, Gao & Zhu, 2002
8. *Pholcus ningan* Yao & Li, 2018
9. *Pholcus phoenixus* Zhang & Zhu, 2009
10. *Pholcus phungiformes* Oligier, 1983
11. *Pholcus sublingulatus* Zhang & Zhu, 2009
12. *Pholcus tongi* Yao & Li, 2012
13. *Pholcus wangi* Yao & Li, 2012
14. *Pholcus wangtian* Tong & Ji, 2010
15. *Pholcus xianrendong* Liu & Tong, 2015

The Mountain Taebaek:

1. *Pholcus acutulus* Paik, 1978
2. *Pholcus cheongogensis* Kim & Ye, 2015
3. *Pholcus chiakensis* Seo, 2014
4. *Pholcus crassus* Paik, 1978
5. *Pholcus extumidus* Paik, 1978
6. *Pholcus gajiensis* Seo, 2014
7. *Pholcus gosuensis* Kim & Lee, 2004
8. *Pholcus joreongensis* Seo, 2004
9. *Pholcus juwangensis* Seo, 2014
10. *Pholcus kwanaksanensis* Namkung & Kim, 1990

11. *Pholcus kwangkyosanensis* Kim & Park, 2009
12. *Pholcus montanus* Paik, 1978
13. *Pholcus nodong* Huber, 2011
14. *Pholcus okgye* Huber, 2011
15. *Pholcus palgongensis* Seo, 2014
16. *Pholcus parkyeonensis* Kim & Yoo, 2009
17. *Pholcus pojeonensis* Kim & Yoo, 2008
18. *Pholcus simbok* Huber, 2011
19. *Pholcus socheunensis* Paik, 1978
20. *Pholcus sokkrisanensis* Paik, 1978
21. *Pholcus woongil* Huber, 2011
22. *Pholcus yeongwol* Huber, 2011

Acknowledgements

The manuscript benefited greatly from comments by Gergin Blagoev, Marc Milne, and John TD Caleb. We are grateful to Tongyao Jiang for providing the ecological photographs and specimens. Abid Ali kindly checked the English. This study was supported by the National Natural Science Foundation of China (NSFC-31872193) and the Liaoning Revitalization Talents Program (XLYC1907150). Part of the laboratory work was supported by the Research Fund for the Doctoral Program of Shenyang Normal University (BS201841) and the Shenyang Youth Science and Technology Project (RC200183).

References

- Dimitrov D, Astrin JJ, Huber BA (2013) Pholcid spider molecular systematics revisited, with new insights into the biogeography and the evolution of the group. *Cladistics* 29(2): 132–146. <https://doi.org/10.1111/j.1096-0031.2012.00419.x>
- Eberle J, Dimitrov D, Valdez-Mondragón A, Huber BA (2018) Microhabitat change drives diversification in pholcid spiders. *BMC Evolutionary Biology* 18: 1–141. <https://doi.org/10.1186/s12862-018-1244-8>
- Fuesslin JC (1775) Verzeichnis der ihm Bekannten Schweizerischen Insekten, mit Einer Ausgemahlten Kupfertafel: Nebst der Ankündigung Eines Neuen Insekten Werkes. Zürich and Winterthur, 62 pp. [Araneae pp: 60–61] <https://doi.org/10.5962/bhl.title.65772>
- Huber BA (2005) The pholcid spiders of Africa (Araneae: Pholcidae): state of knowledge and directions for future research. In: Huber BA, Sinclair BJ, Lampe KH (Eds) *African Biodiversity: Molecules, Organisms, Ecosystems*. Springer Verlag, The Netherlands, 181–186. https://doi.org/10.1007/0-387-24320-8_15
- Huber BA (2011a) Phylogeny and classification of Pholcidae (Araneae): an update. *Journal of Arachnology* 39: 211–222. <https://doi.org/10.1636/CA10-57.1>

- Huber BA (2011b) Revision and cladistic analysis of *Pholcus* and closely related taxa (Araneae, Pholcidae). *Bonner Zoologische Monographien* 58: 1–509.
- Huber BA, Eberle J, Dimitrov D (2018) The phylogeny of pholcid spiders: a critical evaluation of relationships suggested by molecular data (Araneae, Pholcidae). *ZooKeys* 789: 51–101. <https://doi.org/10.3897/zookeys.789.22781>
- Khmelik VV, Kozub D, Glazunov A (2005) *Helicon Focus* 3.10.3. <https://www.heliconsoft.com/heliconsoft-products/helicon-focus/> [accessed 10 January 2020]
- Kim J, Ye S (2015) A new species of the genus *Pholcus* Walckenaer, 1805 (Araneae: Pholcidae) from Korea. *Korean Arachnology* 31(2): 73–80.
- Koch CL (1850) Übersicht des Arachnidensystems. Heft 5. Lotzbeck JL, Nürnberg, 77 pp. <https://doi.org/10.5962/bhl.title.39561>
- Li S (2020) Spider taxonomy for an advanced China. *Zoological Systematics* 45(2): 73–77. <https://doi.org/10.11865/zs.202011>
- Liu Y, Tong Y (2015) A new species of the genus *Pholcus* Walckenaer, 1805 from Xianrendong National Forest Park, Liaoning Province, China (Araneae: Pholcidae). *Acta Arachnologica Sinica* 24(1): 31–34. <https://doi.org/10.3969/j.issn.1005-9628.2015.01.008>
- Oligier TI (1983) New species of spider families (Pholcidae, Clubionidae, Agelenidae) from the Lazovsky State Reserve. *Zoologicheskii Zhurnal* 62: 627–629.
- Peng Y, Zhang F (2011) Two new species of the genus *Pholcus* (Araneae: Pholcidae) from Taihang Mountains, China, with first report of the female of *Pholcus oculosus*. *Entomologica Fennica* 22: 78–84. <https://doi.org/10.33338/ef.84546>
- Peng Y, Zhang F (2013) Two new *Pholcus* species from northern China (Araneae: Pholcidae). *Acta Arachnologica* 62(2): 75–80. <https://doi.org/10.2476/asjaa.62.75>
- Seo BK (2014) Four new species of the genus *Pholcus* (Araneae: Pholcidae) from Korea. *Korean Journal of Applied Entomology* 53(4): 399–408. <https://doi.org/10.5656/KSAE.2014.10.0.059>
- Simon E (1890) Études Arachnologiques. 22e Mémoire. XXXIV. Étude sur les arachnides de l'Yemen. *Annales de la Société Entomologique de France* (6) 10: 77–124.
- Simon E (1893) *Histoire Naturelle des Araignées* (Vol. 1). Roret, Paris, 256–488. <https://doi.org/10.5962/bhl.title.51973>
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S (2011) MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* 28(10): 2731–2739. <https://doi.org/10.1093/molbev/msr121>
- Walckenaer CA (1805) *Tableau des Aranéides ou Caractères Essentiels des Tribus, Genres, Familles et Races que Renferme le Genre Aranea de Linné, Avec la Désignation des Espèces Comprises dans Chacune de ces Divisions*. Dentu, Paris, 88 pp.
- World Spider Catalog (2020) *World Spider Catalog, Version 21.5*. Natural History Museum Bern. <http://wsc.nmbe.ch> [accessed 22 August 2020]
- Yao Z, Dong T, Zheng G, Fu J, Li S (2016) High endemism at cave entrances: a case study of spiders of the genus *Uthina*. *Scientific Reports* 6(35757): 1–9. [I–LII] <https://doi.org/10.1038/srep35757>

- Yao Z, Li S (2012) New species of the spider genus *Pholcus* (Araneae: Pholcidae) from China. *Zootaxa* 3289: 1–271. <https://doi.org/10.11646/zootaxa.3289.1.1>
- Yao Z, Pham DS, Li S (2012) A new species of the genus *Pholcus* (Araneae, Pholcidae) from Vietnam. *Acta Zootaxonomica Sinica* 37: 313–318.
- Yao Z, Pham DS, Li S (2015) Pholcid spiders (Araneae: Pholcidae) from northern Vietnam, with descriptions of nineteen new species. *Zootaxa* 3909(1): 1–82. <https://doi.org/10.11646/zootaxa.3909.1.1>
- Zhang B, Zhang F, Liu J (2016) Three new spider species of the genus *Pholcus* from the Taihang Mountains of China (Araneae, Pholcidae). *ZooKeys* 600: 53–74. <https://doi.org/10.3897/zookeys.600.7924>
- Zhu K, Yao Z, Bai X, Li S (2018) Five new species of the spider genus *Pholcus* Walckenaer, 1805 (Araneae: Pholcidae) from China and Thailand. *Zootaxa* 4377(2): 235–253. <https://doi.org/10.11646/zootaxa.4377.2.5>