

Research article

Open Access

Membrane-associated prostaglandin E synthase-1 is upregulated by proinflammatory cytokines in chondrocytes from patients with osteoarthritisFumiaki Kojima¹, Hiroaki Naraba², Satoshi Miyamoto³, Moroe Beppu³, Haruhito Aoki³ and Shinichi Kawai¹¹Institute of Medical Science, St Marianna University School of Medicine, Kawasaki, Japan²National Cardiovascular Center Research Institute, Osaka, Japan³Department of Orthopedic Surgery, St Marianna University School of Medicine, Kawasaki, JapanCorresponding author: Shinichi Kawai, s2kawai@marianna-u.ac.jp

Received: 27 Dec 2003 Revisions requested: 21 Jan 2003 Revisions received: 1 Apr 2004 Accepted: 12 May 2004 Published: 8 Jun 2004

Arthritis Res Ther 2004, **6**:R355-R365 (DOI 10.1186/ar1195)© 2004 Kojima *et al.*; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.**Abstract**

Prostaglandin E synthase (PGES) including isoenzymes of membrane-associated PGES (mPGES)-1, mPGES-2, and cytosolic PGES (cPGES) is the recently identified terminal enzyme of the arachidonic acid cascade. PGES converts prostaglandin (PG)H₂ to PGE₂ downstream of cyclooxygenase (COX). We investigated the expression of PGES isoenzyme in articular chondrocytes from patients with osteoarthritis (OA). Chondrocytes were treated with various cytokines and the expression of PGES isoenzyme mRNA was analyzed by the reverse transcription-polymerase chain reaction and Northern blotting, whereas Western blotting was performed for protein expression. The subcellular localization of mPGES-1 was determined by immunofluorescent microscopy. Conversion of arachidonic acid or PGH₂ to PGE₂ was measured by enzyme-linked immunosorbent assay. Finally, the expression of mPGES-1 protein in OA articular cartilage was assessed by immunohistochemistry. Expression of mPGES-1 mRNA in chondrocytes was significantly induced by interleukin (IL)-1 β or

tumor necrosis factor (TNF)- α , whereas other cytokines, such as IL-4, IL-6, IL-8, IL-10, and interferon- γ , had no effect. COX-2 was also induced under the same conditions, although its pattern of expression was different. Expression of cPGES, mPGES-2, and COX-1 mRNA was not affected by IL-1 β or TNF- α . The subcellular localization of mPGES-1 and COX-2 almost overlapped in the perinuclear region. In comparison with 6-keto-PGF_{1 α} and thromboxane B₂, the production of PGE₂ was greater after chondrocytes were stimulated by IL-1 β or TNF- α . Conversion of PGH₂ to PGE₂ (PGES activity) was significantly increased in the lysate from IL-1 β -stimulated chondrocytes and it was inhibited by MK-886, which has an inhibitory effect on mPGES-1 activity. Chondrocytes in articular cartilage from patients with OA showed positive immunostaining for mPGES-1. These results suggest that mPGES-1 might be important in the pathogenesis of OA. It might also be a potential new target for therapeutic strategies that specifically modulate PGE₂ synthesis in patients with OA.

Keywords: chondrocytes, interleukin-1 β , osteoarthritis, prostaglandin E synthase, tumor necrosis factor- α **Introduction**

Osteoarthritis (OA) is the most common chronic articular condition and is primarily characterized by the progressive destruction of articular cartilage. Although OA is traditionally defined as a non-inflammatory arthropathy, many reports have suggested that proinflammatory cytokines such as interleukin (IL)-1 β and tumor necrosis factor (TNF)-

α might be important in this disease [1-3]. Stimulation of chondrocytes by proinflammatory cytokines increases the production of major cartilage-degrading enzymes, called matrix metalloproteinases [4-6], inhibits the synthesis of cartilage proteoglycans and/or type II collagen [7-9], stimulates the production of reactive oxygen species such as nitric oxide [10-12], and increases the production of

COX = cyclooxygenase; cPGES = cytosolic prostaglandin E synthase; DMEM = Dulbecco's modified Eagle's medium; ELISA = enzyme-linked immunosorbent assay; FBS = fetal bovine serum; HRP = horseradish peroxidase; IL = interleukin; mPGES = membrane-associated prostaglandin E synthase; OA = osteoarthritis; PBS = phosphate-buffered saline; PG = prostaglandin; PGES = prostaglandin E synthase; RT-PCR = reverse transcription-polymerase chain reaction; SDS = sodium dodecyl sulfate; TBS = Tris-buffered saline; TBS-T = TBS containing 0.1% Tween-20; TNF = tumor necrosis factor; TX = thromboxane.

prostaglandin (PG) E_2 [13,14]. It was recently shown that high levels of PGE $_2$ production by OA tissues are observed after stimulation by IL-1 β , which might have a role in the degeneration of bone and cartilage associated with OA [15]. It was also reported that the nitric oxide-induced death of OA chondrocytes was enhanced through the production of PGE $_2$ by the chondrocytes themselves [16].

In the PGE $_2$ synthesis pathway, cyclooxygenase (COX) is a key enzyme that metabolizes arachidonic acid to PGG $_2$ and PGH $_2$. There are two isoforms of COX, which are designated COX-1 and COX-2 [17,18]. COX-1 is expressed constitutively in various cells and tissues, and is important in maintaining homeostasis. In contrast, COX-2 is induced in inflammatory cells and tissues by various stimuli including cytokines, suggesting that COX-2 has a key role in the process of inflammation. Increased PGE $_2$ production in response to stimulation by proinflammatory cytokines coincides with the upregulation of COX-2 expression.

PGE synthase (PGES) acts downstream of COX to catalyze the conversion of PGH $_2$ into PGE $_2$ [19,20]. At least three forms of human PGES have been cloned and characterized, including membrane-associated PGES (mPGES)-1 [21], mPGES-2 [22], and cytosolic PGES (cPGES) [23]. Among these, mPGES-1 was originally known as microsomal glutathione S-transferase 1-like 1 (MGST1-L1), and it is a glutathione-dependent enzyme that shows coordinated induction with COX-2 by inflammatory stimuli in various cells and tissues [21,24]. We and others have reported that mPGES-1 was coordinately induced with COX-2 by IL-1 β in synovial fibroblasts from patients with rheumatoid arthritis [25-27]. It was also reported that mPGES-1 is expressed in joint tissues isolated from animals with arthritis [28,29]. Moreover, a recent study demonstrated that mPGES-1-deficient mice showed a significant reduction of the inflammatory features and histopathological changes such as pannus formation and joint erosion of experimental inflammatory arthritis [30]. Although the significance of PGE $_2$ in the pathophysiology of OA has been demonstrated, the role of each PGES isoenzyme has not yet been determined. Accordingly, we studied the effects of various inflammatory cytokines on the expression of PGES isoenzymes in OA chondrocytes.

Materials and methods

Materials

Rabbit anti-human mPGES-1 polyclonal antibody, rabbit anti-human cPGES polyclonal antibody, rabbit anti-human COX-2 polyclonal antibody, PGE $_2$, PGH $_2$, and enzyme-linked immunosorbent assay (ELISA) kits for PGE $_2$, 6-keto-PGF $_{1\alpha}$, and thromboxane (TX) B_2 were all purchased from Cayman Chemical Co. (Ann Arbor, MI). Mouse anti-human COX-1 monoclonal antibody and MK-886 were purchased from Wako Pure Chemical Industries (Osaka, Japan). Goat

anti-human COX-2 polyclonal antibody was obtained from Santa Cruz Biotechnology (Santa Cruz, CA). Horseradish peroxidase (HRP)-conjugated goat anti-rabbit IgG, HRP-conjugated goat anti-mouse IgG, tetramethylrhodamine β -isothiocyanate-conjugated donkey anti-goat IgG, and fluorescein isothiocyanate-conjugated donkey anti-rabbit IgG were obtained from Jackson ImmunoResearch (West Grove, PA). Dulbecco's modified Eagle's medium (DMEM), fetal bovine serum (FBS), and TRIzol reagent were from Invitrogen (Carlsbad, CA). Recombinant human IL-1 β , TNF- α , IL-4, IL-6, IL-8, IL-10, and interferon- γ were from BD Biosciences (San Diego, CA). The Megaprime DNA labeling system, [α - 32 P]dCTP, nylon membrane (Hybond N+), and enhanced chemiluminescence reagent were purchased from Amersham Pharmacia Biotech (Little Chalfont, Bucks., UK), and poly(vinylidene difluoride) membrane (Immobilon-P) was obtained from Millipore (Bedford, MA).


Preparation of cells

Cartilage samples were obtained at the time of joint replacement surgery from patients with OA who fulfilled the American College of Rheumatology criteria for this disease [31]. These experiments were performed in accordance with a protocol approved by the ethics committee of St Marianna University, and all patients gave written consent to the use of their tissues for this research.

Chondrocytes were prepared as described elsewhere [32]. In brief, cartilage samples were minced and digested with 1 mg/ml collagenase in DMEM for 18 h at 37°C, filtered through a nylon mesh, and washed extensively. Then the isolated chondrocytes were seeded at a high density in tissue culture flasks coated with type I collagen and were incubated in DMEM containing 10% FBS, 100 units/ml penicillin, and 100 μ g/ml streptomycin at 37°C under an atmosphere of 5% CO $_2$. At confluence, the cells were detached and passaged, and the second-passage cells were used in subsequent experiments. The chondrocytes' phenotype was assessed by analyzing the expression of type II collagen with the reverse transcription-polymerase chain reaction (RT-PCR) (Fig. 1). OA synovial fibroblasts and normal dermal fibroblasts were prepared as described previously [25,33].

RT-PCR

Cells were seeded into the wells of a six-well plate coated with type I collagen at a density of 5×10^4 cells/cm 2 in culture medium containing 1% FBS, and then were cultured under various conditions. RNA from the cells was extracted with TRIzol reagent in accordance with the manufacturer's instructions. Reverse transcription was performed with a SuperScript preamplification system (Invitrogen, Carlsbad, CA), in accordance with the manufacturer's instructions, with 1 μ g of total RNA from the cells as a template. Equal amounts of each reverse-transcription product were

Figure 1

Phenotypic features of chondrocytes. **(a)** Morphology of chondrocytes and synovial fibroblasts from patients with osteoarthritis. **(b)** Chondrocytes from four patients with osteoarthritis were incubated for 12 h. Total RNA from the cells was subjected to reverse transcription–polymerase chain reaction (RT–PCR) for type I collagen, type II collagen, and glyceraldehyde-3-phosphate dehydrogenase (GAPDH), as described in Materials and methods. Synovial fibroblasts and dermal fibroblasts were used as controls. **(c)** Chondrocytes from osteoarthritis patients were incubated with IL-1 β (1 ng/ml) or tumor necrosis factor- α (TNF α ; 10 ng/ml) for 12 h. Total RNA from the cells was subjected to RT–PCR for type I collagen, type II collagen, and GAPDH, as described in Materials and methods. Representative results from three patients are shown here.

amplified by PCR with HotStarTaq polymerase (Qiagen, Valencia, CA). After initial denaturation at 95°C for 15 min, PCR involved amplification cycles of 30 s at 95°C, 30 s at 56°C, and 45 s at 72°C, followed by elongation for 5 min at 72°C. The sequences of the primers, the size of the PCR products, and the number of amplification cycles are shown in Table 1. The amplified cDNA fragments were resolved by electrophoresis on 1.5% (w/v) agarose gel, and were detected under ultraviolet after staining of the gel with ethidium bromide.

Northern blot analysis

Cells were seeded into the wells of a six-well plate coated with type I collagen at a density of 5×10^4 cells/cm² in culture medium containing 1% FBS, and were then cultured under various conditions. Northern blotting was performed as described previously [25]. Total RNA (4 μ g) was extracted from the cells with TRIzol reagent and was sub-

jected to electrophoresis on 1% formaldehyde/agarose gel, followed by transfer to a Hybond N+ membrane and hybridization with cDNA probes that had been labeled with [α -³²P]dCTP by random priming. After hybridization, the membranes were washed and the RNA bands were detected by autoradiography.

Western blot analysis

Cells (at a density of 5×10^4 cells/cm²) were cultured under various conditions in medium containing 1% FBS. Subsequently, the cells were lysed in Tris-buffered saline (TBS) containing 0.1% sodium dodecyl sulfate (SDS), and the protein content of the lysates was determined with the bicinchoninic acid protein assay reagent (Pierce, Rockford, IL), with bovine serum albumin as the standard. Cell lysates were adjusted to 5 μ g of protein and were then applied to SDS-polyacrylamide gels (10–20%) for electrophoresis, as reported previously [25]. Next, the proteins were electroblotted onto Immobilon-P poly(vinylidene difluoride) membranes with a semidry blotter (Atto, Tokyo, Japan). After the membranes had been blocked in 10 mM TBS containing 0.1% Tween-20 (TBS-T) and 5% skimmed milk, the primary antibody (rabbit anti-human mPGES-1 antibody, rabbit anti-human cPGES antibody, mouse anti-human COX-1 antibody, or rabbit anti-human COX-2 antibody) was added at a dilution of 1:500 (mPGES-1 and cPGES) or 1:200 (COX-1 and COX-2) in TBS-T, and incubation was performed for 1.5 h. After the membranes had been washed with TBS-T, the secondary antibody (HRP-conjugated goat anti-rabbit antibody or HRP-conjugated goat anti-mouse antibody) was added (at a dilution of 1:10,000 or 1:5,000, respectively, in TBS-T) and incubation was performed for 1 h. After further washing with TBS-T, protein bands were detected with an enhanced chemiluminescence Western blot analysis system.

Immunofluorescent microscopy

Immunofluorescent microscopy was performed as reported previously [25]. In brief, cells were seeded at a density of 2.5×10^4 cells/cm² onto culture slides coated with type I collagen and were incubated with or without 1 ng/ml IL-1 β for 48 h. After removal of the supernatants, the cells were fixed with 4% (w/v) paraformaldehyde in phosphate-buffered saline (PBS) for 30 min at 4°C. Then the cells were permeabilized with 0.1% Triton-X in PBS for 3 min. After blocking with 1% (w/v) bovine serum albumin in PBS for 1 h at 20–25°C, rabbit anti-human mPGES-1 polyclonal antibody and goat anti-human COX-2 polyclonal antibody (premixed so that the final dilutions were, respectively, 1:100 and 1:250 in PBS containing 1% albumin) were added and incubation was done for 1.5 h at 20–25°C. After washing with PBS, tetramethylrhodamine β -isothiocyanate-conjugated and fluorescein isothiocyanate-conjugated secondary antibodies were added at 1:50 dilution in PBS containing 1% albumin, and incubated for 1 h. Then the

Table 1**Oligonucleotide primers used in the reverse transcription-polymerase chain reaction**

Target gene encoding	Product length (bp)	Primer (5' → 3') ^a	Cycle
Type I collagen	614	CAA CGG TGC TCC TGG TGA AG GCT GGT CAG CCC TGT AGA AG	32
Type II collagen	507	TCA ACA ACC AGA TTG AGA GCA ACG TGA ACC TGC TAT TGC CCT	32
mPGES-1	459	ATG CCT GCC CAC AGC CTG GT TCA CAG GTG GCG GGC CGC TT	26
mPGES-2	502	GCA GCT GAC CCT GTA CCA GT CTC GCG GAC AAT GTA GTC AA	32
cPGES	485	ATG CAG CCT GCT TCT GCA AA CCT TAC TCC AGA TCT GGC AT	30
COX-1	401	CTT TCT CCA ACG TGA GCT ATT GTG CAT CAA CAC AGG CGC CTC TTC	32
COX-2	1157	AGA CAG ATC ATA AGC GAG GGC C ACT TGC ATT GAT GGT GAC TGT	28
GAPDH	598	CCA CCC ATG GCA AAT TCC ATG GCA TCT AGA CGG CAG GTC AGG TCC ACC	26

COX, cyclooxygenase; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; mPGES, membrane-associated prostaglandin E synthase; cPGES, cytosolic prostaglandin E synthase. ^aFor each target gene, the upper primer is the forward primer and the lower primer is the reverse primer.

glass slides were sealed under coverslips using Vectashield Mounting Medium (Vector Laboratories, Burlingame, CA) and examined under an IX71 microscope connected to a Cool SNAP HQ digital camera (Olympus, Tokyo, Japan).

Conversion of arachidonic acid to PGE₂, 6-keto-PGF_{1α}, and TXB₂

Cells stimulated with IL-1β or TNF-α were washed with DMEM and incubated for a further 30 min with 3 μM arachidonic acid, as reported previously [25,34]. The culture medium was then harvested, and the concentrations of PGE₂, 6-keto-PGF_{1α} (a stable metabolite of PGI₂), and TXB₂ (a stable metabolite of TXA₂) were measured by ELISA.

Measurement of PGES activity

PGES activity was measured by assessment of the conversion of PGH₂ to PGE₂, as reported previously [25,35]. In brief, cells (at a density of 5 × 10⁴ cells/cm²) were cultured under various conditions in medium containing 1% FBS. The cells were then scraped off the dish and lysed by sonication (twice for 10 s each at 1 min intervals) in 100 μl of 1 M Tris-HCl (pH 8.0). After centrifugation of the lysate at 15,000 g for 1 min at 4°C, the supernatant was used to measure PGES activity. For assessment of PGES activity, an aliquot of each supernatant equivalent to 50 μg of protein was incubated with 2 μg of PGH₂ for 30 s at 24°C in 0.1 ml of 1 M Tris-HCl containing 2 mM glutathione, in the absence or presence of MK-886. The reaction was terminated by the addition of 100 mM FeCl₂; the reaction mixture was then left to stand at 20–25°C for 15 min. After

centrifugation of the reaction mixture, the PGE₂ concentration in the supernatant was measured with an ELISA kit.

Measurement of prostaglandins in culture medium

Cells were seeded at a density of 5 × 10⁴ cells/cm² in culture medium containing 1% FBS and were incubated with IL-1β for 48 h. In experiments involving treatment with MK-886, it was added 30 min before IL-1β. The culture supernatant was then harvested and the PGE₂ concentration was measured with an ELISA kit.


Immunohistochemical analysis

Tissue sections were incubated for 5 min with 3% (v/v) H₂O₂. After blocking with 1% (w/v) bovine serum albumin in PBS for 30 min at 20–25°C, rabbit anti-human mPGES antibody or goat anti-human COX-2 antibody (diluted 1:100 or 1:200, respectively, in PBS containing 1% albumin) was added and incubation was performed for 1.5 h at 20–25°C. After being washed, the sections were incubated with the biotinylated secondary antibody, followed by streptavidin-HRP complex and 3,3'-diaminobenzidine tetrahydrochloride in accordance with the manufacturer's protocol (LSAB staining kit; Dako, Carpinteria, CA). As adsorption controls, primary antibodies against mPGES-1 and COX-2 were used that had been previously reacted with mPGES-1 (10 μg/ml) or COX-2 peptide (10 μg/ml), respectively. In addition, rabbit anti-human mPGES-1 was replaced with normal rabbit IgG as a negative control.

Statistical analysis

Results are expressed as means ± SD for triplicate experiments, and representative results from three independent

Figure 2



Effects of cytokines on the expression of prostaglandin E synthase (PGES) and cyclooxygenase (COX) isoenzyme mRNAs in chondrocytes. **(a)** Chondrocytes from patients with osteoarthritis were incubated with interleukin (IL)-1 β (1 ng/ml), tumour necrosis factor (TNF α ; 10 ng/ml), IL-4 (10 ng/ml), IL-6 (10 ng/ml), IL-8 (10 ng/ml), IL-10 (10 ng/ml), or interferon- γ (IFN γ ; 100 ng/ml) for 12 h. Total RNA from the cells was subjected to reverse transcription-polymerase chain reaction (RT-PCR) for membrane-associated PGES (mPGES)-1, COX-2, and glyceraldehyde-3-phosphate dehydrogenase (GAPDH), as described in Materials and methods. Representative results from three patients are shown here. **(b)** Chondrocytes from four patients with osteoarthritis were incubated with IL-1 β (1 ng/ml) or TNF- α (10 ng/ml) for 12 h. Total RNA from the cells was subjected to RT-PCR for mPGES-1, mPGES-2, cytosolic PGES (cPGES), COX-1, COX-2, and GAPDH, as described in Materials and methods. Representative results are shown here.

patients are shown. Statistical analysis for triplicate experiments was performed with Tukey's multiple comparison test; $P < 0.05$ was considered significant.

Results

Phenotypic features of chondrocytes

Morphology and collagen mRNA expression were examined for assessment of the chondrocyte phenotype. The morphology of chondrocytes used in this study was obviously different from that of synovial fibroblasts (Fig. 1a). Type I collagen mRNA was expressed in chondrocytes as well as synovial fibroblasts and dermal fibroblasts. In contrast, expression of type II collagen mRNA was detected only in chondrocytes, as shown in Fig. 1b. Moreover, expression of type II collagen mRNA as well as type I colla-

gen mRNA in chondrocytes was maintained after stimulation of IL-1 β or TNF- α for 12 h (Fig. 1c).


Effect of cytokines on expression of mRNAs for PGES and COX isoenzymes in chondrocytes

The expression of mRNA for mPGES-1 and COX-2 in chondrocytes from patients with OA was assessed by RT-PCR after the cells had been treated with various cytokines for 12 h. As shown in Fig. 2a, TNF- α or IL-1 β induced the expression of mPGES-1 and COX-2 mRNA in chondrocytes. In contrast, IL-4, IL-6, IL-8, IL-10, and interferon- γ all had no effect on the expression of mPGES-1 or COX-2 mRNA.

We next examined the induction of mRNA for PGES and COX isoenzymes by IL-1 β or TNF- α in chondrocytes from four patients with OA (Fig. 2b). Expression of mPGES-1 and COX-2 mRNA was significantly induced by IL-1 β or TNF- α in chondrocytes from all four patients. In contrast, mRNA for mPGES-2, cPGES, and COX-1 was constitutively expressed in unstimulated chondrocytes, and exposure to IL-1 β or TNF- α had little effect on the expression of these mRNAs. The pattern of mRNA expression for each enzyme was similar in all four patients.


Subsequently, we examined the effect of various concentrations of IL-1 β and TNF- α on mPGES-1 and COX-2 mRNA by Northern blot analysis. As shown in Fig. 3, expression of mPGES-1 and COX-2 mRNA was induced by IL-1 β in a concentration-dependent manner. TNF- α also induced expression of mPGES-1 mRNA, but had a weaker effect than IL-1 β . The effect of TNF- α on the expression of COX-2 mRNA was also weaker than that of IL-1 β .

Figure 3



Expression of membrane-associated prostaglandin E synthase-1 (mPGES-1) and cyclooxygenase-2 (COX-2) mRNA in cytokine-stimulated chondrocytes. Chondrocytes from patients with osteoarthritis were incubated with interleukin-1 β (IL-1 β ; 0.01–10 ng/ml) or tumor necrosis factor- α (TNF- α ; 0.01–10 ng/ml) for 12 h. Total RNA from the cells was subjected to Northern blot analysis for mPGES-1, COX-2, and glyceraldehyde-3-phosphate dehydrogenase (GAPDH), as described in Materials and methods. Representative results from three patients are shown.

Figure 4



Expression of membrane-associated prostaglandin E synthase (mPGES-1) and cyclooxygenase-2 (COX-2) protein in cytokine-stimulated chondrocytes. **(a)** Chondrocytes from osteoarthritis patients were incubated for 48 h with interleukin-1 β (IL-1 β ; 0.01–10 ng/ml) or tumor necrosis factor- α (TNF α ; 0.01–10 ng/ml). Protein from the cells was subjected to Western blot analysis for mPGES-1 and COX-2. **(b)** Chondrocytes from osteoarthritis patients were incubated for 48 h with IL-1 β (1 ng/ml) or TNF- α (10 ng/ml). The cells were then double-immunostained with tetramethylrhodamine β -isothiocyanate for mPGES-1 (upper row) and fluorescein isothiocyanate for COX-2 (lower row), and were examined under a Fluoview laser fluorescence microscope. Representative findings from three patients are shown here.


Expression of mPGES-1 and COX-2 protein in cytokine-stimulated chondrocytes

Expression of mPGES-1 and COX-2 protein in chondrocytes stimulated with IL-1 β or TNF- α was investigated by Western blot analysis and immunofluorescent microscopy (Fig. 4). As shown in Fig. 4a, both mPGES-1 and COX-2 protein were induced by IL-1 β and TNF- α , similar to the results for mRNA expression shown in Fig. 3. Colocalization of these enzymes was observed in the perinuclear region of IL-1 β -stimulated chondrocytes (Fig. 4b).

Production of PGE₂, 6-keto-PGF_{1 α} , and TXB₂ in cytokine-stimulated chondrocytes

To determine the profile of PG synthesis in OA chondrocytes, cells were exposed to various concentrations of IL-1 β (0.01–10 ng/ml) or TNF- α (0.01–10 ng/ml) for 24 h and the conversion of exogenous arachidonic acid to PGE₂, 6-keto-PGF_{1 α} (a stable metabolite of PGI₂), and TXB₂ (a stable metabolite of TXA₂) was measured by ELISA. As shown in Fig. 5a, PGE₂ synthesis was markedly increased by IL-1 β or TNF- α (to a smaller extent) in a concentration-dependent manner. The pattern of PGE₂ production was similar to results shown in Fig. 2. In contrast, the changes in 6-keto-PGF_{1 α} (Fig. 5b) and TXB₂ (Fig. 5c) after the addition of IL-

Figure 5



Production of prostaglandin E₂ (PGE₂) **(a)**, 6-keto-PGF_{1 α} **(b)**, and thromboxane (TX)₂ **(c)** in cytokine-stimulated chondrocytes. Chondrocytes from osteoarthritis patients were incubated for 24 h with interleukin-1 β (IL-1 β ; 0.01–10 ng/ml, filled circles) or tumor necrosis factor- α (0.01–10 ng/ml, open circles). After being washed, the cells were incubated for a further 30 min with 3 μ M arachidonic acid; the PGE₂, 6-keto-PGF_{1 α} (a stable metabolite of PGI₂), and TXB₂ (a stable metabolite of TXA₂) concentrations in the medium were then measured by enzyme-linked immunosorbent assay. Data are means \pm SD for triplicate cultures, and representative results from three patients are shown. * P < 0.01 compared with untreated control cells.

1 β or TNF- α were almost negligible compared with the effect on PGE₂.


Time course of the expression of PGES and COX isoenzymes in IL-1 β -stimulated chondrocytes

The expression of mRNA (Fig. 6a) and protein (Fig. 6b) for PGES and COX isoenzymes at various times after stimulation with IL-1 β was assessed by Northern and Western blotting, respectively. Expression of mPGES-1 mRNA showed a gradual increase, reaching a maximum after 24–48 h of exposure to IL-1 β . Expression of mPGES-1 protein showed a slight increase at 12 h, and then gradually increased until at least 72 h. In contrast, COX-2 mRNA increased rapidly after the start of IL-1 β stimulation, and its expression remained maximal until 48 h. The COX-2 protein level also showed a rapid increase within 6 h and expression was maximal between 12 and 48 h; it decreased again at 72 h. Expression of mPGES-1 was delayed compared with that of COX-2. In contrast, the levels of cPGES and COX-1 mRNA and protein were not affected by IL-1 β .

Induction of PGES activity in IL-1 β -stimulated chondrocytes

To investigate whether PGES enzymatic activity was upregulated by IL-1 β , we measured the conversion of

Figure 6




Time course of the expression of prostaglandin E synthase (PGES) and cyclooxygenase (COX) isoenzymes in chondrocytes stimulated with interleukin-1 β (IL-1 β). Chondrocytes from osteoarthritis patients were incubated with IL-1 β (1 ng/ml) for the indicated durations. **(a)** Total RNA from the cells was subjected to Northern blot analysis for membrane-associated PGES-1 (mPGES-1), cytosolic PGES (cPGES), COX-1, COX-2, and glyceraldehyde-3-phosphate dehydrogenase (GAPDH). **(b)** Protein from the cells was subjected to Western blot analysis for mPGES-1, cPGES, COX-1, and COX-2. Representative results from three patients are shown here.

PGH₂ to PGE₂ in lysates prepared from IL-1 β -stimulated chondrocytes at various times (Fig. 7a). PGES activity increased gradually until 72 h after the start of IL-1 β stimulation. These results matched the pattern of mPGES-1 protein expression (Fig. 6b).

It was reported that MK-886 could inhibit mPGES-1 activity in mPGES-1 transfected cells [28]. To evaluate the contribution of mPGES-1 to the IL-1 β -induced increase of PGES activity in chondrocytes, we examined the effect of MK-886 on PGES activity in chondrocyte lysates. As shown in Fig. 7b, PGES activity in lysates from unstimulated chondrocytes was very low, and it was not affected by MK-886 alone. In contrast, the increased level of PGES activity in lysates prepared from IL-1 β -stimulated chondrocytes was inhibited by MK-886 in a concentration-dependent manner. These results suggested that the induction of PGES activ-

Figure 7



Induction of prostaglandin E synthase (PGES) activity in chondrocytes stimulated with IL-1 β (IL-1 β). **(a)** For estimation of the PGES activity that produces prostaglandin (PG)E₂ from PGH₂, exogenous PGH₂ (2 μg) was added to lysates of chondrocytes that had been incubated with (filled circles) or without (open circles) IL-1 β (1 ng/ml) for the indicated durations. The lysates were then incubated at 24°C for 30 s in the presence of reduced glutathione, the reaction was stopped with 100 mM FeCl₂, and the PGE₂ concentration was measured by enzyme-linked immunosorbent assay. Spontaneous conversion of PGH₂ to PGE₂ in the absence of cell lysates was subtracted from each result. Data are means \pm SD for triplicate cultures; representative results from three patients are shown. * P < 0.01 and * P < 0.05 compared with 0 h. **(b)** Chondrocytes were incubated with or without IL-1 β (1 ng/ml) for 48 h. Then the PGES activity of the cell lysates in the presence or absence of MK-886 (1–100 μM) was measured as described in Materials and methods, and calculated as a percentage of the control value (IL-1 β alone). Data are means \pm SD for triplicate experiments; representative results from three patients are shown. # P < 0.01, untreated control compared with IL-1 β alone; * P < 0.01, IL-1 β alone compared with IL-1 β plus MK-886.

ity by IL-1 β in chondrocytes was dependent mainly on the enzymatic activity of inducible mPGES-1 rather than on constitutive mPGES-2 and cPGES.


Effect of MK-886 on PGE₂ production in IL-1 β -stimulated chondrocytes

We next examined the effect of MK-886 on endogenous PGE₂ production in IL-1 β -stimulated chondrocytes. Chondrocytes were incubated with IL-1 β for 48 h in the absence or presence of MK-886. As shown in Fig. 8, endogenous PGE₂ production by OA chondrocytes was significantly increased after stimulation with IL-1 β . When chondrocytes were cultured with IL-1 β in the presence of MK-886, there was significant inhibition of PGE₂ production in a concentration-dependent manner. Microscopy of the cells revealed severe toxicity of MK-886 at 30 μM (data not shown).

Expression of mPGES-1 protein in articular cartilage from patients with OA

Articular cartilage obtained from patients with OA was fixed in formaldehyde, embedded in paraffin, and cut into serial sections that were immunostained with antibody against mPGES-1 or COX-2. Chondrocytes in the articular cartilage from patients with OA were stained by the anti-

Figure 8




Effect of MK-886 on endogenous prostaglandin E₂ (PGE₂) production in chondrocytes stimulated with IL-1β (IL-1β). Chondrocytes from patients with osteoarthritis were incubated with IL-1β in the presence of MK-886 (1 or 10 μM) for 48 h. The PGE₂ concentration of the medium was then measured by enzyme-linked immunosorbent assay and calculated as a percentage of the control value (IL-1β alone). Data are means ± SD for triplicate experiments; representative results from three patients are shown. #*P* < 0.01, untreated control compared with IL-1β alone; **P* < 0.01, IL-1β alone compared with IL-1β plus MK-886.

body against mPGES-1 (Fig. 9a). When the absorption control using mPGES-1 peptide was tested (Fig. 9b), mPGES-1 immunostaining was almost completely abolished. In addition, no mPGES-1 immunostaining was observed when rabbit anti-human mPGES-1 was replaced by normal rabbit IgG (Fig. 9c). COX-2 immunostaining was also positive in the chondrocytes (Fig. 9d), whereas the absorption control using COX-2 peptide (Fig. 9e) showed no COX-2 immunostaining.

Discussion

In this study we demonstrated that mPGES-1 is a key enzyme in the regulation of PGE₂ production in cytokine-stimulated chondrocytes from patients with OA. Although several studies on the induction of COX-2 and its regulatory mechanism in chondrocytes have already been published [16,36,37], this is the first report that mPGES-1, located downstream of COX-2, is induced in chondrocytes by cytokine stimulation. Among the various cytokines that we tested, IL-1β and TNF-α caused the induction of both mPGES-1 expression and COX-2 expression. In addition we demonstrated mPGES-1 and COX-2 immunostaining of chondrocytes in articular cartilage from patients with OA. There have been reports of a high level of IL-1β and TNF-α expression on the surface of OA cartilage, and the production of IL-1β in OA tissues seems to be correlated with the extent of cartilage damage [4,38-40]. It was also reported that IL-1β expression is increased in synovial membrane tissues obtained from patients with OA at all stages of the disease [5,41]. Our results suggested that

Figure 9



Expression of membrane-associated prostaglandin E synthase-1 (mPGES-1) and cyclooxygenase-2 (COX-2) protein in articular cartilage sections from osteoarthritis patients. Paraffin-embedded section of cartilage from osteoarthritis patients were subjected to immunostaining for mPGES-1 (a) and COX-2 (d). As adsorption controls for mPGES-1 (b) and COX-2 (e), primary antibodies that had been reacted with mPGES-1 and COX-2 peptide (10 μg/ml) were used, respectively. As a negative control for mPGES-1, rabbit anti-human mPGES-1 was replaced with normal rabbit IgG (c). Representative findings from three patients are shown here.

expression of mPGES-1 and COX-2 in chondrocytes from patients with OA might be promoted by endogenous IL-1β and TNF-α released from cartilage and/or synovial tissue.

Long-term culture and/or multiple passages of chondrocytes generally induce dedifferentiation of the cells. The existence of type II collagen, which is highly specific to chondrocytes, is one of the major phenotypic characterizations of this cell type; dedifferentiated chondrocytes lose the expression of type II collagen [42]. In our study, the chondrocytes that we used expressed type II collagen and were clearly distinguished from synovial fibroblasts and dermal fibroblasts. Thus, we estimated that the chondrocytes in our study were maintained in a differentiated condition.

We also studied the profile of PG synthesis in chondrocytes. PGE₂ synthesis was markedly induced by IL-1β or TNF-α, whereas 6-keto-PGF_{1α} and TXB₂ synthesis were almost negligible in comparison. These results suggest that terminal PG synthase downstream of COX is important for deciding which PG is synthesized. In this study we showed that the expression of mPGES-1 and COX-2 was markedly induced by IL-1β in parallel with an increase in PGE₂ synthesis. In contrast, PGI synthase and TX synthase mRNAs were detectable in chondrocytes by RT-PCR but were not induced by IL-1β (data not shown). Moreover, MK-886, which has an inhibitory effect on mPGES-1 activity [28], significantly inhibited IL-1β-induced PGES activity in lysates from IL-1β-stimulated chondrocytes. This inhibitor also reduced PGE₂ production by intact chondrocytes after stimulation by IL-1β. These results suggest that the induction of mPGES-1 might be necessary for substantial PGE₂ production by chondrocytes in response to proinflammatory

tory stimuli. However, mPGES-1 is a member of the MAPEG (membrane-associated proteins involved in eicosanoid and glutathione metabolism) family, which also contains 5-lipoxygenase-activating protein and leukotriene C₄ synthase. It has been shown that MK-886 inhibits not only mPGES-1 but also 5-lipoxygenase-activating protein [43] and LTC₄ synthase [44]. Alternative specific methods for inhibiting mPGES-1 therefore remain to be studied.

Immunohistochemistry revealed that mPGES-1 and COX-2 protein colocalized to the perinuclear region of IL-1 β -stimulated chondrocytes. We recently reported that mPGES-1 and COX-2 also showed perinuclear colocalization in IL-1 β -stimulated rheumatoid arthritis synovial fibroblasts [25]. Murakami and colleagues showed that mPGES-1 and COX-2 are functionally linked by using an mPGES-1 and COX-2 transfected cell line [24]. Colocalization of mPGES-1 and COX-2 might be related to the preferential increase of PGE₂ production after IL-1 β stimulation due to functional linkage between these two enzymes in chondrocytes.

The present study showed that the induction of mPGES-1 expression in chondrocytes was delayed compared with that of COX-2 and that it persisted for longer after stimulation with IL-1 β . We previously found that increased expression of mPGES-1 was delayed in a similar fashion in IL-1 β -stimulated rheumatoid arthritis synovial fibroblasts [25]. This coordinated expression of mPGES-1 and COX-2 over time might sustain higher PGE₂ production in OA chondrocytes. The different profiles of mPGES-1 and COX-2 expression also imply that regulation of these enzymes occurs through different pathways in chondrocytes. It was recently shown that mPGES-1 expression in response to various stimuli is regulated by a transcription factor, early growth response factor-1 [45], which was reported to bind to the proximal GC box in the mPGES-1 gene promoter. If different transcription factors are associated with mPGES-1 and COX-2, this might help to explain the different time course of the expression of these enzymes in IL-1 β -stimulated chondrocytes.

Cartilage damage is considered to be a major cause of OA, and chondrocytes seem to have a fundamental role in the pathogenesis of this condition [46]. It was reported that cartilage is the main target of cytokines such as IL-1 β and TNF- α in OA, and it seems that activated chondrocytes might be a major source of various inflammatory mediators associated with the progressive destruction of articular cartilage [47,48]. Accordingly, the induction of mPGES-1 in cytokine-activated chondrocytes and the subsequent increase of PGE₂ production might be important in the pathogenesis of OA. mPGES-1 is a specific enzyme that only catalyzes PGE₂ production from PGH₂. mPGES-1 might therefore be a potential new target for therapeutic

strategies to specifically control the PGE₂ synthesis associated with inflammation in patients with OA and other inflammatory diseases.

Conclusion

mPGES-1 was upregulated by proinflammatory cytokines in OA chondrocytes and was one of the key enzymes in the regulation of PGE₂ production. mPGES-1 might be important in articular inflammation with cytokine-stimulated conditions in patients with OA.

Competing interests

None declared.

Author contributions

FK participated mainly in the experiments and drafted the manuscript.

HN suggested the methodology of the experiments and assessed the results.

SM, MB, and HA provided samples from the patients and assessed the results.

SK conceived the study and participated in its design and coordination.

All authors read and approved the final manuscript.

Acknowledgements

We thank Mr Soichiro Kato, Ms Toshiko Mogi, Ms Hiroe Ogasawara, and Ms Natsuko Kusunoki for their support, and Ms Sonoko Sakurai for secretarial assistance. We also thank Dr Masako Mizoguchi and Dr Emiko Yamasaki for providing dermal fibroblasts. This work was partly supported by research grants from The Ministry of Education, Culture, Sports, Science and Technology of Japan.

References

1. Pelletier JP, DiBattista JA, Roughley P, McCollum R, Martel-Pelletier J: **Cytokines and inflammation in cartilage degradation.** *Rheum Dis Clin North Am* 1993, **19**:545-568.
2. Martel-Pelletier J, Alaaeddine N, Pelletier JP: **Cytokines and their role in the pathophysiology of osteoarthritis.** *Front Biosci* 1999, **4**:D694-D703.
3. Fernandes JC, Martel-Pelletier J, Pelletier JP: **The role of cytokines in osteoarthritis pathophysiology.** *Biorheology* 2002, **39**:237-246.
4. Martel-Pelletier J, McCollum R, Fujimoto N, Obata K, Cloutier JM, Pelletier JP: **Excess of metalloproteases over tissue inhibitor of metalloprotease may contribute to cartilage degradation in osteoarthritis and rheumatoid arthritis.** *Lab Invest* 1994, **70**:807-815.
5. Chevalier X: **Upregulation of enzymatic activity by interleukin-1 in osteoarthritis.** *Biomed Pharmacother* 1997, **51**:58-62.
6. Tetlow LC, Adlam DJ, Woolley DE: **Matrix metalloproteinase and proinflammatory cytokine production by chondrocytes of human osteoarthritic cartilage: associations with degenerative changes.** *Arthritis Rheum* 2001, **44**:585-594.
7. Dodge GR, Poole AR: **Immunohistochemical detection and immunochemical analysis of type II collagen degradation in human normal, rheumatoid, and osteoarthritic articular cartilages and in explants of bovine articular cartilage cultured with interleukin-1.** *J Clin Invest* 1989, **83**:647-661.

8. Reboul P, Pelletier JP, Tardif G, Cloutier JM, Martel-Pelletier J: **The new collagenase, collagenase-3, is expressed and synthesized by human chondrocytes but not by synoviocytes. A role in osteoarthritis.** *J Clin Invest* 1996, **97**:2011-2019.
9. Shinmei M, Masuda K, Kikuchi T, Shimomura Y, Okada Y: **Production of cytokines by chondrocytes and its role in proteoglycan degradation.** *J Rheumatol* 1991, **Suppl 27**:89-91.
10. Amin AR, Abramson SB: **The role of nitric oxide in articular cartilage breakdown in osteoarthritis.** *Curr Opin Rheumatol* 1998, **10**:263-268.
11. Melchiorri C, Meliconi R, Frizziero L, Silvestri T, Pulsatelli L, Mazzetti I, Borzi RM, Uguccioni M, Facchini A: **Enhanced and coordinated *in vivo* expression of inflammatory cytokines and nitric oxide synthase by chondrocytes from patients with osteoarthritis.** *Arthritis Rheum* 1998, **41**:2165-2174.
12. Studer R, Jaffurs D, Stefanovic-Racic M, Robbins PD, Evans CH: **Nitric oxide in osteoarthritis.** *Osteoarthr Cartilage* 1999, **7**:377-379.
13. Tawara T, Shingu M, Nobunaga M, Naono T: **Effects of recombinant human IL-1 beta on production of prostaglandin E₂, leukotriene B₄, NAG, and superoxide by human synovial cells and chondrocytes.** *Inflammation* 1991, **15**:145-157.
14. Jacques C, Sautet A, Moldovan M, Thomas B, Humbert L, Berenbaum F: **Cyclooxygenase activity in chondrocytes from osteoarthritic and healthy cartilage.** *Rev Rhum Engl Ed* 1999, **66**:701-704.
15. Hardy MM, Seibert K, Manning PT, Currie MG, Woerner BM, Edwards D, Koki A, Tripp CS: **Cyclooxygenase 2-dependent prostaglandin E₂ modulates cartilage proteoglycan degradation in human osteoarthritis explants.** *Arthritis Rheum* 2002, **46**:1789-1803.
16. Notoya K, Jovanovic DV, Reboul P, Martel-Pelletier J, Mineau F, Pelletier JP: **The induction of cell death in human osteoarthritis chondrocytes by nitric oxide is related to the production of prostaglandin E₂ via the induction of cyclooxygenase-2.** *J Immunol* 2000, **165**:3402-3410.
17. Xie WL, Chipman JG, Robertson DL, Erikson RL, Simmons DL: **Expression of a mitogen-responsive gene encoding prostaglandin synthase is regulated by mRNA splicing.** *Proc Natl Acad Sci USA* 1991, **88**:2692-2696.
18. Kujubu DA, Fletcher BS, Varnum BC, Lim RW, Herschman HR: **TIAS10, a phorbol ester tumor promoter-inducible mRNA from Swiss 3T3 cells, encodes a novel prostaglandin synthase/cyclooxygenase homologue.** *J Biol Chem* 1991, **266**:12866-12872.
19. Urade Y, Watanabe K, Hayaishi O: **Prostaglandin D, E, and F synthases.** *J Lipid Mediat Cell Signal* 1995, **12**:257-273.
20. Kudo I, Murakami M: **Diverse functional coupling of prostanoid biosynthetic enzymes in various cell types.** *Adv Exp Med Biol* 1999, **469**:29-35.
21. Jakobsson PJ, Thoren S, Morgenstern R, Samuelsson B: **Identification of human prostaglandin E synthase: a microsomal, glutathione-dependent, inducible enzyme, constituting a potential novel drug target.** *Proc Natl Acad Sci USA* 1999, **96**:7220-7225.
22. Tanikawa N, Ohmiya Y, Ohkubo H, Hashimoto K, Kangawa K, Kojima M, Ito S, Watanabe K: **Identification and characterization of a novel type of membrane-associated prostaglandin E synthase.** *Biochem Biophys Res Commun* 2002, **291**:884-889.
23. Tanioka T, Nakatani Y, Semmyo N, Murakami M, Kudo I: **Molecular identification of cytosolic prostaglandin E₂ synthase that is functionally coupled with cyclooxygenase-1 in immediate prostaglandin E₂ biosynthesis.** *J Biol Chem* 2000, **275**:32775-32782.
24. Murakami M, Naraba H, Tanioka T, Semmyo N, Nakatani Y, Kojima F, Ikeda T, Fueki M, Ueno A, Oh S, Kudo I: **Regulation of prostaglandin E₂ biosynthesis by inducible membrane-associated prostaglandin E₂ synthase that acts in concert with cyclooxygenase-2.** *J Biol Chem* 2000, **275**:32783-32792.
25. Kojima F, Naraba H, Sasaki Y, Okamoto R, Koshino T, Kawai S: **Coexpression of microsomal prostaglandin E synthase with cyclooxygenase-2 in human rheumatoid synovial cells.** *J Rheumatol* 2002, **29**:1836-1842.
26. Kojima F, Naraba H, Sasaki Y, Beppu M, Aoki H, Kawai S: **Prostaglandin E₂ is an enhancer of interleukin-1beta-induced expression of membrane-associated prostaglandin E synthase in rheumatoid synovial fibroblasts.** *Arthritis Rheum* 2003, **48**:2819-2828.
27. Stichtenoth DO, Thoren S, Bian H, Peters-Golden M, Jakobsson PJ, Crofford LJ: **Microsomal prostaglandin E synthase is regulated by proinflammatory cytokines and glucocorticoids in primary rheumatoid synovial cells.** *J Immunol* 2001, **167**:469-474.
28. Mancini JA, Blood K, Guay J, Gordon R, Claveau D, Chan CC, Riendeau D: **Cloning, expression, and up-regulation of inducible rat prostaglandin E synthase during lipopolysaccharide-induced pyresis and adjuvant-induced arthritis.** *J Biol Chem* 2001, **276**:4469-4475.
29. Claveau D, Sirinyan M, Guay J, Gordon R, Chan CC, Bureau Y, Riendeau D, Mancini JA: **Microsomal prostaglandin E synthase-1 is a major terminal synthase that is selectively up-regulated during cyclooxygenase-2-dependent prostaglandin E₂ production in the rat adjuvant-induced arthritis model.** *J Immunol* 2003, **170**:4738-4744.
30. Trebino CE, Stock JL, Gibbons CP, Naiman BM, Wachtmann TS, Umland JP, Pandher K, Lapointe JM, Saha S, Roach ML, Carter D, Thomas NA, Durtschi BA, McNeish JD, Hambor JE, Jakobsson PJ, Carty TJ, Perez JR, Audoly LP: **Impaired inflammatory and pain responses in mice lacking an inducible prostaglandin E synthase.** *Proc Natl Acad Sci USA* 2003, **100**:9044-9049.
31. Altman R, Asch E, Bloch D, Bole G, Borenstein D, Brandt K, Christy W, Cooke TD, Greenwald R, Hochberg M, Howell D, Kaplan D, Koopman W, Longley S, Mankin H III, McShane DJ, Medsger T, Meenan R Jr, Mikkelsen W, Moskowitz R, Murphy W, Rothschild B, Segal M, Sokoloff L, Wolfe F: **Development of criteria for the classification and reporting of osteoarthritis. Classification of osteoarthritis of the knee. Diagnostic and Therapeutic Criteria Committee of the American Rheumatism Association.** *Arthritis Rheum* 1986, **29**:1039-1049.
32. Yuan GH, Masuko-Hongo K, Sakata M, Tsuruha J, Onuma H, Nakamura H, Aoki H, Kato T, Nishioka K: **The role of C-C chemokines and their receptors in osteoarthritis.** *Arthritis Rheum* 2001, **44**:1056-1070.
33. Inoue H, Ohshima H, Kono H, Yamanaka M, Kubota T, Aihara M, Hiroi T, Yago N, Ishida H: **Suppressive effects of tranilast on the expression of inducible cyclooxygenase (COX2) in interleukin-1beta-stimulated fibroblasts.** *Biochem Pharmacol* 1997, **53**:1941-1944.
34. Kawai S, Nishida S, Kato M, Furumaya Y, Okamoto R, Koshino T, Mizushima Y: **Comparison of cyclooxygenase-1 and -2 inhibitory activities of various nonsteroidal anti-inflammatory drugs using human platelets and synovial cells.** *Eur J Pharmacol* 1998, **347**:87-94.
35. Matsumoto H, Naraba H, Murakami M, Kudo I, Yamaki K, Ueno A, Oh-ishi S: **Concordant induction of prostaglandin E₂ synthase with cyclooxygenase-2 leads to preferred production of prostaglandin E₂ over thromboxane and prostaglandin D₂ in lipopolysaccharide-stimulated rat peritoneal macrophages.** *Biochem Biophys Res Commun* 1997, **230**:110-114.
36. Thomas B, Thirion S, Humbert L, Tan L, Goldring MB, Berezziat G, Berenbaum F: **Differentiation regulates interleukin-1beta-induced cyclo-oxygenase-2 in human articular chondrocytes: role of p38 mitogen-activated protein kinase.** *Biochem J* 2002, **362**:367-373.
37. Thomas B, Berenbaum F, Humbert L, Bian H, Berezziat G, Crofford L, Olivier JL: **Critical role of C/EBPdelta and C/EBPbeta factors in the stimulation of the cyclooxygenase-2 gene transcription by interleukin-1beta in articular chondrocytes.** *Eur J Biochem* 2000, **267**:6798-6809.
38. Towle CA, Hung HH, Bonassar LJ, Treadwell BV, Mangham DC: **Detection of interleukin-1 in the cartilage of patients with osteoarthritis: a possible autocrine/paracrine role in pathogenesis.** *Osteoarthritis Cartilage* 1997, **5**:293-300.
39. Moos V, Fickert S, Muller B, Weber U, Sieper J: **Immunohistological analysis of cytokine expression in human osteoarthritic and healthy cartilage.** *J Rheumatol* 1999, **26**:870-879.
40. Barakat AF, Elson CJ, Westacott CI: **Susceptibility to physiological concentrations of IL-1beta varies in cartilage at different anatomical locations on human osteoarthritic knee joints.** *Osteoarthr Cartilage* 2002, **10**:264-269.
41. Smith MD, Triantafyllou S, Parker A, Youssef PP, Coleman M: **Synovial membrane inflammation and cytokine production in patients with early osteoarthritis.** *J Rheumatol* 1997, **24**:365-371.

42. Schnabel M, Marlovits S, Eckhoff G, Fichtel I, Gotzen L, Vecsei V, Schlegel J: **Dedifferentiation-associated changes in morphology and gene expression in primary human articular chondrocytes in cell culture.** *Osteoarthr Cartilage* 2002, **10**:62-70.
43. Reid GK, Kargman S, Vickers PJ, Mancini JA, Leveille C, Ethier D, Miller DK, Gillard JW, Dixon RA, Evans JF: **Correlation between expression of 5-lipoxygenase-activating protein, 5-lipoxygenase, and cellular leukotriene synthesis.** *J Biol Chem* 1990, **265**:19818-19823.
44. Lam BK, Penrose JF, Freeman GJ, Austen KF: **Expression cloning of a cDNA for human leukotriene C4 synthase, an integral membrane protein conjugating reduced glutathione to leukotriene A4.** *Proc Natl Acad Sci USA* 1994, **91**:7663-7667.
45. Naraba H, Yokoyama C, Tago N, Murakami M, Kudo I, Fueki M, Ohishi S, Tanabe T: **Transcriptional regulation of the membrane-associated prostaglandin E₂ synthase gene. Essential role of the transcription factor Egr-1.** *J Biol Chem* 2002, **277**:28601-28608.
46. Berenbaum F: **Osteoarthritis: epidemiology, pathology, and pathogenesis.** In *Primer on the Rheumatic Diseases* 12th edition. Edited by: Klippel JH, Crofford LJ, Stone JH, Weyand CM. Atlanta: Arthritis Foundation; 2001:285-289.
47. Ruggeri R, Pulsatelli L, Melchiorri C, Da Re R, Focherini MC, Veronesi M, Facchini A: **Differential expression of IL-1 and TNF-receptors in inflammatory arthritis and osteoarthritis.** *Boll Soc Ital Biol Sper* 1996, **72**:15-20.
48. Melchiorri C, Meliconi R, Frizziero L, Silvestri T, Pulsatelli L, Mazzetti I, Borzi RM, Uguccioni M, Facchini A: **Enhanced and coordinated *in vivo* expression of inflammatory cytokines and nitric oxide synthase by chondrocytes from patients with osteoarthritis.** *Arthritis Rheum* 1998, **41**:2165-2174.