

2-(Furan-2-yl)-1,3-bis(furan-2-ylmethyl)-1*H*-benzimidazol-3-ium chloride mono-hydrate

David K. Geiger,* H. Cristina Geiger and Jared M. Deck

Department of Chemistry, State University of New York-College at Geneseo,
1 College Circle, Geneseo, NY 14454, USA
Correspondence e-mail: geiger@geneseo.edu


Received 22 August 2012; accepted 21 September 2012

Key indicators: single-crystal X-ray study; $T = 200$ K; mean $\sigma(C-C) = 0.004$ Å;
disorder in main residue; R factor = 0.045; wR factor = 0.129; data-to-parameter
ratio = 12.6.

The title hydrated salt, $C_{21}H_{17}N_2O_3^+\cdot Cl^- \cdot H_2O$, exhibits disorder in one of the furan rings. The major and minor components have a refined occupancy ratio of 0.844 (19):0.156 (19). The structure displays intermolecular hydrogen bonding involving the water molecule and the chloride anion. Close intermolecular C—H \cdots Cl and C—H \cdots (furan ring) interactions complete the hydrogen bonding.

Related literature

For examples of the synthesis of substituted benzimidazolium salts, see: Wang & Chang (2006); Hoesl *et al.* (2004); Rivas *et al.* (2001, 2002). For examples of structures of other trisubstituted benzimidazolium salts, see: Ennajih *et al.* (2009, 2010); Akkurt *et al.* (2008, 2004); Smith & Luss (1975). For the structure of 1,3-bis(furan-2-ylmethyl)-1*H*-benzo[d]imidazol-3-ium chloride, see: Akkurt *et al.* (2009). For other related literature, see: Costache *et al.* (2007); Elmali *et al.* (2005); Hayakawa *et al.* (1996); Horton *et al.* (2003); Nahlé *et al.* (2012); Welton (1999).



Experimental

Crystal data

$C_{21}H_{17}N_2O_3^+\cdot Cl^- \cdot H_2O$	$\gamma = 93.547$ (4) $^\circ$
$M_r = 398.83$	$V = 976.3$ (2) Å 3
Triclinic, $P\bar{1}$	$Z = 2$
$a = 9.4723$ (12) Å	Mo $K\alpha$ radiation
$b = 9.9129$ (14) Å	$\mu = 0.23$ mm $^{-1}$
$c = 11.2779$ (16) Å	$T = 200$ K
$\alpha = 97.980$ (5) $^\circ$	$0.30 \times 0.07 \times 0.07$ mm
$\beta = 110.359$ (4) $^\circ$	

Data collection

Bruker SMART X2S benchtop diffractometer	9477 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2009)	3419 independent reflections
$T_{\min} = 0.91$, $T_{\max} = 0.98$	2732 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.036$

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.62$ e Å $^{-3}$
 $\Delta\rho_{\min} = -0.21$ e Å $^{-3}$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$
$wR(F^2) = 0.129$
$S = 1.05$
3419 reflections
272 parameters
10 restraints

Table 1
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
OW—HO2 \cdots Cl1	0.85 (1)	2.35 (1)	3.189 (2)	174 (5)
OW—HO1 \cdots Cl1 ⁱ	0.84 (1)	2.34 (1)	3.183 (2)	176 (4)
C9—H9 \cdots O3	0.95	2.49	3.278 (7)	141
C12—H12A \cdots Cl1 ⁱⁱ	0.99	2.77	3.694 (2)	155
C12—H12B \cdots O1	0.99	2.39	3.098 (6)	127
C17—H17B \cdots Cl1	0.99	2.61	3.594 (2)	176
C21—H21 \cdots OW ⁱⁱⁱ	0.95	2.29	3.168 (4)	153
C15—H15 \cdots O3 ^{iv}	0.95	2.62	3.436 (3)	143
C15—H15 \cdots C18 ^{iv}	0.95	2.85	3.788 (4)	170
C15—H15 \cdots C21 ^{iv}	0.95	2.87	3.498 (4)	124

Symmetry codes: (i) $-x + 1, -y + 2, -z + 1$; (ii) $-x, -y + 1, -z + 1$; (iii) $x, y, z + 1$; (iv) $-x, -y + 1, -z + 2$.

Data collection: *APEX2* (Bruker, 2010); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XSHELL* (Bruker, 2004) and *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *pubLCIF* (Westrip, 2010).

This work was supported by a Congressionally directed grant from the US Department of Education (grant No. P116Z100020) for the X-ray diffractometer and a grant from the Geneseo Foundation.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GG2097).

References

- Akkurt, M., Karaca, S., Küçükbay, H., Şireci, N. & Büyükgüngör, O. (2008). *Acta Cryst. E64*, o809.
- Akkurt, M., Özтурk, S., Küçükbay, H., Orhan, E. & Büyükgüngör, O. (2004). *Acta Cryst. E60*, o219–o221.

- Akkurt, M., Sireci, N., Deniz, S., Küçükbay, H. & Büyükgüngör, O. (2009). *Acta Cryst. E* **65**, o2037–o2038.
- Bruker (2004). *XSCHELL*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2009). *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2010). *APEX2*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Costache, M., Heidecker, M., Manias, E., Gupta, R. K. & Wilkie, C. (2007). *Polym. Degrad. Stabil.* **92**, 1753–1762.
- Elmali, A., Elerman, Y., Eren, G., Gümüş, R. & Svoboda, I. (2005). *Z. Naturforsch. Teil B* **60**, 164–168.
- Ennajih, H., Bouhfid, R., Zouihri, H., Essassi, E. M. & Ng, S. W. (2009). *Acta Cryst. E* **65**, o2321.
- Ennajih, H., Bouhfid, R., Zouihri, H., Essassi, E. M. & Ng, S. W. (2010). *Acta Cryst. E* **66**, o455.
- Hayakawa, Y., Kataoka, M. & Noyori, R. (1996). *J. Org. Chem.* **61**, 7996–7997.
- Hoesl, C., Nefzi, A. & Houghten, R. (2004). *J. Combin. Chem.* **6**, 220–223.
- Horton, D. A., Bourne, G. T. & Smythe, M. L. (2003). *Chem. Rev.* **103**, 893–930.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Nahle, A., Abu-Abdoun, I. & Abdel-Rahman, I. (2012). *Int. J. Corros.* Article ID 246013.
- Rivas, F., Giessert, A. & Diver, S. (2002). *J. Org. Chem.* **67**, 1708–1711.
- Rivas, F., Riaz, U., Giessert, A., Smulik, J. & Diver, S. (2001). *Org. Lett.* **3**, 2673–2676.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Smith, D. L. & Luss, H. R. (1975). *Acta Cryst. B* **31**, 402–408.
- Wang, S. & Chang, Y.-T. J. (2006). *J. Am. Chem. Soc.* **128**, 10380–10381.
- Welton, T. (1999). *Chem. Rev.* **99**, 2071–2083.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2012). E68, o3006–o3007 [doi:10.1107/S1600536812040135]

2-(Furan-2-yl)-1,3-bis(furan-2-ylmethyl)-1*H*-benzimidazol-3-i^{um} chloride monohydrate

David K. Geiger, H. Cristina Geiger and Jared M. Deck

Comment

Benzimidazolium salts have uses as room-temperature ionic liquids (Welton, 1999), surfactants (Costache *et al.*, 2007), inhibitors of corrosion (Nahlé *et al.*, 2012), anti-tumor (Elmali *et al.*, 2005) and other pharmacological agents (Horton *et al.*, 2003) and in nucleotide synthesis (Hayakawa *et al.*, 1996).

A perspective view of the benzimidazolium cation of the title compound is shown in Figure 1. The benzimidazole moiety is planar with the largest deviation being 0.011 (1) Å (N1). The furan rings are also planar with the largest deviations being 0.004 (3) Å (C11, major contributor to the disorder structure), 0.034 (13) Å (O1', minor contributor to the disordered structure), 0.011 (1) Å (C16), and 0.003 (2) Å (C20). The major component of the disordered 2-furan substituent is canted 45.1 (2)° from the mean plane of the benzimidazole ring system and the two components of the disordered furan form an angle of 22.0 (2)°. The two 2-furanyl methyl rings are on the same side of the benzimidazole with an angle of 82.79 (10)° between them.

Figure 2 shows the unit cell of the title compound. The hydrogen-bonding network involving the chloride anion and water are displayed. In addition to the O—H···Cl hydrogen bonding network, weak C—H···Cl interactions are observed. The H12A···Cl1 distance is 2.77 Å with C12···Cl1 = 3.627 (2) Å. Also, there is a weak C—H···aromatic interaction involving H15 and the furan ring containing O3. The closest approach is with O3 with H15···O3 = 2.63 Å. H15 is 2.602 (3) Å from the furan mean plane. These interactions result in chains of molecules parallel to the (0 0 1) plane as shown in Figure 3.

Experimental

A single-crystal of the title compound was obtained during the attempted crystallization of 2-(2-furanyl)-1-(furan-ylmethyl)-1*H*-benzimidazole prepared *via* the aluminium trichloride catalyzed reaction of 1,2-diaminobenzene and furan-2-carbaldehyde in refluxing dichloromethane. The product was isolated by column chromatography on silica gel using an eluant that varied from 1:2 ethylacetate:hexanes to pure ethyl acetate.

The crystal used for the diffraction study was obtained by vapor diffusion of heptane into a methanol solution of the chromatographed product. Based on ¹H NMR spectroscopy, the title compound makes up less than 5% of the bulk product.

Refinement

All hydrogen atoms were observed in difference fourier maps. The H atoms were refined using a riding model with a C—H distance of 0.99 Å for the methylene carbon atoms and 0.95 Å for the phenyl and furan carbon atoms. All C—H hydrogen atom thermal parameters were set using the approximation $U_{\text{iso}} = 1.2U_{\text{eq}}$.


The water oxygen atom was refined anisotropically. During the course of the refinement, *SHELXL* suggested that the oxygen atom of the water could be split into two atoms. However, attempts to refine the water assuming disorder result in no improvement in the GOF or the *R* values. The O—H distances were constrained to ~0.84 Å using *DFIX* and the H—O—H angles were restrained to ~104° using a DANG value of 1.34 Å between corresponding H atoms. The O—H hydrogen atom thermal parameters were set using the approximation $U_{\text{iso}} = 1.2U_{\text{eq}}$.

During the later stages of refinement, elongated thermal ellipsoids were noted for one of the furan rings. A disorder model was developed in which the minor component of the furan ring was modeled using the metrics of the major components as a guide. The pivot atom (C8) was assumed to have full occupancy and so was not included in the disorder model. The minor four-atom components (O1', C9', C10' and C11') were constrained to planarity using FLAT. Corresponding bond distances of the minor component and major component were set equal using SADI and corresponding thermal parameters were held the same using EADP. All atoms were refined anisotropically with hydrogen atoms in calculated positions using a riding model. With these constraints, the site occupancy of the major component refined to 0.84 (2). Based on this model, the angle between the mean planes of the major and minor components of the disordered furan ring is 22.2(2)°.


Improvement in the usual indicators occurred with the introduction of the disorder model. The *S* improved from 1.06 to 1.04. The *R* decreased from 0.046 to 0.045 and *wR* decreased from 0.132 to 0.129. However, an unusually large residual peak (0.61 e/Å³) that is 1.00 Å from a hydrogen atom exists. The hydrogen atom in question (H12A) is involved in a close interaction with the chloride ion (2.78 Å). The next largest residual peak is 0.37 e/Å³ and it is located 1.29 Å from H15, which is involved in a weak C—H···aromatic ring hydrogen bonding interaction (the closest approach is to O3 at 2.63 Å).

Computing details


Data collection: *APEX2* (Bruker, 2010); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT* (Bruker, 2009); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XSHELL* (Bruker, 2004) and *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *pubCIF* (Westrip, 2010).

**Figure 1**

Perspective view of the benzimidazolium ion showing the atom labeling scheme. Displacement ellipsoids of non-hydrogen atoms are drawn at the 25% probability level. Only the major contributor to the disordered furan ring is shown.

**Figure 2**


View of the unit cell showing the hydrogen bonding network. Only the major component of the disordered furan is shown. Ellipsoids are drawn at the 25% probability level.

**Figure 3**

Close intermolecular $C-H\cdots Cl$ and $C-H\cdots O(\text{furan})$ interactions resulting in a chain structure.

2-(Furan-2-yl)-1,3-bis(furan-2-ylmethyl)-1*H*-benzimidazol-3-ium chloride monohydrate

Crystal data



$M_r = 398.83$

Triclinic, $P\bar{1}$

$a = 9.4723 (12)$ Å

$b = 9.9129 (14)$ Å

$c = 11.2779 (16)$ Å

$\alpha = 97.980 (5)$ °

$\beta = 110.359 (4)$ °

$\gamma = 93.547 (4)$ °

$V = 976.3 (2)$ Å³

$Z = 2$

$F(000) = 416$

$D_x = 1.357 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3551 reflections
 $\theta = 2.4\text{--}24.8^\circ$
 $\mu = 0.23 \text{ mm}^{-1}$

$T = 200 \text{ K}$
 Needle, clear colourless
 $0.30 \times 0.07 \times 0.07 \text{ mm}$

Data collection

Bruker SMART X2S benchtop diffractometer
 Radiation source: XOS X-beam microfocus source, Bruker SMART X2S benchtop Doubly curved silicon crystal monochromator
 Detector resolution: 8.3330 pixels mm^{-1}
 ω scans
 Absorption correction: multi-scan (*SADABS*; Bruker, 2009)

$T_{\min} = 0.91, T_{\max} = 0.98$
 9477 measured reflections
 3419 independent reflections
 2732 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$
 $\theta_{\max} = 25.2^\circ, \theta_{\min} = 2.0^\circ$
 $h = -11 \rightarrow 11$
 $k = -11 \rightarrow 11$
 $l = -13 \rightarrow 13$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.129$
 $S = 1.05$
 3419 reflections
 272 parameters
 10 restraints
 Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0633P)^2 + 0.4042P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.62 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.21 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^* / U_{\text{eq}}$	Occ. (<1)
C11	0.44526 (6)	0.74517 (6)	0.50520 (6)	0.0460 (2)	
OW	0.6147 (4)	0.9554 (2)	0.3947 (2)	0.0967 (9)	
HO1	0.597 (5)	1.033 (2)	0.423 (4)	0.145*	
HO2	0.575 (5)	0.902 (3)	0.429 (4)	0.145*	
C1	0.1566 (2)	0.5684 (2)	0.68521 (18)	0.0318 (5)	
C2	0.0112 (2)	0.5062 (2)	0.66052 (18)	0.0311 (5)	
C3	-0.0290 (3)	0.3649 (2)	0.62089 (19)	0.0365 (5)	
H3	-0.1283	0.3225	0.6042	0.044*	
C4	0.0851 (3)	0.2905 (2)	0.6075 (2)	0.0435 (6)	
H4	0.0638	0.1938	0.5817	0.052*	
C5	0.2319 (3)	0.3538 (3)	0.6310 (2)	0.0449 (6)	
H5	0.3062	0.2986	0.6197	0.054*	

C6	0.2706 (3)	0.4929 (2)	0.6698 (2)	0.0398 (5)	
H6	0.3695	0.5354	0.6853	0.048*	
N1	-0.07374 (18)	0.61074 (18)	0.68249 (15)	0.0303 (4)	
C7	0.0161 (2)	0.7312 (2)	0.72021 (18)	0.0312 (5)	
N2	0.15618 (19)	0.70877 (18)	0.72206 (15)	0.0316 (4)	
C8	-0.0297 (2)	0.8639 (2)	0.7526 (2)	0.0371 (5)	
C9	0.0340 (8)	0.9713 (6)	0.8502 (5)	0.0481 (14)	0.844 (19)
H9	0.1283	0.9784	0.9192	0.058*	0.844 (19)
C10	-0.0713 (7)	1.0725 (7)	0.8273 (7)	0.0587 (14)	0.844 (19)
H10	-0.0589	1.16	0.8787	0.07*	0.844 (19)
C11	-0.1853 (5)	1.0220 (6)	0.7243 (8)	0.0689 (16)	0.844 (19)
H11	-0.2715	1.0677	0.6884	0.083*	0.844 (19)
O1	-0.1656 (4)	0.8939 (6)	0.6734 (6)	0.0664 (14)	0.844 (19)
C9'	0.059 (5)	0.970 (3)	0.843 (3)	0.0481 (14)	0.156 (19)
H9'	0.1661	0.9845	0.887	0.058*	0.156 (19)
C10'	-0.057 (4)	1.053 (5)	0.850 (4)	0.0587 (14)	0.156 (19)
H10'	-0.0414	1.1381	0.9055	0.07*	0.156 (19)
C11'	-0.187 (3)	0.992 (3)	0.770 (3)	0.0689 (16)	0.156 (19)
H11'	-0.2783	1.0324	0.7541	0.083*	0.156 (19)
O1'	-0.1809 (18)	0.862 (2)	0.708 (3)	0.0664 (14)	0.156 (19)
C12	-0.2298 (2)	0.5859 (2)	0.6823 (2)	0.0389 (5)	
H12A	-0.29	0.5136	0.6089	0.047*	
H12B	-0.2793	0.6709	0.6721	0.047*	
C13	-0.2264 (2)	0.5425 (2)	0.8045 (2)	0.0378 (5)	
C14	-0.2923 (3)	0.4323 (3)	0.8311 (2)	0.0524 (7)	
H14	-0.3555	0.3564	0.7712	0.063*	
C15	-0.2481 (4)	0.4523 (3)	0.9676 (3)	0.0634 (8)	
H15	-0.278	0.393	1.0161	0.076*	
C16	-0.1564 (3)	0.5706 (3)	1.0140 (2)	0.0570 (7)	
H16	-0.1082	0.6076	1.1027	0.068*	
O2	-0.14159 (19)	0.63101 (18)	0.91660 (15)	0.0492 (4)	
C17	0.2888 (2)	0.8126 (2)	0.7527 (2)	0.0387 (5)	
H17A	0.2543	0.9046	0.7483	0.046*	
H17B	0.337	0.7931	0.6881	0.046*	
C18	0.4023 (3)	0.8125 (2)	0.8827 (2)	0.0424 (6)	
C19	0.5399 (3)	0.7716 (3)	0.9248 (3)	0.0663 (8)	
H19	0.5931	0.7321	0.8738	0.08*	
C20	0.5922 (4)	0.7991 (4)	1.0628 (3)	0.0774 (10)	
H20	0.6864	0.7807	1.1206	0.093*	
C21	0.4837 (4)	0.8549 (3)	1.0937 (3)	0.0652 (8)	
H21	0.4882	0.884	1.1789	0.078*	
O3	0.3639 (2)	0.86463 (18)	0.98518 (16)	0.0523 (5)	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0399 (4)	0.0385 (3)	0.0571 (4)	0.0009 (2)	0.0175 (3)	0.0011 (3)
OW	0.197 (3)	0.0528 (13)	0.0828 (16)	0.0360 (17)	0.0950 (18)	0.0222 (12)
C1	0.0345 (12)	0.0383 (12)	0.0231 (10)	0.0040 (9)	0.0104 (9)	0.0074 (8)
C2	0.0344 (12)	0.0379 (12)	0.0215 (9)	0.0063 (9)	0.0091 (8)	0.0084 (8)

C3	0.0400 (13)	0.0381 (12)	0.0290 (11)	0.0011 (10)	0.0095 (9)	0.0075 (9)
C4	0.0601 (16)	0.0352 (12)	0.0345 (12)	0.0095 (11)	0.0155 (11)	0.0055 (10)
C5	0.0477 (15)	0.0508 (15)	0.0397 (13)	0.0198 (12)	0.0173 (11)	0.0086 (11)
C6	0.0360 (12)	0.0494 (14)	0.0363 (12)	0.0098 (11)	0.0145 (10)	0.0086 (10)
N1	0.0274 (9)	0.0353 (10)	0.0271 (9)	0.0025 (8)	0.0082 (7)	0.0067 (7)
C7	0.0326 (11)	0.0373 (12)	0.0233 (10)	0.0021 (9)	0.0091 (8)	0.0075 (8)
N2	0.0295 (10)	0.0366 (10)	0.0290 (9)	0.0019 (8)	0.0110 (7)	0.0067 (7)
C8	0.0356 (12)	0.0401 (13)	0.0367 (12)	0.0067 (10)	0.0136 (10)	0.0080 (10)
C9	0.041 (3)	0.0451 (15)	0.0502 (16)	-0.0033 (18)	0.0118 (16)	-0.0023 (12)
C10	0.075 (3)	0.033 (3)	0.071 (3)	0.0086 (18)	0.032 (2)	0.002 (2)
C11	0.072 (2)	0.055 (2)	0.078 (4)	0.037 (2)	0.020 (2)	0.012 (2)
O1	0.0532 (14)	0.059 (2)	0.067 (2)	0.0210 (14)	-0.0006 (13)	-0.0014 (16)
C9'	0.041 (3)	0.0451 (15)	0.0502 (16)	-0.0033 (18)	0.0118 (16)	-0.0023 (12)
C10'	0.075 (3)	0.033 (3)	0.071 (3)	0.0086 (18)	0.032 (2)	0.002 (2)
C11'	0.072 (2)	0.055 (2)	0.078 (4)	0.037 (2)	0.020 (2)	0.012 (2)
O1'	0.0532 (14)	0.059 (2)	0.067 (2)	0.0210 (14)	-0.0006 (13)	-0.0014 (16)
C12	0.0333 (12)	0.0452 (13)	0.0379 (12)	0.0045 (10)	0.0119 (10)	0.0084 (10)
C13	0.0338 (12)	0.0430 (13)	0.0342 (12)	0.0038 (10)	0.0101 (9)	0.0043 (10)
C14	0.0618 (17)	0.0512 (16)	0.0424 (14)	-0.0091 (13)	0.0191 (12)	0.0079 (12)
C15	0.083 (2)	0.0618 (18)	0.0477 (15)	-0.0101 (16)	0.0261 (15)	0.0175 (14)
C16	0.0727 (19)	0.0634 (18)	0.0328 (13)	0.0045 (15)	0.0153 (12)	0.0126 (12)
O2	0.0529 (11)	0.0516 (10)	0.0393 (9)	-0.0029 (8)	0.0134 (8)	0.0083 (8)
C17	0.0362 (13)	0.0388 (13)	0.0432 (12)	-0.0026 (10)	0.0176 (10)	0.0089 (10)
C18	0.0373 (13)	0.0388 (13)	0.0461 (13)	-0.0051 (10)	0.0116 (11)	0.0044 (10)
C19	0.0443 (16)	0.072 (2)	0.071 (2)	0.0068 (14)	0.0087 (14)	0.0062 (16)
C20	0.0531 (19)	0.074 (2)	0.075 (2)	-0.0040 (17)	-0.0146 (16)	0.0171 (17)
C21	0.070 (2)	0.0564 (18)	0.0466 (16)	-0.0170 (16)	-0.0038 (14)	0.0111 (13)
O3	0.0532 (11)	0.0541 (11)	0.0432 (10)	-0.0020 (9)	0.0109 (8)	0.0085 (8)

Geometric parameters (\AA , $\text{^{\circ}}$)

OW—HO1	0.844 (10)	C9'—C10'	1.432 (18)
OW—HO2	0.847 (10)	C9'—H9'	0.95
C1—C2	1.391 (3)	C10'—C11'	1.297 (19)
C1—N2	1.397 (3)	C10'—H10'	0.95
C1—C6	1.397 (3)	C11'—O1'	1.388 (17)
C2—C3	1.395 (3)	C11'—H11'	0.95
C2—N1	1.398 (3)	C12—C13	1.489 (3)
C3—C4	1.384 (3)	C12—H12A	0.99
C3—H3	0.95	C12—H12B	0.99
C4—C5	1.410 (4)	C13—C14	1.345 (3)
C4—H4	0.95	C13—O2	1.383 (3)
C5—C6	1.373 (3)	C14—C15	1.429 (4)
C5—H5	0.95	C14—H14	0.95
C6—H6	0.95	C15—C16	1.338 (4)
N1—C7	1.347 (3)	C15—H15	0.95
N1—C12	1.483 (3)	C16—O2	1.365 (3)
C7—N2	1.352 (3)	C16—H16	0.95
C7—C8	1.449 (3)	C17—C18	1.486 (3)
N2—C17	1.482 (3)	C17—H17A	0.99

C8—O1'	1.341 (15)	C17—H17B	0.99
C8—C9	1.357 (4)	C18—C19	1.333 (4)
C8—C9'	1.358 (18)	C18—O3	1.373 (3)
C8—O1	1.366 (4)	C19—C20	1.439 (5)
C9—C10	1.444 (5)	C19—H19	0.95
C9—H9	0.95	C20—C21	1.322 (5)
C10—C11	1.290 (5)	C20—H20	0.95
C10—H10	0.95	C21—O3	1.369 (3)
C11—O1	1.371 (4)	C21—H21	0.95
C11—H11	0.95		
HO1—OW—HO2	103 (2)	C8—C9'—H9'	130.8
C2—C1—N2	106.81 (18)	C10'—C9'—H9'	130.8
C2—C1—C6	121.7 (2)	C11'—C10'—C9'	109 (4)
N2—C1—C6	131.4 (2)	C11'—C10'—H10'	125.5
C1—C2—C3	122.2 (2)	C9'—C10'—H10'	125.5
C1—C2—N1	106.73 (18)	C10'—C11'—O1'	114 (3)
C3—C2—N1	131.0 (2)	C10'—C11'—H11'	122.8
C4—C3—C2	115.7 (2)	O1'—C11'—H11'	122.8
C4—C3—H3	122.1	C8—O1'—C11'	97.6 (17)
C2—C3—H3	122.1	N1—C12—C13	110.44 (17)
C3—C4—C5	122.0 (2)	N1—C12—H12A	109.6
C3—C4—H4	119.0	C13—C12—H12A	109.6
C5—C4—H4	119.0	N1—C12—H12B	109.6
C6—C5—C4	122.0 (2)	C13—C12—H12B	109.6
C6—C5—H5	119.0	H12A—C12—H12B	108.1
C4—C5—H5	119.0	C14—C13—O2	110.3 (2)
C5—C6—C1	116.3 (2)	C14—C13—C12	133.2 (2)
C5—C6—H6	121.8	O2—C13—C12	116.5 (2)
C1—C6—H6	121.8	C13—C14—C15	106.2 (2)
C7—N1—C2	108.79 (17)	C13—C14—H14	126.9
C7—N1—C12	126.83 (18)	C15—C14—H14	126.9
C2—N1—C12	123.78 (18)	C16—C15—C14	106.9 (2)
N1—C7—N2	109.02 (18)	C16—C15—H15	126.6
N1—C7—C8	125.66 (19)	C14—C15—H15	126.6
N2—C7—C8	125.32 (19)	C15—C16—O2	110.8 (2)
C7—N2—C1	108.65 (17)	C15—C16—H16	124.6
C7—N2—C17	127.40 (18)	O2—C16—H16	124.6
C1—N2—C17	123.92 (18)	C16—O2—C13	105.8 (2)
O1'—C8—C9	109.7 (10)	N2—C17—C18	111.56 (18)
O1'—C8—C9'	120 (2)	N2—C17—H17A	109.3
C9—C8—O1	109.0 (4)	C18—C17—H17A	109.3
C9'—C8—O1	116 (2)	N2—C17—H17B	109.3
O1'—C8—C7	112.0 (9)	C18—C17—H17B	109.3
C9—C8—C7	133.6 (4)	H17A—C17—H17B	108.0
C9'—C8—C7	126 (2)	C19—C18—O3	109.8 (2)
O1—C8—C7	117.5 (2)	C19—C18—C17	133.6 (3)
C8—C9—C10	105.8 (5)	O3—C18—C17	116.7 (2)
C8—C9—H9	127.1	C18—C19—C20	106.6 (3)

C10—C9—H9	127.1	C18—C19—H19	126.7
C11—C10—C9	107.3 (6)	C20—C19—H19	126.7
C11—C10—H10	126.3	C21—C20—C19	106.7 (3)
C9—C10—H10	126.3	C21—C20—H20	126.6
C10—C11—O1	111.1 (5)	C19—C20—H20	126.6
C10—C11—H11	124.5	C20—C21—O3	110.3 (3)
O1—C11—H11	124.4	C20—C21—H21	124.9
C8—O1—C11	106.8 (3)	O3—C21—H21	124.9
C8—C9'—C10'	98 (3)	C21—O3—C18	106.7 (2)
N2—C1—C2—C3	179.89 (18)	C9—C10—C11—O1	-0.7 (6)
C6—C1—C2—C3	1.0 (3)	O1'—C8—O1—C11	96 (2)
N2—C1—C2—N1	0.3 (2)	C9—C8—O1—C11	-0.3 (4)
C6—C1—C2—N1	-178.61 (18)	C9'—C8—O1—C11	-11 (2)
C1—C2—C3—C4	-0.1 (3)	C7—C8—O1—C11	178.9 (3)
N1—C2—C3—C4	179.42 (19)	C10—C11—O1—C8	0.6 (5)
C2—C3—C4—C5	-0.7 (3)	O1'—C8—C9'—C10'	-4 (2)
C3—C4—C5—C6	0.7 (3)	C9—C8—C9'—C10'	-33 (9)
C4—C5—C6—C1	0.2 (3)	O1—C8—C9'—C10'	24 (2)
C2—C1—C6—C5	-1.0 (3)	C7—C8—C9'—C10'	-166.6 (15)
N2—C1—C6—C5	-179.6 (2)	C8—C9'—C10'—C11'	-1 (2)
C1—C2—N1—C7	-0.6 (2)	C9'—C10'—C11'—O1'	5 (3)
C3—C2—N1—C7	179.8 (2)	C9—C8—O1'—C11'	12.4 (16)
C1—C2—N1—C12	-172.23 (17)	C9'—C8—O1'—C11'	6 (2)
C3—C2—N1—C12	8.2 (3)	O1—C8—O1'—C11'	-80 (2)
C2—N1—C7—N2	0.7 (2)	C7—C8—O1'—C11'	171.4 (13)
C12—N1—C7—N2	172.00 (17)	C10'—C11'—O1'—C8	-7 (3)
C2—N1—C7—C8	-179.78 (18)	C7—N1—C12—C13	-92.9 (2)
C12—N1—C7—C8	-8.5 (3)	C2—N1—C12—C13	77.2 (2)
N1—C7—N2—C1	-0.5 (2)	N1—C12—C13—C14	-124.4 (3)
C8—C7—N2—C1	179.96 (18)	N1—C12—C13—O2	57.0 (3)
N1—C7—N2—C17	177.42 (17)	O2—C13—C14—C15	0.5 (3)
C8—C7—N2—C17	-2.1 (3)	C12—C13—C14—C15	-178.2 (3)
C2—C1—N2—C7	0.1 (2)	C13—C14—C15—C16	-1.4 (3)
C6—C1—N2—C7	178.9 (2)	C14—C15—C16—O2	1.8 (4)
C2—C1—N2—C17	-177.90 (17)	C15—C16—O2—C13	-1.5 (3)
C6—C1—N2—C17	0.9 (3)	C14—C13—O2—C16	0.6 (3)
N1—C7—C8—O1'	-17.5 (16)	C12—C13—O2—C16	179.5 (2)
N2—C7—C8—O1'	161.9 (15)	C7—N2—C17—C18	105.0 (2)
N1—C7—C8—C9	134.8 (4)	C1—N2—C17—C18	-77.4 (2)
N2—C7—C8—C9	-45.8 (5)	N2—C17—C18—C19	108.4 (3)
N1—C7—C8—C9'	147 (2)	N2—C17—C18—O3	-71.7 (3)
N2—C7—C8—C9'	-34 (2)	O3—C18—C19—C20	0.2 (3)
N1—C7—C8—O1	-44.2 (5)	C17—C18—C19—C20	-180.0 (3)
N2—C7—C8—O1	135.2 (5)	C18—C19—C20—C21	-0.4 (4)
O1'—C8—C9—C10	-26.4 (14)	C19—C20—C21—O3	0.4 (4)
C9'—C8—C9—C10	127 (10)	C20—C21—O3—C18	-0.3 (3)
O1—C8—C9—C10	-0.1 (4)	C19—C18—O3—C21	0.1 (3)
C7—C8—C9—C10	-179.1 (4)	C17—C18—O3—C21	-179.8 (2)

C8—C9—C10—C11 0.5 (5)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
OW—HO2···Cl1	0.85 (1)	2.35 (1)	3.189 (2)	174 (5)
OW—HO1···Cl1 ⁱ	0.84 (1)	2.34 (1)	3.183 (2)	176 (4)
C9—H9···O3	0.95	2.49	3.278 (7)	141
C12—H12A···Cl1 ⁱⁱ	0.99	2.77	3.694 (2)	155
C12—H12B···O1	0.99	2.39	3.098 (6)	127
C17—H17B···Cl1	0.99	2.61	3.594 (2)	176
C21—H21···OW ⁱⁱⁱ	0.95	2.29	3.168 (4)	153
C15—H15···O3 ^{iv}	0.95	2.62	3.436 (3)	143
C15—H15···C18 ^{iv}	0.95	2.85	3.788 (4)	170
C15—H15···C21 ^{iv}	0.95	2.87	3.498 (4)	124

Symmetry codes: (i) $-x+1, -y+2, -z+1$; (ii) $-x, -y+1, -z+1$; (iii) $x, y, z+1$; (iv) $-x, -y+1, -z+2$.