



HHS Public Access

Author manuscript

Nat Neurosci. Author manuscript; available in PMC 2009 September 01.

Published in final edited form as:

Nat Neurosci. 2009 March ; 12(3): 295–302. doi:10.1038/nn.2258.

Intra-Retinal Visual Cycle Required for Rapid and Complete Cone Dark Adaptation

Jin-Shan Wang¹, Maureen E. Estevez², M. Carter Cornwall², and Vladimir J. Kefalov¹

¹Department of Ophthalmology and Visual Sciences, Washington University School of Medicine, 660 South Euclid, Saint Louis, MO 63110

²Department of Physiology and Biophysics, Boston University School of Medicine, 715 Albany Street, Boston, MA 02118

Summary

Daytime vision is mediated by retinal cones which, unlike rods, remain functional even in bright light and dark-adapt rapidly. These cone properties are enabled by rapid regeneration of their pigment. This in turn requires rapid chromophore recycling which may not be achieved by the canonical retinal pigment epithelium visual cycle. Recent biochemical studies have suggested the presence of a second, cone-specific visual cycle, although its physiological function remains to be established. Here we report that the Müller cells within the salamander neural retina promote cone-specific pigment regeneration and dark adaptation that are independent of the pigment epithelium. Without this pathway, dark adaptation of cones is slow and incomplete. Interestingly, the rates of cone pigment regeneration by the retina and pigment epithelium visual cycles are essentially identical suggesting a possible common rate-limiting step. Finally, we also observed cone dark adaptation in the isolated mouse retina.

Introduction

Phototransduction in rods and cones begins with the light-triggered isomerization of the visual pigment chromophore from 11-cis to all-trans retinal¹. Eventually, the photoactivated pigment dissociates into free opsin and all-trans retinal², which is then reduced to all-trans retinol³. Dark-adaptation of both rods and cones requires regeneration of the visual pigment from opsin and 11-cis retinal^{4, 5}. However, the speed of pigment regeneration and hence sensitivity recovery is very different in rods and cones with full recovery requiring only about 5 minutes in cones and up to an hour in rods^{6, 7}. The fast turnover of cone visual pigment required for cones to rapidly dark-adapt and to remain functional in bright light imposes the need for rapid recycling of their chromophore from all-trans retinol back into 11-cis retinal. The canonical pathway for chromophore recycling⁸ involves the pigment epithelium, where all-trans retinol is converted into 11-cis retinal via a series of enzymatic

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:http://www.nature.com/authors/editorial_policies/license.html#terms

Address for Editorial Correspondence: Dr. Vladimir Kefalov, Ophthalmology & Visual Sciences, Washington University School of Medicine, 660 S. Euclid Ave, Box 8096, Saint Louis, MO 63110, USA, Tel: 314-362-4376, FAX: 314-362-3720, E-Mail:kefalov@wustl.edu.

reactions and then transported back to the photoreceptors for incorporation into opsin. There are reasons to think that, apart from the epithelial pathway common for rods and cones, a separate, cone-specific, chromophore-recycling pathway may exist. First, while rods are non-functional in bright light, their pigment continues to cycle through repetitive bleaching and regeneration, acting as a sink for 11-cis retinal. Thus, even saturated, rods continue to use 11-cis retinal thereby reducing its availability to cones. As in the mouse and human retina, cones constitute only 3–5% of all photoreceptors, cone opsin has to compete with overwhelming levels of rod opsin for recycled 11-cis retinal. Second, unlike the extremely stable rod pigment, cone pigment can dissociate spontaneously into opsin and 11-cis retinal^{9, 10}. Thus, even after regeneration cone pigment might lose its chromophore to rod pigment². Finally, the rate of pigment regeneration required for sustaining cone function in bright light exceeds the maximal reported rate of chromophore recycling by the pigment epithelium¹¹.

Recent biochemical studies have established a series of enzymatic reactions within the retina consistent with a chromophore-recycling pathway^{11–15}. This pathway is possibly cone-specific because it is characterized in cone-dominant retinas such as those of chicken and ground squirrel. The emerging notion¹¹ is that the all-trans retinol released from cones is converted into 11-cis retinol within the retina independently of the pigment epithelium. The 11-cis retinol is then used by the cones, which, unlike rods, can convert 11-cis retinol into 11-cis retinal (at least in salamander)⁴. This pathway is capable of turning over chromophore 20-fold faster than the canonical pigment epithelium pathway¹¹. While these biochemical studies are widely accepted, the functional validation of this separate pathway *in situ* has not been carried out and it is not known whether this pathway can promote cone pigment regeneration or dark adaptation. Equally importantly, the question remains whether such a visual cycle exists in rod-dominant retinas. In fact, two recent studies, albeit indirect, failed to find any evidence for chromophore recycling within the rod-dominant mouse retina^{16, 17}. Finally, it is not known what, if any, role this putative retina visual cycle plays in the dark adaptation of cones. We have now successfully addressed these questions by combining microspectrophotometric measurements with single-cell and whole-retina recordings from amphibian and mouse photoreceptors *in situ* in the retina.

Results

The Retina Promotes Cone Pigment Regeneration

We used the rod-dominant salamander retina, which has 35% cones¹⁸, to investigate whether a cone-specific visual cycle is present in the vertebrate retina and is functional under physiological conditions. Following bleaching by bright light, such a pathway would be expected to promote regeneration exclusively of cone pigment, independently of the pigment epithelium. Using single-cell microspectrophotometry, we compared the pigment content in dark-adapted cones and cones that had been bleached and then allowed to recover for two hours in darkness without pigment epithelium. We measured the pigment recovery in cones that had been dissociated from the retina prior to the bleach and in cones bleached while still within the intact retina. All measurements were performed at the end of the recovery period from individual isolated cells. We found that exposure of dissociated red

cones to 40 s white bleaching light induced a 4.7-fold decrease in their optical density (Fig. 1a, compare left and middle panels), corresponding to a loss of 79% of the cone pigment (Table 1). The bleached cone pigment could be partially regenerated with exogenous 11-cis retinal chromophore (Fig. 1a, right panel). In contrast, identical bleach of cones from whole isolated retina produced only 4.3% decrease in their optical density (Fig. 1b, compare left and middle panels). Thus, the retina was able to regenerate 95% (74.7%/79%) of all bleached cone pigment, whereas pigment regeneration in isolated cones was insignificant. Treating such cones with exogenous 11-cis retinal fully restored their pigment content (Fig. 1b, right panel). Thus, the incomplete recovery of cone pigment content in isolated retina was due to insufficient recycled chromophore. Possibly, some chromophore released from the bleached photoreceptors was lost to the bath; alternatively, retina integrity was compromised during the dissection reducing the efficiency of its visual cycle. The ability of the retina to promote pigment regeneration was restricted to cones as we found no pigment regeneration in rods from intact retina (Fig. 1c).

The Retina Promotes Cone Dark Adaptation

Using single-cell suction recordings, we investigated whether the salamander retina could promote cone dark adaptation independently of the pigment epithelium. To allow comparison with our microspectrophotometric results, these experiments were performed under identical conditions. As expected, in the absence of pigment regeneration in dissociated cones, the bleach induced a substantial persistent desensitization (Fig. 2a). Photosensitivity decreased from $10.4 \pm 1.5 \times 10^{-3}$ pA photon⁻¹ μm² (mean ± s.e.m., $n = 19$) in dark-adapted cones to $2.0 \pm 0.4 \times 10^{-4}$ pA photon⁻¹ μm² ($n = 20$) in cones dissociated from the retina before the bleach. Using the relation between percentage of bleached cone pigment and decrease in sensitivity of salamander red cones⁹, this decrease in sensitivity corresponds to bleaching 83% of the cone visual pigment (Table 1), a value similar to the 79% pigment loss from microspectrophotometric measurements. Consistent with the adaptation induced by the bleaching light, the average integration time of the cone dim flash response was also accelerated from 445 ± 42 ms ($n=19$) for dark-adapted cones to 183 ± 15 ms ($n=20$) for bleached cones. Exogenous 11-cis retinal restored the sensitivity of bleached dissociated cones to $8.2 \pm 2.0 \times 10^{-3}$ pA photon⁻¹ μm² ($n = 7$) (Fig. 2a). Similar results were obtained with 11-cis retinol (not shown), confirming that salamander cones can oxidize it to 11-cis retinal for pigment regeneration⁴. In contrast, cones from whole isolated retina recovered their sensitivity following identical bleach in the absence of pigment epithelium and without exogenous chromophore (Fig. 2b). On average, the flash sensitivity changed from $7.3 \pm 0.8 \times 10^{-3}$ pA photon⁻¹ μm² ($n = 14$) for dark-adapted cones to $4.8 \pm 0.4 \times 10^{-3}$ pA photon⁻¹ μm² ($n = 14$) for cones bleached in the intact retina. This represents a 1.5-fold decrease in sensitivity, corresponding to a loss of only 7.5% of the pigment. Consistent with the substantial recovery of cone sensitivity in isolated retina following a bleach, the flash response acceleration was also largely reversed and the integration time of the dim flash response changed from 496 ± 48 ms ($n=14$) for dark-adapted cones to 451 ± 39 ms ($n=14$) for bleached cones. Exogenous 11-cis retinal produced only a small further increase in sensitivity to $8.5 \pm 1.1 \times 10^{-3}$ pA photon⁻¹ μm² ($n = 7$), just higher than the initial dark-adapted level (Fig. 2b, bottom panel) consistent with the presence of free opsin in dark-adapted salamander cones⁹. Thus, the intact amphibian retina, removed from the pigment

epithelium, was able to reverse the effects of bleaching in cones and promote their dark adaptation. The ability of the retina to promote dark adaptation was restricted to cones as we found no dark adaptation in rods from intact retina (Fig. 2c).

Müller Cells are Part of the Retina Visual Cycle

We next investigated the role of Müller cells in the regeneration of cone pigment using the Müller cell-specific gliotoxin α -aminoadipic acid (L- α -AAA)¹⁹. L- α -AAA inhibits cysteine uptake through the cysteine/glutamate antiporter, primarily localized in glial cells, leading to depletion of the antioxidant glutathione and eventually causing cellular damage²⁰. As in the adult retina L- α -AAA does not affect retinal neurons, including photoreceptors^{21, 22}, it is widely used for selectively abolishing Müller cells. As expected from the lack of uptake of L- α -AAA by neurons²³, 48-hour incubation of isolated salamander retina in 10 mM L- α -AAA prior to recordings did not affect significantly its morphology (Supplementary Fig. 1 online) or the function of cones (but see Discussion). However, a subsequent bleach induced a significant decrease in sensitivity, from $5.7 \pm 0.9 \times 10^{-3}$ pA photon⁻¹ μm^2 ($n = 12$) in dark-adapted cones to $3.3 \pm 0.6 \times 10^{-4}$ pA photon⁻¹ μm^2 ($n = 15$) following the bleach (Fig. 3a). This 17-fold decrease in sensitivity corresponds to loss of 68% of the pigment (Table 1), 9 times more than following identical bleach in control solution. In parallel microspectrophotometry measurements, the bleach induced a loss of 69% of the pigment (Table 1). L- α -AAA did not significantly affect cone pigment content or sensitivity in darkness. Furthermore, both cone pigment loss (not shown) and desensitization (Fig. 3a) in bleached L- α -AAA-treated retinas were readily reversed by exogenous 11-cis retinal confirming that L- α -AAA did not affect the function of dark-adapted cones while largely inhibiting cone pigment regeneration and dark adaptation in isolated retina. In contrast, in eyecup with the retina attached to the pigment epithelium, L- α -AAA did not block recovery of sensitivity of bleached cones (Fig. 3b) and had no effect on the dark adaptation of rods (Fig. 5b). Thus, the Müller cell gliotoxin inhibited specifically the retina visual cycle and not the canonical pigment epithelium visual cycle.

Role of the Retina Visual Cycle in Cone Dark Adaptation

To observe the recovery of cone sensitivity in real time and determine the kinetics of chromophore recycling by the cone-specific retina visual cycle, we recorded rod and cone photoresponses from whole salamander retina. We isolated the photoreceptor component (a-wave) of isolated retina electroretinogram (ERG) responses by pharmacologically blocking synaptic transmission (see Methods). The photoreceptor response from salamander retina contains both rod (slow) and cone (fast) components (Fig. 4a, left inset). Using backgrounds that were bright enough to saturate the rods while minimally affecting the cone component of the response (Fig. 4a, right inset), we were able to record cone test-flash responses from intact retina (Fig. 4c, e). By subtracting this cone response from the rod-and-cone response elicited by identical flash in darkness, we were able, in turn, to obtain the rod component of the retina responses (Fig. 4b, d). While the bleach reduced the rod sensitivity by over three orders of magnitude, its effect on the sensitivity of cones from the same retina was a reduction of only 1.7-fold, corresponding to a loss of only 8.0% cone pigment (Table 1), a value similar to the 7.5% pigment loss from single-cell recordings.

Using ERG recordings from isolated salamander retina we were able to observe in real time the recovery of cone sensitivity following a bleach (Fig. 5a, black circles). Consistent with our single-cell results, this recovery was inhibited by 48-hour incubation in L- α -AAA (Fig. 5a, red circles) while similar incubation in control solution had no effect on recovery (not shown). In contrast, L- α -AAA did not affect recycling of chromophore by the pigment epithelium as demonstrated by the recovery of rod (Fig. 5b) and cone (Fig. 5c, red) sensitivity in retina from eyecup, still attached to the pigment epithelium. Importantly, the recovery of cone sensitivity was fastest in eyecup, where both retina and pigment epithelium recycled chromophore for cones (Fig. 5c, black), so that 5 minutes following a bleach cones regained 32% of their dark-adapted sensitivity (pigment epithelium and retina) as opposed to only 8% (pigment epithelium alone). Thus, the addition of chromophore recycled within the retina significantly accelerated the dark adaptation of cones.

The pigment epithelium alone was not sufficient for complete cone pigment regeneration: in the L- α -AAA-treated eyecup, cone sensitivity following bleach recovered to 92% (Fig 5c, red), corresponding to regeneration of 98% of cone pigment (Fig 5d, red). The cone pigment regeneration driven by the isolated retina was somewhat lower at 93% (64% sensitivity recovery, Fig. 5a), most likely due to dissection damage to the retina and loss of chromophore to the bath. However, combining the pigment epithelium and the retina visual cycles in the bleached intact eyecup fully restored cone sensitivity indicating 100% pigment regeneration (Fig. 5c and 5d, black, respectively). Thus, both visual cycles were required for full recovery of cone sensitivity following a bleach. Interestingly, the initial rates of pigment regeneration by the two visual cycles were similar (Fig. 5d, see Discussion).

A Retina Visual Cycle is Functional in the Mouse Retina

Finally, to determine whether a visual cycle functions in the mammalian retina, we recorded ERG responses from isolated mouse retina. As in the case of salamander, we used the difference in light adaptation between rods and cones to isolate their responses within the same retina. For simplicity, we used 500 nm light both for bleaching and for test-flash stimulation, effectively excluding mouse cone S-opsin and limiting our studies to mouse cone M-opsin. This allowed us to investigate the possible function of a visual cycle in the mouse retina without considering the coexpression of two cone pigments in mouse cones.

As expected, the bleach induced significant reduction in the rod response amplitude (Fig. 6, compare a and c) and 140-fold (n=16) decrease in rod sensitivity. In contrast, following an identical bleach, the amplitude of the cone response recovered fully and the cone sensitivity recovered to 1/3 (n=8) of its prebleach dark-adapted level (Fig. 6, compare b and d) indicating substantial pigment regeneration in cones from isolated retina. Thus, similarly to the case in salamander, the mouse retina was able to promote dark adaptation independently of the pigment epithelium in cones, but not in rods.

Discussion

Functional Cone-Specific Visual Cycle in Vertebrate Retina

We investigated whether the vertebrate neural retina is able to recycle chromophore and promote cone pigment regeneration independently of the pigment epithelium. Using microspectrophotometry, we observed cone-specific pigment regeneration in the isolated salamander retina following exposure to bright bleaching light. In parallel physiological experiments using single-cell and whole-retina recordings, we observed cone-specific dark-adaptation in the same conditions. Recordings from mouse photoreceptors revealed that the isolated mouse retina is also able to promote cone-specific dark adaptation. Together, these results demonstrate that a functional visual cycle is present within the vertebrate retina where it provides recycled chromophore, presumably 11-cis retinol, exclusively to cones and independently of the pigment epithelium (Supplementary Fig. 3 online). We also found that both the canonical pigment epithelium visual cycle and the retina visual cycle are capable of independently promoting pigment regeneration in cones. Thus, while rods rely on chromophore recycled solely in the pigment epithelium, cones utilize chromophore supplied by both the pigment epithelium and the retina. Our results help explain the long-standing but controversial observation of recovery of cone early receptor potential following a bleach in isolated frog retina²⁴. The implications of the addition of a retina visual cycle for cone function are discussed below.

How is the cone-specificity of the retina visual cycle achieved? Previous physiological studies from salamander photoreceptors provide two possible mechanisms. First, exogenous 11-cis retinol can produce dark-adaptation only in cones, but not rods⁴. As 11-cis retinol is the presumptive form of recycled chromophore produced by the retina, this will provide one mechanism for the cone specificity of the retina visual cycle. Second, chromophore can diffuse from the inner to the outer segment only in cones but not in rods²⁵. As Müller cell processes surround photoreceptor cell bodies but do not reach their outer segments (Ref. 26; see also Supplementary Fig. 1 online), the recycled chromophore released from Müller cells will diffuse to the outer segment and combine with opsin only in cones but not in rods.

As a first step in dissecting the physiology of the retina visual cycle, we examined whether Müller cells play a role in chromophore recycling. First, we found that the retina visual cycle functions only in intact retina, where contacts between cones and Müller cells are preserved. Even simply drawing the outer segment of a single cone, still attached to a small piece of retina, into the recording electrode was sufficient to prevent retina-driven pigment regeneration (Supplementary Fig. 2 online). Thus, preserving retina morphology and proper contact between cones and adjacent inner neurons and Müller cells is paramount for the function of the retina visual cycle. Second, we found that preincubation of the retina in the Müller cells-specific gliotoxin L- α -AAA selectively blocks the retina-driven cone pigment regeneration and dark adaptation (Fig. 4a and Fig. 5a). In the adult frog and chicken retina, treatment with L- α -AAA at concentrations between 60 and 300 mM (higher than the one used in this study) has no effect on retinal elements other than Müller cells²². Indeed, we found that treatment of salamander retina with L- α -AAA has no effect on photoreceptor morphology (Supplementary Fig. 1 online) or on the ability of bleached cones to regenerate

their pigment and dark-adapt using exogenous 11-cis retinal. Together, these results indicate that Müller cells are essential part of the retina visual cycle.

The inhibition of chromophore recycling by L- α -AAA resulted in ca. 50% reduction in the sensitivity of dark-adapted cones. This desensitization likely reflects the gradual accumulation of free opsin produced due to the high rate of spontaneous thermal activation of pigment in salamander red cones⁹. Indeed, subsequent treatment of such cells with exogenous 11-cis retinal resulted in significant overshoot of sensitivity (Fig. 3, bottom panels) indicating a larger than normal levels of free opsin. As L- α -AAA did not completely block the retina visual cycle, some chromophore released from the gradual “thermal bleach” of cone pigment would have been recycled to regenerate cone pigment. These results indicate that, similarly to activation by a photon, thermal activation of the cone pigment likely results in its decay to free opsin.

Functional Significance of the Retina Visual Cycle

What are the advantages of having two separate sources of chromophore for cones but only one for rods? Rapid cone pigment regeneration is crucial for the rapid dark adaptation of cones and for their persistent function in bright light, where cone pigment is continuously bleached at a high rate. The rapid turnover of cone pigment facilitated by its rapid formation and decay² also imposes the need for rapid recycling of chromophore.

Our results demonstrate that, while both the pigment epithelium and retina visual cycles are able to promote substantial cone pigment regeneration, the combined action of the two visual cycles is required for the complete dark adaptation of cones. Furthermore, the addition of the retina visual cycle to the canonical pigment epithelium visual cycle accelerates 4-fold cone dark adaptation (Fig. 5c), indicating that that the retina visual cycle is required for the rapid and complete dark adaptation of cones. Rod pigment regeneration and dark adaptation, on the other hand, are not affected by the retina visual cycle and the action of the pigment epithelium visual cycle alone is sufficient for complete rod pigment regeneration and dark adaptation (Fig. 5b).

Comparable Rates of the Retina and Pigment Epithelium Visual Cycles

Previous biochemical experiments have suggested that the retina might be able to recycle chromophore up to 20 times faster than the pigment epithelium¹¹. Our whole-retina ERG recordings allowed for a direct comparison of the rates of the two visual cycles under physiological conditions. Since chromophore loss and retina damage reduce the efficiency of the retina visual cycle as measured from isolated retina, we used instead measurements from intact eyecup to estimate the rate of the retina visual cycle. By subtracting the fractional pigment regenerated by the pigment epithelium alone (eyecup + L- α -AAA) from the pigment regenerated by both pigment epithelium and retina (eyecup), we obtained the fractional pigment regenerated by the retina alone (Fig. 5d, open circles). This estimate is based on the lack of effect by L- α -AAA on the pigment epithelium cycle (Fig. 5b). Surprisingly, we find that the initial rates of cone pigment regeneration by the pigment epithelium and by the retina are essentially equal. Thus, the additional action of the retina visual cycle doubles the amount of recycled chromophore available to cones during dark

adaptation. Because of the non-linear relation between bleached pigment and desensitization⁹, this results in a 4-fold increase in cone sensitivity and significantly accelerates cone dark adaptation.

How can we explain the discrepancy between the biochemical estimate of the maximum rate of enzymatic recycling of chromophore in ground squirrel retina¹¹ and our physiological measurement of pigment regeneration kinetics in amphibian retina? A simple explanation could be the differences between species. More likely, however, cone pigment regeneration via the two cycles could be rate-limited not by their enzymatic reactions but rather by a common step such as the delivery of recycled chromophore to cones, as recently proposed for rods²⁷.

Ruling out Potential Contamination from the Pigment Epithelium

We took extreme care to remove all visible pigment epithelium from the retina and performed multiple control experiments demonstrating that residual pigment epithelium was not the source of chromophore in isolated retina. First, only cones in close proximity to a piece of pigment epithelium would be expected to undergo pigment regeneration and dark adaptation. Instead, we found uniform pigment regeneration (Fig. 1b) and dark adaptation (Fig. 2b) in individual cones and almost complete recovery of whole-retina cone sensitivity (Fig. 5a) indicating that all bleached cones underwent dark adaptation. Second, pigment epithelium contamination would be expected to promote both rod and cone pigment regeneration. However, we found no rod recovery of pigment content (Fig. 1c) or single-cell (Fig. 2c) and whole-retina (Fig. 4d) sensitivity in bleached isolated retina indicating that rods did not receive chromophore. Third, the Müller cell inhibitor L- α -AAA blocked cone pigment regeneration and single-cell (Fig. 3a) and whole-retina (Fig. 5a) cone dark adaptation. This treatment specifically affected chromophore recycling within the retina and had no effect on the pigment epithelium as shown by the dark adaptation of rods and cones in eyecup (Fig. 3b, 5b, and 5c). Fourth, cone pigment regeneration (Fig. 1a) and dark adaptation (Fig. 2a) were not observed in dissociated solitary cones or in cones from retina where proper contact with Müller cells was disturbed by a suction electrode (Supplementary Fig. 2 online). Thus, if any pigment epithelium contaminants were present in our retina preparation, their role in cone pigment regeneration was negligible.

Interaction between the Retina and the Pigment Epithelium Visual Cycles

Recently, two independent studies of *RPE65*^{-/-} *Nrl*^{-/-} mice found no evidence for chromophore recycling within the retina, raising doubts about the functional presence of such a pathway in rod-dominant species^{16, 17}. Both studies found that ablation of RPE65, a chromophore-binding protein essential for the pigment epithelium visual cycle²⁸, leads to the absence of 11-cis retinal in the *Nrl*^{-/-} retina populated exclusively by *Nrl* cones. The conclusion from these results was that the pigment epithelium is the only significant source of chromophore in the rod-dominant retina ruling out chromophore recycling in the retina. However, as our findings presented here clearly demonstrate the function of a visual cycle in the mouse retina (Fig. 6), the results from these studies most likely reflect an interaction between the two visual cycles. Specifically, as chromophore is not synthesized *de novo* in the eye, the pigment epithelium pathway is likely required for its initial uptake from the

circulation. Furthermore, the lack of chromophore in *RPE65^{-/-} Nr1^{-/-}* mice is not surprising if one considers that, as we show here, within the retina chromophore is recycled in Müller cells and supplied only to a small fraction of all photoreceptors (cones). Thus, simply increasing the mouse cone population would not be expected to increase the chromophore recycling capacity of the retina. Lastly, if RPE65 itself is expressed in cones (see Ref. 17 for discussion), its deletion might directly impair the retina visual cycle. Our results demonstrating the function of a cone visual cycle in the rod-dominant amphibian and mouse retinas settle this controversy.

Finally, unlike in primates, the salamander and mouse retinas do not have central cone-rich foveal region. Interestingly, while the inner retina neurons are sparse in the central primate retina, presumably to minimize light scattering, the ratio of cones to Müller cells in the fovea of primates is 1:129 indicating that they play an important function in the cone-rich central area of the retina. Experiments to determine whether a similar retina visual cycle is present in human and sub-human primates are currently underway.

Methods

Electrophysiology

Single-cell and whole-retina ERG recordings were done as previously described^{9, 30}. Briefly, dark-adapted larval tiger salamanders (*Ambystoma tigrinum*) were decapitated in dim red light, double-pithed, and the eyes were enucleated and hemisected. The retina was removed from the eyecup and any visible residual pigment epithelium was cleaned with forceps. For single-cell recordings, the retina was chopped with a razor blade and a small aliquot of cell suspension was placed in the recording chamber. Under infrared illumination, the outer segment of an isolated single cone or rod photoreceptor was drawn into the tip of a tight-fitting glass pipette for recording. For ERG recordings, the retina was transferred to a recording chamber on filter paper (photoreceptor side up). In both cases, tissue was perfused with amphibian Ringer solution containing 110 mM NaCl, 2.5 mM KCl, 1.6 mM MgCl₂, 1.0 mM CaCl₂, 10 mM dextrose, 10 mM HEPES, pH 7.8, and bovine serum albumin (100 mg/l)⁹. For ERG recordings, to isolate the photoreceptor component (a-wave) of the retina, the solution also contained 5 μM L-AP4 to block on-bipolar cell signals³¹, 5 μM NBQX to block AMPA/kainate signals³² and 50 μM D-AP5 to block NMDA signals³³. ERG recordings were done between a microelectrode built into the bottom of the chamber and a capillary microelectrode positioned above the retina. To suppress any glial components of the retina response, 10 mM barium chloride, which blocks potassium channels on Müller cells³⁴, was added to the chamber electrode. ERG recordings from retina in eyecup were performed in a modified recording chamber containing a small well in which the eyecup was placed. For mouse ERG recordings, C57/BL6 mice were euthanized by CO₂ asphyxiation/cervical dislocation and the retina was prepared as described above for salamander. The mouse retina was perfused using 36–38 °C bicarbonate-buffered solution containing 140 mM NaCl, 3.6 mM KCl, 2.4 mM MgCl₂, 1.2 mM CaCl₂, 3 mM HEPES (pH 7.4), 0.02 mM EDTA, 10 mM glucose, 0.1 × MEM amino acids, 0.1 × MEM Vitamins, 5 μM APB, 2 μM NBQX, and 10 μM D-AP5.

Photoreceptors were stimulated with calibrated 20-ms flashes. The signals were amplified, low-pass filtered at 20 Hz (8-pole Bessel) and digitized at 100 Hz for further analysis. Photosensitivity was calculated from the linear region of the intensity-response curve as the ratio of response amplitude and flash intensity. Integration time was calculated as the integral of the dim-flash response with the transient-peak amplitude normalized to unity.

Calculation of Pigment Content from Electrophysiology

We used the previously derived dependence of flash sensitivity on the percentage of pigment bleached by light for salamander red cones (Ref. 9, Fig. 4c) to estimate pigment content from the level of cone desensitization. This method was used for both single-cell and whole-retina ERG recordings. The pigment content values derived from sensitivity measurements matched very closely those measured directly using microspectrophotometry (Table 1).

Microspectrophotometry

Isolated photoreceptors were prepared as described above for electrophysiology. Microspectrophotometric measurements were made as previously described^{35, 36}. The optical density (O.D.) of a rod or cone outer segment was measured over the wavelength range of 400–700 nm with a rectangular slit smaller than the outer segment, where $O.D. = \log_{10}(I_o/I_t)$, where I_o is the transmitted light in the absence of a sample and I_t is the transmitted light through the outer segment. The collected O.D. spectra for each photoreceptor were fitted by the 11-cis A1 and A2 retinal spectral templates derived by Govardovskii and colleagues³⁷. Ringer solutions containing chromophore were prepared daily from 300 μ g dry aliquots of retinoid by dissolving the chemical in 0.1% ethanol as described previously⁹.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We thank Dr. Rosalie Crouch for a generous gift of 11-cis retinal and 11-cis retinol. We also thank Petri Ala-Laurila for his expert assistance with the analysis of microspectrophotometric data. We thank King-Wai Yau, Peter Lukasiewicz, Joseph Corbo, Rosalie Crouch, and Petri Ala-Laurila for comments on the manuscript. Supported by Career Development Award from Research to Prevent Blindness, Karl Kirchgessner Foundation, and EY 019312 (V.J.K), EY 01157 (M.C.C.), and EY 02687 (Department of Ophthalmology & Visual Sciences at Washington University).

REFERENCES

1. Ebrey T, Koutalos Y. Vertebrate photoreceptors. *Prog Retin Eye Res.* 2001; 20:49–94. [PubMed: 11070368]
2. Wald G, Brown PK, Smith fv. Iodopsin. *J Gen Physiol.* 1955; 38:623–681. [PubMed: 14367777]
3. Palczewski K, et al. Rod outer segment retinol dehydrogenase: substrate specificity and role in phototransduction. *Biochemistry.* 1994; 33:13741–13750. [PubMed: 7947785]
4. Jones GJ, Crouch RK, Wiggert B, Cornwall MC, Chader GJ. Retinoid requirements for recovery of sensitivity after visual-pigment bleaching in isolated photoreceptors. *Proc Natl Acad Sci U S A.* 1989; 86:9606–9610. [PubMed: 2594788]

5. Pepperberg DR, et al. Interphotoreceptor retinoid-binding protein (IRBP). Molecular biology and physiological role in the visual cycle of rhodopsin. *Mol Neurobiol.* 1993; 7:61–85. [PubMed: 8318167]
6. Hecht S, Haig C, Chase AM. Rod and cone dark adaptation. *J. Gen. Physiol.* 1937; 20:831–850. [PubMed: 19873031]
7. Rushton WA. Visual Adaptation. *Proc R Soc Lond B Biol Sci.* 1965; 162:20–46. [PubMed: 14296430]
8. Saari JC. Biochemistry of visual pigment regeneration: the Friedenwald lecture. *Invest Ophthalmol Vis Sci.* 2000; 41:337–348. [PubMed: 10670460]
9. Kefalov VJ, et al. Breaking the Covalent Bond- A Pigment Property that Contributes to Desensitization in Cones. *Neuron.* 2005; 46:879–890. [PubMed: 15953417]
10. Matsumoto H, Tokunaga F, Yoshizawa T. Accessibility of the iodopsin chromophore. *Biochim Biophys Acta.* 1975; 404:300–308. [PubMed: 1182163]
11. Mata NL, Radu RA, Clemmons RC, Travis GH. Isomerization and oxidation of vitamin A in cone-dominant retinas: a novel pathway for visualpigment regeneration in daylight. *Neuron.* 2002; 36:69–80. [PubMed: 12367507]
12. Das SR, Bhardwaj N, Kjeldbye H, Gouras P. Muller cells of chicken retina synthesize 11-cis-retinol. *Biochem J.* 1992; 285(Pt 3):907–913. [PubMed: 1497628]
13. Bustamante JJ, Ziari S, Ramirez RD, Tsini AT. Retinyl ester hydrolase and the visual cycle in the chicken eye. *Am J Physiol.* 1995; 269:R1346–R1350. [PubMed: 8594936]
14. Trevino SG, Villazana-Espinoza ET, Muniz A, Tsini AT. Retinoid cycles in the cone-dominated chicken retina. *J Exp Biol.* 2005; 208:4151–4157. [PubMed: 16244173]
15. Villazana-Espinoza ET, Hatch AL, Tsini AT. Effect of light exposure on the accumulation and depletion of retinyl ester in the chicken retina. *Exp Eye Res.* 2006; 83:871–876. [PubMed: 16780835]
16. Feathers KL, et al. Nrl-Knockout Mice Deficient in Rpe65 Fail to Synthesize 11-cis Retinal and Cone Outer Segments. *Invest. Ophthalmol. Vis. Sci.* 2008; 49:1126–1135. [PubMed: 18326740]
17. Wenzel A, et al. RPE65 is essential for the function of cone photoreceptors in NRL-deficient mice. *Invest Ophthalmol Vis Sci.* 2007; 48:534–542. [PubMed: 17251447]
18. Mariani AP. Photoreceptors of the larval tiger salamander retina. *Proc R Soc Lond B Biol Sci.* 1986; 227:483–492. [PubMed: 2873578]
19. Jablonski MM, Iannaccone A. Targeted disruption of Muller cell metabolism induces photoreceptor dysmorphogenesis. *Glia.* 2000; 32:192–204. [PubMed: 11008218]
20. Kato S, Ishita S, Sugawara K, Mawatari K. Cystine/glutamate antiporter expression in retinal Muller glial cells: implications for DL-alpha-amino acid toxicity. *Neuroscience.* 1993; 57:473–482. [PubMed: 7906874]
21. Pedersen OO, Karlsen RL. Destruction of Muller cells in the adult rat by intravitreal injection of D,L-alpha-amino acid. An electron microscopic study. *Exp Eye Res.* 1979; 28:569–575. [PubMed: 446576]
22. Bonaventure N, Roussel G, Wioland N. Effects of DL-alpha-amino adipic acid on Muller cells in frog and chicken retinae in vivo: relation to ERG b wave, ganglion cell discharge and tectal evoked potentials. *Neurosci Lett.* 1981; 27:81–87. [PubMed: 7329627]
23. Pow DV. Visualising the activity of the cystine-glutamate antiporter in glial cells using antibodies to amino acid, a selectively transported substrate. *Glia.* 2001; 34:27–38. [PubMed: 11284017]
24. Goldstein EB. Cone pigment regeneration in the isolated frog retina. *Vision Res.* 1970; 10:1065–1068. [PubMed: 5492790]
25. Jin J, Jones GJ, Cornwall MC. Movement of retinal along cone and rod photoreceptors. *Vis Neurosci.* 1994; 11:389–399. [PubMed: 8003460]
26. Sarantis M, Mobbs P. The spatial relationship between Muller cell processes and the photoreceptor output synapse. *Brain Res.* 1992; 584:299–304. [PubMed: 1325248]
27. Lamb TD, Pugh EN Jr. Dark adaptation and the retinoid cycle of vision. *Prog Retin Eye Res.* 2004; 23:307–380. [PubMed: 15177205]

28. Redmond TM, et al. Rpe65 is necessary for production of 11-cis-vitamin A in the retinal visual cycle. *Nat Genet.* 1998; 20:344–351. [PubMed: 9843205]
29. Ahmad KM, Klug K, Herr S, Sterling P, Schein S. Cell density ratios in a foveal patch in macaque retina. *Vis Neurosci.* 2003; 20:189–209. [PubMed: 12916740]
30. Green DG, Guo H, Pillers DA. Normal photoresponses and altered b-wave responses to APB in the mdx(Cv3) mouse isolated retina ERG supports role for dystrophin in synaptic transmission. *Vis Neurosci.* 2004; 21:739–747. [PubMed: 15683561]
31. Thoreson WB, Ulphani JS. Pharmacology of selective and non-selective metabotropic glutamate receptor agonists at L-AP4 receptors in retinal ON bipolar cells. *Brain Res.* 1995; 676:93–102. [PubMed: 7796182]
32. Yu W, Miller RF. NBQX, an improved non-NMDA antagonist studied in retinal ganglion cells. *Brain Res.* 1995; 692:190–194. [PubMed: 8548303]
33. Coleman PA, Miller RF. Do N-methyl-D-aspartate receptors mediate synaptic responses in the mudpuppy retina? *J Neurosci.* 1988; 8:4728–4733. [PubMed: 2904492]
34. Green DG, Kapousta-Bruneau NV. A dissection of the electroretinogram from the isolated rat retina with microelectrodes and drugs. *Vis Neurosci.* 1999; 16:727–741. [PubMed: 10431921]
35. Levine, JS.; MacNichol, EF, Jr. Microspectrophotometry of primate photoreceptors: art, artifact, and analysis. In: Fein, A.; Levine, JS., editors. *The visual system.* New York: Liss; 1985. p. 73-78.
36. MacNichol, EF, Jr. A photon-counting microspectrophotometer for the study of single vertebrate photoreceptor cells. In: Cool, SJ.; Smith, EL., editors. *Frontiers of visual science.* Berlin: Springer; 1978. p. 194-208.
37. Govardovskii VI, Fyhrquist N, Reuter T, Kuzmin DG, Donner K. In search of the visual pigment template. *Vis Neurosci.* 2000; 17:509–528. [PubMed: 11016572]




Figure 1. Effect of bleach on pigment content in salamander photoreceptors in dissociated and intact retina

Shown are average absorbance spectra of cones from dissociated retina (**a**, $n = 20$), cones from intact retina (**b**, $n = 20$), and rods from intact retina (**c**, $n = 10$) under three different conditions: in dark-adapted state (left), 2 hours after a bleach (middle), and following treatment with exogenous 11-cis retinal (right). In all cases photoreceptors were bleached by identical 40 s white light. Optical density in dark, bleached and 11-cis treated conditions are 0.049 ± 0.004 , 0.010 ± 0.002 , and 0.042 ± 0.002 respectively for cones of dissociated retina, 0.040 ± 0.002 , 0.039 ± 0.002 , and 0.040 ± 0.002 for cones of intact retina, and 0.130 ± 0.009 , 0.024 ± 0.003 , and 0.149 ± 0.006 for rods of intact retina. The fraction of bleached pigment was 82% in rods of intact retina and 79% in cones from dissociated retina. Note the recovery of pigment content after bleach of cones from intact retina (**b**) but not of cones from dissociated retina (**a**) or rods from intact retina (**c**). Error bars give s.e.m.




Figure 2. Effect of bleach on sensitivity in salamander photoreceptors in dissociated and intact retina

Suction recordings of flash intensity-response families from single dissociated cones (**a**), cones from intact retina (**b**), and rods from intact retina (**c**). Cells were stimulated at time 0 with 20-ms flashes of intensity increasing in 0.5 log unit steps. Top panels show test flash responses from cells in dark-adapted state (left), following a 40 s white light bleach (middle), and after treatment with exogenous 11-cis retinal (right). For cones (**a** and **b**), red traces represent photoresponses to $6,550 \text{ photons } \mu\text{m}^{-2}$, 620 nm. For rods (**c**), red traces represent photoresponses to $119 \text{ photons } \mu\text{m}^{-2}$, 520 nm. Bottom panels show the corresponding intensity-response relation for each cell, fit with Michaelis-Menten function $R/R_{max} = I/(I+I_O)$, where R/R_{max} is the normalized response amplitude, I is the flash intensity, and I_O is the intensity required to produce half-saturating response. Note the recovery of sensitivity of bleached cones from intact retina (**b**) but not of cones from dissociated retina (**a**) or rods from intact retina (**c**).




Figure 3. Effect of the Müller cell inhibitor L- α -AAA on the recovery of cone sensitivity following a bleach

Recordings from cones bleached in isolated intact retina (**a**) and in eyecup, with retina still attached to the pigment epithelium (**b**). All retinas were treated for 48 hours with 10 mM L- α -AAA and then transferred to Ringer prior to recordings. Red traces represent photoresponses to 6,550 photons μm^{-2} , 620 nm. Inhibiting the function of Müller cells blocked the recovery of sensitivity of cones from isolated retina but not of cones from eyecup following exposure to 40 s white bleaching light.




Figure 4. Rod and cone responses from salamander whole-retina ERG recordings

Background adaptation of isolated salamander retina (**a**), showing distinct rod and cone components. Data is fit with the Weber-Fechner relation $S/S_{DA} = (1 + I_B/I_O)^{-1}$, where S is the light-adapted sensitivity, S_{DA} is the dark-adapted sensitivity, I_B is the intensity of the background, and I_O is the background that reduced sensitivity to $0.5 S_{DA}$. I_O was $0.94 \text{ photons } \mu\text{m}^{-2} \text{ s}^{-1}$ for rods and $4,840 \text{ photons } \mu\text{m}^{-2} \text{ s}^{-1}$ for cones. Insets show a combined rod and cone response in darkness (left), and a cone-only response in background saturating the rods (right). Time course of test flash and background under each trace. Bottom panels

show rod (**b, d**) and cone (**c, e**) responses from one retina in darkness (top) and following a 40 s white light bleach (bottom). For each trace, a 20-ms flash was delivered at $t = 0$. Red traces represent photoresponses to 530 photons μm^{-2} , 520 nm for rods and 2,100 photons μm^{-2} , 620 nm for cones. Note the significant desensitization of rods and the recovery of sensitivity in cones from the same retina following the bleach. Error bars give s.e.m.




Figure 5. Kinetics of cone dark adaptation from whole-retina ERG recordings

(a) Recovery of salamander cone sensitivity driven by isolated retina in Ringer (black, $n = 4$) or following 48 hour treatment with L- α -AAA (red, $n = 3$). Cone sensitivity in Ringer recovered to 64% of its dark-adapted value, corresponding to regeneration of 93% cone pigment. In contrast, bleached cones from retina treated with L- α -AAA recovered only 3.2% of their dark-adapted sensitivity. (b) Recovery of rod sensitivity in eyecup in Ringer (black, $n = 4$) and following 48 hour treatment with L- α -AAA (red, $n = 4$). L- α -AAA did not affect the rate or final level of recovery of sensitivity of rods driven by the pigment epithelium. (c) Recovery of cone sensitivity in eyecup in Ringer (black, $n = 5$), driven by both retina and pigment epithelium visual cycles, or following 48 hour treatment with L- α -AAA (red, $n = 5$), driven by the pigment epithelium alone. The recovery of cone sensitivity was significantly accelerated and driven to completion by the addition of the retina visual cycle. (d) Recovery of cone pigment content, estimated from (c), using the relation between cone pigment loss and desensitization⁹. The initial rate of pigment regeneration by the retina visual cycle (open circles) was estimated by subtracting the pigment regenerated by the pigment epithelium (red) from the total regenerated pigment (black). Note the comparable

rates of pigment regeneration by the two cycles. In all cases identical 40 s white bleaching light was delivered at $t = 0$. Error bars give s.e.m.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 6. Rod and cone responses from mouse whole-retina ERG recordings

Shown are rod (a, c) and cone (b, d) responses from one retina in darkness (top) and following a 40 s 9.7×10^6 photons $\mu\text{m}^{-2} \text{s}^{-1}$ 500 nm bleach (bottom). Test flashes of intensity increasing in 0.5 log unit steps were delivered at $t = 0$. Red traces represent photoresponses to 1,977 photons μm^{-2} for rods and 22,850 photons μm^{-2} for cones, both at 500 nm. Note the significant desensitization of rods and the recovery of sensitivity in cones from the same retina following the bleach.

Table 1
Percentage of cone pigment content following a bleach

For microspectrophotometry, cone pigment content was estimated from optical density. For single-cell and whole-retina ERG recordings, cone pigment content was estimated from the bleach-induced desensitization using the relation between pigment content and sensitivity in salamander red cones (see text for details). All values are the mean \pm s.e.m.

	Dissociated Retina	Intact Retina	Intact Retina + L- α -AAA
Microspectrophotometry	21 \pm 4 (<i>n</i> = 20)	96 \pm 5 (<i>n</i> = 20)	31 \pm 5 (<i>n</i> = 15)
Single-cell Recordings	17 \pm 2 (<i>n</i> = 20)	93 \pm 2 (<i>n</i> = 14)	32 \pm 3 (<i>n</i> = 15)
Whole-retina ERG	--	92 \pm 2 (<i>n</i> = 11)	23 \pm 2 (<i>n</i> = 4)

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript