

CASE REPORT

Cutis verticis gyrata: Three cases illustrating three different etiologies

Amal Chamli^{1,2}  | Meriem Jones^{1,2} | Takwa Bacha^{1,2} | Nouredine Litaïem^{1,2} | Faten Zegloui^{1,2}

¹Department of Dermatology, Charles Nicolle Hospital, Tunis, Tunisia

²Faculty of Medicine of Tunis, University of Tunis El Manar, Tunis, Tunisia

Correspondence

Amal Chamli, Department of Dermatology, Charles Nicolle Hospital, Tunis, Tunisia.
Email: amal.fmt@gmail.com

Funding information

None

Abstract

Cutis Verticis gyrata is an uncommon neurocutaneous syndrome characterized by excessive growth of the skin of the scalp or the face, forming folds of similar aspect to cerebral cortex gyri. Three categories have been individualized: the primary form, essential or non-essential, and the secondary form.

KEYWORDS

cutis verticis gyrata, folds, hypertrophy, neurocutaneous syndrome

1 | INTRODUCTION

Cutis verticis gyrata (CVG) is a rare skin disease, affecting 1 of 100,000 males and 0.026 of 100,000 females in the general population.¹ CVG is characterized by excessive growth of the skin of the scalp or the face, forming folds of similar aspect to cerebral cortex gyri. The following three forms have been individualized: the primary form, essential or non-essential, and the secondary form. The main complaint of the patients is usually cosmetic. The treatment, if requested, is surgical. We report three cases that illustrate one of each of these three categories.

2 | OBSERVATIONS

2.1 | Case 1

A 28-year-old male patient presented with multiple trichilemmal cysts of the scalp. Physical examination revealed, in addition, a thickened scalp with multiple symmetric skin folds from vertex to occipital area (Figure 1).

The systemic examination did not reveal any neurologic, ophthalmologic, or metabolic anomalies. Laboratory investigations were within the normal range limit. The diagnosis of primary essential CVG was retained and the patient required only excision of the cysts.

2.2 | Case 2

A 32-year-old male patient with a history of epilepsy was referred to our consultation for a drug reaction secondary to phenobarbital. On examination, apart from the skin lesions consistent with the morbilliform drug reaction (Figure 2), the patient had cerebriform folds on his forehead. The furrows were running horizontally and had a width of 1–2 cm (Figure 3). He did not have any lesions on the scalp. No other anomalies were found. Laboratory investigations including growth hormone and insulin-like growth factor levels were within normal range. Computerized tomography of the brain was unremarkable. The diagnosis of primary non-essential CVG associated with epilepsy and involving only the forehead was retained.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. *Clinical Case Reports* published by John Wiley & Sons Ltd.



FIGURE 1 Multiple symmetric skin folds extending from the vertex to the occipital area in a 28-year-old male patient



FIGURE 2 Morbiliform drug eruption: red papules and macules on the trunk

2.3 | Case 3

A 54-year-old female patient treated for hypertension, dyslipidemia, and a heart rhythm disorder, presented to our department with an occipital tumefaction evolving for a few years. The examination revealed cerebriform transverse occipital skin folds. The remaining physical

examination was otherwise unremarkable. Given her chronic metabolic pathologies, secondary CVG was diagnosed.

3 | DISCUSSION

CVG is a rare benign condition first described by Alibert in 1837 and characterized by proliferation and hypertrophy of the skin of the scalp or forehead. The affected skin then exhibits folds and furrows resembling the cerebral cortex convolutions. Unna introduced the term *cutis veritica gyrata* in 1907.²

This entity was afterward divided into different subtypes: a secondary and a primary form, then the primary form was subdivided into primary essential and non-essential. The primary essential form is the rarest form, characterized by an isolated cutaneous involvement with no associated pathology. It usually begins in adolescence and is more common in men than in women. Typically, as seen in case 1, primary CVG presents as symmetric scalp folds that usually extend anteroposteriorly from the vertex to the occiput and transversally in the occipital region. Terminal hair density is reduced on the folds, but not in the furrows. The primary non-essential form has been described in association with several neuropsychiatric pathologies like mental retardation, epilepsy, microcephaly, schizophrenia, encephalopathy, and other neurological malformations in addition to ophthalmic pathologies such as blindness, strabismus, and congenital cataract, the latter being the most frequent association. In primary non-essential CVG, the associated conditions do not have a clear pathophysiological link with the CVG.

Secondary CVG, on the other hand, is associated with conditions that may be implicated in modifications of the trophicity of the skin. It is slightly more common than the primary form. It can occur at any age without gender predominance. It is associated with a variety of underlying disorders. It may be caused by general diseases comprise as liver affections, paraneoplastic syndromes, and pachydermoperiostosis or also associated with inflammatory dermatoses most commonly psoriasis and eczema.³ In our second case, the absence of digital clubbing, and evident enlargement of bones in clinical examination allowed us to rule out the diagnosis of pachydermoperiostosis. Furthermore, various associated genetic conditions have been described such as neurofibromatosis, Noonan, Turner, Klinefelter and Ehlers-Danlos syndrome. It may also occur in association with endocrine etiologies, as seen in case 3, such as diabetes mellitus, myxedema, and the most described in the literature is acromegaly.³ Among iatrogenic causes, treatment with minoxidil or misuses of anabolic substances were reported.^{4,5}



FIGURE 3 Cerebriform folds on the forehead of a 32-year-old male patient

The pathophysiology of CVG remains unclear. On a structural level, it is suggested that the folds are caused by connective tissue septa formation between the skin and the galea. These septa prevent the expansion of the skin which bulges into folds.⁶ In the primary form, an endocrine origin has been postulated as the disease usually manifests in postpubertal males. Cases reported in the literature are sporadic although some familial cases are described suggesting possible genetic factors in primary CVG. The physiopathology in the secondary form remains also unclear, especially in general and inflammatory diseases. However, in the physiopathology of endocrine etiology, trophic hormones such as growth hormone (GH) and insulin growth factor 1 (IGF 1) were incriminated to be responsible for skin modifications with collagen thickening and hypertrophy of sebaceous glands.

When performed, histopathology of primary CVG shows normal skin. Thickened connective tissue with hypertrophy or hyperplasia of adnexal structures has also been described. Secondary CVG has variable histology representing the underlying disease.

No treatment has proven its effectiveness in this entity. Abstention is the rule in the primary form. In all cases, regular rigorous hygiene of the skin folds and furrows is recommended to avoid maceration and infection. If treatment is requested by the patient because of an anesthetic complaint, medical and/or surgical treatment may be

suggested. Surgical treatment seems to be the best option with a regularly reported efficacy. The undermining and relaxation incisions alter the relationship between the skin and the deeper structures and so result in the disappearance of the folds and furrows.⁷ Medical treatment includes isotretinoin or systemic steroids. No treatment can prevent a recurrence.

In summary, CVG is a rare disease with mainly anesthetic impact. A rigorous clinical examination with a minimal workup is, however, required to eliminate a secondary or a primary non-essential form. If esthetic prejudice is important, surgical treatment can be suggested.

AUTHOR CONTRIBUTIONS

Drs Amal Chamli and Meriem Jones contributed to the first draft of the manuscript. Drs Amal Chamli, Meriem Jones, Takwa Bacha, and Noureddine Litaïem contributed to the literature search, analysis, and interpretation of the data. Dr Faten Zeglaoui critically revised the manuscript and gave final approval. All authors read and approved the final manuscript and agree to be finally accountable for ensuring the integrity of and accuracy of the work.

ACKNOWLEDGMENT

None.

CONFLICT OF INTEREST

None declared.

DATA AVAILABILITY STATEMENT

Data sharing was not applicable to this article as no datasets were generated or analyzed during the current study.

ETHICAL APPROVAL

Appropriate consent has been obtained, prior to submission, for the publication of images and data.

CONSENT

Written informed consent was obtained from the patient to publish this report in accordance with the journal's patient consent policy.

ORCID

Amal Chamli  <https://orcid.org/0000-0002-7927-9088>

REFERENCES

1. Harish V, Clarke F. Isolated cutis verticis gyrata of the glabella and nasal bridge: a case report and review of the literature. *J Plast Reconstr Aesthet Surg*. 2013;66(10):1421-1423.
2. Samira B, Meriem D, Oumkeltoum E, Driss E, Yassine B, Saloua E. Primitive essential Cutis verticis gyrata, a rare skin disease: case report and review of literature. *Pan Afr Med J*. 2014;19:345.
3. Chentli F, Terki B, Azzoug S. Cutis verticis gyrata and acromegaly. *Presse Med*. 2015;44(9):958-961.

4. Nguyen KH, Marks JG. Pseudoacromegaly induced by the long-term use of minoxidil. *J Am Acad Dermatol.* 2003;48(6):962-965.
5. Sommer A, Gambichler T, Altmeyer P, Kreuter A. A case of cutis verticis gyrata, induced by misuse of anabolic substances? *Clin Exp Dermatol.* 2006;31(1):134-136.
6. Tani T, Miyamoto Y, Kurozumi N. Surgical treatment of cutis verticis gyrata: a report of 2 cases. *Br J Plast Surg.* 1977;30(3):235-239.
7. El-Husseiny M, Yarrow J, Moiemmen N. Primary cutis gyrata: review of literature and a successful new surgical approach. *Eur J Plast Surg.* 2010;33(3):153-157.

How to cite this article: Chamli A, Jones M, Bacha T, Litaïem N, Zeglaoui F. Cutis verticis gyrata: Three cases illustrating three different etiologies. *Clin Case Rep.* 2022;10:e05889. doi:[10.1002/ccr3.5889](https://doi.org/10.1002/ccr3.5889)