



## Neurological manifestations by sex and age group in COVID-19 in-hospital patients

Daniella Nunes Pereira<sup>a,\*</sup>, Maria Aparecida Camargos Bicalho<sup>a,b</sup>, Alzira de Oliveira Jorge<sup>a,c</sup>, Angélica Gomides dos Reis Gomes<sup>d</sup>, Alexandre Vargas Schwarzbald<sup>e</sup>, Anna Luiza Homan Araújo<sup>f</sup>, Christiane Corrêa Rodrigues Cimini<sup>g,w,x</sup>, Daniela Ponce<sup>h</sup>, Danyelle Romana Alves Rios<sup>i,y</sup>, Genna Maira Santos Grizende<sup>j</sup>, Euler Roberto Fernandes Manenti<sup>k</sup>, Fernando Anschau<sup>l</sup>, Fernando Graça Aranha<sup>m</sup>, Frederico Bartolazzi<sup>n</sup>, Joanna d'Arc Lyra Batista<sup>o,z,aa</sup>, Julia Teixeira Tupinambás<sup>p</sup>, Karen Brasil Ruschel<sup>q</sup>, Maria Angélica Pires Ferreira<sup>r</sup>, Pedro Gibson Paraíso<sup>s</sup>, Sílvia Ferreira Araújo<sup>t</sup>, Antonio Lucio Teixeira<sup>u,ab</sup>, Milena Soriano Marcolino<sup>v,x</sup>

<sup>a</sup> Faculdade de Medicina, Universidade Federal de Minas Gerais, Brazil

<sup>b</sup> Fundação Hospitalar do Estado de Minas Gerais (Fhemig), Germany

<sup>c</sup> Hospital Risoleta Tolentino Neves, Brazil

<sup>d</sup> Hospitais da Rede Mater Dei, Brazil

<sup>e</sup> Hospital Universitário de Santa Maria/EBSERH; Universidade Federal de Santa Maria, Brazil

<sup>f</sup> Hospital Universitário Ciências Médicas, Brazil

<sup>g</sup> Hospital Santa Rosália, Brazil

<sup>h</sup> Hospital das Clínicas da Faculdade de Medicina de Botucatu, Brazil

<sup>i</sup> Complexo da Saúde São João de Deus, Universidade Federal de São João del-Rei, Brazil

<sup>j</sup> Santa Casa de Misericórdia de Belo Horizonte, Brazil

<sup>k</sup> Hospital Mãe de Deus, Brazil

<sup>l</sup> Grupo Hospitalar Conceição, Brazilian National Health System, Hospital Nossa Senhora da Conceição and Hospital Cristo Redentor, Brazil

<sup>m</sup> Hospital SOS Córdio, Brasil

<sup>n</sup> Hospital Santo Antônio, Brasil

<sup>o</sup> Universidade Federal Fronteira do Sul, Brazil

<sup>p</sup> Hospital Metropolitano Odilon Behrens, Brazil

<sup>q</sup> Hospital Mãe de Deus, Hospital Universitário de Canoas, Universidade Federal do Rio Grande do Sul e Instituto de Avaliação de Tecnologia em Saúde (IATS/CNPQ), Brazil

<sup>r</sup> Hospital de Clínicas de Porto Alegre; IATS/CnPq, Brazil

<sup>s</sup> Orizonti- Instituto Oncomed de Saúde e Longevidade, Brazil

<sup>t</sup> Hospital Semper, Brazil

<sup>u</sup> Faculdade Santa Casa BH, Brazil

<sup>v</sup> Department of Internal Medicine, Medical School, and Telehealth Center, University Hospital, Universidade Federal de Minas Gerais, Brazil

<sup>w</sup> Universidade Federal dos Vales do Jequitinhonha e Mucuri (UFVJM), Brazil

<sup>x</sup> Institute for Health Technology Assessment (IATS/ CNPq), Brazil

<sup>y</sup> Campus Centro Oeste Dona Lindu, Universidade Federal de São João del-Rei, Brazil

<sup>z</sup> Instituto de Avaliação de Tecnologia em Saúde (IATS/CNPQ), Brazil

<sup>aa</sup> Hospital Regional do Oeste, Brazil

<sup>ab</sup> Neuropsychiatry Program, Department of Psychiatry and Behavioral Sciences, UT Health Houston, Brazil

### ARTICLE INFO

**Keywords:**  
COVID-19

### ABSTRACT

**Introduction:** Neurological manifestations have been associated with a poorer prognosis in COVID-19. However, data regarding their incidence according to sex and age groups is still lacking.

**Abbreviations:** ACE2, Angiotensin-converting enzyme 2; AD, Alzheimers disease; BMI, Body mass index; CI, Confidence interval; CNS, Central nervous system; COPD, Chronic obstructive pulmonary disease; IQR, Interquartile range; OR, Odds ratio.

\* Corresponding author.

E-mail addresses: [danielanp@ufmg.br](mailto:danielanp@ufmg.br) (D.N. Pereira), [milenamarc@ufmg.br](mailto:milenamarc@ufmg.br) (M.S. Marcolino).

<https://doi.org/10.1016/j.ensci.2022.100419>

Received 15 April 2022; Received in revised form 15 July 2022; Accepted 24 July 2022

Available online 30 July 2022

2405-6502/© 2022 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Neurological manifestations  
Age  
Sex  
Delirium

**Methods:** This retrospective multicentric cohort collected data from 39 Brazilian hospitals from 17 cities, from adult COVID-19 admitted from March 2020 to January 2022. Neurological manifestations presented at hospital admission were assessed according to incidence by sex and age group.

**Results:** From 13,603 COVID-19 patients, median age was 60 years old and 53.0% were men. Women were more likely to present with headaches (22.4% vs. 17.7%,  $p < 0.001$ ; OR 1.36, 95% confidence interval [CI] 1.22–1.52) than men and also presented a lower risk of having seizures (OR 0.43, 95% CI 0.20–0.94). Although delirium was more frequent in women (6.6% vs. 5.7%,  $p = 0.020$ ), sex was not associated with delirium in the multivariable logistic regression analysis. Delirium, syncope and coma increased with age (1.5% [18–39 years] vs. 22.4% [80 years or over],  $p < 0.001$ , OR 1.07, 95% CI 1.06–1.07; 0.7% vs. 1.7%,  $p = 0.002$ , OR 1.01, 95% CI 1.00–1.02; 0.2% vs. 1.3%  $p < 0.001$ , OR 1.04, 95% CI 1.02–1.06), while, headache (26.5% vs. 7.1%, OR 0.98, 95% CI 0.98–0.99), anosmia (11.4% vs. 3.3%, OR 0.99, 95% CI] 0.98–0.99 and ageusia (13.1% vs. 3.5%, OR 0.99, CI 0.98–0.99) decreased ( $p < 0.001$  for all).

**Conclusion:** Older COVID-19 patients were more likely to present delirium, syncope and coma, while the incidence of anosmia, ageusia and headaches decreased with age. Women were more likely to present headache, and less likely to present seizures.

## 1. Introduction

Neurological manifestations have been frequently associated with COVID-19 [1]. Symptoms may differ among patients, including symptoms and signs of both peripheral and central nervous system (CNS) dysfunction [1–3]. Such neurological manifestations, especially clinically-defined neurological syndromes (e.g. delirium), have been associated with a poorer COVID-19 prognosis, even when controlling for age, sex and number of medical comorbidities [4,5].

Previous studies have shown that age and sex are important prognostic factors in COVID-19 patients. Children and young adults are not as prone to severe forms of COVID-19 compared to older adults [6–9]. Elderly patients are more likely to progress to severe disease and mortality due to age-related immunosenescence and comorbidities [10–12]. The reasons for sex differences are believed to be related to distinct immune response profiles [13] and gender inequalities, including socioeconomic status and occupational exposure [14–16]. Age and sex-related differences in the efficiency of the immune response may influence disease presentation and outcomes [17].

In this context, there is a dearth of data regarding incidence of neurological manifestations in COVID-19 patients according to sex and age groups [18], especially in Latin America. Therefore, our aim was to fill this gap in knowledge, assessing a large database of Brazilian patients.

## 2. Methods

This study is part of a multicentric cohort, Brazilian COVID Registry Project, which collected data from 39 Brazilian hospitals from 17 Brazilian cities, described in detail elsewhere [19]. It was approved by the National Commission for Research Ethics (CAAE 30350820.5.1001.0008). Due to the worldwide concern on the pandemic and the urgency for studies of COVID-19 and to the fact that data was collected solely by review of medical records, individual informed consent was waived by the National Commission for Research Ethics.

Consecutive adult patients with laboratory confirmed diagnosis of COVID-19 [20], who were admitted to the participating hospitals from March 2020 to January 2022, were enrolled in the study. Data on patient's demographic and clinical characteristics, laboratory findings and outcomes were collected through the revision of medical records by trained hospital staff using Research Electronic Data Capture (REDCap) tools [21]. Neurological manifestations caused by COVID-19 presented at hospital presentation systematically assessed in this study included stroke, delirium, coma, seizures, manifestations of the peripheral nervous system (e.g. hypo/anesthesia, dysesthesia, pain), anosmia, ageusia, headaches and syncope.

For data analysis, categorical data was expressed by proportions and absolute numbers, meanwhile continuous variables were presented as

medians and interquartile ranges. To compare distribution of categorical and continuous variables, the Fisher Exact test and the Kruskal-Wallis tests were used, respectively. The analysis was conducted evaluating neurological manifestations by sex and age group: i) 18–39; ii) 40–59; iii) 60–79; iv) 80 years old or over.

To assess age and sex as predictors of each neurological manifestation, stepwise multivariate logistic regression analysis was performed. In addition to age and sex, other variables assessed at hospital admission which have shown to be prognostic predictors in a previous analysis were also included as covariates [22]. For the analysis of self-reported symptoms (ageusia, anosmia, headache), patients with coma, delirium and dementia were not included in the models given the possibility of reporting bias.

All statistical analyses were performed using the R software (version 4.0.2), and its packages data.table, tidyverse, gtsummary and gt. A  $p$ -value below 0.05 was considered as statistically significant.

## 3. Results

The current study included 13,603 COVID-19 patients (median age 60 years old, interquartile range [IQR] 47–71, 53.0% were men). Hypertension (53.0%) was the most common medical comorbidity, followed by diabetes (27.1%) and obesity (17.8%) (Table 1).

When comparing demographic data regarding both sexes, it was observed that women were slightly older than men (median age of 61 vs. 58 years old). Also, the prevalence of hypertension, heart failure, asthma, diabetes, psychiatric conditions, obesity and dementia were higher in women, meanwhile coronary artery disease, cirrhosis and chronic kidney disease were more prevalent in men (Table 1).

When stratifying groups according to sex, women had a higher frequency of delirium (6.6% vs. 5.7%,  $p = 0.020$ ) and headache (22.4% vs. 17.7%,  $p < 0.001$ ) when compared to men (Table 2). However, sex was not associated with delirium in the multivariable logistic regression analysis. Women had a higher risk of headaches (OR 1.36, 95% confidence interval [CI] 1.22–1.52), and lower risk of seizures (OR 0.43, 95% CI 0.20–0.94) (Table 3).

As shown on Table 4, incidence of delirium, syncope and coma increased with age; older age was associated with delirium and coma in the multivariable logistic regression analysis. Conversely, headache, anosmia and ageusia decreased with increased age, findings confirmed in the multivariate analysis. (Tables 3 and 4).

## 4. Discussion

The Brazilian COVID Registry Project is one of the largest cohorts of COVID-19 patients which assessed neurological manifestations in detail. We observed that older patients presented a higher incidence of neurological manifestations associated with worse prognosis, such as delirium and coma, when compared to other age groups. Conversely,

**Table 1**

Demographic and clinical characteristics according to sex in the Brazilian COVID Registry Project, 2020–2022 (n = 13,603).

Characteristics	Overall <sup>a</sup> (n = 13,603)	Men <sup>a</sup> (n = 7211)	Women <sup>a</sup> (n = 6392)	p-value <sup>b</sup> c
Age	60.0 (47.0, 71.0)	58.0 (46.0, 70.0)	61.0 (48.0, 73.0)	<0.001
Atrial flutter/fibrillation	379 (2.8%)	197 (2.7%)	182 (2.8%)	0.722
Hypertension	7205 (53.0%)	3486 (48.3%)	3719 (58.2%)	<0.001
Coronary artery disease	683 (5.0%)	421 (5.8%)	262 (4.1%)	<0.001
Heart failure	745 (5.5%)	348 (4.8%)	397 (6.2%)	<0.001
Stroke	458 (3.4%)	226 (3.1%)	232 (3.6%)	0.121
Chagas disease	45 (0.3%)	21 (0.3%)	24 (0.4%)	0.481
Asthma	797 (5.9%)	256 (3.6%)	541 (8.5%)	<0.001
COPD	697 (5.1%)	350 (4.9%)	347 (5.4%)	0.139
Pulmonary fibrosis	55 (0.4%)	26 (0.4%)	29 (0.5%)	0.472
Diabetes mellitus	3689 (27.1%)	1790 (24.8%)	1899 (29.7%)	<0.001
Obesity (BMI > 30 kg/m <sup>2</sup> )	2417 (17.8%)	1059 (14.7%)	1358 (21.2%)	<0.001
Cirrhosis	76 (0.6%)	50 (0.7%)	26 (0.4%)	0.034
Psychiatric diseases	1018 (7.5%)	354 (4.9%)	664 (10.4%)	<0.001
Chronic renal disease	674 (5.0%)	382 (5.3%)	292 (4.6%)	0.055
Dementia	298 (2.2%)	111 (1.5%)	187 (2.9%)	<0.001

<sup>a</sup> Statistics are presented as n (%).<sup>b</sup> Statistical tests performed: chi-square test of independence; Fisher's exact test.<sup>c</sup> COPD: chronic obstructive pulmonary disease; BMI: body mass index.**Table 2**

Incidence of neurological manifestations according to sex in the Brazilian COVID Registry Project, 2020–2022 (n = 13,603).

Neurological manifestation	Overall <sup>a</sup> (N = 13,603) n(%)	Men <sup>a</sup> (N = 7211) n(%)	Women <sup>a</sup> (N = 6392) n(%)	p-value <sup>b</sup>
Headache	2705 (19.9%)	1275 (17.7%)	1430 (22.4%)	<0.001
Ageusia	1275 (9.4%)	643 (8.9%)	632 (9.9%)	0.056
Anosmia	1110 (8.2%)	561 (7.8%)	549 (8.6%)	0.091
Delirium	834 (6.1%)	409 (5.7%)	425 (6.6%)	0.020
Syncope	133 (1.0%)	69 (1.0%)	64 (1.0%)	0.861
Coma	68 (0.5%)	36 (0.5%)	32 (0.5%)	>0.999
Stroke	27 (0.2%)	16 (0.2%)	11 (0.2%)	0.647
Seizures	39 (0.3%)	26 (0.4%)	13 (0.2%)	0.121
Peripheral neuropathy	10 (0.1%)	8 (0.1%)	2 (0.0%)	0.116

<sup>a</sup> Statistics are presented as n (%).<sup>b</sup> Statistical tests performed: chi-square test of independence; Fisher's exact test.

younger patients were more likely to present mild neurological symptoms, such as anosmia, ageusia and headache. Women were more likely to present headache, and less likely to have seizures.

COVID-19 clinical presentations vary widely from asymptomatic cases and mild respiratory symptoms to severe pulmonary, neurological and cardiovascular manifestations [23]. When infecting the human body, SARS-CoV-2 binds to Angiotensin-Converting Enzyme 2 (ACE2) [24] receptors present in different cell types of the CNS [25]. These CNS cells expressing ACE2 receptors are mainly present in the posterior cortex, posterior cingulate cortex, medial temporal gyrus and the olfactory bulb [25]. Besides a potential direct effect of SARS-CoV-2 on neurons and glial cells, the elicited systemic inflammatory cascade can lead to neuroinflammation and, therefore, contribute to CNS dysfunction and related symptoms and signs [26–28]. This unregulated inflammatory response is more common in elderly people, what justifies a higher incidence of symptoms related to a poorer prognosis and are more likely to present severe COVID-19 neurological presentation,

**Table 3**

Prediction models for each neurological manifestation, taking account clinical features obtained at hospital presentation.

Variable	OR (CI 95%)	p-value
<b>Ageusia</b>		
(Intercept)	0.636	0.102
Age	0.987 (0.982–0.992)	<0.001
Female sex	1.124 (0.978–1.292)	0.100
Urea	0.995 (0.992–0.998)	0.001
C-reactive protein	0.998 (0.997–0.999)	<0.001
Platelet count	0.999 (0.998–1.000)	0.002
Heart rate	0.996 (0.992–1.001)	0.112
<b>Anosmia</b>		
(Intercept)	0.460	0.004
Age	0.987 (0.982–0.992)	<0.001
Urea	0.997 (0.994–1.000)	0.035
C-reactive protein	0.999 (0.998–1.000)	0.020
Heart rate	0.994 (0.990–0.999)	0.016
<b>Coma</b>		
(Intercept)	0.002	<0.001
Age	1.039 (1.020–1.058)	<0.001
Mechanical ventilation	12.109 (5.411–27.098)	<0.001
Oxygen saturation	0.994 (0.991–0.996)	<0.001
<b>Delirium</b>		
(Intercept)	0.000	<0.001
Age	1.070 (1.064–1.077)	<0.001
Mechanical ventilation	0.000 (0.000)	0.957
Urea	1.006 (1.004–1.008)	<0.001
Oxygen saturation	0.999 (0.998–1.000)	0.020
Heart rate	1.012 (1.007–1.017)	<0.001
<b>Headache</b>		
(Intercept)	1.477	0.082
Age	0.981 (0.978–0.985)	<0.001
Female sex	1.363 (1.224–1.517)	<0.001
Urea	0.993 (0.990–0.995)	<0.001
Number of comorbidities	0.920 (0.871–0.971)	0.003
C-reactive protein	0.999 (0.998–1.000)	0.004
SF ratio	1.000 (1.000–1.001)	0.055
Platelets	0.999 (0.999–1.000)	0.027
Heart rate	0.998 (0.994–1.001)	0.147
<b>Peripheral neuropathy</b>		
(Intercept)	0.001	<0.001
Number of comorbidities	0.657 (0.338–1.275)	0.214
<b>Seizure</b>		
(Intercept)	0.008	<0.001
Female sex	0.433 (0.199–0.942)	0.035
C-reactive protein	0.991 (0.984–0.998)	0.009
<b>Stroke</b>		
(Intercept)	0.001	<0.001
Platelets	1.004 (1.001–1.007)	0.016
<b>Syncope</b>		
(Intercept)	0.005	0 < 0.001
Age	1.012 (1.000–1.025)	0.059
Number of comorbidities	1.157 (0.984–1.361)	0.078
C-reactive protein <sup>a</sup>	0.998 (0.995–1.000)	0.077

<sup>a</sup> OR: odds ratio; CI: confidence interval

including delirium and coma. This finding corroborated previous studies showing delirium as the most frequent neurological symptom in older adults with COVID-19 [29,30]. Delirium has even been described as an atypical presentation of COVID-19, particularly in frailty older adults and those with dementia [31].

Recent systematic reviews observed that anosmia was more common in less-severe cases of COVID-19, but information on anosmia and ageusia in older patients was still limited [32,33]. The current study addressed this gap showing that older patients are less likely to present anosmia. Studies with non-COVID-19 upper respiratory tract infections have already shown that women present a higher prevalence of anosmia than men. Nonetheless, the prevalence of anosmia in these respiratory infections was higher in older than younger patients [34]. In COVID-19, younger patients have a higher incidence of anosmia and ageusia. Since delirium is frequent in older patients with COVID-19, it is possible that these patients with delirium did not report anosmia and ageusia. The same trend of decreasing typical manifestations and increasing

**Table 4**  
Incidence of neurological manifestations according to age group in the Brazilian COVID Registry Project, 2020–2022 (n = 13,603).

Manifestation	Age group (years-old) <sup>a</sup>				p-value <sup>b</sup>
	18–39 N = 1840 n(%)	40–59 N = 4923 n(%)	60–79 N = 5277 n(%)	80 or over N = 1563 n(%)	
Delirium	27 (1.5%)	107 (2.2%)	350 (6.6%)	350 (22.4%)	<0.001
Coma	4 (0.2%)	17 (0.3%)	26 (0.5%)	21 (1.3%)	<0.001
Seizure	6 (0.3%)	13 (0.3%)	15 (0.3%)	5 (0.3%)	0.924
Stroke	3 (0.2%)	7 (0.1%)	14 (0.3%)	3 (0.2%)	0.570
Anosmia	210 (11.4%)	473 (9.6%)	375 (7.1%)	52 (3.3%)	<0.001
Ageusia	241 (13.1%)	565 (11.5%)	414 (7.8%)	55 (3.5%)	<0.001
Peripheral neuropathy	2 (0.1%)	6 (0.1%)	1 (0.0%)	1 (0.1%)	0.153
Headache	487 (26.5%)	1271 (25.8%)	836 (15.8%)	111 (7.1%)	<0.001
Syncope	12 (0.7%)	36 (0.7%)	58 (1.1%)	27 (1.7%)	0.002

<sup>a</sup> Statistics are presented as n (%).

<sup>b</sup> Statistical tests performed: chi-square test of independence; Fisher's exact test.

frequency of delirium with age has been observed in other diseases, such as in viruses' and bacterial infections [33,35].

The increased incidence of headache in women might reflect a larger epidemiological and biological trend that women are more prone to headache disorders than men [36,37]. A previous outpatient clinic study also observed that headaches were more frequent COVID-19 symptoms in women, as well as olfactory and taste dysfunctions [38]. Although it is considered a mild symptom, post-COVID headaches can be persistent for over three months in up to 19% of cases, impacting directly in the quality of life of post-COVID-19 patients [39].

It is unclear why women had a higher frequency of delirium than men in the study sample. This is a puzzling finding, as male sex is a risk factor for severe COVID-19, and delirium has been associated with worse outcomes, representing a sign of brain/neurological dysfunction [40]. However, in the multivariate analysis, sex was not a predictor for delirium, indicating confounding factors. For instance, the prevalence of dementia was higher in women than men (2.9% vs. 1.5%;  $p < 0.001$ ) and, as a consequence, of delirium [41]. Compelling evidence suggests that women have a greater lifetime risk of developing Alzheimer's disease (AD) or any dementia than men. At age 65, men and women have a lifetime risk of approximately 14% and 20%, respectively, to develop dementia [42,43]. Of note, there is a growing interest in the study of the contribution of social/cultural determinants of health, including social circumstances, environmental characteristics, and early-life exposures, to delirium risk. In this context, it would be tempting to speculate whether these social determinants played any role in the susceptibility of Brazilian women to delirium. However, the matter is still understudied [44,45], and even education level, an important risk factor for dementia, is not properly considered in the delirium literature [45].

The incidence of coma at hospital presentation was low (0.5%, and 1.3% in octogenarians). Results of a registry for neurologic manifestations of 971 patients from 19 countries from an earlier phase of the pandemic also reported that coma corresponded to 23.6% of these manifestations and had a high risk for a worse outcome [46]. As the previously cited study was meant to evaluate only neurological manifestation, a selection bias may have impacted results, since there is no other studies that found such high frequencies of coma in COVID-19 patients.

Overall, the present study allowed us to have a better understanding of neurological manifestations of COVID-19 according to age groups as well as differences of symptoms between the sexes. Even though this study has several strengths, such as its multicentric character, large

sample size, it is the largest Brazilian cohort that evaluated COVID-19 neurological manifestations, and one of the largest cohorts which assessed neurological symptoms worldwide, it has limitations. First, as this was an observational study based on data collected from patient charts, there was no formal checklist for neurological signs or symptoms. However, both the Brazilian Ministry of Health COVID-19 Guidelines and the Guidelines from the Brazilian Society of Infectology, Brazilian Society of Pneumology and Brazilian Medical Association recommended global assessment of COVID-19 patients [47], including formal assessment of neurological manifestations. Moreover, there were several on-line lectures and training in each hospital, to improve and harmonize patient assessment and treatment. Still, we cannot guarantee that the attending physician followed these recommendations, including a standardized assessment of neurological signs and symptoms. Standardized definitions were used for data collectors, who collected data from the medical records [48]. Another limitation is the fact that the current study evaluated neurological manifestations at hospital presentation only, and symptoms emerging later during the hospitalization were not registered for the purpose of the present analysis. At last, it is possible that older people and those who presented more severe neurological symptoms, such as delirium and coma, were not able to report milder symptoms, such as anosmia and ageusia. However, even excluding patients with delirium, coma and dementia from the regression analysis of the self-reported symptoms, younger age was still associated with higher risk of those symptoms.

## 5. Conclusion

In conclusion, elderly COVID-19 patients presented a higher prevalence of delirium, syncope and coma when compared to younger patients, and older age increased risk of delirium and coma. Younger patients were more likely to present less severe symptoms, such as anosmia, ageusia and headaches. Additionally, women were more likely to present with headache.

## Funding

The present study is supported in part by Minas Gerais State Agency for Research and Development (*Fundação de Amparo à Pesquisa do Estado de Minas Gerais - FAPEMIG*) [grant number APQ-01154-21], National Institute of Science and Technology for Health Technology Assessment (*Instituto de Avaliação de Tecnologias em Saúde - IATS*)/ National Council for Scientific and Technological Development (*Conselho Nacional de Desenvolvimento Científico e Tecnológico - CNPq*) [grant number 465518/2014–1], =. MM was supported in part by CNPq (grant 310561/2021–3). DNP was funded by a research scholarship from CNPq [grant number 147122/2021–0].

## Credit author statement

Data collection: DNP, MSM, MACB, AOJ, AGRG, AVS, ALHA, CCRC, DP, DRAR, GMSG, ERFM, FA, FGA, FB, JALB, JTT, KBR, PGP, SFA.

Manuscript idea: MSM, DNP, ALT.

Manuscript elaboration: DNP, MSM, MACB, AOJ, ALT.

Manuscript revision: all authors

## Acknowledgements

We would like to thank the collaborating hospitals that supported the Brazilian COVID Registry Project. We would also like to thank the clinical staff of all participating hospitals, who cared for the patients, and all undergraduate students that aided in data collection.

For this analysis we would like to thank the hospitals: Hospital Nossa Senhora da Conceição, Hospitais da Rede Mater Dei, Hospital Julia Kubitschek, Hospital Eduardo de Menezes, Hospital das Clínicas da UFMG, Hospital de Clínicas de Porto Alegre, Hospital Santo Antônio,



Hospital Tacchini, Hospital Márcio Cunha, Hospital Metropolitanou Doutor Célio de Castro, Hospital Metropolitanou Odilon Behrens, Hospital Risoleta Tolentino Neves, Hospital Santa Rosália, Complexo de Saúde São João de Deus, Hospital Semper, Hospital UNIMED, Hospital Mãe de Deus, Hospital Universitário Canoas, Hospital Universitário de Santa Maria, Hospital Moinhos de Vento, Instituto Mário Penna/ Hospital Luxemburgo, Hospital Nossa Senhora da Conceição, Hospital João XXIII, Hospital Regional Antônio Dias, Hospital Mãe de Deus, Hospital Regional do Oeste, Hospital das Clínicas da Faculdade de Medicina de Botucatu, Hospital das Clínicas da Universidade Federal de Pernambuco, Hospital Universitário Ciências Médicas, Hospital Bruno Born, Hospital SOS Córdio, Hospital São Lucas da PUCRS, Orizonti- Instituto de Saúde e Longevidade, Santa Casa de Misericórdia de Belo Horizonte.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ensci.2022.100419>.

## References

- [1] A. Johansson, M.S. Mohamed, T.C. Moulin, H.B. Schiöth, Neurological manifestations of COVID-19: a comprehensive literature review and discussion of mechanisms, *J. Neuroimmunol.* 358 (2021) 577–658, <https://doi.org/10.1016/j.jneuroim.2021.577658>.
- [2] A.A. Divani, S. Andalib, J. Biller, M. Di Napoli, N. Moghimi, C.A. Rubinos, C. O. Nobleza, P.N. Sylaja, M. Toledano, S. Lattanzi, L.D. McCullough, S. Cruz-Flores, M. Torbey, M.R. Azarpazhooh, Central nervous system manifestations associated with COVID-19, *Curr. Neurol. Neurosci. Rep.* 20 (12) (2020) 60–66, <https://doi.org/10.1007/s11910-020-01079-7>.
- [3] S. Andalib, J. Biller, M. Di Napoli, N. Moghimi, L.D. McCullough, C.A. Rubinos, C. O'Hana Nobleza, M.R. Azarpazhooh, L. Catanese, I. Elicer, M. Jafari, F. Liberati, C. Camejo, M. Torbey, A.A. Divani, Peripheral nervous system manifestations associated with COVID-19, *Curr. Neurol. Neurosci. Rep.* 21 (3) (2021) 9, <https://doi.org/10.1007/s11910-021-01102-5>.
- [4] M.S. Marcolino, F. Anschau, L. Kopitke, M.C. Pires, I.G. Barbosa, D.N. Pereira, et al., Research Square. Frequency and Burden of Neurological Manifestations in Hospitalized Patients with COVID-19: Findings from a Large Brazilian Cohort, 2022, <https://doi.org/10.21203/rs.3.rs-1231708/v1>. Jan 5.
- [5] S.H.Y. Chou, E. Beghi, R. Helbok, E. Moro, J. Sampson, V. Altamirano, S. Mainali, C. Bassetti, J.I. Suarez, M. McNett, GCS-NeuroCOVID Consortium and ENERGY Consortium, Global incidence of neurological manifestations among patients hospitalized with COVID-19: a report for the GCS-NeuroCOVID consortium and the ENERGY consortium, *JAMA Netw. Open* 4 (5) (2021) 112–131, <https://doi.org/10.1001/jamanetworkopen.2021.12131>.
- [6] T.C. Jones, G. Biele, B. Mühlemann, T. Veith, J. Schneider, J. Beheim-Schwarzbach, et al., Estimating infectiousness throughout Sars-Cov-2 infection course, *Science* 373 (2021) eabi5273, <https://doi.org/10.1126/science.abi5273>.
- [7] P.-I. Lee, Y.-L. Hu, P.-Y. Chen, Y.-C. Huang, P.-R. Hsueh, Are children less susceptible to COVID-19? *J. Microbiol. Immunol. Infect.* 53 (2020) 371–372, <https://doi.org/10.1016/j.jmii.2020.02.011>.
- [8] J.P. Dudley, N.T. Lee, Disparities in age-specific morbidity and mortality from Sars-Cov-2 in China and the Republic of Korea, *Clin. Infect. Dis.* 71 (2020) 863–865, <https://doi.org/10.1093/cid/ciaa354>.
- [9] K. Liu, Y. Chen, R. Lin, K. Han, Clinical features of COVID-19 in elderly patients: A comparison with young and middle-aged patients, *J. Infect.* 80 (2020) 14–18, <https://doi.org/10.1016/j.jinf.2020.03.005>.
- [10] J. Wang, X. Zhu, Z. Xu, G. Yang, G. Mao, Y. Jia, et al., Clinical and Ct findings of COVID-19: differences among three age groups, *BMC Infect. Dis.* 20 (2020) 1–11, <https://doi.org/10.1186/s12879-020-05154-9>.
- [11] X. Yang, Y. Yu, J. Xu, H. Shu, H. Liu, Y. Wu, et al., Clinical course and outcomes of critically ill patients with Sars-Cov-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study, *Lancet Respir. Med.* 8 (2020) 475–481, [https://doi.org/10.1016/S2213-2600\(20\)30079-5](https://doi.org/10.1016/S2213-2600(20)30079-5).
- [12] F. Zhou, T. Yu, R. Du, G. Fan, Y. Liu, Z. Liu, et al., Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study, *Lancet* 395 (2020) 1054–1062, [https://doi.org/10.1016/S0140-6736\(20\)30566-3](https://doi.org/10.1016/S0140-6736(20)30566-3).
- [13] C.C. Whitacre, S.C. Reingold, P.A. O'Looney, E. Blankenhorn, F. Brinley, E. Collier, et al., A gender gap in autoimmunity: task force on gender, multiple sclerosis and autoimmunity, *Science* 283 (1999) 1277–1278, <https://doi.org/10.1126/science.283.5406.1277>.
- [14] C.A. Roberts, M.E. Lewis, P. Boocock, Infectious disease, sex, and gender: the complexity of it all, *Sex Gender Paleopathol. Perspect.* (1998) 93–113.
- [15] R. Tolhurst, K. De Koning, J. Price, J. Kemp, S. Theobald, S. Squire, The challenge of infectious disease: time to take gender into account, *J. Health Manag.* 4 (2002) 135–151, <https://doi.org/10.1177/097206340200400204>.
- [16] R. Morgan, S.L. Klein, The intersection of sex and gender in the treatment of influenza, *Curr. Opin. Virol.* 35 (2019) 35–41, <https://doi.org/10.1016/j.coviro.2019.02.009>.
- [17] E.J. Williamson, A.J. Walker, K. Bhaskaran, S. Bacon, C. Bates, C.E. Morton, et al., Factors associated with COVID-19-related death using open-safely, *Nature* 584 (2020) 430–436, <https://doi.org/10.1038/s41586-020-2521-4>.
- [18] M. Shubham, K. Kavitha, P. Many, R. Divya, T.T. Kiran, S. Tom, et al., Frequency of neurologic manifestations in COVID-19: A systematic review and meta-analysis, *Neurology* 97 (23) (2021) e2269–e2281, <https://doi.org/10.1212/WNL.00000000000012930>. Dec.
- [19] M.S. Marcolino, P.K. Ziegelmann, M.V.R. Souza-Silva, I.J.B. Nascimento, L. M. Oliveira, L.S. Monteiro, Brazilian COVID-19 Registry Investigators, et al., Clinical characteristics and outcomes of patients hospitalized with COVID-19 in Brazil: results from the Brazilian COVID-19 registry, *Int. J. Infect. Dis.* 107 (2021) 300–310, <https://doi.org/10.1016/j.ijid.2021.01.019>. Jun.
- [20] World Health Organization, Diagnostic Testing for SARS-CoV-2. Interim Guidance, Online. 11 September. Available at, <https://www.who.int/publications/i/item/diagnostic-testing-for-sars-cov-2>, 2020.
- [21] P.A. Harris, R. Taylor, B.L. Minor, V. Elliott, M. Fernandez, L. O'Neal, et al., The REDCap consortium: building an international community of software platform partners, *J. Biomed. Inform.* 95 (2019), 103208.
- [22] M.S. Marcolino, M.C. Pires, L.E.F. Ramos, R.T. Silva, L.M. Oliveira, R.L. R. Carvalho, et al., ABC2-SPH risk score for in-hospital mortality in COVID-19 patients: development, external validation and comparison with other available scores, *Int. J. Infect. Dis.* 110 (2021) 281–308, <https://doi.org/10.1016/j.ijid.2021.07.049>.
- [23] Q. Bi, Y. Wu, S. Mei, C. Ye, et al., Epidemiology and transmission of COVID-19 in 391 cases and 1286 of their close contacts in Shenzhen, China: a retrospective cohort study, *Lancet Infect. Dis.* 20 (8) (2020) 911–919, 08.
- [24] F. Salamanna, M. Maglio, M.P. Landini, M. Fini, Body localization of ACE-2: on the trail of the keyhole of SARS-CoV-2, *Front. Med. (Lausanne)* 7 (2020), 594495.
- [25] R. Chen, K. Wang, J. Yu, D. Howard, et al., The spatial and cell-type distribution of SARS-CoV-2 receptor ACE2 in the human and mouse brains, *Front. Neurol.* 11 (2020), 573095.
- [26] S.F. Farhadian, D. Seilhean, S. Spudich, Neuropathogenesis of acute coronavirus disease 2019, *Curr. Opin. Neurol.* 34 (3) (2021) 417–422, 06 01.
- [27] D. Kumar, S. Jahan, A. Khan, A.J. Siddiqui, et al., Neurological manifestation of SARS-CoV-2 induced inflammation and possible therapeutic strategies against COVID-19, *Mol. Neurobiol.* 58 (7) (Mar 2021) 3417–3434.
- [28] J. Matschke, M. Lütgehetmann, C. Hagem, J.P. Sperhake, A.S. Schröder, C. Edler, H. Mushumba, A. Fitzek, L. Allweiss, M. Dandri, M. Dottermusch, A. Heinemann, S. Pfeiffer, M. Schwabenland, D. Sumner Magruder, S. Bonn, M. Prinz, C. Gerloff, K. Püschel, S. Krasemann, M. Aepfelbacher, M. Glatzel, Neuropathology of patients with COVID-19 in Germany: a post-mortem case series, *Lancet Neurol.* 19 (11) (2020) 919–929, [https://doi.org/10.1016/S1474-4422\(20\)30308-2](https://doi.org/10.1016/S1474-4422(20)30308-2).
- [29] C.M. Romero-Sánchez, I. Díaz-Maroto, E. Fernández-Díaz, Á. Sánchez-Larsen, A. Layos-Romero, J. García-García, E. González, I. Redondo-Peñas, A.B. Perona-Moratalla, J.A. Del Valle-Pérez, J. Gracia-Gil, L. Rojas-Bartolomé, I. Ferial-Vilar, M. Monteagudo, M. Palao, E. Palazón-García, C. Alcahut-Rodríguez, D. Sopenana-Garay, Y. Moreno, J. Ahmad, T. Segura, Neurologic manifestations in hospitalized patients with COVID-19: The ALBA COVID registry, *Neurology* 95 (8) (2020) e1060–e1070, <https://doi.org/10.1212/WNL.0000000000009937>.
- [30] S. Misra, K. Kolappa, M. Prasad, D. Radhakrishnan, K.T. Thakur, T. Solomon, B. D. Michael, A.S. Winkler, E. Beghi, A. Guekht, C.A. Pardo, G.K. Wood, S. Hsiang-Yi Chou, E.L. Fink, E. Schmutzhard, A. Kheradmand, F.K. Hoo, A. Kumar, A. Das, A. K. Srivastava, A. Agarwal, T. Dua, K. Prasad, Frequency of neurologic manifestations in COVID-19: A systematic review and meta-analysis, *Neurology* 97 (23) (2021) e2269–e2281, <https://doi.org/10.1212/WNL.00000000000012930>. Dec 7. (Epub 2021 Oct 11).
- [31] M.B. Zazzara, R.S. Penfold, A.L. Roberts, K.A. Lee, H. Dooley, C.H. Sudre, C. Welch, R.C.E. Bowyer, A. Visconti, M. Mangino, M.B. Freidin, J.S. El-Sayed Moustafa, K. S. Small, B. Murray, M. Modat, M.S. Graham, J. Wolf, S. Ourselin, F.C. Martin, C. J. Steves, M.N. Lochlainn, Probable delirium is a presenting symptom of COVID-19 in frail, older adults: a cohort study of 322 hospitalised and 535 community-based older adults, *Age Ageing* 50 (1) (2021) 40–48, <https://doi.org/10.1093/ageing/afaa223>. Jan 8.
- [32] K.J. Foster, E. Jauregui, B. Tajudeen, F. Bishehsari, M. Mahdavinia, Smell loss is a prognostic factor for lower severity of coronavirus disease 2019, *Ann. Allergy Asthma Immunol.* 125 (4) (2020) 481–483, <https://doi.org/10.1016/j.anaai.2020.07.023>.
- [33] J. Porta-Etessam, I.J. Núñez-Gil, N. González García, et al., COVID-19 anosmia and gustatory symptoms as a prognosis factor: a subanalysis of the HOPE COVID-19 (health outcome predictive evaluation for COVID-19) registry, *Infection* 49 (4) (2021) 677–684.
- [34] A.F. Temmel, C. Quint, B. Schickinger-Fischer, L. Klimek, E. Stoller, T. Hummel, Characteristics of olfactory disorders in relation to major causes of olfactory loss, *Arch. Otolaryngol. Head Neck Surg.* 128 (6) (2002) 635–641, <https://doi.org/10.1001/archotol.128.6.635>.
- [35] P. Rebora, R. Rozzini, A. Bianchetti, P. Blangiardo, A. Marchegiani, et al., Delirium in patients with SARS-CoV-2 infection: a multicenter study, *J. Am. Geriatr. Soc.* 69 (2) (2021) 293–299.
- [36] M. Ashina, Migraine, *N. Engl. J. Med.* 383 (19) (2020) 1866–1876, <https://doi.org/10.1056/NEJMr1915327>.
- [37] Z. Delaruelle, T.A. Ivanova, S. Khan, A. Negro, R. Ornello, B. Raffaelli, A. Terrin, D. D. Mitsikostas, U. Reuter, European Headache Federation School of Advanced Studies (EHF-SAS), Male and female sex hormones in primary headaches, *J. Headache Pain* 19 (1) (2018) 117, <https://doi.org/10.1186/s10194-018-0922-7>.
- [38] A. Biadsee, A. Biadsee, F. Kassem, O. Dagan, S. Masarwa, Z. Ormianer, Olfactory and oral manifestations of COVID-19: sex-related symptoms—a potential pathway to

- early diagnosis, *Otolaryngol. Head Neck Surg.* 163 (4) (2020) 722–728, <https://doi.org/10.1177/0194599820934380>.
- [39] D. García-Azorin, A. Layos-Romero, J. Porta-Etessam, J.A. Membrilla, E. Caronna, A. Gonzalez-Martinez, Á.S. Mencia, T. Segura, N. Gonzalez-García, J. Díaz-de-Terán, V.J. Gallardo, A.B. Gago-Veiga, A. Ballvé, J. Trigo López, M. Sastre-Real, A. Llauradó, A. Cornejo, Í. de Lorenzo, Á. Guerrero-Peral, P. Pozo-Rosich, Post-COVID-19 persistent headache: a multicentric 9-months follow-up study of 905 patients, *Cephalalgia*. 15 (2022 Feb) 804–809, <https://doi.org/10.1177/03331024211068074>.
- [40] J.W. Devlin, Y. Skrobik, C. Gelinas, D.M. Needham, A.J.C. Slooter, P.P. Pandharpande, P.L. Watson, G.L. Weinhouse, M.E. Nunnally, B. Rochweg, M.C. Balas, M. van den Boogaard, K.J. Bosma, N.E. Brummel, G. Chanques, L. Denehy, X. Drouot, G.L. Fraser, J.E. Harris, A.M. Joffe, M.E. Kho, J.P. Kress, J.A. Lanphere, S. McKinley, K.J. Neufeld, M.A. Pisani, J.F. Payen, B.T. Pun, K.A. Puntillo, R. Riker, B.R.H. Robinson, Y. Shehabi, P.M. Szumita, C. Winkelman, J. E. Centofanti, C. Price, S. Nikayin, C.J. Misak, P.D. Flood, K. Kiedrowski, W. Alhazzani, Clinical practice guidelines for the prevention and management of pain, agitation/sedation, delirium, immobility, and sleep disruption in adult patients in the ICU, *Crit. Care Med.* 46 (2018) e825–e873.
- [41] J.H. Han, J. Suyama, Delirium and Dementia, *Clin. Geriatr. Med.* 34 (3) (2018) 327–354, <https://doi.org/10.1016/j.cger.2018.05.001>. Aug. (PMID: 30031420).
- [42] S. Seshadri, P.A. Wolf, Lifetime risk of stroke and dementia: current concepts, and estimates from the Framingham study, *Lancet Neurol.* 6 (12) (2007) 1106–1114, [https://doi.org/10.1016/S1474-4422\(07\)70291-0](https://doi.org/10.1016/S1474-4422(07)70291-0). Dec. (PMID: 18031707).
- [43] C. Hasselgren, H. Ekbrand, B. Halleröd, et al., Sex differences in dementia: on the potentially mediating effects of educational attainment and experiences of psychological distress, *BMC Psychiatry* 20 (2020) 434, <https://doi.org/10.1186/s12888-020-02820-9>.
- [44] F. Arias, M. Alegria, A.J. Kind, R.N. Jones, T.G. Trivison, E.R. Marcantonio, E. M. Schmitt, T.G. Fong, S.K. Inouye, A framework of social determinants of health for delirium tailored to older adults, *J. Am. Geriatr. Soc.* 70 (1) (2022) 235–242, <https://doi.org/10.1111/jgs.17465>. Jan.
- [45] T.T. Wu, M. Zegers, R. Kookan, J.L. Griffith, B.E. Molnar, J.W. Devlin, M. van den Boogaard, Social determinants of health and delirium occurrence and duration in critically ill adults, *Crit. Care Explor.* 3 (9) (2021) 05–32, <https://doi.org/10.1097/CCE.0000000000000532>.
- [46] F.B. Garcez, M.J.R. Aliberti, P.C.E. Poco, M. Hiratsuka, S.F. Takahashi, V. A. Coelho, et al., Delirium and adverse outcomes in hospitalized patients with COVID-19, *J. Am. Geriatr. Soc.* 68 (11) (2020) 2440–2446.
- [47] Ministério da Saúde, Diretrizes da Atenção Especializada no Contexto da Pandemia de COVID-19, Gov. Brasil, 2021.
- [48] Associação Médica Brasileira, Sociedade Brasileira de Infectologia, Sociedade Brasileira de Pneumologia e Tisiologia, Manejo Pré-Hospitalar da COVID-19 (Prevenção e Tratamento de Pacientes com Sintomas Leves), *A. Med. Brasileira* (2021) 01–47.