

CASE REPORT

Open Access



Hypokalaemic paralysis and metabolic alkalosis in a patient with Sjögren syndrome: a case report and literature review

Rasika Ranaweerage*, Shehan Perera and Aruna Gunapala

Abstract

Background: Acquired Gitelman syndrome is a very rare disorder reported in association with autoimmune disorders, mostly Sjögren syndrome. It is characterized by the presence of hypokalaemic metabolic alkalosis, hypocalciuria, hypomagnesaemia and hyper-reninaemia, in the absence of typical genetic mutations associated with inherited Gitelman syndrome.

Case presentation: A 20 year old woman who was previously diagnosed with primary Sjögren syndrome and autoimmune thyroiditis presented with two week history of lower limb weakness and salt craving. Examination revealed upper limb and lower limb muscle weakness with muscle power of 3/5 on MRC scale and diminished deep tendon reflexes. On evaluation, she had hypokalaemia with high trans-tubular potassium gradient, metabolic alkalosis and hypocalciuria, features suggestive of Gitelman syndrome. New onset hypokalaemic alkalosis in a previously normokalaemic patient with Sjögren syndrome strongly favored a diagnosis of acquired Gitelman syndrome. Daily potassium supplementation and spironolactone resulted in complete clinical recovery.

Conclusions: Acquired Gitelman syndrome associated with Sjögren syndrome is rare. It should be considered as a differential diagnosis during evaluation of acute paralysis and hypokalaemic metabolic alkalosis in patients with autoimmune disorders, especially Sjögren syndrome.

Keywords: Acquired Gitelman syndrome, Sjögren syndrome, Hypokalaemia, Metabolic alkalosis

Background

Sjögren syndrome is a systemic autoimmune disease primarily affecting the exocrine glands. Renal involvement in Sjögren syndrome usually manifest as chronic interstitial nephritis, distal renal tubular acidosis and nephrogenic diabetes insipidus. Sjögren syndrome presenting with features of Gitelman syndrome is rare and named as acquired Gitelman syndrome. In medical literature only eight such cases have been reported. Awareness of the treating physician regarding this rare clinical

presentation is imperative to avoid fatal outcomes, especially in patients presenting with acute paralysis. Here, we describe a young female with Sjögren syndrome presenting with features of Gitelman syndrome and hypokalaemic paralysis.

Case presentation

A 20 year old woman previously diagnosed with primary Sjögren syndrome and autoimmune thyroiditis with hypothyroidism presented with two week history of muscle cramps, gradual onset, progressive lower limb weakness and salt craving. The weakness was not

* Correspondence: rranaweerage@gmail.com

General medicine, National Hospital of Sri Lanka, Colombo, Sri Lanka



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

associated with ingestion of high carbohydrate meals, diurnal variation of symptoms, easy fatigability or other neurological symptoms including diplopia, bulbar involvement and paresthesia. There was no preceding respiratory tract infection or gastroenteritis. She was on Thyroxin 50 µg daily and denied the use of diuretics or laxatives. There was no personal or family history of similar episodes. Her blood pressure was 100/70 mmHg. Pulse rate was 78 bpm. She was not volume depleted. There was no apparent wasting or tenderness of muscles. Muscle power was 3/5 on MRC scale. Deep tendon reflexes were diminished. Rest of the clinical examination was normal.

Her investigation results revealed that she was having hypokalaemia. ECG revealed T wave flattening and U waves. Imaging did not reveal nephrocalcinosis. Her investigation results are summarized in Table 1.

Previous records revealed that she was diagnosed with primary Sjögren syndrome and autoimmune thyroiditis 2 years ago, when she presented with salivary gland swelling and sicca symptoms. She did not have any other symptoms and signs suggestive of connective tissue disorders. Salivary gland biopsy showed periductal and periacinar lymphoplasmacytic infiltrate that was

compatible with Sjögren syndrome. Rheumatoid factor, anti-Ro (SSA) antibody, anti TPO antibody and Thyroglobulin antibody were positive. ANA titer was positive at 1:320. At that time, serum potassium was noted to be within normal parameters.

Presence of symptomatic hypokalemia with high Trans-tubular Potassium Gradient (TTKG) was suggestive of renal loss of potassium. Coexisting metabolic alkalosis, absence of hypercalciuria, increased renin activity with normal blood pressure in the absence of diuretic or laxative use favored a diagnosis of Gitelman syndrome, either inherited or acquired. Previous records revealed persistent normokalaemia prior to this presentation. New onset hypokalaemic alkalosis in a previously normokalaemic patient with Sjögren syndrome (which is frequently associated with the renal tubular dysfunctions) and autoimmune thyroiditis strongly favored acquired Gitelman syndrome of immune origin. Genetic studies are crucial in differentiating between the inherited and acquired forms of Gitelman syndrome, but we did not have access to genetic studies to confirm our diagnosis.

She was treated with intravenous and oral potassium salts which resulted in improvement of symptoms. At discharge, maintenance potassium salts and spironolactone were prescribed which she tolerated well. Two months later she had a relapse due to poor compliance. After ensuring the compliance she had no further relapses and remained symptoms free thereafter.

Table 1 Summary of laboratory investigations

Investigation	Result	Normal range
Haemoglobin	12 g/dL	12.0–15.5 g/dL
Platelets	200,000 per microl	150–450 per microl
Serum investigations		
Sodium	135 mmol/L	135–145 mmol/L
Potassium	2.5 mmol/L	3.5–5 mmol/L
Magnesium	1.8 mg/dL	1.7–2.30 mg/dL
Chloride	91 mmol/L	95–111 mmol/L
Creatinine	0.8 mg/dL	0.6–1.2 mg/dL
Arterial pH	7.5	7.35–7.45
PaCO ₂	43 mmHg	38–42 mmHg
Bicarbonate	28 mmol/L	22–28 mmol/L
Renin activity	60.6 ng/mL/h	5.4–34.5 ng/mL/h
TSH	3.872 mIU/L	0.5 to 5.0 mIU/L
Free T ₄	1.29 ng/dL	0.9–2.3 ng/dL
9 a.m. Serum cortisol	550 nmol/L	140 to 690 nmol/L
Urinary investigations		
Urinary potassium	40 mmol/L	< 20 mmol/L
Urinary chloride	64 mmol/L	55–125 mmol/L
Trans-Tubular Potassium Gradient (TTKG)	13	< 3
Spot urine potassium/creatinine ratio	3.1 mmol/mmol	> 2 mmol/mmol
Spot urine calcium/creatinine ratio	0.01 g calcium /g creatinine	< 0.2 g calcium /g creatinine

Discussion and conclusion

Primary Sjögren syndrome is a systemic autoimmune disease characterized by lymphoplasmacytic infiltration of lacrimal and salivary glands and multiple extraglandular tissues including skin and joints, lungs, heart, gastrointestinal tract, kidneys, bladder, gynecological system and nervous system [1, 2]. It is commonly seen in females with typical onset in fourth to fifth decade of life. The diagnosis of Sjögren syndrome is made in the presence of typical clinical signs, serological and/or histological evidence after the exclusion of conditions that may manifest similar to Sjögren syndrome [3].

Renal involvement is a frequent manifestation of Sjögren syndrome. It is rarely overt and may precede onset sicca symptoms [4]. The manifestations vary from chronic interstitial nephritis, distal tubular acidosis, and nephrogenic diabetes insipidus to symptomatic hypokalaemia due to tubular injury [4, 5]. The commonest presentation is chronic interstitial nephritis [6]. Distal renal tubular acidosis (RTA) may occur in up to 25 % of the patients with Sjögren syndrome. It is characterized by normal anion gap metabolic acidosis and urinary potassium wasting leading to hypokalaemia muscle paralysis with respiratory arrest which has been the presenting

symptom in some cases of Sjögren syndrome. Glomerular involvement is noted to be rare and carries high mortality [4].

Gitelman syndrome is a renal tubular disorder characterized by hypokalemia, metabolic alkalosis, hypomagnesaemia, hypocalciuria, and hyper-reninaemia with normal blood pressure. Gitelman syndrome is an autosomal recessively inherited genetic disorder caused due to the biallelic inactivating mutations of SLC12A3 gene encoding for thiazide-sensitive Sodium chloride cotransporter (NCCT) in distal tubule of the kidney. Gitelman syndrome is usually not diagnosed until late childhood or adulthood [7]. More common disorders that have a similar presentation and must be excluded are surreptitious vomiting, surreptitious diuretic use and hypokalaemic periodic paralysis.

A rare, phenotypically similar clinical presentation has been described in patients with autoimmune disorders such as Sjögren syndrome, Systemic lupus erythematosus and Systemic sclerosis, which has been named as “acquired Gitelman syndrome” [8–11]. There are only eight such cases have been reported in patients with primary Sjögren syndrome and three of these patients presented with hypokalemic paralysis [12–15]. It should be considered as a differential diagnosis during evaluation of hypokalaemic metabolic alkalosis in patients with autoimmune disorders, especially Sjögren syndrome.

Pathogenesis of acquired Gitelman syndrome associated with Sjögren syndrome is being evaluated. Kim et al. reported the presence of a circulating autoantibody against NCCT in a patient presenting with acquired Gitelman syndrome secondary to Sjögren syndrome. This was demonstrated by incubation of serum of a patient with acquired Gitelman syndrome with normal mouse kidney tissue, which showed similar staining patterns of NCCT of distal tubules compared to the incubation of normal mouse kidney with rabbit polyclonal anti-NCCT antibody [8, 14]. Genetic studies are crucial in differentiating between the genetic and the acquired form of Gitelman syndrome. In comparison, type 1 renal tubular acidosis (RTA) secondary to Sjögren syndrome was found to be associated with tubular interstitial nephritis with lymphocytic infiltration and reduced expression of both anion exchanger 1 (AE1) and hydrogen-ATPase located in the α -intercalated cells in absence of autoantibodies against these transporters [16]. In recent studies, a subset of patients with RTA with anti-carbonic anhydrase antibodies has been identified [17]. These antibodies are thought to be contributory to the pathogenesis of RTA in Sjögren syndrome.

Hypomagnesaemia is a characteristic manifestation of classic Gitelman syndrome, but magnesium level can even be normal in some patients [7]. The mechanism of hypomagnesaemia has been attributed to down

regulation TRPM6 magnesium-permeable channels that are located at the apical domain of the distal convoluted tubules and brush border of the duodenal magnesium-transporter cells [18]. In our patient magnesium level was normal.

Our patient demonstrated a very good clinical response to potassium supplementation, but she needed very high doses of potassium chloride to maintain normokalaemia. Similar clinical responses to potassium and/or magnesium and/or spironolactone was noted among other cases as well. In five case reports, steroids were administered with above treatment [13–15]. Four patients demonstrated reduced renal potassium wasting while one patient continued to have renal potassium wasting while on steroids. One patient was given cyclophosphamide [19]. None of these patients had relapses while on treatment. Three patients who had apparently responded to steroids were also documented to be on spironolactone at the same time. Hence the observed reduction of renal potassium wasting cannot be attributed to action of steroids alone.

In comparison, the effects of immunosuppression with steroids in type 1 renal tubular acidosis secondary to Sjögren syndrome has been extensively evaluated [20]. Immunosuppression with steroids has been found to effective in slowing the progression of renal disease in majority of these patients. It is our opinion that use of steroids in management of acquired Gitelman syndrome in Sjögren syndrome needs further evaluation as a potential therapeutic modality. Presence of lymphoplasmacytic cell infiltration in renal biopsy may also assist in deciding on immunosuppressive therapy.

Further studies focused on the pathogenesis of renal tubular dysfunctions in Sjögren syndrome will be immensely beneficial in establishing the prognosis and effective therapeutic interventions in this rare and possibly underdiagnosed manifestation of Sjögren syndrome. Paucity of diagnosed patients with acquired Gitelman syndrome appears to be a major limiting factor in studying pathogenesis and treatment responsiveness. Establishing a global disease registry and long term follow up of the diagnosed patients may help to alleviate this issue.

Abbreviations

bpm: beats per minute; MRC scale: Medical research council scale; NCCT: Sodium chloride co-transporter; SLC12A3: Solute carrier family 12 member 3; TRPM6: Transient receptor potential melastatin 6; RTA: Renal tubular acidosis

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12882-021-02371-5>.

Additional file 1.

Acknowledgements

The authors would like to acknowledge all the staff of ward 46, National Hospital, Sri Lanka for the assistance provided in investigation and management of this patient.

Authors' contributions

All authors were involved in the management of the patient. RR did the initial literature review and wrote the first draft. SP also reviewed literature and corrected the first draft. AG did the final correction before submission. All authors have read and approved the final manuscript.

Funding

Authors did not receive any funding during the management of this patient or in writing this case report.

Availability of data and materials

All necessary data and material are provided.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Informed written consent was taken from the patient to publish the details relevant to the disease for this scientific publication. The copy of consent form is available for review if deemed necessary.

Competing interests

The authors declare that they do not have any competing interests.

Received: 11 December 2020 Accepted: 22 April 2021

Published online: 30 April 2021

References

- Ramos-Casals M, Tzioufas AG, Font J. Primary Sjögren's syndrome: new clinical and therapeutic concepts. *Annals of the rheumatic diseases*. 2005 Mar 1;64(3):347–54.
- ASMUSSEN K, ANDERSEN V, BENDIXEN G, SCHIØDT M, OXHOLM P. A new model for classification of disease manifestations in primary Sjögren's syndrome: evaluation in a retrospective long-term study. *Journal of internal medicine*. 1996 Jun;239(6):475–82.
- Shiboski CH, Shiboski SC, Seror R, Criswell LA, Labetoulle M, Lietman TM, Rasmussen A, Scofield H, Vitali C, Bowman SJ, Mariette X. 2016 ACR-EULAR classification criteria for primary Sjögren's syndrome: a consensus and data-driven methodology involving three international patient cohorts. *Arthritis & rheumatology* (Hoboken). 2017 Jan;69(1):p. 35.
- Goules A, Masouridi S, Tzioufas AG, Ioannidis JP, Skopouli FN, Moutsopoulos HM. Clinically significant and biopsy-documented renal involvement in primary Sjögren syndrome. *Medicine*. 2000 Jul 1;79(4):241–9.
- Nagayama Y, Shigeno M, Nakagawa Y, Suganuma A, Takeshita A, Fujiyama K, Ashizawa K, Kiriya T, Yokoyama N, Nagataki S. Acquired nephrogenic diabetes insipidus secondary to distal renal tubular acidosis and nephrocalcinosis associated with Sjögren's syndrome. *J Endocrinol Investig*. 1994 Sep;17(8):659–63.
- Maripuri S, Grande JP, Osborn TG, Fervenza FC, Matteson EL, Donadio JV, Hogan MC. Renal involvement in primary Sjögren's syndrome: a clinicopathologic study. *Clinical Journal of the American Society of Nephrology*. 2009 Sep 1;4(9):1423–31.
- Blanchard A, Bockenauer D, Bolignano D, Calo LA, Cosyns E, Devuyst O, Ellison DH, Frankl FE, Knoers NV, Konrad M, Lin SH. Gitelman syndrome: consensus and guidance from a kidney disease: improving global outcomes (KDIGO) controversies conference. *Kidney international*. 2017 Jan 1;91(1):24–33.
- Kim YK, Song HC, Kim YS, Choi EJ. Acquired gitelman syndrome. *Electrolyte Blood Press*. 2009 Jun;7(1):5–8.
- Barathidasan GS, Krishnamurthy S, Karunakar P, Rajendran R, Ramya K, Dhandapany G, Ramamoorthy JG, Ganesh RN. Systemic lupus erythematosus complicated by a Gitelman-like syndrome in an 8-year-old girl. *CEN case reports*. 2019 Dec 18:1–4.
- Masab M, Goyal A, Abrol S, Rangaswami J. Acquired Gitelman Syndrome associated with systemic sclerosis. *Cureus*. 2019 Jan;11(1).
- Casatta L, Ferraccioli GF, Bartoli E. Hypokalaemic alkalosis, acquired Gitelman's and Bartter's syndrome in chronic sialoadenitis. *British journal of rheumatology*. 1997 Oct 1;36(10):1125–8.
- Chen YC, Yang WC, Yang AH, Lin SH, Li HY, Lin CC. Primary Sjögren's syndrome associated with Gitelman's syndrome presenting with muscular paralysis. *American journal of kidney diseases*. 2003 Sep 1;42(3):586–90.
- Schwarz C, Barisani T, Bauer E, Druml W. A woman with red eyes and hypokalemia: a case of acquired Gitelman syndrome. *Wiener klinische Wochenschrift*. 2006 May;118(7):239–42.
- Kim YK, Song HC, Kim WY, Yoon HE, Choi YJ, Ki CS, Park CW, Yang CW, Kim J, Kim YS, Choi EJ. Acquired Gitelman syndrome in a patient with primary Sjögren syndrome. *American Journal of Kidney Diseases*. 2008 Dec 1;52(6):1163–7.
- Kulkarni M, Kadri P, Pinto R. A case of acquired Gitelman syndrome presenting as hypokalemic paralysis. *Indian J Nephrol*. 2015 Jul 1;25(4):246–7.
- Walsh S, Turner CM, Toye A, Wagner C, Jaeger P, Laing C, Unwin R. Immunohistochemical comparison of a case of inherited distal renal tubular acidosis (with a unique AE1 mutation) with an acquired case secondary to autoimmune disease. *Nephrology Dialysis Transplantation*. 2007 Mar 1;22(3):807–12.
- Pertovaara M, Bootorabi F, Kuuslahti M, Pasternack A, Parkkila S. Novel carbonic anhydrase autoantibodies and renal manifestations in patients with primary Sjögren's syndrome. *Rheumatology*. 2011 Aug 1;50(8):1453–7.
- McCormick JA, Ellison DH. Distal convoluted tubule. *Comprehensive Physiology*. 2011 Jan 17;5(1):45–98.
- Gu X, Su Z, Chen M, Xu Y, Wang Y. Acquired Gitelman syndrome in a primary Sjögren syndrome patient with a SLC12A3 heterozygous mutation: a case report and literature review. *Nephrology*. 2017 Aug;22(8):652–5.
- Kaufman I, Schwartz D, Caspi D, Paran D. Sjögren's syndrome—not just Sicca: renal involvement in Sjögren's syndrome. *Scandinavian journal of rheumatology*. 2008 Jan 1;37(3):213–8.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

