

Review Article

Camel semen collection, viscosity, and cryopreservation: a review

Niasari-Naslaji, A.

Department of Theriogenology, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran

Correspondence: A. Niasari-Naslaji, Department of Theriogenology, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran. E-mail: niasari@ut.ac.ir



10.22099/IJVR.2023.44924.6624

(Received 1 Oct 2022; revised version 19 Dec 2022; accepted 2 Jan 2023)

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Abstract

Artificial insemination in camels remains undeveloped due to the difficulties in semen collection, semen viscosity, and semen cryopreservation. The semen collection procedure has been facilitated to some extent using camel phantom and/or possibly an intravaginal condom. Main reasons for semen viscosity in camelids have been unraveled and different mechanical and enzymatic approaches were used to alleviate this problem; however, there is still no conclusive protocol to safely remove semen viscosity completely. It seems that along with the problem of semen viscosity, semen cryopreservation in camels remains unresolved. As a result, there is no convincing report on successful and repeatable pregnancies following insemination with frozen semen in camel. This review gathered most of the information that appeared in the peer reviewed journals to highlight major problems in camel semen technology, including semen collection, semen viscosity, and semen cryopreservation.

Key words: Camel semen, semen collection, semen viscosity, sperm cryopreservation

Introduction

There is a great lag in the development of reproductive technologies in camels compared to other domestic animals. Semen technology, as the first generation of reproductive technologies, remained behind other generations of reproductive technologies like *in vitro* and *in vivo* production of embryos and cloning in camel. While AI with fresh and frozen semen is widely used in domestic animals, the progress of AI in camels is very slow. There is not a single camel bull stud in the world to collect and process semen for routine AI and cryopreservation successfully. Development of an artificial insemination network in camel depends on the ability:

- To collect clean and viable semen safely and easily
- To remove viscosity of semen without affecting sperm viability
- To extend the non-viscous semen in a suitable extender for performing artificial insemination and/or semen cryopreservation

The present review has been prepared to elaborate on the current status of semen technology with respect to semen collection, viscosity, and cryopreservation in camel.

Semen collection

The first step in developing AI in camels is the ability to collect clean and viable semen. Semen collection in camel is a challenging procedure due to the mating in sternal recumbency, prolonged time of copulation, and possible injuries to the operator (Tibary and Anouassi, 1997). There are four main approaches for collecting semen in camels, including artificial vagina (Tibary and Anouassi, 1997; Mosafari *et al.*, 2005), electroejaculation (Tingari *et al.*, 1986), phantom or dummy (Ziapour *et al.*, 2014), and intravaginal condoms (Tibary and Anouassi, 2018; Mansour, 2022). These methods have advantages and disadvantages among which semen quality and animal welfare are the main challenges. Artificial vagina (AV) is the most common approach for semen collection in camels (Skidmore *et al.*, 2020). Although AV simulates natural mating, the corresponding procedure in camel is tiring and hazardous to the operator mainly due to the prolonged time of semen collection and the position required collecting semen. More importantly, due to the several backward and forward movements of a bull during semen collection, the specimen is more likely to be contaminated (Ziapour *et al.*, 2014). Electroejaculation is

another approach for collecting semen from camel bulls (Tingari *et al.*, 1986). This method is not recommended for routine semen collection from valuable males. The procedure requires sedation or even anesthesia, and potentiates some risks to the life and welfare of the animal. Moreover, the quality of semen collected by an electroejaculator varies in volume and concentration (Tibary and Anouassi, 2018). Phantom could be a suitable replacement for live female camels for semen collection. Phantom eliminates the risk for the operator, facilitates semen collection procedure, and provides more natural conditions to collect good quality and clean specimens (Ziapour *et al.*, 2014; Panahi *et al.*, 2017). The main point in using phantom for semen collection in camels is the patience required for training the bull. Once a single bull is trained, it is possible to use this bull to stimulate others. Several tactics could be taken to train the bull to accept the camel dummy. To stimulate the bull camels, complete isolation of them from she-camels throughout the season, spraying urine of estrous she-camel around the back and perineal region of the camel dummy, and playing the recorded sound of she-camel during mating could be helpful. Recently, using intravaginal condoms was revisited for semen collection in camels (Tibary and Anouassi, 2018; Mansour, 2022). Major constraints in using condoms for semen collection in camels include problems associated with the installation of the device inside the vagina and its fixation around the vulva, and removing the installed condom due to clockwise and anti-clockwise movement of the penis. Moreover, due to causing the unpleasant condition for she-camel, using the condom device does not seem to receive animal ethics approval.

Semen viscosity

Camel semen is highly viscous by nature (Deen *et al.*, 2003), with mass vibration at the initial raw semen assessment (Panahi *et al.*, 2017) and prolonged liquefaction time of 18-41 h (Mal *et al.*, 2016). Proteins belonging to the β -nerve growth factors family might be responsible for the liquefaction of camel semen (Mal *et al.*, 2016). The viscous nature of the ejaculates makes the processing and pipetting of raw or even diluted semen very difficult. Besides, semen cryopreservation becomes unsuccessful because the viscous seminal plasma prevents penetration of cryoprotectant into the spermatozoa. The role of camel semen viscosity is not fully understood. It is postulated that semen viscosity might be required to prevent loss of sperm from the female reproductive tract (Kershaw-Young and Maxwell, 2012). Regardless, semen processing, evaluation, cryopreservation, and even AI require complete liquefaction. The causes of semen viscosity and its elimination have been a challenging subject in the world of camelid research. During the early sixties, it was suggested that mucopolysaccharide secreted by bulbourethral gland might be responsible for camel semen viscosity (Perk, 1962). However, recent studies suggested that these chemicals, renamed

glycosaminoglycans, may not be the main source of semen viscosity in camelids (Kershaw-Young and Maxwell, 2012; Kershaw-Young *et al.*, 2013). There is sufficient evidence to accept that proteins within seminal plasma such as mucin may be responsible for viscosity in camelid semen (Kershaw-Young and Maxwell, 2012). The viscosity of semen can be reduced by adding cysteine protease, such as papain, present in papaya (Kershaw-Young *et al.*, 2013, 2017; Monaco *et al.*, 2016; El-Bahrawy *et al.*, 2017) or ficin, present in fig (Keshavarz *et al.*, 2016) to semen. Following ficin treatment and centrifugation, round pellet forms at the bottom of the conical tube in association with the ability to separate supernatant from the pellet. However, in an untreated-centrifugated specimen, an oblique and sticky pellet forms at the wall of the tube and is removed with the supernatant during withdrawal (Keshavarz *et al.*, 2016). The positive and negative effects of enzyme treatment on semen occur fairly quickly. In fact, the amount of time required to remove semen viscosity by the enzyme is fairly similar to the time that the adverse effect on sperm viability occurs by the enzyme. Attempts have been conducted to neutralize the enzyme following partial semen liquefaction (Kershaw-Young *et al.*, 2017; Malo *et al.*, 2017b); however, there is no convincing so far report on the benefit of such treatment for semen cryopreservation and successful pregnancy following AI.

Apart from the enzymatic approach to improving the rheological characteristics of camel semen, mechanical and ultrasonic approaches were also investigated. Stirring of camel semen at a very low speed (150 rpm) for 15 min (Mosafari *et al.*, 2005), gentle pipetting of diluted semen (Morton *et al.*, 2008), and passage of semen back and forth through a needle could also be used to reduce semen viscosity (Santiani *et al.*, 2005). Ultrasound wave (40 kHz) was also imposed to Dromedary camel semen for 2 min, interspersed for 2 min, and repeated 4 times to reduce semen viscosity (Rateb, 2016; El-Bahrawy *et al.*, 2017). However, the safety and feasibility of such a method were not confirmed in other studies. While all of these methods could be simple and relatively effective, they do not completely eliminate semen viscosity. It is suggested that camel sperm could be extracted from seminal plasma without enzymatic extraction using a combination of pipetting and colloid single layer centrifugation without detrimental effect on sperm quality (Malo *et al.*, 2017a, 2018a; Morrell *et al.*, 2021). This method could be helpful to achieve a low sample of good quality sperm for *in vitro* embryo production; however, it did not provide a promising result in terms of semen cryopreservation.

Semen cryopreservation

The progress in camel semen cryopreservation has been slow compared to other livestock species. This, in turn, prevents extensive use of artificial insemination in camels. To date, there is only one report on pregnancy following AI with frozen-thawed semen in dromedary

camel (Deen *et al.*, 2003) and some reports in Bactrian camel (Chen *et al.*, 1984, 1985; Zhao *et al.*, 1994, 1996). There are no other reports on successful pregnancy in camels. Deen *et al.* (2003) reported a pregnancy rate of 0/10, 1/13, and 4/10 following insemination with liquid semen, frozen-thawed semen, and whole semen in dromedary camel, respectively. The later investigators considered 20% post-thaw motility as successful cryopreservation; whereas, it is commonly accepted that 40% post-thawed progressively motile sperm could be considered as a minimum standard for frozen semen to be used in extensive AI programs.

If a clean, viable, and non-viscous semen sample is an initial step, a suitable extender is the next step toward the development of extensive AI programs. There are few suitable and chemically defined extenders for semen preservation in camel. Lactose and sucrose have been used to preserve Bactrian camel semen (Chen *et al.*, 1984, 1985; Zhao *et al.*, 1994, 1996; Niasari-Naslaji *et al.*, 2006a, b, 2007a). These two extenders were not efficient even for chilled storage of Bactrian camel semen (Niasari-Naslaji *et al.*, 2006a, b, 2007a); however, they were reported to be good for cryopreservation of Bactrian camel semen, resulting in pregnancy rates of 86-100% following insemination with frozen semen (Chen *et al.*, 1984, 1985; Zhao *et al.*, 1994, 1996). Laiciphos, Androhep, and glucose-EDTA have also been investigated for the preservation of camel semen (Sieme *et al.*, 1990), with no exact quantification of the sperm viability parameters. Tris extender was suggested to be better than lactose (Vyas *et al.*, 1998) and Bicephos (Deen *et al.*, 2004) for the chilled storage of camel semen. Green buffer (Niasari-Naslaji *et al.*, 2006a; Skidmore *et al.*, 2013) and INRA 96 (Morton *et al.*, 2013; Malo *et al.*, 2020) were successfully used to preserve camel semen. Chemically defined tris-based extender named "SHOTOR" diluent was used successfully to store Bactrian camel semen in chilled and frozen states (Niasari-Naslaji *et al.*, 2006a, 2007b, 2008). However, the post-thaw motility of Bactrian camel sperm using SHOTOR diluent did not reach 40% to produce a successful pregnancy. We found that, SHOTOR diluent was not a proper extender to preserve Dromedary camel semen. Therefore, we introduced new extender named "HASHI" diluent for chilled storage of Dromedary camel semen (Panahi *et al.*, 2017). HASHI diluent consists of 60% SHOTOR diluent, 20% pigeon plasma egg yolk, and 20% camel skim milk (Panahi *et al.*, 2017).

Several experiments were conducted to determine the type (Malo *et al.*, 2017b) and concentration (Niasari-Naslaji *et al.*, 2007b; Malo *et al.*, 2017b) of cryoprotectants, dilution rate (Malo *et al.*, 2017a), equilibration time (Malo *et al.*, 2017b), the speed of cooling from room to 4°C (Niasari-Naslaji *et al.*, 2007b), freezing rate (Malo *et al.*, 2018b), thawing rates (Malo *et al.*, 2018b), addition of antioxidants (Medan *et al.*, 2008; Malo *et al.*, 2019, 2020), and surfactants (Niasari-Naslaji *et al.*, 2008) to cryopreserve camel semen. Unfortunately, the main obstacle to get meaningful and

conclusive results in the majority of studies was the inability to safely remove the viscosity of camel semen completely. Moreover, in several studies, the investigators reported the total motility and or very low progressive forward motility of sperm, which is not a good indicator of successful cryopreservation or successful AI using frozen semen.

Conclusion

Although there is considerable progress toward collecting good quality semen in camel using phantom or intravaginal condom and providing a sound extender to preserve liquid stored camel semen, to date, to the best of my knowledge, there is no solid, convincing, and repeatable result on the production of healthy and live birth of camel calf following insemination with frozen-thawed semen. It seems that still the main constraint preventing successful semen cryopreservation in camels is semen viscosity. Therefore, several investigations with conclusive outcomes are required to solve the problem of semen viscosity prior to further attempts for cryopreservation of camel semen.

Acknowledgement

The author wishes to dedicate this review article to his mentor, Dr Lulu Skidmore, the scientific director of Camel Reproduction Center, Dubai, UAE, whom I learned the first steps in understanding the reproduction of camel in 1999. Since then, we became close friend. I wish her a very happy, prosperous life.

References

- Chen, BX; Yuen, ZX and Pan, CW (1984). Factors inducing ovulation in the Bactrian camel. In: Cockrill, WR (Ed.), *The Camelid and all-purpose animal*. Uppsala, Sweden, Scandinavian Institute for African Studies. PP: 387-398.
- Chen, BX; Yuen, ZX and Pan, GW (1985). Semen induced ovulation in the Bactrian camel (*Camelus bactrianus*). *J. Reprod. Fertil.*, 74: 335-339.
- Deen, A; Vyas, S; Jain, M and Sahani, MS (2004). Refrigeratory preservation of camel semen. *J. Camel Pract. Res.*, 11: 137-139.
- Deen, A; Vyas, S and Sahani, MS (2003). Semen collection, cryopreservation and artificial insemination in the dromedary camel. *Anim. Reprod. Sci.*, 77: 223-233.
- El-Bahrawy, K; Rateb, S; Khalifa, M; Monaco, D and Lacalandra, G (2017). Physical and kinematic properties of cryopreserved camel sperm after elimination of semen viscosity by different techniques. *Anim. Reprod. Sci.*, 187: 100-108.
- Kershaw-Young, CM; Evans, G; Rodney, R and Maxwell, WMC (2017). Papain and its inhibitor E-64 reduce camelid semen viscosity without impairing sperm function and improve post-thaw motility rates. *Reprod. Fertil. Dev.*, 29: 1107-1114.
- Kershaw-Young, CM and Maxwell, WMC (2012). Seminal plasma components in camelids and comparisons with other species. *Reprod. Domest. Anim.*, (Suppl. 4), 47: 369-375.

- Kershaw-Young, CM; Stuart, C; Evans, G and Maxwell, WMC** (2013). The effect of glycosaminoglycan enzymes and proteases on the viscosity of alpaca seminal plasma and sperm function. *Anim. Reprod. Sci.*, 138: 261-267.
- Keshavarz, M; Niasari-Naslaji, A; Zare, H; Ziapour, S; Mirtavoosi, M; Omid, M; Kalantari, A and Moosavi-Movahedi, AA** (2016). Effect of Ficin enzyme on semen viscosity in dromedary camel. *J. Camel Pract. Res.*, 23: 219-222.
- Mal, G; Vyas, S; Srinivasan, A; Patil, NVR and Pathak, KML** (2016). Studies on liquefaction time and proteins involved in the improvement of seminal characteristics in dromedary camels (*Camelus dromedarius*). *Scientifica 1-6*. <https://doi.org/10.1155/2016/4659358>.
- Malo, C; Crichton, EG; Morrell, JM; Pukazhenth, BS; Johannisson, A; Splan, R and Skidmore, JA** (2018a). Colloid centrifugation of fresh semen improves post-thaw quality of cryopreserved dromedary camel spermatozoa. *Anim. Reprod. Sci.*, 192: 28-34.
- Malo, C; Crichton, EG; Morrell, JM; Pukazhenth, BS and Skidmore, JA** (2017a). Single layer centrifugation of fresh dromedary camel semen improves sperm quality and *in vitro* fertilization capacity compared with simple sperm washing. *Reprod. Domest. Anim.*, 52: 1097-1103.
- Malo, C; Crichton, EG and Skidmore, JA** (2017b). Optimization of the cryopreservation of dromedary camel semen: Cryoprotectants and their concentration and equilibration times. *Cryobiology*. 74: 141-147.
- Malo, C; Crichton, EG and Skidmore, JA** (2020). Preservation of the spermatozoa of the dromedary camel (*Camelus dromedarius*) by chilling and freezing: The effects of cooling time, extender composition and catalase supplementation. *Theriogenology*. 153: 9-18.
- Malo, C; Elwing, B; Soederstroem, L; Lundeheim, N; Morrell, JM and Skidmore, JA** (2018b). Effect of different freezing rates and thawing temperatures on cryosurvival of dromedary camel spermatozoa. *Theriogenology*. 125: 43-48.
- Malo, C; Grundin, J; Morrell, JM and Skidmore, JA** (2019). Individual male dependent improvement in post-thaw dromedary camel sperm quality after addition of catalase. *Anim. Reprod. Sci.*, 209: 106168.
- Mansour, N** (2022). Semen collection from dromedary camel bulls, methodology and a new patented approach. In: *Proceedings of ICAR 2020+2 Satellite Meeting on Camelid Reproduction, Bologna, Italy*. PP: 40-44.
- Medan, MS; Absy, G; Zeidan, AE; Khalil, MH; Khalifa, HH; AbdelSalaam, AM and Abdel-Khalek, TM** (2008). Survival and fertility rate of cooled dromedary camel spermatozoa supplemented with catalase enzyme. *J. Reprod. Dev.*, 54: 84-89.
- Monaco, D; Fatnassi, M; Padalino, B; Hammadi, M; Khorchani, T and Lacalandra, GM** (2016). Effect of α -Amylase, Papain, and Spermfluid[®] treatments on viscosity and semen parameters of dromedary camel ejaculates. *Res. Vet. Sci.*, 105: 5-9.
- Morrell, JM; Warring, SK; Norrestam, E; Malo, C and Huanca, W** (2021). Non-enzymatic extraction of spermatozoa from alpaca ejaculates by pipetting followed by colloid centrifugation. *Livest. Sci.*, 251: 104627.
- Morton, KM; Billah, M and Skidmore, JA** (2013). Effect of sperm diluent and dose on the pregnancy rate in dromedary camels after artificial insemination with fresh and liquid-stored semen. *J. Camelid Sci.*, 6: 49-62.
- Morton, KM; Vaughan, JL and Maxwell, WMC** (2008). The continued development of artificial insemination technology in Alpacas. *Rural Industries Research and Development Corporation: Kingston, ACT*. <https://www.semanticscholar.org/paper/The-continued-development-of-artificial-in-alpacas-Morton/1becbca8d69d6513a5880d260992e9e4326710ca>.
- Mosaferi, S; Niasari-Naslaji, A; Abarghani, A; Gharahdaghi, AA and Gerami, A** (2005). Biophysical and biochemical characteristics of Bactrian camel semen collected by artificial vagina. *Theriogenology*. 63: 92-101.
- Niasari-Naslaji, A; Gharahdaghi, AA; Mosaferi, S; Bahmani, N; Abarghani, A and Gerami, A** (2007a). Effect of sucrose extender with different levels of osmolality on the viability of spermatozoa in Bactrian camel (*Camelus bactrianus*). *Pajoujesh-va-Sazandeghi*. 75: 112-117.
- Niasari-Naslaji, A; Mosaferi, S; Abarghani, A; Gharahdaghi, AA; Bahmani, N; Ghanbari, A and Gerami, A** (2006a). Effectiveness of tris-based extender (SHOTOR diluent) for the preservation of Bactrian camel (*Camelus bactrianus*) semen. *Cryobiology*. 53: 12-21.
- Niasari-Naslaji, A; Mosaferi, S; Bahmani, N; Abarghani, A; Gharahdaghi, AA and Gerami, A** (2006b). Effect of lactose extender with different levels of osmolality and pH on the viability of Bactrian camel (*Camelus bactrianus*) spermatozoa. *Iran. J. Vet. Res.*, 7: 14-22.
- Niasari-Naslaji, A; Mosaferi, S; Bahmani, N; Gerami, A; Gharahdaghi, AA; Abarghani, A and Ghanbari, A** (2007b). Cryopreservation of semen in Bactrian camel (*Camelus bactrianus*) using SHOTOR diluent: effects of cooling rates and glycerol concentrations. *Theriogenology*. 68: 618-625.
- Niasari-Naslaji, A; Mosaferi, S; Moghiseh, A; Nikjou, D; Bahmani, N; Gerami, A; Moosavi-Movahedi, AA; Gharahdaghi, AA and Abarghani, A** (2008). Effects of equex on the survival of Bactrian camel (*Camelus bactrianus*) sperm in chilled and frozen conditions. *J. Camel Pract. Res.*, 15: 1-8.
- Panahi, F; Niasari-Naslaji, A; Seyedasgari, F; Ararooti, T; Razavi, K and Moosavi-Movahedi, AA** (2017). Supplementation of tris-based extender with plasma egg yolk of six avian species and camel skim milk for chilled preservation of dromedary camel semen. *Anim. Reprod. Sci.*, 184: 11-19.
- Perk, K** (1962). Seasonal changes in the glandula bulbourethralis of the camel. *Bull. Res. Council. Isr.*, 10: 37-44.
- Rateb, SA** (2016). Ultrasound-assisted liquefaction of dromedary camel semen. *Small Rumin. Res.*, 141: 48-55.
- Santiani, A; Huanca, W; Sapana, R; Huanca, T; Sepulveda, N and Sanchez, R** (2005). Effects on the quality of frozen-thawed alpaca (*Lama pacos*) semen using two different cryoprotectants and extenders. *Asian J. Androl.*, 7: 303-309.
- Sieme, H; Merkt, H; Musa, B; Hago, B and Willem, T** (1990). Liquid and deep freeze preservation of camel semen using different extenders and methods. In: *Proceedings of the Workshop "Is it possible to improve the reproductive performance of the camel"*. Paris. PP: 273-283.
- Skidmore, JA; Crichton, EG; Malo, CM; Vaughan, JL; Wani, NA and Herrid, M** (2020). Reproductive technologies in camelids. In: *Presicce, GA (Ed.), Reproductive technologies in animals*. (1st Edn.), Rome, Italy, Academic Press. PP: 119-134. <https://doi.org/10.1016/C2018-0-01374-2>.
- Skidmore, JA; Morton, KM and Billah, M** (2013). Artificial insemination in dromedary camels. *Anim. Reprod. Sci.*, 136: 178-186.
- Tibary, A and Anouassi, A** (1997). Artificial breeding and

- manipulation of reproduction in Camelidae. In: Tibary, A and Anouassi, A (Eds.), *Theriogenology in Camelidae: Anatomy, physiology, BSE, pathology and artificial breeding*. (1st Edn.), Ins. Agron. Vet. Hassan II, Rabat, Maroc. PP: 413-452.
- Tibary, A and Anouassi, A** (2018). Challenges in the development of artificial insemination in the dromedary camel. *Rev. Mar. Sci. Agron. Vét.*, 6: 178-188.
- Vyas, S; Goswami, P; Raj, AK and Khanna, ND** (1998). Use of tris and lactose extenders in preservation of camel semen at refrigerated temperature. *Indian Vet. J.*, 75: 810-812.
- Zhao, XX; Chen, BX and Huang, YM** (1996). Artificial insemination with deep-frozen semen in the Bactrian camel (*Camelus bactrianus*). *Research Progress in Animal Industry and Animal Products Processing*. Compiled by the Chinese Association of Agricultural Sciences. Presented at the 1st China International Annual Meeting on Agricultural Science and Technology, March 19-22, 1996. Chinese Agriculture Press, Beijing, PP: 237-242.
- Zhao, XX; Huang, YM and Chen, BX** (1994). Artificial insemination and pregnancy diagnosis in the Bactrian camel (*Camelus bactrianus*). *J. Arid Environ.*, 26: 61-65.
- Ziapour, S; Niasari-Naslaji, A; Mirtavousi, M; Keshavarz, M; Kalantari, A and Adel, A** (2014). Semen collection using phantom in dromedary camel. *Anim. Reprod. Sci.*, 151: 15-21.