

OPEN

Serum Bone Morphogenic Protein-4 Contributes to Discriminating Coronary Artery Disease Severity

Chul Soo Park, MD, PhD, Oak-Kee Hong, PhD, Mee Kyoung Kim, MD, PhD, Woo Baek Chung, MD, Yun Seok Choi, MD, PhD, Ki-Hyun Baek, MD, PhD, Ki-Ho Song, MD, PhD, Man Young Lee, MD, PhD, and Hyuk-Sang Kwon, MD, PhD

Abstract: Bone morphogenic protein 4 (BMP-4) is a known pro-inflammatory and pro-atherogenic cytokine. Here, we investigated whether the serum BMP-4 level predicts coronary artery disease (CAD) severity in humans.

We measured serum BMP-4 concentrations in 1044 consecutive patients who underwent elective coronary angiography and percutaneous coronary intervention. CAD severity was estimated by the number of diseased vessels showing $\geq 50\%$ diameter stenosis.

Among males, the serum BMP-4 level was significantly lower in patients with multivessel disease (MVD) compared with those with single-vessel disease (SVD) (16.3 ± 22.6 vs. 22.0 ± 28.4 pg/mL, $P < 0.01$). After adjustment for other cardiovascular risk factors, a high serum BMP-4 level was an independent predictor for a decreased risk of MVD (odds ratio, 0.992; 95% confidence interval [CI], 0.985–0.998; $P = 0.01$) and patients in the lower tertile were 1.55-fold more likely to have MVD compared with upper tertile patients. Receiver-operating characteristic curve analysis demonstrated that the serum BMP-4 level had a 54% sensitivity and 54% specificity for predicting MVD (area under the curve [AUC], 56.5%; 95% CI, 51.9–61.0%; $P < 0.01$). Serum BMP-4 improved the predictive capability of risk factors for MVD (AUC with and without BMP-4: 64.9 and 63.6%, respectively). Considering the likelihood ratio and number of parameters, adding the serum BMP-4 level provided a better-fit model for predicting MVD compared with the model consisting of conventional risk factors

(likelihood ratio $\chi^2 = 6.20$, $P = 0.01$). However, an association between serum BMP-4 and CAD was not observed in females.

Serum BMP-4 levels are independently associated with CAD severity and contribute to discriminating CAD severity in males.

(*Medicine* 94(39):e1530)

Abbreviations: AUC = area under the curve, BMP-4 = bone morphogenic protein 4, CAD = coronary artery disease, CI = confidence interval, DM = diabetes mellitus, HDL = high-density lipoprotein, LDL = low-density lipoprotein, MVD = multivessel disease, OR = odds ratio, PCI = percutaneous coronary intervention, SVD = single-vessel disease.

INTRODUCTION

Atherosclerosis is an inflammatory disease,^{1,2} associated with multiple risk factors and a complex pathophysiology. It is well known that vessel bifurcations or branched regions are prone to atherosclerosis.^{3–5} In these regions, a disruption in flow induces bone morphogenic protein-4 (BMP-4), which causes endothelial dysfunction and inflammation in a nuclear factor- κ B- and nicotinamide adenine dinucleotide phosphate oxidase-dependent manner.^{6–8} However, inhibition of BMP-4 has been shown to restore endothelial function.⁹ BMP-2 and BMP-4 expression is increased in human atherosclerotic plaques and the overlying endothelium of early atherosclerotic lesions within the coronary artery.^{8,10} Moreover, in a diabetic animal model, hyperglycemia reportedly induced BMP-4 expression, resulting in vascular calcification.¹¹

Although BMP-4 has been proposed as a proinflammatory and proatherosclerotic cytokine, previous *in vitro* or animal studies focused on the role of BMP-4 in early atherogenesis, such as initiation of endothelial dysfunction,^{7–9,12} and monocyte adhesion.⁷ There is a paucity of data regarding the association between serum BMP-4 level and the progression of atherosclerosis, especially in humans. We previously reported contradictory data on the association between serum BMP-4 level and atherosclerotic parameters in different study subjects. In a general population of individuals without diabetes, serum BMP-4 level was positively correlated with adiposity and insulin resistance,¹³ while it was inversely correlated with intima-media thickness and cardio-ankle vascular index in patients with type 2 diabetes.¹⁴ In a recent small study,¹⁵ serum BMP-4 levels were increased in subjects with chronic kidney disease and coronary artery disease (CAD) compared with those without CAD, but there was no difference among patients without chronic kidney disease. These human studies raise the possibility that BMP-4 might have a different role in the context of disease type or progression of atherosclerosis.

Editor: Hsueh Wang.

Received: August 3, 2015; revised: August 10, 2015; accepted: August 13, 2015.

From the Department of Internal Medicine, College of Medicine (CSP, O-KH, MKK, WBC, YSC, K-HB, K-HS, YL, HSK); Division of Cardiology, Department of Internal Medicine, Yeouido St. Mary's Hospital, College of Medicine (CSP, WBC, YSC, MYL); and Division of Endocrinology and Metabolism, Department of Internal Medicine, Yeouido St. Mary's Hospital, College of Medicine, The Catholic University of Korea, Seoul, Republic of Korea (MKK, K-HB, K-HS, HSK).

Correspondence: Hyuk Sang Kwon, Division of Endocrinology, Department of Internal Medicine, The Catholic University of Korea, Yeouido St. Mary's Hospital, 62, Yoido-dong, Youngdeungpo-gu, Seoul 150-713, Korea (e-mail: drkwon@catholic.ac.kr).

Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (NRF-2012R1A1A207098) and the Grant from Daewoong Pharmaceutical Company (2014-21).

This study was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (NRF-2012R1A1A207098) and the Grant from Daewoong Pharmaceutical Company (2014-21). The granting agencies had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. No additional external funding was received for this study.

Copyright © 2015 Wolters Kluwer Health, Inc. All rights reserved.

This is an open access article distributed under the Creative Commons Attribution-NonCommercial License, where it is permissible to download, share and reproduce the work in any medium, provided it is properly cited. The work cannot be used commercially.

ISSN: 0025-7974

DOI: 10.1097/MD.0000000000001530

TABLE 1. Clinical Characteristics of Patients With SVD or MVD

Characteristics	Male			Female		
	SVD (n = 254)	MVD (n = 375)	P Value	SVD (n = 166)	MVD (n = 249)	P Value
Demographic						
Age (yr)	60 ± 9	64 ± 10	<0.01	66 ± 9	66 ± 9	0.82
BMI (kg/m ²)	25.0 ± 2.7	24.8 ± 2.9	0.55	25.2 ± 3.1	25.5 ± 3.2	0.29
Risk factor						
Hypertension (%)	143 (56)	234 (62)	0.13	125 (75)	184 (74)	0.75
DM (%)	66 (26)	141 (38)	<0.01	56 (34)	123 (49)	<0.01
Current smoker (%)	79 (31)	124 (33)	0.61	7 (4)	10 (4)	0.92
Laboratory findings						
FBG (mg/dL)	119 ± 47	122 ± 37	0.38	115 ± 34	126 ± 52	0.02
HbA1c (%)	6.3 ± 1.4	6.4 ± 1.3	0.33	6.2 ± 1.2	6.9 ± 1.6	<0.01
WBC count (10 ³ /mm ³)	6.88 ± 2.24	6.89 ± 1.98	0.95	6.84 ± 2.91	7.06 ± 2.21	0.38
Creatinine (mg/dL)	1.30 ± 1.16	1.32 ± 1.21	0.88	1.10 ± 1.04	1.06 ± 0.87	0.71
Total cholesterol (mg/dL)	161 ± 38	166 ± 40	0.16	168 ± 34	183 ± 46	<0.01
Triglyceride (mg/dL)	152 ± 95	162 ± 111	0.24	135 ± 74	164 ± 90	<0.01
HDL (mg/dL)	43 ± 11	41 ± 10	0.12	47 ± 11	45 ± 11	0.01
LDL (mg/dL)	88 ± 35	93 ± 37	0.13	95 ± 34	105 ± 43	0.01
hs-CRP (mg/dL)	0.56 ± 2.32	0.49 ± 1.08	0.67	1.35 ± 0.11	1.95 ± 0.13	0.46
BMP-4 (pg/mL)	22.0 ± 28.4	16.3 ± 22.6	<0.01	18.4 ± 23.7	19.8 ± 25.2	0.57

Data are mean ± SD and number (%) of patients. BMI = body mass index, BMP-4 = bone morphogenic protein 4, DM = diabetes mellitus, FBG = fasting blood glucose, HDL = high-density lipoprotein, hs-CRP = high-sensitivity C-reactive protein, LDL = low-density lipoprotein, MVD = multivessel disease, SVD = single-vessel disease, WBC = white blood cell.

CAD is one of the leading causes of death worldwide,¹⁶ and there are many branching regions of the coronary arteries vulnerable to BMP-4 expression-related atherosclerosis. Therefore, we measured serum BMP-4 concentrations in subjects who underwent elective coronary angiography and investigated whether the serum BMP-4 level was associated with CAD severity and contributed to CAD discrimination in humans.

METHODS

Subjects and Coronary Angiography

We included 1044 consecutive subjects who underwent elective coronary angiography and percutaneous coronary intervention at Yeouido Saint Mary's Hospital (Seoul, Korea) between March 2007 and March 2009. We excluded patients with myocardial infarction defined as having an elevated troponin T level (>0.1 µg/L) with or without ST elevation, as well as patients who underwent a previous revascularization, who were on dialysis, had cancer, or had an infectious disease. Elective coronary angiography was performed in all subjects. CAD severity was estimated by the number of vessels showing ≥50% diameter stenosis on quantitative coronary analysis. We classified the subjects with CAD according to the number of diseased vessels into single-vessel disease (SVD) and multivessel disease (MVD) groups. All laboratory and clinical parameter data were obtained from a careful review of the patients' medical records at index admission. This study protocol complied with the principles of the Declaration of Helsinki revised in 2000, and ethical approval was obtained from the Institutional Review Board of Yeouido Saint Mary's Hospital, the Catholic University of Korea (Seoul,

Korea). Written informed consent was obtained from all subjects prior to participation.

Blood Sampling and Measurement of Serum BMP-4 Concentration

After an overnight fast, arterial blood was obtained via an arterial sheath just before the routine coronary angiography procedure. The serum was separated and stored at -70°C. Serum BMP-4 concentrations were measured using a Quantikine Human BMP-4 Immunoassay Kit (R&D Systems, Minneapolis, MN) with intra- and interassay coefficients of variation (CVs) of 5.3 and 5.8%, respectively. Serum BMP-4 measurements were performed in duplicate by researchers who were blinded to the coronary angiography results.

Statistical Analysis

Since the association between serum BMP-4 and CAD severity was significantly different in males and females, all analyses were performed separately by gender. All data are presented as means ± standard deviation (SD) or frequency and percentage. We compared the clinical characteristics and serum BMP-4 concentrations between the SVD and MVD groups using an independent t-test or χ^2 test. One-way analysis of variance (ANOVA) with a post-hoc and χ^2 test was used to compare clinical characteristics across the groups divided by serum BMP-4 level tertile. To estimate the association between serum BMP-4 level and CAD severity, we used multiple logistic regression analysis to discriminate MVD from SVD after adjustment for traditional cardiovascular risk factors defined as age, diabetes mellitus (DM), hypertension, smoking, LDL.




FIGURE 1. Serum BMP-4 levels grouped by coronary artery disease severity in males and females. Data are expressed as means ± standard deviation (SD). BMP-4 = bone morphogenic protein-4, MVD = multivessel disease, SVD = single-vessel disease.

significant at $P < 0.05$. All statistical analyses were performed using SPSS 19.0 and R statistical language v. 2.15.0.

RESULTS

Baseline Clinical Characteristics

The clinical characteristics of SVD and MVD are described in Table 1. Among males, MVD patients were older with a higher prevalence of DM. However, there were no significant differences in other laboratory or demographic parameters. Among females, MVD patients showed higher fasting blood glucose, hemoglobin (Hb) A1c, total cholesterol, and low-density lipoprotein (LDL) levels with a higher prevalence of DM. The serum BMP-4 concentration was significantly decreased in MVD compared with SVD patients among males (16.3 ± 22.6 vs. 22.0 ± 28.4 pg/mL, $P < 0.01$), but not among females (19.8 ± 25.2 vs. 18.4 ± 23.7 pg/mL, $P = 0.57$, Figure 1).

Association of Serum BMP-4 With Cardiovascular Risk Factors

Since BMP-4 is a pro-inflammatory cytokine, we investigated the relationship between serum BMP-4 level and cardiovascular risk factors. As shown in Table 2, among males and females, there were no significant differences in age, prevalence of DM, hypertension, smoking status, creatinine, lipid profile, and inflammatory markers, such as white blood cell count and

We evaluated improvements in the discrimination after adding serum BMP-4 level to a model consisting of traditional cardiovascular risk factors with an increased area under the curve (AUC) in receiver-operating characteristic analysis. We assessed whether the serum BMP-4 level provided a better-fit model to discriminate CAD severity compared with known conventional risk factors using the likelihood ratio χ^2 test for nested models. The results were considered statistically

TABLE 2. Association Between Serum BMP 4 Tertile and Clinical Characteristics

Characteristics	Male				Female			
	T1 (n = 210)	T2 (n = 210)	T3 (n = 209)	P Value	T1 (n = 140)	T2 (n = 138)	T3 (n = 137)	P value
Demographic								
Age (yr)	63 ± 10	63 ± 10	61 ± 9	0.13	66 ± 9	67 ± 9	67 ± 9	.61
BMI (kg/m ²)	25.2 ± 3.0	24.7 ± 2.7	24.9 ± 2.8	0.22	25.6 ± 3.5	25.5 ± 3.2	25.0 ± 2.8	.29
Risk factor								
Hypertension (%)	124 (59)	130 (62)	123 (59)	0.78	104 (74)	105 (76)	100 (73)	.84
DM (%)	66 (31)	80 (38)	61 (29)	0.13	64 (46)	56 (41)	59 (43)	.69
Current smoker (%)	70 (33)	63 (30)	70 (34)	0.69	5 (4)	7 (5)	5 (4)	.78
Laboratory findings								
FBG (mg/dL)	118 ± 34	122 ± 37	121 ± 52	0.58	121 ± 42	120 ± 49	124 ± 46	.83
HbA1c (%)	6.3 ± 1.2	6.5 ± 1.5	6.3 ± 1.4	0.23	6.6 ± 1.5	6.7 ± 1.5	6.7 ± 1.5	.92
WBC count (10 ³ /mm ³)	6.94 ± 2.03	6.81 ± 1.97	6.90 ± 2.26	0.83	7.22 ± 2.40	6.90 ± 2.75	6.79 ± 2.37	.35
Creatinine (mg/dL)	1.45 ± 1.67	1.24 ± 0.78	1.24 ± 0.91	0.13	1.14 ± 1.10	1.07 ± 0.87	1.01 ± 0.83	.51
Total cholesterol (mg/dL)	167 ± 38	162 ± 36	164 ± 43	0.49	175 ± 40	180 ± 43	176 ± 40	.65
Triglyceride (mg/dL)	153 ± 88	153 ± 107	167 ± 117	0.30	160 ± 92	157 ± 100	140 ± 76	.12
HDL (mg/dL)	41 ± 10	43 ± 11	42 ± 10	0.38	46 ± 12	45 ± 11	46 ± 10	.74
LDL (mg/dL)	95 ± 33	89 ± 35	88 ± 40	0.12	96 ± 41	103 ± 40	103 ± 39	.31
hs-CRP (mg/dL)	0.73 ± 2.67	0.40 ± 0.86	0.41 ± 0.85	0.09	0.59 ± 1.47	0.37 ± 0.63	0.68 ± 1.74	.31
CAD severity								
SVD (%)	68 (32)	92 (44)	94 (45)	0.02	59 (42)	52 (38)	55 (40)	.75
MVD (%)	142 (68)	118 (56)	115 (55)		81 (58)	86 (62)	82 (60)	
BMP-4 (pg/mL)	3.2 ± 1.5	8.7 ± 1.9	44.0 ± 30.4	<0.01	3.0 ± 1.6	8.9 ± 2.2	45.2 ± 27.2	<.01

Data are mean ± SD and number (%) of patients. BMI = body mass index, BMP-4 = bone morphogenic protein 4, CAD = coronary artery disease, DM = diabetes mellitus, FBG = fasting blood glucose, HDL = high-density lipoprotein, hs-CRP = high-sensitivity C-reactive protein, LDL = low-density lipoprotein, MVD = multivessel disease, SVD = single-vessel disease, WBC = white blood cell.

TABLE 3. Logistic Regression Analysis to Predict Multivessel Disease in Male

Variables	Univariate		Multivariate*	
	OR (95% CI)	P Value	OR (95% CI)	P Value
Age (yr)	1.038 (1.020–1.056)	<0.01	1.042 (1.023–1.062)	<0.01
BMI (kg/m ²)	0.982 (0.928–1.040)	0.54		
DM	1.716 (1.210–2.435)	<0.01	1.669 (1.152–2.419)	<0.01
Hypertension	1.288 (0.931–1.782)	0.13	1.184 (0.835–1.678)	0.34
Smoking	1.094 (0.777–1.540)	0.61	1.307 (0.902–1.895)	0.16
Total cholesterol (mg/dL)	1.003 (0.999–1.007)	0.16		
Triglyceride (mg/dL)	1.001 (0.999–1.003)	0.24	1.005 (1.000–1.009)	0.06
HDL(mg/dL)	0.988 (0.973–1.003)	0.12		
LDL (mg/dL)	1.003 (0.999–1.008)	0.14	0.992 (0.985–0.998)	0.01
hs-CRP (mg/dL)	0.979 (0.889–1.078)	0.67		
BMP-4 (pg/mL)	0.991 (0.985–0.997)	<0.01		

BMI = body mass index, BMP-4 = bone morphogenic protein-4, CI = confidence interval, DM = diabetes mellitus, HDL = high-density lipoprotein, hs-CRP = high-sensitive creatinine protein, LDL = low-density lipoprotein, OR = odds ratio.

*Adjusted for age, DM, hypertension, smoking status, LDL-cholesterol, and BMP-4.

high-sensitivity C-reactive protein level, across the serum BMP-4 level tertiles.

Association Between Serum BMP-4 Concentration and CAD Severity

Table 3 shows that in males, age (odds ratio [OR], 1.038; 95% confidence interval [CI], 1.020–1.056, $P < 0.01$), DM (OR, 1.716; 95% CI, 1.210–2.435, $P < 0.01$), and serum BMP-4 level (OR, 0.991; 95% CI, 0.985–0.997; $P < 0.01$) were predictors for MDV in univariate analysis. A high serum BMP-4 level was an independent predictor for a decreasing risk of MVD after adjusting for age, DM, hypertension, smoking, and LDL (OR, 0.992; 95% CI, 0.985–0.998; $P = 0.01$). Patients with a lower serum BMP-4 tertile showed a higher risk of MVD compared with upper tertile patients (OR, 1.556; 95% CI, 1.024–2.364; $P = 0.03$, Figure 2). However, there was no association between serum BMP-4 and CAD severity in females (data not shown).

Contribution of Serum BMP-4 Level in Discriminating CAD Severity in Males

Receiver-operating characteristic analysis determined that the serum BMP-4 level had a 56.5% AUC (95% CI, 51.9–61.0%, $P < 0.01$) with a 54% sensitivity and 54% specificity for predicting MVD (Figure 3A). In contrast, the model consisting of conventional risk factors such as age, DM, hypertension, smoking, and LDL showed a 63.6% AUC (95% CI, 59.1–68.0%, $P < 0.01$) with a 62% sensitivity and 60% specificity (Figure 3B). A combined model consisting of serum BMP-4 level and conventional risk factors showed a 64.9% AUC (95% CI, 60.4–69.3%, $P < 0.01$) with a 63% sensitivity and 61% specificity (Figure 3C). According to the likelihood ratio test, the combined model was a better-fit for predicting MVD compared with the model consisting of conventional risk factors only (likelihood ratio $\chi^2 = 6.20$, $P = 0.01$).

DISCUSSION

In the present study, we evaluated a potential link between serum BMP-4 level and CAD severity. To our knowledge, this is the first study to demonstrate an association between serum

BMP-4 level and CAD severity in a relatively large human study population. We found that the serum BMP-4 concentration significantly decreased in the MVD group compared with the SVD group. On multivariate analysis, the lower serum BMP-4 level was associated with MVD independently of traditional cardiovascular risk factors in males, while no association was observed in females.

Although BMP-4 signaling has been associated with endothelial dysfunction,^{7–9,12} monocyte adhesion,⁷ and foam cell formation,¹⁷ which is an early step in atherogenesis, it is unclear whether the atherogenic effect of BMP-4 signaling is maintained and actively involved in the progression of atherosclerosis. Kim et al¹⁸ recently reported that knockdown of the BMP receptor II (BMPRII) markedly elevated monocyte adhesion independently of BMP-4, and that BMPRII was down-regulated by pro-atherogenic risk factors such as disturbed flow or hypercholesterolemia. These results suggested a limited role for BMP-4 in atherogenesis.




FIGURE 2. Adjusted odds ratios for multivessel disease by serum BMP-4 tertile level. The upper BMP-4 tertile level is the reference, and 1 bars represent the 95% confidence interval. BMP-4 = bone morphogenic protein-4.




FIGURE 3. Receiver-operating characteristic curves of BMP-4, conventional risk factors, and combined model for predicting multivessel disease. BMP-4 = bone morphogenic protein-4.

The decrease in serum BMP-4 levels following progression of atherosclerosis observed in the current study can be explained in several ways. First, as Kim et al¹⁸ reported, BMP-4 is primarily involved in early atherogenesis, and the progression to advanced atherosclerosis might be dependent on other mechanisms. Second, irrespective of the atherogenic effect of BMP-4, its serum level might simply decrease as a result of BMP-4 deposition into the atherosclerotic vascular tissue as demonstrated in previous studies.^{8,10} Third, it also might be due to a possible inhibitory mechanism by BMP-4 antagonists such as gremlin, matrix Gla protein, follistatin, and noggin.^{6,11,19,20} Although we do not know the exact mechanisms underlying the decrease in serum BMP-4 during CAD progression, this might suggest that the active atherogenic role of BMP-4 in the progression to advanced CAD is not important.

We found that a model incorporating the serum BMP-4 level produced a better fit in the prediction of MVD compared with the model consisting only of established cardiovascular risk factors. The capability to predict MVD estimated by the AUC was also modestly improved from 63.6 to 64.9% by the addition of the serum BMP-4 level to the model consisting of conventional cardiovascular risk factors. Even though this improvement is modest, patients in the lower serum BMP-4 tertile were 1.55-fold more likely to have MVD compared with upper tertile patients.

In our study, there was no correlation between serum BMP-4 concentration and CAD severity in females. As a regulator of BMP-4 expression, estrogen facilitates osteoblast differentiation by up-regulating BMP-4 signaling²¹ and prevents cardiac hypertrophy by inhibiting BMP-4 expression in cardiomyocytes.²² Hence, in females, BMP-4 expression might be more influenced by this hormonal change than by the development of atherosclerosis. This sex-based difference in BMP-4 expression should be considered in future studies.

This study has several limitations. First, although we created a model to discriminate CAD according to severity, we could not validate our model in a different test set. Further studies are warranted in larger study populations. Second, because there were no data on the subjects without CAD, we could not estimate the role of serum BMP-4 level to predict CAD. Because previous human studies^{13,14,23} reported contradictory data in different clinical settings, it would be interesting to evaluate subjects with and without CAD. Third, no association was seen between serum BMP-4 level and CAD severity in females in the present study. However, there could be an association in menopausal females if this lack of an association is due to the effect of estrogen. We did not analyze postmenopausal females in this study due to the lack of information about menopausal status and the small number of females included.

In conclusion, here we demonstrated that serum BMP-4 concentration is independently associated with CAD severity and contributes to its discrimination.

ACKNOWLEDGMENT

This study was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (NRF-2012RIA1A207098) and the Grant from Daewoong Pharmaceutical Company (2014-21).

REFERENCES

- Libby P. Inflammation in atherosclerosis. *Nature*. 2002;420:868–874.
- Ross R. Atherosclerosis: an inflammatory disease. *N Engl J Med*. 1999;340:115–126.
- Roger VL, Go AS, Lloyd-Jones DM, et al. Heart disease and stroke statistics: 2011 update: a report from the American Heart Association. *Circulation*. 2011;123:e18–e209.
- VanderLaan PA, Reardon CA, Getz GS. Site specificity of atherosclerosis: site-selective responses to atherosclerotic modulators. *Arterioscler Thromb Vasc Biol*. 2004;24:12–22.
- Zarins CK, Giddens DP, Bharadvaj BK, et al. Carotid bifurcation atherosclerosis. Quantitative correlation of plaque localization with flow velocity profiles and wall shear stress. *Circ Res Oct*. 1983;53:502–514.
- Chang K, Weiss D, Suo J, et al. Bone morphogenic protein antagonists are coexpressed with bone morphogenic protein 4 in endothelial cells exposed to unstable flow in vitro in mouse aortas and in human coronary arteries: role of bone morphogenic protein antagonists in inflammation and atherosclerosis. *Circulation*. 2007;116:1258–1266.
- Sorescu GP, Song H, Tressel SL, et al. Bone morphogenic protein 4 produced in endothelial cells by oscillatory shear stress induces monocyte adhesion by stimulating reactive oxygen species production from a nox1-based NADPH oxidase. *Circ Res*. 2004;95:773–779.
- Sorescu GP, Sykes M, Weiss D, et al. Bone morphogenic protein 4 produced in endothelial cells by oscillatory shear stress stimulates an inflammatory response. *J Biol Chem*. 2003;278:31128–31135.
- Zhang Y, Liu J, Tian XY, et al. Inhibition of bone morphogenic protein 4 restores endothelial function in db/db diabetic mice. *Arterioscler Thromb Vasc Biol*. 2014;34:152–159.
- Dhore CR, Cleutjens JP, Lutgens E, et al. Differential expression of the bone matrix regulatory proteins in human atherosclerotic plaques. *Arterioscler Thromb Vasc Biol*. 2001;21:1998–2003.
- Bostrom KI, Jumabay M, Matveyenko A, et al. Activation of vascular bone morphogenic protein signaling in diabetes mellitus. *Circ Res*. 2011;108:446–457.

12. Wong WT, Tian XY, Chen Y, et al. Bone morphogenic protein-4 impairs endothelial function through oxidative stress-dependent cyclooxygenase-2 upregulation: implications on hypertension. *Circ Res*. 2010;107:984–991.
13. Son JW, Kim MK, Park YM, et al. Association of serum bone morphogenetic protein 4 levels with obesity and metabolic syndrome in non-diabetic individuals. *Endocr J*. 2011;58:39–46.
14. Son JW, Jang EH, Kim MK, et al. Serum BMP-4 levels in relation to arterial stiffness and carotid atherosclerosis in patients with Type 2 diabetes. *Biomark Med*. 2011;5:827–835.
15. Stahls PF3rd, Lightell DJ Jr, Moss SC, et al. Elevated serum bone morphogenetic protein 4 in patients with chronic kidney disease and coronary artery disease. *J Cardiovasc Transl Res*. 2013;6:232–238.
16. Go AS, Mozaffarian D, Roger VL, et al. Heart disease and stroke statistics: 2014 update: a report from the American Heart Association. *Circulation Jan*. 2014;129:e28–e292.
17. Feng J, Gao J, Li Y, et al. BMP4 enhances foam cell formation by BMPR-2/Smad1/5/8 signaling. *Int J Mol Sci*. 2014;15:5536–5552.
18. Kim CW, Song H, Kumar S, et al. Anti-inflammatory and antiatherogenic role of BMP receptor II in endothelial cells. *Arterioscler Thromb Vasc Biol*. 2013;33:1350–1359.
19. Sun J, Zhuang FF, Mullersman JE, et al. BMP4 activation and secretion are negatively regulated by an intracellular gremlin-BMP4 interaction. *J Biol Chem*. 2006;281:29349–29356.
20. Sweatt A, Sane DC, Hutson SM, et al. Matrix Gla protein (MGP) and bone morphogenetic protein-2 in aortic calcified lesions of aging rats. *J Thromb Haemost*. 2003;1:178–185.
21. Matsumoto Y, Otsuka F, Takano-Narazaki M, et al. Estrogen facilitates osteoblast differentiation by upregulating bone morphogenetic protein-4 signaling. *Steroids*. 2013;78:513–520.
22. Wang YC, Xiao XL, Li N, et al. Oestrogen inhibits BMP4-induced BMP4 expression in cardiomyocytes: a potential mechanism of oestrogen-mediated protection against cardiac hypertrophy. *Br J Pharmacol*. 2014doi: 10.1111/bph.12983.
23. Stahls PF3rd, Lightell DJ Jr, Moss SC, et al. Elevated serum bone morphogenetic protein 4 in patients with chronic kidney disease and coronary artery disease. *J Cardiovasc Transl Res*. 2013;6:232–238.