

Randomized phase III study comparing the efficacy and safety of irinotecan plus S-1 with S-1 alone as first-line treatment for advanced gastric cancer (study GC0301/TOP-002)

Hiroyuki Narahara · Hiroyasu Iishi · Hiroshi Imamura · Akira Tsuburaya · Keisho Chin · Haruhiko Imamoto · Taito Esaki · Hiroshi Furukawa · Chikuma Hamada · Yuh Sakata

Received: 30 July 2010/Accepted: 31 October 2010/Published online: 23 February 2011
© The Author(s) 2011. This article is published with open access at Springerlink.com

Abstract

Background Irinotecan hydrochloride and S-1, an oral fluoropyrimidine, have shown antitumor activity against advanced gastric cancer as single agents in phase I/II studies. The combination of irinotecan and S-1 (IRI-S) is also active against advanced gastric cancer. This study was conducted to compare the efficacy and safety of IRI-S versus S-1 monotherapy in patients with advanced or recurrent gastric cancer.

Methods Patients were randomly assigned to oral S-1 (80 mg/m² daily for 28 days every 6 weeks) or oral S-1 (80 mg/m² daily for 21 days every 5 weeks) plus irinotecan (80 mg/m² by intravenous infusion on days 1 and 15 every 5 weeks) (IRI-S). The primary endpoint was overall survival. Secondary endpoints included the time to treatment failure, 1- and 2-year survival rates, response rate, and safety.

Results The median survival time with IRI-S versus S-1 monotherapy was 12.8 versus 10.5 months ($P = 0.233$), time to treatment failure was 4.5 versus 3.6 months ($P = 0.157$), and the 1-year survival rate was 52.0 versus 44.9%, respectively. The response rate was significantly higher for IRI-S than for S-1 monotherapy (41.5 vs. 26.9%, $P = 0.035$). Neutropenia and diarrhea occurred more frequently with IRI-S, but were manageable. Patients treated with IRI-S received more courses of therapy at a relative dose intensity similar to that of S-1 monotherapy.

Conclusions Although IRI-S achieved longer median survival than S-1 monotherapy and was well tolerated, it did not show significant superiority in this study.

Keywords Irinotecan S-1 · Gastric cancer · Phase III · Randomized controlled trial

H. Narahara (✉)

Division of Clinical Oncology, Hiroshima University Graduate School of Biomedical Sciences, 1-2-3 Kasumi, Minami-ku, Hiroshima 734-8551, Japan
e-mail: narahara@hiroshima-u.ac.jp

H. Iishi

Department of Gastrointestinal Oncology, Osaka Medical Center for Cancer and Cardiovascular Diseases, Osaka, Japan

H. Imamura · H. Furukawa

Department of Surgery, Sakai Municipal Hospital, Osaka, Japan

A. Tsuburaya

Department of Gastrointestinal Surgery,
Kanagawa Cancer Center, Yokohama, Japan

K. Chin

Department of Medical Oncology, Cancer Institute Hospital, Tokyo, Japan

H. Imamoto

Department of Surgery, Kinki University School of Medicine, Osaka-sayama, Japan

T. Esaki

Department of Medical Oncology, National Kyushu Cancer Center, Fukuoka, Japan

C. Hamada

Faculty of Engineering, Tokyo University of Science, Tokyo, Japan

Y. Sakata

Department of Internal Medicine, Misawa Municipal Hospital, Misawa, Japan

Introduction

Gastric cancer is the second leading cause of cancer-related deaths after lung cancer in Japan, and it was responsible for approximately 50,000 deaths in 2005 [1]. While surgery and appropriate adjuvant chemotherapy have resulted in superior stage-by-stage survival when compared with that in other parts of the world [2], the prognosis of unresectable or recurrent gastric cancer remains dismal. The development of more effective chemotherapeutic regimens is therefore warranted.

In Western countries where a combination of 5-fluorouracil (5-FU) and cisplatin (CDDP) [3] has served as a reference arm in several phase III studies [4–6], triplets employing epirubicin [7] or docetaxel [5] in addition to this combination are the current standards, with modifications such as the replacement of CDDP with oxaliplatin and the replacement of infusional 5-FU with oral agents such as capecitabine [8]. Failure with the first-line treatment usually denotes the termination of chemotherapy, and second-line treatments are rarely considered outside of clinical trials. In Japan, where a phase III study (JCOG9205) failed to show superiority of a 5-FU/CDDP combination over 5-FU alone [9], the 5-FU monotherapy remained a standard of care, and other cytotoxic agents were usually delivered sequentially as second-line and third-line therapies rather than concurrently as combination therapy. With this strategy, the median survival time (MST) of patients with advanced gastric cancer whose treatment started with infusional 5-FU alone actually reached 10.8 months [9].

In the 1990s, S-1 (TS-1; Taiho Pharmaceutical, Tokyo, Japan), an oral derivative of 5-FU, was developed for the treatment of gastric cancer [10–12]. With an exceptionally high response rate of 46% as a single agent, this drug rapidly established itself as a community standard in Japan and was used widely in clinical practice. Phase III trials eventually proved the non-inferiority of S-1 when compared with infusional 5-FU in the advanced/metastatic setting [13], along with the superiority of S-1 monotherapy over observation alone in the postoperative adjuvant setting [14]. In addition, S-1 was found to be a unique cytotoxic drug, in that Japanese patients tolerated higher doses than Western patients, due to differences in the gene polymorphism of relevant enzymes [15]. Thus, the development of novel chemotherapeutic regimens in Japan during the 2000s has inevitably centered around this drug.

The establishment of doublets to enhance response rates and improve on survival was the next important step, and several phase I/II studies were performed to explore combinations of S-1 with other cytotoxic drugs such as CDDP [16], docetaxel [17], paclitaxel [18], and irinotecan (Yakult Honsha, Tokyo, Japan; Daiichi Sankyo, Tokyo, Japan) [19]. All these combinations were found to be promising,

with response rates of around 50% and relatively favorable safety profiles. A series of phase III trials comparing these doublets with S-1 monotherapy were subsequently planned and conducted to seek optimal first-line treatments. Of these, a phase III trial to explore S-1/CDDP was the first to complete accrual, and a significant improvement in MST of this combination over S-1 monotherapy was proven [20]. The present study, entitled GC0301/TOP-002, represents another of these attempts, exploring the efficacy of a combination of S-1 and irinotecan (IRI-S). The dose and schedule for this combination had been established by a phase I trial [21], and treatment at the recommended dose has shown a response rate of 47.8% [95% confidence interval (CI) 27.4–68.2%] with an MST of 394 days in a phase II study [19]. Given these earlier results and the synergistic effect of irinotecan and 5-FU observed in pre-clinical studies, the results of this present trial have been eagerly awaited.

Patients and methods

Eligibility

The eligibility criteria were histologically and cytologically confirmed unresectable or recurrent gastric adenocarcinoma; oral food intake possible; age between 20 and 75 years; no prior radiotherapy or chemotherapy; expected survival for ≥ 12 weeks; Eastern Cooperative Oncology Group (ECOG) performance status of 0–2; and adequate major organ function before chemotherapy (leukocyte count of 4,000–12,000/mm³, hemoglobin ≥ 8.0 g/dl, platelet count $\geq 100,000/\text{mm}^3$, total bilirubin ≤ 1.5 mg/dl, aspartate aminotransferase ≤ 100 IU/l, alanine aminotransferase ≤ 100 IU/l, creatinine ≤ 1.2 mg/dl). The main exclusion criteria were massive ascites, active concomitant malignancy, uncontrolled diabetes mellitus, and pregnancy or breast-feeding. Written informed consent was obtained from each patient. Institutional review board approval was obtained at each participating institution. An independent data monitoring committee evaluated safety throughout this study. The study was performed in accordance with the Declaration of Helsinki and Good Clinical Practice Guidelines. This trial was registered with the Japan Pharmaceutical Information Center (JapicCTI-050083).

Treatment schedule

In the S-1 monotherapy group, patients received oral S-1 twice daily for 28 days every 6 weeks. In the IRI-S group, S-1 (80 mg/m²) was given orally for 21 days and irinotecan (80 mg/m²) was infused intravenously on days 1 and 15 every 5 weeks. In both groups, the dose of S-1 was

based on body surface area: 40 mg if the area was <1.25 m²; 50 mg for 1.25–1.5 m², and 60 mg for ≥1.5 m². Dose modification criteria were defined in the protocol. Treatment was discontinued if there was documented disease progression, unacceptable toxicity, or withdrawal of consent.

Assessment of response and toxicity

All patients who had at least one measurable lesion were evaluated for tumor response according to the Response Evaluation Criteria in Solid Tumors (RECIST) [22]. All radiologic assessments were confirmed by extramural review. Toxicity was evaluated according to the National Cancer Institute Common Toxicity Criteria (version 2.0).

Statistical analysis

Eligible patients were registered with the data center and randomized by centralized dynamic allocation with stratification for advanced/recurrent disease (with or without adjuvant chemotherapy), performance status (0/1/2), and institution. The full analysis set was defined as all patients who received treatment at least once and met all inclusion criteria. The per-protocol set was defined as all patients who received treatment at least once and had no major protocol violations.

The primary endpoint was overall survival, which was compared between groups using the stratified log-rank test. Secondary endpoints were the time to treatment failure (TTF), the 1- and 2-year survival rates, the response rate, and safety. Overall survival time was defined as the interval from the date of registration to the date of death (patients who remained alive at the final follow-up were censored at that time). Survival curves were estimated by the Kaplan–Meier method, and differences were analyzed with the stratified log-rank test. Hazard ratios (HRs) for various prognostic factors were calculated using a stratified Cox proportional hazards model. TTF was defined as the time from the date of registration to the date of detection of progressive disease, death, or treatment discontinuation.

In addition, subset analyses were conducted, using the Cox proportional hazards model, to identify factors that influenced overall survival in each group. As well as the predetermined variables such as gender, age, performance status, and disease status (whether the disease was unresectable or recurrent), subset analyses were conducted for 6 additional variables; the presence or absence of a measurable lesion by the RECIST, hepatic metastasis, peritoneal metastasis, existent of primary focus, metastasis the number of metastatic foci, and tumor histology. All analyses were performed using SAS system version 8.2 (SAS Institute, Cary, NC, USA).

This study was designed to detect a 40% improvement in MST at a two-tailed significance level of $P \leq 0.05$ with 80% power. The MST for S-1 monotherapy was assumed to be 8.5 months, based on the results of previous phase I/II studies [12, 23]. A total of 142 patients per group were required according to calculations made with nQuery Advisor version 4.0 (Statistical Solutions, Boston, MA, USA), and the sample size was set as 300 (150 patients per group).

We initially planned to continue follow-up for ≥1.5 years after the registration of all patients, with a cut-off date of April 2007. However, an unexpectedly high survival rate of 22% (68 of 315 patients) at the cut-off date prompted the Coordinating Committee, the medical expert, and the biostatistician to advise the sponsor to continue follow-up for a further year before performing the final analysis. Thus, the MST was also calculated using 2.5-year follow-up data.

Results

Patient characteristics

Between June 2004 and November 2005, a total of 326 patients (S-1 monotherapy, $n = 162$; IRI-S, $n = 164$) were enrolled from 54 institutions and randomized (Fig. 1). Seven patients were subsequently found to be ineligible or withdrew before receiving any treatment. Another 4 patients were found to be ineligible after starting treatment and were not included in the analysis. Therefore, 315 patients (S-1 monotherapy, $n = 160$; IRI-S, $n = 155$) were evaluable and were included in the full analysis set to assess overall survival and TTF. In addition, 187 patients were evaluable for tumor response. Baseline patient characteristics are shown in Table 1.

Treatments given

The median number of treatment courses was three (range 1–19) for S-1 monotherapy whose duration was 6 weeks, and four (range 1–25) for IRI-S whose duration was 5 weeks. The main reasons for treatment discontinuation were disease progression [S-1 monotherapy vs. IRI-S, 116/160 (72.5%) vs. 89/155 (57.4%)], adverse events [12/160 (7.5%) vs. 23/155 (14.8%)], attending physician's decision [18/160 (11.3%) vs. 18/155 (11.6%)], and consent withdrawal [11/160 (6.9%) vs. 17/155 (11.0%)]. The median TTF was 3.6 months (95% CI 2.9–4.1) and 4.5 months (95% CI 3.7–5.3), respectively ($P = 0.157$). The relative dose intensity was 88.9% for S-1 monotherapy, versus 90.0% for S-1 and 86.2% for irinotecan




Fig. 1 Patient disposition. *FAS* Full analysis set, *IRI-S* S-1 plus irinotecan, *PPS* per-protocol set, *TTF* time to treatment failure

among those treated with IRI-S. Most patients in both groups received the scheduled dose of chemotherapy.

Second-line chemotherapy was administered to 240 patients (76%; S-1 monotherapy, $n = 112$; IRI-S, $n = 128$) (Table 2). The most common second-line therapy in both groups was a taxane alone (S-1 monotherapy, 26.9%; IRI-S, 40.6%). Among patients initially treated with S-1, 13 received crossover treatment with IRI-S, while 31 patients originally treated with IRI-S received second-line S-1 monotherapy.

Response and survival

The overall response rate was determined in 187 patients evaluable by the RECIST, and was significantly higher with IRI-S than with S-1 monotherapy (39/94, 41.5% vs. 25/93, 26.9%; $P = 0.035$) (Table 3).

The MST at the predetermined cut-off date was 12.8 months with IRI-S compared with 10.5 months with S-1 monotherapy (HR 0.856, $P = 0.233$) (Fig. 2), but the difference was not statistically significant. The 1-year survival rates were 44.9% [95% CI 37.2–52.6%] with S-1

Table 1 Baseline characteristics and prior therapy

Characteristic	Treatment					
	S-1		IRI-S		Total	
	n	%	n	%	n	%
Patients randomized	162		164		326	
Patients receiving at least one dose of study medication (full analysis set)	160		155		315	
Sex						
Male	127	79	110	71	237	75
Female	33	21	45	29	78	25
Age (years)						
Median	63		63		63	
Range	27–75		33–75		27–75	
ECOG performance status						
0	109	68	102	66	211	67
1	46	29	48	31	94	30
2	5	3	5	3	10	3
Tumor histology						
Intestinal	71	44	61	39	132	42
Diffuse	88	55	93	60	181	57
Other	1	1	1	1	2	1
Resection of primary tumor						
+	93	58	93	60	186	59
-	67	42	62	40	129	41
Advanced	133	83	129	83	262	83
Recurrent						
Adjuvant chemotherapy (+)	5	3	5	3	10	3
Adjuvant chemotherapy (-)	22	14	21	14	43	14

IRI-S S-1 plus irinotecan, ECOG Eastern Cooperative Oncology Group

monotherapy and 52.0% (95% CI 44.1–59.9%) with IRI-S, while the 2-year survival rates were 19.5% (95% CI 12.6–26.4%) and 18.0% (95% CI 11.2–24.8%), respectively.

MST was additionally calculated as an exploratory analysis after 2.5 years of follow-up, but the result was identical to the initial analysis at 12.8 months for IRI-S and at 10.5 months for S-1 monotherapy (HR 0.927; log-rank test $P = 0.536$). Again, the difference was not statistically significant.

Prognostic factors of all patients and factors that favored treatment with IRI-S

Baseline risk factors with a significant influence on the overall survival of all patients accrued ($P < 0.05$) were performance status (HR 1.348, 95% CI 1.079–1.686, Wald test $P = 0.009$), tumor histology (HR 1.720, 95% CI

Table 2 Second-line chemotherapy

Regimen	S-1 (n = 160)		IRI-S (n = 155)	
	n	%	n	%
IRI-S	13	8.1	—	—
Irinotecan-based regimen ^a	27	16.9	4	2.6
S-1 alone	—	—	31	20.0
S-1-based regimen ^b	9	5.6	11	7.1
Taxane alone	43	26.9	63	40.6
Others	20	12.5	19	12.3
None	48	30.0	27	17.4

IRI-S S-1 plus irinotecan

^a Irinotecan/cisplatin, irinotecan/taxane

^b S-1/cisplatin, S-1/taxane

Table 3 Response to treatment

	S-1 (n = 93)		IRI-S (n = 94)	
	n	%	n	%
Complete response	0	0	0	0
Partial response	25	27	39	41
Stable disease	35	38	40	43
Progressive disease	30	32	12	13
Not assessable	3	3	3	3
Overall response rate	26.9		41.5*	
95% CI	18.2–37.1		31.4–52.1	

CI confidence interval

* P = 0.035 (χ^2 test)

1.161–2.548, P = 0.007), target lesion (HR 1.525, 95% CI 1.164–1.999, P = 0.002), and surgery for the primary tumor (HR 0.698, 95% CI 0.538–0.906, P = 0.007).

Stratified analysis according to baseline patient characteristics (Fig. 3) showed that IRI-S was significantly more effective than S-1 monotherapy for patients with diffuse-type histology (HR 0.632, 95% CI 0.454–0.880) and for those with an ECOG performance status of 1 or 2 (HR 0.614, 95% CI 0.401–0.940). No differences were observed for the other factors assessed.

Safety

Adverse events that occurred in each group are listed in Table 4. The incidence of major hematological toxicities was higher with IRI-S than with S-1 monotherapy. Grade 3 or 4 neutropenia was observed in 10.6% of patients treated with S-1 monotherapy versus 27.1% of patients treated with IRI-S, while the corresponding incidences of infection/febrile neutropenia were 3.8 versus 1.9%. The most common grade 3 or 4 non-hematological toxicities were diarrhea (S-1 monotherapy vs. IRI-S, 5.6 vs. 16.1%),




Fig. 2 Kaplan–Meier estimates of overall survival (a) and time to treatment failure (b) for 315 evaluable patients treated with S-1 monotherapy or S-1 plus irinotecan (IRI-S). MST Median survival time, TTF time to treatment failure, CI confidence interval

anorexia (18.8 vs. 17.4%), nausea (5.6 vs. 7.1%), and vomiting (1.9 vs. 3.2%). Hand-foot skin reaction, a characteristic adverse event associated with some oral fluoropyrimidines, was confined to grade 2 or less and was observed in only 4.4 and 5.2% of patients treated with S-1 monotherapy and IRI-S, respectively. There were no treatment-related deaths among patients treated with S-1 monotherapy, whereas two patients in the IRI-S died of potentially treatment-related conditions (severe bone marrow dysfunction, multiple organ failure that was probably associated with multiple duodenal ulcers).

Discussion

This study was conducted to determine whether IRI-S could prolong MST compared with S-1 monotherapy. Basic studies have indicated that irinotecan has a multifactorial synergistic effect with the anti-tumor activity




Fig. 3 Subset analysis of overall survival stratified by baseline patient characteristics. CI Confidence interval, ECOG Eastern Cooperative Oncology Group, RECIST Response Evaluation Criteria in Solid Tumors

of 5-FU [24, 25]. In addition, several trials exploring combinations of S-1 and irinotecan have reported promising response rates [19, 23, 26, 27]; the dose and schedule in the present study was selected based on the lower incidence of grade 3 neutropenia and gastrointestinal toxicity evidenced from phase II studies among these trials.

Although the combination therapy in the present study achieved a significantly higher response rate, the initial expectation that the addition of irinotecan would improve the MST by 40% was not met. Thus, the combination of S-1 and CDDP remains the first-line chemotherapy that can be recommended for Japanese patients, while patients who are frail or those who wish to refrain from the short stay in the hospital required for hydration could turn to S-1 monotherapy. Another standard treatment could be available pending the results of a phase III trial comparing S-1 with an S-1/docetaxel combination [17]. A combination of CDDP with 5-FU or its derivative capecitabine has been used as a platform for molecularly targeting agents in recent international trials [28]; however, the place of

platinum agents in the first-line treatment of gastric cancer would seem indispensable at present.

Irinotecan has often been delivered in combination with CDDP for gastric cancer in the West [29]. This combination was also explored in Japan in a phase II trial [30] and subsequently in a phase III trial [13], but failed to show statistically significant superiority over infusional 5-FU alone. Irinotecan was more recently found to be similarly effective to CDDP when delivered with 5-FU [31], with benefit in terms of a more favorable toxicity profile. The combination then went on to be compared with a 5-FU/CDDP combination [4], but, again, failed to show a survival advantage. With similar results obtained from the present study, irinotecan-based chemotherapy would no longer be expected to surpass 5-FU or its derivatives with or without CDDP in the first-line setting.

Our stratified analysis revealed that IRI-S had a significant effect on overall survival in patients with diffuse-type histology and an ECOG performance status of 1 or 2 (Fig. 3). IRI-S was more effective in symptomatic patients. This finding may be related to its higher response rate,

Table 4 Summary of adverse events

	S-1 (<i>n</i> = 160)				IRI-S (<i>n</i> = 155)			
	All events		Grade 3/4		All events		Grade 3/4	
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%
Anemia	83	51.9	19	11.5	113	72.9	24	15.5
Leukopenia	83	51.9	5	3.1	115	74.2	18	11.6
Neutropenia	86	53.8	17	10.6	113	72.9	42	27.1
Infection/febrile neutropenia	28	17.5	6	3.8	40	25.8	3	1.9
Thrombocytopenia	18	11.3	6	3.8	17	11.0	2	1.3
Increased AST	75	46.9	8	5.0	69	44.5	5	3.2
Increased ALT	58	36.3	3	1.9	69	44.5	3	1.9
Increased bilirubin	74	46.3	9	5.6	56	36.1	5	3.2
Increased creatinine	17	10.6	2	1.3	19	12.3	3	1.9
Fatigue	101	63.1	12	7.5	123	79.4	10	6.5
Alopecia	13	8.1	0	0.0	87	56.1	0	0.0
Anorexia	104	65.0	30	18.8	125	80.6	27	17.4
Diarrhea	63	39.4	9	5.6	103	66.5	25	16.1
Nausea	84	52.5	9	5.6	115	74.2	11	7.1
Vomiting	60	37.5	3	1.9	68	43.9	5	3.2
Stomatitis/pharyngitis	27	16.9	2	1.3	34	21.9	4	2.6
Hand-foot skin reaction	7	4.4	0	0.0	8	5.2	0	0.0
Pigmentation changes	74	46.3	0	0.0	77	49.7	0	0.0

Adverse events were graded according to National Cancer Institute Common Toxicity Criteria, version 2.0

ALT alanine aminotransferase, AST aspartate aminotransferase, IRI-S S-1 plus irinotecan

resulting from tumor shrinkage, with subsequent attenuation of clinical symptoms, possibly leading to enhanced survival time. The effect of IRI-S in cancer with diffuse-type histology was in line with the finding of the subset analysis of another phase III study that an irinotecan/CDDP combination improved the survival of patients with undifferentiated gastric cancer [13]. However, these data are contradictory to data from a phase II study of the combination of S-1 and irinotecan [19], where a higher response rate was observed for intestinal-type histology. It would not seem feasible at this time, therefore, to attempt to identify patients who may benefit from the IRI-S, using clinicopathologic factors that are easily accessible.

As mentioned previously, cytotoxic drugs tend to be used sequentially as second-line and third-line therapies in some countries, including Japan. Recently, Thuss-Patienc et al. [32] reported on second-line treatment for metastatic gastric cancer, and stated that irinotecan monotherapy significantly extended survival compared with best supportive care. A retrospective study exploring a combination of irinotecan and CDDP for patients who failed first-line therapy with S-1 has shown a promising response rate of

28.6% and a MST of 9.4 months from the first day of the second-line treatment [33]. Another retrospective study, also in the second-line setting, has shown promising MSTs, ranging from 9.5 to 10.1 months [34]. These studies suggest a role for irinotecan after the failure of a 5-FU-based first-line treatment, provided that the patients retain sufficient performance status to tolerate this drug. Because definite evidence remains unavailable, further prospective studies in the second-line and third-line settings are warranted to confirm the place of irinotecan in the treatment of gastric cancer. IRI-S uses up one of promising drug combination for the second line treatment without sufficient prolongation of TTF when compared with S-1 monotherapy. It could partially explain why the combination failed to attain significant gain in MST in the present study.

IRI-S was generally well tolerated in the present study. The dose intensity of S-1 in patients treated with IRI-S was equivalent to that in patients receiving S-1 monotherapy, demonstrating the good tolerability of the IRI-S. The most common grade 3 or 4 adverse events associated with this regimen included neutropenia (27.1%) and diarrhea (16.1%), both of these being more frequent than in patients receiving S-1 monotherapy. IRI-S appears to be better tolerated than either the S-1/CDDP or irinotecan/CDDP regimens explored in other phase III studies [13, 20]. Grade 3 or 4 neutropenia was less common with IRI-S than with the S-1/CDDP and irinotecan/CDDP regimens (27 vs. 40% and 65%, respectively), as was anorexia (17 vs. 30% and 33%) and nausea (7 vs. 12% and 21%). Only diarrhea was more common with IRI-S than with the S-1/CDDP and irinotecan/CDDP regimens (16 vs. 4% and 9%, respectively) [13, 20]. However, it is of note that, in the present study, two patients who received IRI-S died of potentially treatment-related conditions. The evaluation of uridine 5'-diphospho-glucuronosyl-transferase gene polymorphism, which had not been approved at the time the trial was conducted, could now identify a small number of patients who may suffer from overt adverse reactions to IRI-S [35].

Although manageable in most cases, the IRI-S was found to be more toxic than S-1 monotherapy. To conclude, the improvement in the response rate observed with the IRI-S did not translate into the predicted prolongation of MST.

Acknowledgments This study was supported by Yakult Honsha Co., Ltd. and Daiichi Sankyo Co., Ltd. (GC0301/TOP002). The funding source had no role in the study design, data collection, data analysis, or interpretation. We thank the patients, clinicians, and support staff who participated in this study. We also thank Y. Takada, K. Miyakawa, and K. Tamura for performing extramural review to assess objective responses, as well as T. Taguchi, S. Yoshida, and H. Origasa for their helpful advice.

Conflict of interest Chikuma Hamada has received advisory fees from Yakult Honsha Co., Ltd. (Tokyo, Japan) and Daiichi Sankyo

Co., Ltd. (Tokyo, Japan). Yuh Sakata has received advisory fees and honoraria from Yakult Honsha Co., Ltd. (Tokyo, Japan), Daiichi Sankyo Co., Ltd. (Tokyo, Japan), and Taiho Pharmaceutical Co., Ltd. (Tokyo, Japan). The other authors have no conflicts of interest to declare.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- Foundation for Promotion of Cancer Research. Cancer Statistics in Japan 2007. Available from URL http://ganjoho.ncc.go.jp/public/statistics/backnumber/2007_en.html. Accessed June 21, 2010.
- Maruyama K, Kaminishi M, Hayashi K, Isobe Y, Honda I, Katai H, et al. Gastric cancer treated in 1991 in Japan: data analysis of nationwide registry. *Gastric Cancer*. 2006;9:51–66.
- Vanhoefer U, Rougier P, Wilke H, Dureux MP, Lacave AJ, Van Cutsem E, et al. Final results of a randomized phase III trial of sequential high-dose methotrexate, fluorouracil, and doxorubicin versus etoposide, leucovorin, and fluorouracil versus infusional fluorouracil and cisplatin in advanced gastric cancer: a trial of the European Organization for Research and Treatment of Cancer Gastrointestinal Tract Cancer Cooperative Group. *J Clin Oncol*. 2000;18:2648–57.
- Dank M, Zaluski J, Barone C, Valvere V, Yalcin S, Peschel C, et al. Randomized phase III study comparing irinotecan combined with 5-fluorouracil and folinic acid to cisplatin combined with 5-fluorouracil in chemotherapy naive patients with advanced adenocarcinoma of the stomach or esophagogastric junction. *Ann Oncol*. 2008;19:1450–7.
- Ajani JA, Moiseyenko VM, Tjulandin S, Majilis A, Constenla M, Boni C, et al. Clinical benefit with docetaxel plus fluorouracil and cisplatin compared with cisplatin and fluorouracil in a phase III trial of advanced gastric or gastroesophageal adenocarcinoma: the V-325 Study Group. *J Clin Oncol*. 2007;25:3205–9.
- Ajani JA, Rodriguez W, Bodoky G, Moiseyenko CM, Lichinitser M, Gorbunova V, et al. Multicenter phase III comparison of cisplatin/S-1 with cisplatin/infusional fluorouracil in advanced gastric or gastroesophageal adenocarcinoma study: the FLAGS trial. *J Clin Oncol*. 2010;28:1547–53.
- Waters JS, Norman A, Cunningham D, Scarffe JH, Webb A, Harper P, et al. Long-term survival after epirubicin, cisplatin and fluorouracil for gastric cancer: results of a randomized trial. *Br J Cancer*. 1999;80:269–72.
- Cunningham D, Starling N, Rao S, Iveson T, Nicolson M, Coxon F, et al. Capecitabine and oxaliplatin for advanced esophagogastric cancer. *N Engl J Med*. 2008;358:36–46.
- Ohtsu A, Shimada Y, Shirao K, Boku N, Hyodo I, Saito H, et al. Randomized phase III trial of fluorouracil alone versus fluorouracil plus cisplatin versus uracil and tegafur plus mitomycin in patients with unresectable, advanced gastric cancer: the Japan Clinical Oncology Group Study (JCOG9205). *J Clin Oncol*. 2003;21:54–9.
- Fukushima M, Satake H, Uchida J, Shimamoto Y, Kato T, Takechi T, et al. Preclinical antitumor efficacy of S-1: a new oral formulation of 5-fluorouracil on human tumor xenografts. *Int J Oncol*. 1998;13:693–8.
- Koizumi W, Kurihara M, Nakano S, Hasegawa K. Phase II study of S-1, a novel oral derivative of 5-fluorouracil, in advanced gastric cancer. For the S-1 Cooperative Gastric Cancer Study Group. *Oncology*. 2000;58:191–7.
- Sakata Y, Ohtsu A, Horikoshi N, Sugimachi K, Mitachi Y, Taguchi T. Late phase II study of novel oral fluoropyrimidine anticancer drug S-1 (1 M tegafur-0.4 M gimestat-1 M ostadt potassium) in advanced gastric cancer patients. *Eur J Cancer*. 1998;34:1715–20.
- Boku N, Yamamoto S, Fukuda H, Shirao K, Doi T, Sawaki A, et al. Fluorouracil versus combination of irinotecan plus cisplatin versus S-1 in metastatic gastric cancer: a randomised phase 3 study. *Lancet Oncol*. 2009;10(11):1027–8.
- Sakuramoto S, Sasako M, Yamaguchi T, Kinoshita T, Fujii M, Nashimoto A, et al. Adjuvant chemotherapy for gastric cancer with S-1, an oral fluoropyrimidine. *N Engl J Med*. 2007;357:1810–20.
- Ajani JA, Faust J, Ikeda K, Yao JC, Anbe H, Carr KL, et al. Phase I pharmacokinetic study of S-1 plus cisplatin in patients with advanced gastric carcinoma. *J Clin Oncol*. 2005;23:6957–65.
- Koizumi W, Tanabe S, Saigenji K, Ohtsu A, Boku N, Nagashima F, et al. Phase I/II study of S-1 combined with cisplatin in patients with advanced gastric cancer. *Br J Cancer*. 2003;89:2207–12.
- Yoshida K, Ninomiya M, Takakura N, Hirabayashi N, Takiyama W, Sato Y, et al. Phase II study of docetaxel and S-1 combination therapy for advanced or recurrent gastric cancer. *Clin Cancer Res*. 2006;12:3402–7.
- Narahara H, Fujitani K, Takiuchi H, Sugimoto N, Inoue K, Uedo N, et al. Phase II study of a combination of S-1 and paclitaxel in patients with unresectable or metastatic gastric cancer. *Oncology*. 2008;74:37–41.
- Uedo N, Narahara H, Ishihara R, Takiuchi H, Goto M, Fujitani K, et al. Phase II study of a combination of irinotecan and S-1 in patients with advanced gastric cancer (OGSG0002). *Oncology*. 2007;73:65–71.
- Koizumi W, Narahara H, Hara T, Takagane A, Akiya T, Takagi M, et al. S-1 plus cisplatin versus S-1 alone for first-line treatment of advanced gastric cancer (SPIRITS trial): a phase III trial. *Lancet Oncol*. 2008;9:2.
- Takiuchi H, Narahara H, Tsujinaka T, Gotoh M, Kawabe S, Katsu K, et al. Phase I study of S-1 combined with irinotecan (CPT-11) in patients with advanced gastric cancer (OGSG 0002). *Jpn J Clin Oncol*. 2005;35:520–5.
- Therasse P, Arbuck SG, Eisenhauer EA, Wanders J, Kaplan RS, Rubinstein L, et al. New guidelines to evaluate the response to treatment in solid tumors. European Organization for Research and Treatment of Cancer, National Cancer Institute of the United States, National Cancer Institute of Canada. *J Natl Cancer Inst*. 2000;92:205–16.
- Inokuchi M, Yamashita T, Yamada H, Kojima K, Ichikawa W, Nihei Z, et al. Phase I/II study of S-1 combined with irinotecan for metastatic advanced gastric cancer. *Br J Cancer*. 2006;94:1130–5.
- Peters GJ, van der Wilt CL, van Moorsel CJ, Kroep JR, Bergman AM, Ackland SP. Basis for effective combination cancer chemotherapy with antimetabolites. *Pharmacol Ther*. 2000;87:227–53.
- Guichard S, Hennebelle I, Bugat R, Canal P. Cellular interactions of 5-fluorouracil and the camptothecin analogue CPT-11 (irinotecan) in a human colorectal carcinoma cell line. *Biochem Pharmacol*. 1998;55:667–76.
- Yamada Y, Yasui H, Goto A, Arai T, Ura T, Hamaguchi T, et al. Phase I study of irinotecan and S-1 combination therapy in patients with metastatic gastric cancer. *Int J Clin Oncol*. 2003;8:374–80.
- Komatsu Y, Yuki S, Miyagishima T, Asaka M. Irinotecan plus oral S-1 in patients with advanced gastric cancer biweekly IRIS regimen. *Gan To Kagaku Ryoho (Cancer and Chemotherapy)*. 2006;33(Suppl 1):131–4. (in Japanese).

28. Van Cutsem E, Kang Y, Chung H, Shen L, Sawaki A, Lordick F, et al. Efficacy results from the ToGA trial: a phase III study of trastuzumab added to standard chemotherapy (CT) in first-line human epidermal growth factor receptor 2 (HER2)-positive advanced gastric cancer (GC) [abstract]. *J Clin Oncol.* 2009;27(Suppl 18S):LBA4509.
29. Enzinger PC, Ilson DH, Saltz LB, O'Reilly EM, Kelsen DP. Irinotecan and cisplatin in upper gastrointestinal malignancies. *Oncology.* 1998;12:110–3.
30. Boku N, Ohtsu A, Shimada Y, Shirao K, Seki S, Saito H, et al. Phase II study of a combination of irinotecan and cisplatin against metastatic gastric cancer. *J Clin Oncol.* 1999;17:319–23.
31. Pozzo C, Barone C, Szanto J, Padi E, Peschel I, Bukki J, et al. Irinotecan in combination with 5-fluorouracil and folinic acid or with cisplatin in patients with advanced gastric or esophageal-gastric junction adenocarcinoma: results of a randomized phase II study. *Ann Oncol.* 2004;15:1773–81.
32. Thuss-Patience PC, Kretzschmar A, Deist T, Hinke A, Bichev D, Lebedinew B, et al. Irinotecan versus best supportive care (BSC) as second-line therapy in gastric cancer: a randomized phase III study of the Arbeitsgemeinschaft Internistische Onkologie (AIO) [abstract]. *J Clin Oncol.* 2009; 27 (Suppl 15S):4540.
33. Takahari D, Shimada Y, Takeshita S, Nishitani H, Takashima A, Okita N, et al. Second-line chemotherapy with irinotecan plus cisplatin after the failure of S-1 monotherapy for advanced gastric cancer. *Gastric Cancer.* 2010;13:186–90.
34. Sakamoto T, Yasui H, Boku N, Onozawa Y, Hironaka S, Fukutomi A, et al. Comparison of combination chemotherapy with irinotecan and cisplatin regimen administered every 2 or 4 weeks in pretreated patients with unresectable or recurrent gastric cancer: retrospective analysis. *Int J Clin Oncol.* 2010;15:287–93.
35. Ando Y, Fujita K, Sasaki Y, Hasegawa Y. UGT1A1*6 and UGT1A1*27 for individualized irinotecan chemotherapy. *Curr Opin Mol Ther.* 2007;9:258–62.