

Research article

The fission yeast COP9/signalosome is involved in cullin modification by ubiquitin-related Ned8p

Chunshui Zhou¹, Volker Seibert¹, Rory Geyer¹, Edward Rhee¹, Svetlana Lyapina², Greg Cope², Raymond J Deshaies^{2,3} and Dieter A Wolf^{f*1}Address: ¹Department of Cancer Cell Biology, Harvard School of Public Health, Boston, MA, USA, ²Division of Biology, California Institute of Technology, CA, USA and ³Howard Hughes Medical Institute, California Institute of Technology, CA, USAE-mail: Chunshui Zhou - czhou@hsph.harvard.edu; Volker Seibert - vseibert@hsph.harvard.edu; Rory Geyer - erhee@hsph.harvard.edu; Edward Rhee - rgeyer@hsph.harvard.edu; Svetlana Lyapina - lyapina@cco.caltech.edu; Greg Cope - cope@caltech.edu; Raymond J Deshaies - dshaies@caltech.edu; Dieter A Wolf* - dewolf@hsph.harvard.edu

*Corresponding author

Published: 18 July 2001

Received: 3 May 2001

BMC Biochemistry 2001, 2:7

Accepted: 18 July 2001

This article is available from: <http://www.biomedcentral.com/1471-2091/2/7>© 2001 Zhou et al; licensee BioMed Central Ltd. Verbatim copying and redistribution of this article are permitted in any medium for any non-commercial purpose, provided this notice is preserved along with the article's original URL. For commercial use, contact info@biomedcentral.com

Abstract

Background: The function of the fission yeast cullins Pcu1p and Pcu4p requires modification by the ubiquitin-related peptide Ned8p. A recent report by Lyapina *et al.* shows that the COP9/signalosome (CSN), a multifunctional eight subunit complex, regulates Ned8p modification of Pcu1p. Disruption of *caal/csn1*, which encodes subunit 1 of the putative *S. pombe* CSN, results in accumulation of Pcu1p exclusively in the modified form. However, it remained unclear whether this reflects global control of all cullins by the entire CSN complex.

Results: We demonstrate that multiple CSN subunits control Ned8p modification of Pcu3p, another fission yeast cullin, which, in complex with the RING domain protein Pip1p, forms a ubiquitin ligase that functions in cellular stress response. Pcu3p is modified by Ned8p on Lys 729 and accumulates exclusively in the neddylated form in cells lacking the CSN subunits 1, 3, 4, and 5. These CSN subunits co-elute with Pcu3p in gel filtration fractions corresponding to ~ 550 kDa and specifically bind both native and Ned8p-modified Pcu3p *in vivo*. While CSN does not influence the subcellular localization of Pcu3p, Pcu3p-associated *in vitro* ubiquitin ligase activity is stimulated in the absence of CSN.

Conclusions: Taken together, our data suggest that CSN is a global regulator of Ned8p modification of multiple cullins and potentially other proteins involved in cellular regulation.

Background

The cullin/RING family of ubiquitin ligases comprises a multifunctional set of enzymes controlling the stability and activity of cell cycle regulators, transcription factors, and signaling proteins (reviewed in ref. [1]). Among the best studied examples of cullin/RING enzymes are the so-called SCF complexes. SCF complexes share homo-

logues of the core components cullin 1 (CUL1), SKP1, and the RING domain protein HRT1/RBX1/ROC1, which associate with different F-box proteins and ubiquitin-conjugating enzymes [1]. F-box proteins specifically bind substrates, following their phosphorylation in response to activation of various signaling pathways. Although substrate phosphorylation is one major determinant of

SCF-mediated ubiquitylation [1], covalent modification of the CUL1 subunit through attachment of the ubiquitin-related peptide NEDD8 also regulates SCF activity. NEDD8 modification stimulates SCF-dependent substrate ubiquitylation *in vitro* [2,3,4], and mutant CUL1 resistant to NEDD8 modification is defective *in vivo* [5]. In addition, fission yeast mutants deficient in the enzymes that attach Ned8p, or in the *ned8* gene itself, are inviable [5].

In addition to CUL1, the human genome contains at least five other cullins, CUL 2, 3, 4A, 4B, and 5. This diversity is partially recapitulated in *S. pombe*, which encodes *pcu3* and *pcu4*, homologues of human CUL3 and CUL4, for which there are no direct correlates in budding yeast [6]. All human cullins interact with HRT1/RBX1/ROC1 [7], are modified by NEDD8 [8], and have ubiquitin ligase activity *in vitro* [9]. Similarly, the fission yeast cullins Pcu1p and Pcu4p are neddylation, and neddylation is critical for their function *in vivo*. [5].

While it is clear that neddylation affects the activity of cullin/RING complexes, mechanisms controlling this modification are just emerging. A recent study by Lyapina *et al.* uncovered a novel connection between the COP9/signalosome (CSN) and cullin neddylation [10]. CSN is a highly conserved multiprotein complex consisting of eight subunits, which have been implicated in a wide variety of regulatory processes, including, cell cycle control [11, 12], signal transduction [13], transcriptional activation [14], and plant photomorphogenesis [15, 16]. To date, no distinct biochemical activity has been identified that can reconcile this diversity in CSN functions. The eight CSN subunits show a one-to-one relationship to components of the 19S proteasome lid complex [17,18,19] and, like *pcu3* and *pcu4*, at least seven of the eight human subunits are conserved in fission yeast, but not in budding yeast. [16, 20].

Lyapina *et al.* recently showed that disruption of *S. pombe* CSN subunit 1 (*caa1/csn1*) results in accumulation of Pcu1p in the neddylation state [10]. However, it remained unclear whether this reflects global control of multiple cullins by Csn1p as part of the entire CSN complex. We have examined this question by determining the effect of disruption of several CSN subunits on Pcu3p neddylation, localization, and ubiquitin ligase activity. Our results indicate that the entire CSN complex is required for proper control of the neddylation state of multiple cullins.

Results

Pcu3p is modified by Ned8p

Immunoblotting of protein lysates prepared from *S. pombe* cells, whose *pcu3* chromosomal locus was modi-

fied to encode Pcu3p with thirteen copies of the Myc epitope attached to its C-terminus, revealed two distinct species of Pcu3p, which migrated on SDS gels as two closely spaced species of roughly equal abundance (Fig. 1A). Several findings indicate that the slower migrating form of Pcu3p arose from covalent modification with the ubiquitin-related protein Ned8p: First, overexpression of a fusion protein of Ned8p and glutathione transferase (GST-Ned8p) converted the faster, unmodified form of Pcu3p into a high molecular weight species that reacted with antibodies to GST (Fig. 1A). Second, replacing lysine 729, which is homologous to the conserved neddylation sites in other cullins, with arginine, completely abolished the slower migrating, modified form of Pcu3p (Fig. 1B). In contrast, mutation of a neighboring conserved lysine (residue 760) did not alter the mobility of Pcu3p. Thus, as with other *S. pombe* cullins [5], Pcu3p undergoes covalent modification by ubiquitin-like Ned8p at lysine 729.

Multiple *csn* mutants accumulate neddylation Pcu3p

Interestingly, fission yeast *caa1/csn1* deletion mutants show a set of defects, which partially overlaps that found in *pcu3* disruptants: slow growth, cell elongation, and sensitivity to UV-irradiation, but not hydroxyurea [6, 21]. With respect to the recent implication of CSN in cullin regulation [10], we considered the possibility that the overlapping phenotypes of *csn1* and *pcu3* mutants could arise from interference with Pcu3p function.

Initial support for this hypothesis was provided by the observation that disruption of *csn1* led to accumulation of Pcu3p in the slower migrating form (Fig. 2A). This shift in migration is due to Pcu3p neddylation as shown by overexpression of GST-Ned8p in *csn1* mutants. This caused an additional supershift, converting a substantial portion of the slower migrating form into a GST-Ned8p modified species (Fig. 2B). Importantly, accumulation of neddylation Pcu3p was also apparent in strains deleted for *csn3*, *csn4*, and *csn5* (Fig. 2A), three genes encoding other putative subunits of the *S. pombe* CSN, which cofractionate with Csn1p upon size exclusion chromatography (see below, Fig. 4). However, unlike *csn1* disruptants, which have a defect in S phase control resulting in slow growth and cell elongation [21], deletion of *csn3*, *csn4*, and *csn5* genes did not cause cell elongation, S phase delay, or any other obvious growth defects (Fig. 2C, 2D, data not shown). This indicates that the described phenotype of *csn1* mutants [21] is unlikely to be a consequence of interference with Pcu3p activity. Pcu1p, another fission yeast cullin also accumulated in the neddylation form in all *csn* mutants [10] (Fig. 2A), suggesting that CSN has a general role in the control of cullin neddylation in fission yeast. The accumulation of neddylation Pcu3p was prevented in all *csn* mutants by providing the missing CSN subunit on a plasmid (Fig. 3).




Figure 1
 Pcu3p is modified by Ned8p on lysine 729. **(a)** Cells containing Pcu3p tagged with 13 Myc epitope tags at the C-terminus (*pcu3-13myc* cells) were transformed with a pRep3-based plasmid driving the overexpression of a fusion of GST with *S. pombe* Ned8p. Pcu3p-13Myc was immunoprecipitated and precipitates were analyzed for the presence of Pcu3p and GST-Ned8p with the antibodies indicated. The lower panel shows GST-Ned8p expression in the total cell lysates. **(b)** N-terminally Myc epitope-tagged wild-type Pcu3p or point mutants where lysines 729 and 760 were replaced by arginine were expressed from pRep81. Cell lysates were blotted with Myc antibodies to reveal the neddylated and unmodified species of Pcu3p. The comparatively low level of Pcu3p neddylated in this experiment may be attributable to overexpression.





Figure 2
 Neddylated Pc3p accumulates in *csn* mutants. **(a)** The indicated *csn* genes were disrupted in a *pcu3-13myc* background. Cell lysates of these strains were probed with antibodies against Myc and Pc1p to reveal neddylated and unmodified forms. **(b)** GST-Ned8p was overexpressed in *csn1* mutants, and modified Pc3p was detected by immunoblotting with Myc antibodies. The star denotes an unidentified cross-reactive band. **(c)** Morphology of *csn* mutants. The indicated strains were fixed in formaldehyde, mounted on microscope slides, and photographed. The elongation phenotype of *csn1* mutants is not apparent in other *csn* mutants. **(d)** DNA content of *csn* mutants. The indicated strains were grown to mid log phase, fixed in ethanol, and analyzed by flow cytometry. Note that the broadening of the 2C peak indicative of the described slow S phase progression of *csn1* mutants [21] is completely absent from other *csn* mutants.

**Figure 3**

Complementation of *csn* mutants. The indicated *csn* mutants in a *pcu3-13myc* background were transformed with pRep81 plasmids driving the expression of the respective 6 × histidine-Myc-tagged *csn* genes. Cells were maintained in the presence (promoter off) or absence of thiamin (promoter on) to regulate expression. Expression of endogenous Pcu3p-13Myc **(a)** and exogenous CSN subunits **(b)** was determined by immunoblotting with Myc antibodies. The weak complementing activity of some of the plasmids even in the repressed state may result from leakiness of the *nmt1* promoter [34].

The conserved "cysteine box" in *Csn5p* is not required for *Pcu3p* deneddylation

Lyapina *et al.* recently demonstrated that partially purified CSN promotes the cleavage of Pcu1p-Ned8p conjugates *in vitro* [10], but it remained unclear whether the enzymatic activity is contained within one of the CSN subunits or a tightly associated peptidase. All peptidases known to cleave ubiquitin- or SUMO-protein conjugates are cysteine-based proteases (reviewed in ref. [22]). *Csn5p* contains a motif closely resembling the conserved cysteine box previously recognized in the catalytic center of deubiquitylating enzymes and in the budding yeast proteasome subunit Rpn11p [23]. The complementation assay shown in Fig. 3 allowed us to determine the potential role of this motif in the control of Pcu3p neddylation

by replacing the critical cysteine residue with alanine. This mutant *Csn5p* retained full activity for preventing Pcu1p and Pcu3p hyperneddylation (Fig. 3A lanes 12-15, and data not shown).

Pcu3p interacts with CSN

We next determined whether the effect of CSN deficiency on Pcu3p neddylation reflects a physical interaction of these proteins. Size exclusion chromatography of lysates prepared from cells harboring epitope-tagged Pcu3p or CSN subunits at the endogenous genomic loci, revealed precise co-elution of Pcu3p with CSN subunits 1, 3, 4, and 5 in high molecular weight fractions corresponding to approximately 550 kDa (Fig. 4A). Interestingly, *Csn5p*, unlike the other CSN subunits, showed a second




Figure 4

Pcú3p interacts with CSN. **(a)** Lysates prepared from cells containing Myc epitope-modified Pcú3p, Csn3p, Csn4p, and Csn5p, or HA-tagged Csn1p were fractionated by gel filtration. Fractions were probed with Myc and HA antibodies to determine the elution profiles of the respective proteins. Size standards are indicated. Pcú3p-Myc was not resolved into the neddylated and deneddylated forms on this gel. **(b)** Lysate from *csn1-3HA* and *pcú3-13myc csn1-3HA* cells, which carry epitope-tagged versions of Pcú3p and Csn1p at the endogenous genomic locus, were immunoprecipitated with Myc antibodies followed by immunoblot analysis with Myc and HA antibodies. Co-precipitated Pcú3p-13Myc and Csn1p-3HA are indicated. Specificity controls for the immunoprecipitation include an irrelevant antibody (left panel) and a strain lacking Pcú3p-Myc (right panel). **(c)** The same strains as used in Fig. 3A were induced to express the indicated 6 × histidine-Myc-tagged CSN subunits. CSN subunits were absorbed on Ni-NTA agarose beads, and co-purification of Pcú3p-13Myc was determined by immunoblotting with Myc antibodies. In the control lane, lysate from nontransformed *pcú3-13myc* cells was used. Note that Csn3p, while inefficiently expressed in the total cell lysate, is equally retained on Ni-NTA beads, probably due to better accessibility of the 6 × histidine tag.

peak in gel filtration eluting with an approximate molecular weight of 200 kDa, suggesting that Csn5p may also be involved in protein complexes separate from the entire CSN (Fig. 4B).

Based on the co-elution of Pcu3p with CSN subunits in gel filtration, we determined whether Pcu3p and Csn1p interact *in vivo* at endogenous expression levels. Pcu3-Myc immunoprecipitated from lysates of cells expressing HA-tagged Csn1p together with Pcu3p-Myc from the endogenous promoters revealed co-precipitation of Csn1p-HA. This interaction was specific as it was not observed with an irrelevant antibody (Fig. 4C left panel) or with lysate from a strain containing untagged Pcu3p (Fig. 4C, right panel). In contrast, immunoprecipitation with HA antibodies failed to precipitate both Csn1p-3HA and Pcu3p-Myc, potentially due to inaccessibility of the C-terminal 3HA tag of Csn1p within the CSN complex (data not shown).

We therefore generated Pcu3p-Myc containing strains overexpressing N-terminally 6xhistidine-Myc-tagged Csn1p, Csn3p, Csn4p, or Csn5p. Binding of each of the four different CSN subunits to nickel beads resulted in co-purification of Pcu3p, indicative of specific *in vivo* interactions (Fig. 4C). Notably, both modified and unmodified Pcu3p interacted with CSN subunits (Fig. 4C). We conclude that, at least a fraction of Pcu3p forms a stable complex with CSN *in vivo*.

Pcu3p localization in *csn* mutants

In *A. thaliana*, CSN was shown to be required for the efficient nuclear accumulation of the putative RING-type

ubiquitin ligase COP1 [24], which mediates ubiquitin-dependent degradation of the photomorphogenic transcription factor HY5 [25]. We therefore considered the possibility that accumulation of neddylated Pcu3p in *csn* mutants is a consequence of defective subcellular targeting. Indirect immunofluorescence staining indicated that endogenous Myc-tagged Pcu3p is present in the cytoplasm, but enriched in the nucleus of wild-type cells (Fig. 5). This localization pattern was fully preserved in all four *csn* mutants, indicating that CSN does not regulate the subcellular localization of Pcu3p.

Mutation of CSN increases Pcu3p-dependent polyubiquitylation activity

All human cullins have been shown to assemble into complexes possessing polyubiquitylation activity *in vitro* [9, 26]. To determine whether Pcu3p is associated with such an activity and how it is affected by CSN, we developed a substrate-independent *in vitro* polyubiquitylation assay, similar to that originally reported by Lyapina et al. [27]. As Pcu3p-associated ubiquitin ligase activity was not described before, we first needed to determine which of the fourteen different ubiquitin-conjugating enzymes (UBCs) present in the fission yeast genome associate with the RING domain protein Pip1p, an essential cofactor of cullin ubiquitin ligases [7, 28,29,30]. Binding assays showed that overexpressed HA-tagged Pip1p specifically associated with co-overexpressed 6xHis-Myc-tagged Ubc1p, Ubc7p, Ubc11p, Ubc12p, Ubc13p, Ubc14p, and Mms2p. In the next step, we confirmed, by immunoprecipitation, that both forms of Pcu3p-Myc bound endogenous Pip1p in a CSN-independent manner (Fig. 6B).




Figure 5

Localization of Pcu3p in *csn* mutants. The indicated strains were fixed and processed for indirect immunofluorescence staining of Pcu3p-Myc with Myc antibodies as described in material and methods.




Figure 6

Pcu3p in vitro ubiquitylation activity. (a) Each of the 14 *S. pombe* UBCs were co-overexpressed as 6 × His-Myc-tagged versions with HA-tagged Pip1p from plasmids. UBCs were bound to Ni-agarose beads and co-purified Pip1p was detected by HA antibodies. (b) Lysates prepared from *pcu3-13myc* and *pcu3-13myc Δcsn1* cells were immunoprecipitated with affinity-purified antisera against Pip1p and precipitates were analyzed for coprecipitation of Pcu3p by immunoblotting with Myc antibodies. (c) Pcu3p-associated ubiquitin ligase activity. Pcu3p complexes were immunoprecipitated from the indicated strains with Myc antibodies and polyubiquitylation assays were performed as described in materials and methods using the UBCs indicated ("3" denotes human UBC3/CDC34). Total lysates used for immunoprecipitation and the resulting Pcu3p immunocomplexes are shown in the lower panels. Pcu3p-Myc is not clearly resolved into the neddylated and unneddylated forms in these gels. (d) Ubiquitylation assay with Pcu3p-Myc complexes isolated from the indicated strains were performed with wild-type ubiquitin or a mutant lacking all lysines (Ub-R).

Pcu3p-Myc complexes prepared by immunoprecipitation with Myc antibodies were then incubated with human E1, ATP, ubiquitin, and the recombinant UBCs that interacted with Pip1p. Polyubiquitylated species were visualized by anti-ubiquitin blot. Pcu3p-Myc complexes isolated from *pcu3-13myc* cells did not show any recognizable polyubiquitylation activity above background with any of the seven UBCs tested (Fig. 6C, lane 2, and data not shown). In contrast, the ubiquitylation assay with Pcu3p-Myc complexes purified from *csn5* mutants led to efficient formation of high molecular weight species reactive with ubiquitin antibodies (Fig. 6C). This activity was specific for Ubc7p (Fig. 6C, compare lanes 3 - 7). Appearance of multiubiquitin chains was inhibited by mutant ubiquitin lacking all lysines, indicating that they were specifically generated in the reaction rather than co-precipitated with Pcu3p-Myc (Fig. 6D). Thus, CSN deficiency appears to stimulate Pcu3p-associated ubiquitin ligase activity *in vitro*. A similar stimulation was obtained for substrate-specific polyubiquitylation of phosphorylated Rum1p by SCF^{POP} purified from *csn5* mutants (unpublished observation).

Discussion

Our data confirm and extend the recently discovered connection between CSN and cullin regulation [10] by showing that deletion of four putative CSN subunits causes accumulation of Pcu3p in the neddylation state (Fig. 2A). Co-elution of these four proteins in a 550 kDa complex (Fig. 4B) strongly suggests that they represent part of the *S. pombe* equivalent of the eight subunit CSN complex found in higher eukaryotes [18,19,20]. Thus, cullin regulation through modulation of the balance of Ned8p modification appears to require the entire CSN complex.

Our study also provides some insight into the role of deneddylation in cullin function, as we were able to create a situation in which Pcu1p and Pcu3p are fully neddylation *in vivo* (ref [10] and data presented here). Deletion of *pcu1* and *pcu3* results either in lethality or in severe growth retardation, respectively [6]. The normal growth behavior of *csn3*, *4*, and *5* mutants (Fig. 2C,2D), where both cullins seem constitutively neddylation, argues against a critical role for dynamic cycles of neddylation and deneddylation in cullin function in fission yeast. Consistent with this notion, the subcellular localization of Pcu3p and Pcu1p were unaffected by mutations in CSN (Fig. 5) [10]. Instead, neddylation Pcu3p and Pcu1p purified from *csn* mutant cell extract exhibit increased ubiquitin ligase activity *in vitro* (Fig. 6B) [10]. Thus, CSN may exert negative control on Pcu1p and Pcu3p by promoting their deneddylation. However, while mutation of *csn5* caused a modest two-fold increase in the steady state fraction of neddylation Pcu3p (Fig. 2A), Pcu3p-associated

ubiquitin ligase activity recovered from this mutant showed a much more dramatic increase (Fig. 6C). This suggests the possibility that CSN represses Pcu3p activity by other mechanisms, in addition to its control of neddylation. Further experiments are required to determine whether recombinant Csn5p or purified CSN complex can inhibit Pcu3p-associated ubiquitin ligase activity.

Lyapina *et al.* recently provided evidence for a cullin deneddylation activity of CSN by demonstrating that a partially purified preparation of the pig CSN can deneddylation the fission yeast cullin Pcu1p *in vitro* [10]. The results presented here eliminate the attractive hypothesis that the cysteine box in Csn5p, which is conserved in numerous deubiquitylation enzymes [23], harbors a critical activity for deneddylation (Fig. 3A). In addition, recombinant Csn5p failed to deneddylation Pcu1p and Pcu3p *in vitro* (unpublished observation). Our finding that hyperneddylation Pcu3p and Pcu1p accumulate in all *csn* mutants tested (Fig. 2A, and data not shown), also suggests that control of cullin neddylation results from cooperation of multiple CSN components rather than from the activity of a single catalytic subunit. The CSN complex may, for example, serve as a platform for recruitment of a separate deneddylation enzyme. A human NEDD8-processing isopeptidase has been described previously [31]. The potential role of CSN as an organizing center for certain enzymatic activities may not be restricted to the control of cullin neddylation, as CSN was also shown to mediate phosphorylation of c-JUN, IκBa, and p53 by a tightly associated kinase(s) [12, 19].

So far, we were unable to obtain deneddylation of Pcu3p-13Myc *in vitro* by partially purified CSN preparations active in deneddylation of Pcu1p-13Myc (data not shown). We can therefore not exclude that CSN's role in the control of Pcu3p neddylation differs from its role established for Pcu1p [10]. For example, binding by CSN could protect Pcu3p from the Ned8p-conjugating enzyme Ubc12p, thus maintaining a fraction of Pcu3p in a deneddylation state. As this shield is experimentally removed through deletion of CSN subunits, Pcu3p may accumulate in the neddylation state. In such a scenario, no enzymatic activity of purified CSN toward neddylation Pcu3p would be expected *in vitro*.

Finally, our data suggest CSN functions in addition to cullin regulation that are not shared by all individual subunits. Out of the four mutants tested, only *csn1* showed the previously described defect in S phase progression [21] (Fig. 2C,2D), thereby underscoring the significance of our finding that all CSN subunits tested here are involved in cullin regulation. It is likely that some subunits have very specialized functions within the complex, while others are required for its broader activities

such as control of cullin neddylation. On the other hand, we found that a sizable fraction of Csn5p is dissociated from the putative CSN holocomplex, but enriched in a 200 kDa assembly (Fig. 4B). It thus appears that part of the diversification of CSN function may arise from formation of smaller subcomplexes.

Conclusions

Our study reveals a general role of the *S. pombe* CSN complex in the control of cullin modification by Ned8p. Regardless of the exact mechanism, CSN may regulate the activity of multiple cullin ubiquitin ligases through counteracting their stimulatory Ned8p modification. Future studies will reveal whether some of the other described functions of CSN also involve post-translational modification of regulatory proteins by ubiquitin-like modifiers.

Materials and methods

Plasmids and yeast strains

S. pombe genes for UBCs and *csn3*, *4*, and *5* were identified in the Sanger Centre *S. pombe* sequence database based on their homology to the respective human and budding yeast proteins. Complementary DNAs were amplified by PCR and subcloned into pRep81.6xHis-Myc, pRep3.6xHis-Myc, or pRep4.HA. Deletion strains and epitope-tagged strains were constructed by one-step gene replacement using PCR-generated fragments containing kanamycin or *ura4* cassettes [32]. Growth media, flow cytometry, and all other relevant *S. pombe* techniques were described previously [33].

Immunological techniques

Protein lysates were prepared by bead lysis in a Fastprep device (Bio 101) in the presence of proteinase inhibitors, followed by boiling in SDS sample buffer. Pip1p and Pcup were detected with affinity-purified rabbit antisera. Epitope-tagged proteins were detected by the monoclonal antibodies 9E10 and 12CA5 prepared from hybridoma supernatants.

Cell lysates for immunoprecipitation were prepared by disrupting cells in immunoprecipitation buffer (20 mM Tris/HCl, pH 7.4; 150 mM NaCl; 0.5 % Triton X-100, 10 µg/ml leupeptin, 10 µg/ml pepstatin, 5 µg/ml aprotinin, 1 mM PMSF). Lysates were cleared and precipitated with the respective antisera. Immunocomplexes were collected by binding to protein A or G beads, washed and analyzed by immunoblotting as described [33].

For indirect immunofluorescence staining, cells were fixed in freshly prepared 3.7% para-formaldehyde for 1h at 30°C. Cells were washed in PEM (100 mM Pipes pH6.9, 1 mM EGTA, 1 mM MgSO₄), and digested with Novozyme (0.5 mg/ml, Sigma) and Zymolyase (0.5 mg/

ml, US Biological) in PEM, 1M sorbitol. Unspecific epitopes were blocked by incubation in PEM-BAL (PEM plus 1% BSA, 0.1% Na-azide, 0.1 M L-lysine HCl) for 1h. Incubation with monoclonal Myc antibodies diluted in PEM-BAL was done overnight, followed by three washes in PEM-BAL. An overnight incubation with TRITC-labeled anti mouse antibodies was followed by washing and mounting on polylysine coated coverslips. Cell nuclei were counterstained with DAPI. Photomicrographs were obtained with a CCD camera mounted on a Nikon E600 epifluorescence microscope. Brightness and contrast were adjusted in Adobe Photoshop 5.0.

In vitro ubiquitylation assay

For ubiquitylation reactions, Pcu3p-Myc complexes were immunoprecipitated from 100 - 200 µg total cell lysates prepared as described above. Precipitates were washed four times in 20 mM Tris/HCl, pH 7.4; 150 mM NaCl; 0.5 % Triton X-100, 10 µg/ml leupeptin, 10 µg/ml pepstatin, 5 µg/ml aprotinin, 1 mM PMS, and equilibrated in 20 mM HEPES, pH 7.4, 100 mM potassium acetate, 1 mM DTT. The reaction was started by the addition of a cocktail of 8 µM ubiquitin, ATP regenerating system (2 mM HEPES at pH 7.4, 1 mM ATP, 30 mM creatine phosphate, 1 mM magnesium acetate, 0.15 mg/ml creatine kinase), reaction buffer (4 mM magnesium acetate, 1 mM DTT), 500 nM bacterially expressed 6xHis-UBCs, 100 nM 6xHis-tagged human E1, and 0.5 µM ubiquitin aldehyde in a volume of 15 µl. After 90 min at 30°C, the reaction was terminated by the addition 5 × SDS sample buffer. Samples were separated on 11 % SDS-polyacrylamide gels and analyzed by immunoblotting with ubiquitin antibodies (Zymed).

Abbreviations

CSN: COP9/signalosome

GST: Glutathione S-transferase

HA: hemagglutinin

SCF: SKP1/Cullin/F-box protein complex

UBC: Ubiquitin-conjugating enzyme

Acknowledgments

We thank F. McKeon for hybridoma lines, R. King for the human E1 plasmid, D. Griffiths for numerous technical suggestions, and C. Maki and Z. Yuan for critical reading of this manuscript. This work is supported by NIH grant GM59780 to DAW.

References

1. Deshaies RJ: **SCF and Cullin/Ring H2-based ubiquitin ligases** *Annu Rev Cell Dev Biol* 1999, **15**:435-467
2. Read MA, Brownell JE, Gladysheva TB, Hottelet M, Parent LA, Coggin MB, et al: **Nedd8 modification of cul-1 activates SCF-beta-TrCP-dependent ubiquitination of IkappaBalpha** *Mol Cell Biol* 2000, **20**:2326-2333

3. Wu K, Chen A, Pan ZQ: **Conjugation of Nedd8 to CUL1 Enhances the Ability of the ROC1-CUL1 Complex to Promote Ubiquitin Polymerization** *J Biol Chem* 2000, **275**:32317-32324
4. Podust VN, Brownell JE, Gladysheva TB, Luo RS, Wang C, Coggins MB, et al: **A Nedd8 conjugation pathway is essential for proteolytic targeting of p27Kip1 by ubiquitination** *Proc Natl Acad Sci U S A* 2000, **97**:4579-4584
5. Osaka F, Saeki M, Katayama S, Aida N, Toh EA, Kominami K, et al: **Covalent modifier NEDD8 is essential for SCF ubiquitin-ligase in fission yeast** *Embo J* 2000, **19**:3475-3484
6. Kominami K, Ochotorena I, Toda T: **Two F-box/WD-repeat proteins Pop1 and Pop2 form hetero- and homo-complexes together with cullin-1 in the fission yeast SCF (Skp1-Cullin-1-F-box) ubiquitin ligase** *Genes Cells* 1998, **3**:721-735
7. Ohta T, Michel JJ, Schottelius AJ, Xiong Y: **ROC1, a homolog of APC11, represents a family of cullin partners with an associated ubiquitin ligase activity** *Mol Cell* 1999, **3**:535-541
8. Hori T, Osaka F, Chiba T, Miyamoto C, Okabayashi K, Shimbara N, et al: **Covalent modification of all members of human cullin family proteins by NEDD8** *Oncogene* 1999, **18**:6829-6834
9. Ohta T, Xiong Y: **Phosphorylation- and Skp1-independent in vitro ubiquitination of E2F1 by multiple ROC-cullin ligases** *Cancer Res* 2001, **61**:1347-1353
10. Lyapina S, Cope G, Shevchenko A, Serino G, Tsuge T, Zhou C, et al: **Promotion of NEDD8-CUL1 conjugate cleavage by COP9 signalosome** *Science* 2001, **292**:1382-1385
11. Tomoda K: **Degradation of the cyclin-dependent-kinase inhibitor p27Kip1 is instigated by Jab1** *Nature* 1999, **398**:160-165
12. Bech-Otschir D, Kraft R, Huang X, Henklein P, Kapelari B, Pollmann C, et al: **COP9 signalosome-specific phosphorylation targets p53 to degradation by the ubiquitin system** *Embo J* 2001, **20**:1630-1639
13. Kleemann R, Hausser A, Geiger G, Mischke R, Burger-Kentscher A, Flieger O, et al: **Intracellular action of the cytokine MIF to modulate AP-1 activity and the cell cycle through Jab1** *Nature* 2000, **408**:211-216
14. Claret FX, Hibi M, Dhut S, Toda T, Karin M: **A new group of conserved coactivators that increase the specificity of AP-1 transcription factors** *Nature* 1996, **383**:453-457
15. Osterlund MT, Ang LH, Deng XW: **The role of COP1 in repression of Arabidopsis photomorphogenic development** *Trends Cell Biol* 1999, **9**:113-118
16. Wei N, Deng XW: **Making sense of the COP9 signalosome. A regulatory protein complex conserved from Arabidopsis to human** *Trends Genet* 1999, **15**:98-103
17. Glickman MH, Rubin DM, Coux O, Wefes I, Pfeifer G, Cjeka Z, et al: **A subcomplex of the proteasome regulatory particle required for ubiquitin-conjugate degradation and related to the COP9-signalosome and eIF3** *Cell* 1998, **94**:615-623
18. Wei N, Tsuge T, Serino G, Dohmae N, Takio K, Matsui M, et al: **The COP9 complex is conserved between plants and mammals and is related to the 26S proteasome regulatory complex** *Curr Biol* 1998, **8**:919-922
19. Seeger M, Kraft R, Ferrell K, Bech-Otschir D, Dumdey R, Schade R, et al: **A novel protein complex involved in signal transduction possessing similarities to 26S proteasome subunits** *Faseb J* 1998, **12**:469-478
20. Deng XW, Dubiel W, Wei N, Hofmann K, Mundt K, Colicelli J, et al: **Unified nomenclature for the COP9 signalosome and its subunits: an essential regulator of development** *Trends Genet* 2000, **16**:289
21. Mundt KE, Porte J, Murray JM, Brikos C, Christensen PU, Caspari T, et al: **The COP9/signalosome complex is conserved in fission yeast and has a role in S phase** *Curr Biol* 1999, **9**:1427-1430
22. Hochstrasser M: **Ubiquitin-dependent protein degradation** *Annu Rev Genet* 1996, **30**:405-439
23. Glickman MH, Rubin DM, Fried VA, Finley D: **The regulatory particle of the Saccharomyces cerevisiae proteasome** *Mol Cell Biol* 1998, **18**:3149-3162
24. Chamovitz DA, Wei N, Osterlund MT, von Arnim AG, Staub JM, Matsui M, et al: **The COP9 complex, a novel multisubunit nuclear regulator involved in light control of a plant developmental switch** *Cell* 1996, **86**:115-121
25. Osterlund MT, Hardtke CS, Wei N, Deng XW: **Targeted destabilization of HY5 during light-regulated development of Arabidopsis** *Nature* 2000, **405**:462-466
26. Lisztwan J, Imbert G, Wirbelauer C, Gstaiger M, Krek W: **The von Hippel-Lindau tumor suppressor protein is a component of an E3 ubiquitin-protein ligase activity** *Genes Dev* 1999, **13**:1822-1833
27. Lyapina SA, Correll CC, Kipreos ET, Deshaies RJ: **Human CUL1 forms an evolutionarily conserved ubiquitin ligase complex (SCF) with SKP1 and an F-box protein** *Proc Natl Acad Sci U S A* 1998, **95**:7451-7456
28. Kamura T, Koepp DM, Conrad MN, Skowyra D, Moreland RJ, Iliopoulos O, et al: **Rbx1, a component of the VHL tumor suppressor complex and SCF ubiquitin ligase** *Science* 1999, **284**:657-661
29. Skowyra D, Koepp DM, Kamura T, Conrad MN, Conaway RC, Conaway JW, et al: **Reconstitution of G1 cyclin ubiquitination with complexes containing SCFGrr1 and rbx1** *Science* 1999, **284**:662-665
30. Seol JH, Feldman RM, Zachariae W, Shevchenko A, Correll CC, Lyapina S, et al: **Cdc53/cullin and the essential Hrt1 RING-H2 subunit of SCF define a ubiquitin ligase module that activates the E2 enzyme Cdc34** *Genes Dev* 1999, **13**:1614-1626
31. Gong L, Kamitani T, Millas S, Yeh ET: **Identification of a novel isopeptidase with dual specificity for ubiquitin- and NEDD8-conjugated proteins** *J Biol Chem* 2000, **275**:14212-14216
32. Bahler J, Wu JQ, Longtine MS, Shah NG, McKenzie A, Steever AB 3rd, et al: **Heterologous modules for efficient and versatile PCR-based gene targeting in Schizosaccharomyces pombe** *Yeast* 1998, **14**:943-951
33. Wolf DA, McKeon F, Jackson PK: **F-box/WD-repeat proteins Pop1p and Sud1p/Pop2p form complexes that bind and direct the proteolysis of Cdc18p** *Curr Biol* 1999, **9**:373-376
34. Forsburg SL: **Comparison of Schizosaccharomyces pombe expression systems** *Nucleic Acids Res* 1993, **21**:2955-2956

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMedCentral will be the most significant development for disseminating the results of biomedical research in our lifetime."

Paul Nurse, Director-General, Imperial Cancer Research Fund

Publish with **BMC** and your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours - you keep the copyright



Submit your manuscript here:

<http://www.biomedcentral.com/manuscript/>

editorial@biomedcentral.com