

RESEARCH

Open Access



# Eutectic mixture of local anaesthetics (EMLA<sup>®</sup>) as a primary dressing on painful chronic leg ulcers: a pilot randomised controlled trial

Anne Purcell<sup>1,2,3\*</sup>, Thomas Buckley<sup>4</sup>, Jennie King<sup>3,4</sup>, Wendy Moyle<sup>1,2</sup> and Andrea P. Marshall<sup>1,2,5,6</sup>

## Abstract

**Background:** The physical, occupational, social and psychological impact of chronic leg ulcers (CLUs) on an individual is considerable. Wound-related pain (WRP), the most common symptom, is frequently reported as moderate to severe and mostly occurs at dressing change. WRP pain may not be alleviated by oral analgesics alone. Persistent poorly controlled leg ulcer pain can negatively impact wound healing and health-related quality of life (HRQoL).

**Methods:** A pilot, parallel group, non-blinded, randomised controlled trial was conducted in six procedure clinics located in a public community nursing service in New South Wales, Australia to evaluate eutectic mixture of local anaesthetics (EMLA<sup>®</sup>) on painful CLUs when used as a primary dressing. The primary objective was to assess feasibility by using pre-determined criteria: at least 80% recruitment rate, 80% retention rate and 80% adherence to the study protocol. Key eligibility criteria were that participants had a painful CLU no larger than 100 cm<sup>2</sup>, a numerical rating scale (NRS) wound-related pain intensity score equal to or greater than 4, low to moderate exudate, no contraindications to EMLA<sup>®</sup> and capacity to consent. One hundred and seven patients with painful CLUs were screened for eligibility; 56% ( $n = 60$ ) were eligible and consented to participate in the study. Participants were randomly assigned to the intervention ( $n = 30$ ) or control ( $n = 30$ ) groups. The intervention group received a measured dose of the topical anaesthetic EMLA<sup>®</sup> 5% cream daily as a primary dressing for 4 weeks followed by usual wound management for a further 8 weeks. The control group received usual wound management. Participants and investigators were not blinded to the treatment. WRP was measured at every dressing change. Wound healing and HRQoL were measured at baseline, 4 and 12 weeks.

**Results:** Recruitment rate was lower than expected which likely meant patients were missed. Study retention rate was 90% ( $n = 54$ ). Intervention fidelity was impacted by availability of resources and patient factors such as increased WRP.

**Conclusion:** This study identified that a larger randomised controlled trial investigating EMLA<sup>®</sup> applied as a primary dressing on painful chronic leg ulcers is feasible with modifications to the study protocol.

**Trial registration:** Australian New Zealand Clinical Trials Register: Registered 16 December, 2009

**Keywords:** Chronic leg ulcers, Wound-related pain, EMLA<sup>®</sup>

\* Correspondence: [anne.purcell@health.nsw.gov.au](mailto:anne.purcell@health.nsw.gov.au)

<sup>1</sup>Menzies Health Institute, Griffith University, Nathan Campus, Building N48, Room 2.10, 170 Kessels Road, Nathan, Queensland 4111, Australia

<sup>2</sup>School of Nursing and Midwifery, Griffith University, Nathan Campus, Building N48, Room 2.06, 170 Kessels Road, Nathan, Queensland 4111, Australia

Full list of author information is available at the end of the article



## Background

Wound-related pain (WRP) is the most common symptom of CLUs with reported prevalence as high as 85% [1, 2]. The most significant pain occurs at dressing change [3, 4]. For many individuals, WRP persists despite the use of conventional pharmacologic strategies such as oral analgesia [4]. Topical analgesia and anaesthetics applied directly to the wound bed is an option for relieving WRP [5]. The eutectic mixture of local anaesthetics cream (EMLA<sup>®</sup>) has been shown to be effective for relieving pain that occurs during debridement of CLUs [5]. Regulations for the use of EMLA<sup>®</sup> on open wounds such as CLUs and its drug schedule status may differ between countries. High quality evidence evaluating topical anaesthetics for managing WRP is still emerging [5–7].

In our published single-case report [8], we suggested that the topical application of EMLA<sup>®</sup> as a primary dressing may be a promising therapy for managing pain associated with CLUs but recognised that this treatment was yet to be empirically tested. Additionally, most individuals with painful CLUs are older with multiple comorbidities so higher rates of non-compliance are more likely. Therefore, in line with current recommendations, we conducted a pragmatic, external pilot, parallel group, randomised controlled trial (RCT) to evaluate the protocol implementation which would inform a larger trial [9]. A pragmatic approach was selected to assess the potential effectiveness of the intervention, the secondary objective, in a routine real-life practice setting. To assess feasibility as the primary outcome of this study, we were guided by the feasibility framework developed by Thabane et al. [10] including the following:

(1) Eligibility, recruitment and retention; (2) resource requirements; (3) human resources and data management; and (4) scientific assessment to identify potential effectiveness and any adverse events resulting from the intervention [10].

Study feasibility was assessed using the following pre-determined criteria for determining success:

- Recruitment of at least 80% of eligible patients within 12 months;
- Retaining 80% of participants in the study;
- Achieving 80% adherence to the intervention protocol

This paper reports on the feasibility findings of this pilot study. Patient-related outcomes were evaluated in this pilot study as secondary outcomes with the findings reported elsewhere [11, 12].

## Methods

### Study design

This feasibility study was a pilot, parallel group, non-blinded, randomised, controlled trial (RCT). The study

protocol was approved by the Northern Sydney Health (AU RED Ref. HREC/09/HARBR/162) and the Griffith University Human Research (GU Ref No: NRS/16/12/HREC) Ethics Committees (HREC), registered with the Australian New Zealand Clinical Trials Register (ACTRN12609001080213) and conducted in accordance with the Declaration of Helsinki (revised 2013); written informed consent was obtained from all participants. The study is reported according to the CONSORT 2010 statement [13] (Fig. 1). In line with Good Clinical Practice (GCP) [14], complications such as adverse reactions to EMLA<sup>®</sup> or wound infection were reported to the Data Safety and Monitoring Board (DSMB) and HREC.

### Setting and sample


Participants considered for inclusion in this study were individuals already referred to a large health district community nursing service in New South Wales, Australia. This study was conducted across six procedure clinics within the service where approximately three-quarters of patients required wound management; 76% had one or more CLUs [15].

Community nurses assisted with preliminary screening of patients and notified the study investigators of any patient with a lower leg ulcer greater than 6 weeks' duration, who required analgesia for WRP and had the capacity to attend the community nursing clinics for wound management. Potential patients were assessed against the inclusion and exclusion criteria (Table 1) by a member of the research team. Eligible patients were informed of the study and consent obtained.

A sample size of 60 was selected for this study as it was thought to be a good representation of the target population, large enough to provide practical information about the feasibility aspects of the study [10]. Although reported sample sizes for pilot studies vary, the median sample size per arm is often 30 [16]. In an external pilot study such as this, there is a 'trade-off' between maximising the precision of estimates of important parameters and sample size which impacts resources, time and costs of a study [17]. To address feasibility, we enrolled 60 patients to accommodate possible attrition throughout the 12-week study period. Attrition rates are often high in wound studies due to comorbidities associated this patient group particularly when the study period is long, where wounds may deteriorate, and the innate difficulty to adhering to the protocol over many weeks [18]. Even so, based on the community service patient profile, we estimated it would take approximately 12 months to enrol 60 patients assuming a 10% drop out rate [19]; a 20% drop-out is considered acceptable [20].

### Randomisation, blinding and allocation concealment

Following eligibility assessment and consent, a simple randomisation method (1:1) was used to randomise



**Fig. 1** Flow of participants through study

participants to either the intervention (EMLA®) or control group. PASS 2008 Power Analysis and Sample Size software (NCSS, Kaysville, UT) was used to generate the allocation sequence by a researcher not involved with screening patients; the investigator was blinded to the allocations. Patients were allocated to study groups by retrieving the next in a series of sequentially numbered, opaque, pre-prepared sealed envelopes. This method for allocation concealment can achieve a low risk of bias [21].

Similar to other wound care studies [18], it was not possible to blind the participants, treating nurses or the investigators to the treatment allocation as the intervention was compared to usual care and not a placebo. However, the statistician was blinded. To minimise bias, the intervention and control groups were treated as equally as possible apart from the intervention itself. Treating clinicians were required to adhere to the Australian Standards for Wound Management [22] and health service policies and procedures.

**Table 1** Inclusion and exclusion criteria and rationale for inclusion in study

	Rationale
<b>Inclusion criteria</b>	
1. One or more chronic lower leg ulcer(s) of at least 6 weeks duration	A chronic wound is defined as one which has not followed the expected path of healing when related to time, appearance and responses to optimum wound management and is often demonstrated when a wound has not shown signs of healing within a 6-week period [54]
2. Wound size up to 100 cm <sup>2</sup> in size in total	The maximum wound size of 100 cm <sup>2</sup> was selected due to the maximum dose/surface area ratio of 1–2 g EMLA® to 10 cm <sup>2</sup> of wound surface area, the recommended maximum dose being 10 g. This dose/surface area ratio is recommended for the application of EMLA® when used for the debridement of non-viable tissue from a leg ulcer [55]
3. Patients with low to moderate wound exudate	This will enable the EMLA® to remain on the wound bed over 24 h and not run off the wound due to excessive wound exudate
4. Numerical rating scale (NRS) pain score ≥ 4 at assessment and/or within the previous week	WRP ≥ NRS of 4 can indicate uncontrolled pain during or after dressing change which may require a change of management [56]
5. Patients currently requiring oral analgesics due to previously reported wound-related pain	Individuals with WRP are often prescribed oral analgesia indicating their level of pain is significant
6. Patients ≥ 18 years of age	The prevalence of CLUs increases with age thus the prevalence in those < 18 years of age is low [57]
7. Patients with the capacity (cognition and/or language)	Participants in this study should be capable of providing well-informed and considered consent
8. Patients have the capacity to attend the CCCNS Procedure Clinics. The exception is participants requiring visits on weekends and public holidays; health centres are closed at these times. This was in accordance with the NSCCH Home Safety Procedures for Community Nursing, CCH (PR2007_016).	Continuity of care
<b>Exclusion criteria</b>	
1. Patients scheduled for leg amputation	Amputation of the lower limb with a CLU would negate the need for wound management
2. History of peripheral sensory neuropathy (PSN) of the feet Modification: After 6 July, 2011, patients with painful peripheral neuropathy were no longer excluded	This study includes painful CLUs. Individuals with PSN often do not experience any peripheral sensation in their feet or lower legs [58] Patients with painful peripheral neuropathy can also experience painful WRP
3. Patients that have had or require the use of EMLA® for debridement of the wound bed within the previous 4 weeks before recruitment Modification: After 9 February 2011, patients were only excluded if they had EMLA® applied for debridement of the wound bed within the previous week	The introduction of EMLA® within 4 weeks may influence baseline data The half-life lignocaine and prilocaine is 65–150 and 10–150 min, respectively, a similar half-life of their intravenous counterparts [55, 59]
4. Patients with suspected wound malignancy or pyoderma gangrenosum confirmed by biopsy	The management of leg ulceration caused by malignancy or pyoderma gangrenosum requires different management than ulcerations due to venous, arterial or diabetic aetiology
5. Patients with diagnosed localised or spreading clinical wound infection	Management of wound infection requires the introduction of strategies that may influence the individual's pain levels beyond the capabilities of the primary dressing
6. End-stage palliative care patients	End-stage palliative patients often require increased use of analgesics for palliation, thus the effectiveness of an intervention may be masked
7. Patients where EMLA® is contraindicated or cautioned	To reduce potential participant harm [55]

WRP wound-related pain, HRQoL health-related quality of life, CLU chronic leg ulcer, EMLA® eutectic mixture of local anaesthetics, NRS numerical rating scale, CCCNS Central Coast Community Nursing Service, CCH Central Coast Health, PSN peripheral sensory neuropathy

### Interventions

An intervention period of 4 weeks was selected as healing rates over a 4-week period can determine intervention effectiveness [23–25]. Patients were followed-up over a 12-week period because wound margin advancement, initial healing rate and percentage of wound surface area reduction are strong indicators of complete

healing at 12 weeks independent of topical dressing used [25].

### Intervention group

Participants in the intervention group received a measured dose of EMLA® daily as a primary dressing to their CLUs for 4 weeks followed by 8 weeks of usual care.

EMLA<sup>®</sup> (Aspen Pharmacare, St Leonards, New South Wales, Australia) is a non-sterile, preservative-free, eutectic mixture of two amide type local anaesthetics lignocaine 2.5% and prilocaine 2.5% and has a good safety profile when used for debridement [26–28]. The EMLA<sup>®</sup> dose of 1–2 g per 10 cm<sup>2</sup> to a maximum dose of 10 g was based on manufacturer recommendations [28]. At each dressing change during the intervention period and following wound cleansing, EMLA<sup>®</sup> was drawn up into a syringe then spread evenly over the wound bed.

To ensure high quality and consistent application of the EMLA<sup>®</sup>, we developed an intervention protocol and provided education sessions and supervision of nurses to promote intervention fidelity. All treatments were performed by community nurses with previous experience in wound management. To assess the timing and dose of the intervention delivery and effect of the intervention on participants, random quality assurance visits to clinics or patient's homes, review of the participant's medical records and follow-up of the data collected were made by the chief investigator.

#### **Control group**

Initially, we considered using a placebo as the control group intervention however, further consideration identified that a placebo would require daily dressing changes to enable a placebo cream to be applied. The daily dressing change would confer no benefit to the participant and had potential to negatively impact wound healing; such an approach was difficult to ethically justify. Participants in the control group received usual care for 12 weeks where the primary dressing and dressing frequency (daily to weekly) were determined by clinical judgement over the treatment period. Usual care primary dressings could change throughout the study period in line with changes in exudate, non-viable tissue and microorganism levels over the course of treatment. Most common primary dressings used were hydrofibre, hydrogel, enzyme alginogel, povidine iodine mesh, cadexamer iodine and silver impregnated dressings.

Both groups received the same secondary dressing, a soft non-woven, hydrophobic polyamide fibre containing cellulose fluff core (Zetuvit<sup>®</sup>) [29]. Participants with low exudating CLUs in the intervention and control groups who experienced an increase in WRP during dressing change from secondary dressing adherence had a triglyceride mesh applied over the primary dressing and under the secondary dressing to prevent adherence. All wounds were cleansed with normal saline 0.9%. If clinically indicated, conservative sharp wound debridement and compression therapy were implemented. Regardless of treatment allocation, and in line with standard practice, all participants received EMLA<sup>®</sup> prior to conservative sharp debridement to eliminate

operative pain. EMLA<sup>®</sup> was removed prior to debridement. Intervention group participants had EMLA<sup>®</sup> re-applied following debridement.

#### **Management of adverse events**

Participants recorded any problems or adverse reactions to the intervention in their Pain Diary which was reviewed at each clinic visit. Treating clinicians observed for known reactions to EMLA<sup>®</sup> such as blanching, erythema, oedema, pruritus, burning, purpura and contact hypersensitivity [28] at each dressing change and documented them in the participant's electronic medical record. Additionally, a Notification of Adverse Event form was completed followed by immediate notification and discussion with the chief investigator (CI). In the event of any adverse reaction in the intervention group, treatment was ceased immediately and the medical team notified. Notifications of adverse events were mandatory and reported promptly to the NSH HREC and the DSMB by the CI.

#### **Measurements used to address study aims**

The feasibility of conducting an RCT in a public community health service to prevent unnecessary spending or wasting of resources in a larger study was measured by assessing quantitative and qualitative data to address key study processes, resources, management and scientific feasibility. Quantitative feasibility outcome measures included validation of the recruitment rate and randomisation processes, consent rate, retention rate, and suitability of the eligibility criteria and, data collection instruments. Additionally, to measure intervention adherence, the number of protocol deviations or violations during the study by treating nurses and participants were identified, and the reasons why these occurred were documented. Further quantitative and qualitative measures included the availability and commitment of human resources, the time to perform study processes, availability and quality of study equipment, data management outcomes and challenges and the cost estimates to conduct such a study. The feasibility of the intervention was measured by evaluating the patient's physiological responses to the intervention and the monitoring of adverse events. Qualitative measures included participant and investigator burden. Scientific feasibility was assessed to address the clinical responses to intervention. Scientific feasibility is reported elsewhere [11, 12].

Additional, patient-specific data were collected. Baseline measurements are detailed in Table 2. Data collection instruments, their psychometric properties, assessment time-points and estimated time to complete are presented in Table 3. When data specific to a leg ulcer were required and if a participant had more than one CLU, the largest ulcer was the reference ulcer; all ulcers were treated as per group allocation.

**Table 2** Baseline measurements

Participant history	Wound-related pain	HRQoL relating to CLU	CLU Characteristics
- Socio-demographic history - Medical and surgical history	- WRP at dressing change -Before -During -After - Pain type - Quality - Location - Triggers - Relievers - Pain medications - Effects on activities	- Social life - Wellbeing - Physical symptoms - Overall HRQoL [60]	- Leg ulcer surface area - Aetiology and duration of leg ulcer - Ankle Brachial Pressure Index - Leg ulcer measurement: -Exudate type and amount -Necrotic tissue type and amount -Granulation type and amount -Condition of wound edges -Peri ulcer viability - Oedema type & location - Assessment of bioburden - Wound-related pain intensity and frequency assessment over previous 24 h - How HRQoL relates to the leg ulcer [61]

WRP wound-related pain, HRQoL health-related quality of life, CLU chronic leg ulcer

### Data analysis

Data were entered and checked for missing and invalid values in Microsoft Excel® then imported into Statistical Analysis for Social Scientists (SPSS Version 22, Chicago, USA) for analysis. A random sample (10%) of the data was verified against the original case report form. Quantitative data were analysed using descriptive statistics.

Qualitative data from field notes were analysed using descriptive content analysis [30].

### Results

Detailed results for key components of the feasibility assessment are provided below.

**Table 3** Data collection instruments

Data collection instrument	Outcome measure	Estimated time of completion	Psychometric properties	Outcome assessment time points
11-point numerical rating scale [62, 63]	Wound-related pain intensity	1 min	Discriminative power relating to chronic pain. Test, re-test reliability high ( $r = 0.96$ ); Construct validity—highly correlated with the visual analogue scale for chronic pain conditions (reported range, 0.86 to 0.95) [64]	Baseline and every dressing
Cardiff Wound Impact Schedule [60]	Health-related quality of life	5 to 10 min	Established reliability with internal consistency subscale scores $> 0.75$ , and good reproducibility [60]	Baseline, 2, 4, 8 and 12 weeks
Leg Ulcer Measurement Tool [61] -Clinical and patient domains	Chronic leg ulcer appearance	5 to 10 min	Concurrent construct validity high ( $r = 0.82$ ) with excellent intra-rater/inter-rater reliability for the total LUMT scores ( $ICC > 0.75$ ) and for many of the 14 domains; some domains were less reproducible; this tool was able to detect change in wound status over time [61]	Baseline, 2, 4, 8 and 12 weeks
Wound-related pain at dressing change assessment tool [56, 65]	Wound-related pain response to intervention over the previous 24 h	5 min	Data collection tool not validated	Baseline and every dressing
Wound photography and 2-dimensional photo-digital planimetry [66]	Chronic leg ulcer measurement over time ( $\text{cm}^2$ )	15 min including download onto computer for measurement	Inter-rater reliability and intra-rater reliability is higher than traditional wound tracing methods (94 and 98.3%, respectively) [66]	Baseline, 2, 4, 8 and 12 weeks
American Geriatric Society Pain Diary [67]	WRP intensity and frequency of pain-relieving medications, mood and response to wound treatments [68, 69]		Data collection tool, not validated	Whenever pain perceived at home

ICC intraclass correlation coefficient,  $r$  correlation coefficient, LUMT Leg Ulcer Measurement Tool

## Study process assessment

### Recruitment

Participants considered for inclusion in this study were individuals already referred to a large health district community nursing service New South Wales, Australia. We screened 107 patients with painful CLUs of whom  $n = 70$  (65%) were eligible. Sixty of the eligible patients (56%) consented to participate in the study. While this met the feasibility objective, the recruitment rate was slower than anticipated and took 30 months (September 2010 to March 2013). In total, 30 patients screened did not meet the eligibility criteria. The most common reasons patients were excluded were insufficient wound-related pain, the presence of wound infection and high wound exudate. Twelve of the screened participants were not eligible due to frailty and chronic disease; seven of these participants were excluded due to the exclusion criteria. In the first 5 months, only three patients were recruited. We recognised that the exclusion criteria were too restrictive and modification took place to improve recruitment rates (Table 1). In addition, the appointment of a research assistant (RA) to assist the chief investigator (CI) also increased the recruitment rate to 2–3 per month. Comparison of groups' socio-demographic and clinical history at baseline is presented in Table 4. Completed follow-up of all participants took 33 months (September 2010 to June 2013). The trial ended once follow-up for all participants was complete.

### Retention

A retention rate of 90% of study participants was achieved. One patient in the intervention group and five in the control group did not complete the study (Fig. 1). One participant required a below knee amputation, another withdrew due to severe back pain and three withdrew as they were unable to maintain a commitment to data collection over the length of the study period. One participant in the control group was lost to follow-up.

### Adherence to study procedures

Prior to commencing the study, all treating nurses were assessed as competent in applying the EMLA<sup>®</sup> dressings. During the study, approximately 30 random quality assurance checks were attended with 100% compliance to intervention protocols. Consistent with current community nursing practice, eight participants performed their own dressing changes (up to three dressings each) when treating nurses or investigators were unavailable during a holiday period. For two participants, EMLA<sup>®</sup> was continued beyond the 4-week intervention period, at the patient's request, for management of ongoing pain. EMLA<sup>®</sup> was ceased for five (16.7%) participants; one participant requested EMLA<sup>®</sup> to be ceased due to participant burden and four participants reported unchanged or increased WRP following application of EMLA<sup>®</sup>. There were confounding factors influencing WRP for the participants who reported unchanged or increased WRP such as

**Table 4** Comparison of Groups' socio-demographic and clinical history at baseline

	Intervention group ( $n = 30$ )	Control group ( $n = 30$ )	<i>P</i>
	Mean (SD)	Mean (SD)	
Age (years)	73.4 (12.5)	73.8 (10.1)	0.89
CLU duration (weeks)	26.4 (26.0)	20.5 (13.4)	0.32
CLU surface area (cm <sup>2</sup> ) at baseline	8.01 (10.4)	9.2 (8.9)	0.48
Sex	<i>n</i> (%)	<i>n</i> (%)	
Male	13 (43.3)	12 (40.0)	0.79
Female	17 (56.6)	18 (60.0)	0.79
Ulcer type			
Venous	18 (60.0)	22 (73.0)	0.27
Arterial	5 (20.0)	3 (10.0)	0.45
Mixed	5 (13.3)	3 (10.0)	0.45
Incompressible	1 (3.3)	2 (6.6)	0.55
Diabetic foot ulcer	1 (3.3)	0	0.31
Pain medications			
Opiates	17 (56.6)	13 (40.0)	0.30
NSAIDS	4 (13.3)	5 (16.6)	0.72
Other pain meds	21 (70.0)	23 (76.6)	0.56

SD standard deviation, *p* *p* value, CLU chronic leg ulcer, NSAIDS non-steroidal anti-inflammatory drugs

compression therapy applied too early and severe arterial disease.

#### Data collection

Data collection instruments and time-points are presented in Table 3. The numerical rating scale (NRS) and the Wound-related Pain at Dressing Change Assessment Tool were quick and easy for both the investigator and the participant to use. The Cardiff Wound Impact Schedule was found to be long and confusing for the frailest of participants. Data from the Pain Diary was inconsistently completed and the diary was not wound specific, thus pain unrelated to CLUs was also documented by the participants. The Leg Ulcer Measurement Tool measured CLU progress; however, there was an overlap with other data collected including WRP intensity, WRP frequency and participant satisfaction with HRQoL. Health service protocols for digital photography were not always adhered to by the community nurses. This contributed to missing data, firstly due to uploading of unclear images, secondly as the computer could not calibrate some images as a ruler was not included in the image thus preventing accurate digital measurement, and finally, some images were not taken at the required time-points (47 out of 266 images). For images that were able to be measured, a specialist wound clinician was required to manually assess accuracy of CLU surface area measurements.

The mean percentage of missing data was 19% (range 8 to 25%) and was missing completely at random [11, 12]. Missing data increased as the study progressed, particularly when nurses trained in data collection were unavailable.

#### Resource assessment

Human resources required to conduct this study were impacted by availability of the CI and RA who had competing full-time clinical roles. The RA was only available 1 day per week for 17 months to support the study; funding was required for the RA position.

The nurse time required for dressing change and pain data collection was the same in both groups (30 min) as was the added time taken for HRQoL and wound photography data collection at the 2, 4, 8, and 12 week time-points (1 h).

Some participants required home visits during business hours and weekends mostly due to participant frailty, lack of transport options, lack of clinic capacity, and the intervention protocol that is, daily dressings for the intervention period (weekdays: intervention group  $n = 9$  (30%); control group:  $n = 6$  (20%); weekends: intervention group:  $n = 30$  (100%); control group  $n = 4$  (33.3%); range: 1 to all visits). Seven participants required all visits in their homes (intervention group:  $n = 5$

(16.6%), range: 24 to 77 visits); control group:  $n = 2$  (6.6%), range: 7 to 24 visits).

A limited economic feasibility assessment was informed by a comparison of cost estimates for the intervention and usual care primary and secondary dressings. The intervention group had higher overall costs over the 12-week study period, with increased costs attributed to increased dressing frequency. Throughout the data collection period of 33 months, the intervention group required almost double the number of dressings compared to the control group (intervention group, 1232 dressings (65.4%); control group, 651 dressings (34.6%)). Daily dressings during the intervention period for the intervention group contributed to the considerable difference between groups (intervention group, 741 dressings (74.6%); control group 252 dressings (25.3%)). The overall cost of dressings per dressing change was less in the intervention group (intervention group, A\$6.03; control group A\$8.73). However, the intervention group had a 13.2% higher overall cost of primary and secondary dressing consumables over the study period (intervention group A\$7441; control group A\$5684), due to the increased frequency of dressings compared to the control group.

#### Management assessment

Participants were initially seen in the community health service clinics. However, as the study progressed, 13 participants had difficulty attending the clinics for treatment. Hire cars were provided for these participants as a short-term solution however, due to budget restrictions, this strategy was not feasible and was discontinued after 20 months. Subsequently, participants unable to attend the clinic were treated in their homes during business hours and weekends as previously reported using existing community nursing resources. This meant that some visits were attended by nurses not educated in the intervention protocols at the beginning of the study; nurse continuity was also an issue. The potential impact on the data collected during home visits was anticipated; the CI attended the visit particularly on weekends or made phone contact with the nurse to explain the protocols and procedures prior to the home visit.

Participant burden was observed in this study. The length of the eligibility interview, randomisation and recruitment processes (1 to 3 h), the length of the intervention (4 weeks) and study periods (12 weeks), the length of some data collection tools (up to 50 questions), the frequency of the data collection (at each dressing change), the average age and health status of the participants (73 years) and requirement of participants to come to the clinics, all contributed to participant burden.



### Scientific assessment

There was no difference in WRP intensity scores between groups before dressing change over the 4-week intervention period (intervention group: mean 4.10 [95% CI 3.55, 4.63] vs control group: mean 4.21 [95% CI 3.66, 4.76]). Nevertheless, there was a statistically significant reduction in WRP for the intervention group compared to the control group during dressing change (intervention group: mean 3.39 [95% CI 2.59, 4.19] vs control group: mean 4.82 [95% CI 3.98, 5.66]) and after dressing change: (intervention group: mean 2.71 [95% CI 1.99, 3.43] vs control group: mean 3.92 [95% CI 3.16, 4.68]) [11]. EMLA<sup>®</sup> was tolerated well for 4 weeks by 83.3% ( $n = 25$ ) of the intervention group. The remaining 16.6% ( $n = 5$ ) of the intervention group had adverse effects from the application of EMLA<sup>®</sup> to their leg ulcers. Erythema, pallor, itching, oedema, purpuric or petechial lesions, or allergic reaction were not reported by the attending clinicians; however, five participants required EMLA<sup>®</sup> to be ceased due to increased or unchanged WRP and increased wound size. Usual care was recommenced on all participants. There were no serious adverse events to the intervention during this study.

Interestingly, two participants required recommencement of EMLA<sup>®</sup> following the 4-week intervention period at their request for the remainder of the 12 week study period due to significant exacerbation of their WRP after cessation of the EMLA<sup>®</sup> and commencement of usual care. Once EMLA<sup>®</sup> was recommenced, WRP was reduced.

### Discussion

This is the first study to investigate EMLA<sup>®</sup> used as a primary dressing for relieving wound-related pain for patients with painful chronic leg ulcers. The pilot study was pivotal to assessing feasibility for a larger clinical trial and to determine potential effectiveness. The identification of potential practical problems that may cause breakdown when implementing the research study protocol into clinical practice is crucial for the success of a larger study [31, 32]. By undertaking this feasibility study, we have been able to identify ways in which the study protocol for a future multicentre randomised controlled trial could be refined although the generalisability of our results may be limited due to participant enrolment from a single site. Solutions to manage any challenges during the study and recommendations for protocol modifications to inform a larger RCT are presented in Table 5. The key learnings from this pilot study related to recruitment and retention of participants, establishing resources required and managing data collection to ensure data accuracy and completeness and are presented below.

In terms of the feasibility objectives, although we were able to recruit 100% of the target sample, it was not

achieved in the predicted timeframe of 12 months. We were able to meet participant retention and intervention adherence targets of 80%, and the study outcomes suggest that it is feasible to proceed to a larger multisite clinical trial to examine EMLA<sup>®</sup> as a primary dressing on painful CLUs. However, modifications to the protocol are recommended.

### Recruitment

Participant and research process factors influenced recruitment rates in this study. This experience is not uncommon in RCTs [33] where up to two-thirds of trials are unable to successfully recruit their original target [34, 35]. Protracted or ineffective recruitment can have undesirable scientific, ethical and economic consequences [36, 37]. Although random assignment may result in refusal to participate in a study [36, 38], this was not the case in this study. We overestimated the pool of patients with CLUs in the community nursing service that would meet the eligibility criteria. Known as 'Lasagna's Law', this phenomenon is a common problem in clinical research with the evidence indicating that the incidence of the disease investigated reduces to 10% of the original estimate once the study starts [38]. This is a common threat to the success of clinical research resulting in increased direct costs and challenges the commitment and morale of research staff and participants [38]. Study processes and under resourcing of research personnel contributed to slower than expected recruitment, and it is likely that eligible patients were missed.

Lack of interest, inability to commit, physical and time limitations, change to daily activities and inability to travel to the community nursing clinics were patient-related factors affecting slow recruitment in this study; all of which are frequently cited in the literature [38]. The biggest obstacle to recruitment was patient eligibility at initial screening; we excluded more than we had predicted (34%), thus potential participants with painful CLUs were omitted from the study which may have negatively affected the generalisability of our results. A literature review found that an average of 30% of patients attending eligibility screening in RCTs are ineligible [34]. The majority of potentially eligible participants in this study were older; this was expected since the older person is more likely to succumb to CLUs [39–41].

Considering the prevalence of chronic leg ulcers in society, most chronic wound trials have small sample sizes reflecting the difficulty in recruiting patients if the eligibility criteria are too restrictive [42]. It became apparent that two of the exclusion criteria were considered unnecessary hence amendments were made to the exclusion criteria to increase the recruitment rate previously described. Based on these findings, we acknowledge the importance of understanding the needs and abilities of

**Table 5** Feasibility challenges, solutions and recommendations

	Challenges	Solutions and recommendations
Feasibility		
Recruitment rate	<ul style="list-style-type: none"> <li>- Recruitment was protracted. The reasons were:               <ul style="list-style-type: none"> <li>- Insufficient dedicated research personnel coupled with competing full-time workloads</li> <li>- Structured screening process prevented identification of all eligible patients</li> <li>- Exclusion criteria too restrictive</li> </ul> </li> </ul>	Solutions: <ul style="list-style-type: none"> <li>- Employment of an RA</li> <li>- Community nursing referral screening tool was developed</li> <li>- Transportation was provided for some participants to clinics</li> <li>- Amendments to some exclusion criteria</li> <li>- Some participants were treated in their homes</li> </ul>
Retention	<ul style="list-style-type: none"> <li>- Participant burden was increased for some frail participants</li> <li>- There was limited availability of transportation to clinics</li> </ul>	Recommendations: <ul style="list-style-type: none"> <li>- Employment of a dedicated trial manager</li> <li>- A comprehensive screening process to identify potential participants at the beginning of the study</li> <li>- Establish centralised intake system to identify potential patients at first CN contact</li> <li>- In-depth review of nursing resources including skill mix prior to commencement of study</li> <li>- Include home visits for treatments in a larger study</li> </ul>
Resources	<ul style="list-style-type: none"> <li>- Insufficient human resources to conduct the research within the timeframe</li> <li>- Poor continuity of nursing services especially for home visits</li> <li>- Some patients could not attend clinics</li> <li>- Re-calibration of photo digital planimetry software required for wound measurement accuracy</li> </ul>	Solutions: <ul style="list-style-type: none"> <li>- Further institution support was acquired during study</li> <li>- Support from experienced clinical nurses to administer intervention and collect data</li> <li>- The application of the intervention was able to be accommodated within existing clinic schedules</li> <li>- Information technology support was acquired</li> </ul> Recommendations: <ul style="list-style-type: none"> <li>- Review treatment location options</li> <li>- Collaboration with health service management regarding staff backfill to promote continuity</li> <li>- Study-specific investigators and support staff</li> <li>- Usual care wound products for multisite larger study as per health service formulary; negotiate for in-kind supply of wound products for a multisite RCT</li> <li>- Use dedicated study equipment</li> </ul>
Management	<ul style="list-style-type: none"> <li>- Oversight of the study was difficult for CI and RA due to competing full-time workload</li> <li>- Participant burden was high</li> <li>- Prolonged consent, randomisation and baseline data collection processes</li> <li>- Poor quality photos of some wounds</li> <li>- Existing resources made available by health service for wound size measurements were insufficient</li> <li>- Research data were collected parallel to health service data resulting in some duplication and extended nurse time</li> <li>- Data collected by clinicians untrained in study processes resulted in higher rates of missing data</li> </ul>	Solutions: <ul style="list-style-type: none"> <li>- Ensured computer software for data entry and analysis available and appropriate</li> <li>- Ensured technology to acquire, store and measure wound photography appropriate</li> <li>- Separated recruitment processes from baseline data collection</li> <li>- Established secure data storage</li> <li>- There was easy access to information technology support</li> <li>- Wound size measurements by wound specialist nurse</li> </ul> Recommendations: <ul style="list-style-type: none"> <li>- Trial-specific investigators, data collectors and administrative team with no competing interests</li> <li>- Shorten recruitment processes</li> <li>- Revise and reduce the size and number of data collection tools and data collection time points</li> </ul>
Scientific	<ul style="list-style-type: none"> <li>- Change to intervention protocol required for some participants due to negative clinical response</li> </ul>	Solutions: <ul style="list-style-type: none"> <li>- Intervention ceased, changed to usual care</li> </ul> Recommendations: <ul style="list-style-type: none"> <li>- Investigate potential use of placebo</li> </ul>

RA research assistant, CI chief investigator, CN community nursing

the prospective study population prior to developing eligibility criteria for a larger study. Prior to recruitment, we over-estimated how many of the potentially eligible patients would qualify for the study. Although recruitment was initially slow, the rate improved when adequate resources for screening and recruitment were in place.

The difficulty in recruiting older individuals has been well identified; however, age itself does not

determine an individual's ability to give consent to research [43]. There are factors however that are associated with age that could impact on an individual's ability to consent effectively such as frailty, fatigue, cognitive impairment, chronic disease and/or feelings of vulnerability [43]. Nevertheless, in this study, we had a good consent rate (86%) when compared to other RCTs relating to CLUs [44–46].

### Community nurses

Most wound care is attended to in the community setting and community nurses are essential for the identification of potential participants in wound care research. They have been described as “effectively the gatekeepers to trial participation” [47]. They are an important link between the investigators and participants and can influence recruitment and retention rates [38]. In this study, community nurses were enthusiastic about being involved in an RCT and could see the benefits to themselves and to the patient. They were provided with criteria to assist them to identify potentially eligible patients however, subsequently, the demands of their clinical workload impacted their ability to undertake patient screening and contributed to delayed or missed participant identification [47].

### Chronic leg ulcer types

Patients with venous, arterial, mixed (arterial/venous) and diabetic foot ulcerations were recruited so outcomes reflected real-world clinical practice. These ulcer types differ in their underlying aetiology and wound characteristics. To increase the ability to meet our required sample size, patients with any of the above CLU types were included. We acknowledge that such heterogeneity can threaten study validity and usefulness of a clinical trial [42]. Subgroup analysis would be a solution however, a much larger sample size would be required [42] which was not realistic for this single-centre pilot study. Additionally, simple randomisation may not be sufficient to provide well-balanced treatment groups regarding confounders in this broadly defined study; a large, multicentre RCT using stratified randomisation may be appropriate [42].

### Attrition

To maintain statistical power, we aimed for less than 20% loss to follow-up. Attrition rates of 20% or more introduces bias and is a serious threat to the internal and external validity by altering the structure of the intervention and control groups [38, 48]. Common predictors for study withdrawal are older age and functional impairment [49]. The attrition rate for this study was only 10% which was encouraging considering the majority of participants were older, frail, in significant pain, had committed to a long study period, were subject to a large amount of data collection and were required to travel to community nursing clinics. To aid retention, some participants required home visits; however, in a larger study, this will require more time allocation plus additional costs [49].

### Participant burden

Participant burden is a subjective, multidimensional construct relating to the perception of the participants

physiological, physical and/or economic adversity with involvement in the research process [50, 51]. Investigators have traditionally addressed participant burden in clinical trials by focusing on direct risks associated with the intervention or data collection procedures. Nevertheless, it is the indirect burden that can vary due to factors such as study duration, intensity and invasiveness of study procedures [50] that needs to be considered. This pragmatic pilot study has identified direct and indirect factors that contributed to participant burden and will be able to inform a larger study to use a more pragmatic approach to reduce participant burden to maximise research participation and response rates (Table 5).

### Missing data

In this study, we had a mean percentage of 19% missing data. The literature does not identify an established cut-off regarding an acceptable percentage of missing data; 5% or less is considered inconsequential, and more than 10% can result in a biased statistical analysis [52]. Additionally, missing data mechanisms and patterns can have a bigger influence on results than the proportion of missing data [52]. To manage missing data in this study, we attempted to follow up all participants, included all available data in the analysis making a plausible assumption about missing data and did a sensitivity analyses that weakened the assumptions about missing data [48]. Missing data was missing completely at random; therefore, systemic attrition did not occur and an unbiased treatment effect estimate was derived from the obtained data [48, 53]. For a definitive effectiveness trial, missing data would need to be minimised to reduce the threat to study validity. Evidence shows that 95% of RCTs report some missing data which can threaten the validity of an RCT, make a true intention to treat analysis difficult to achieve, reduce the power and efficiency of the study and lead to bias [48].

### Strengths of the study

The strengths of the study include the recruitment of 100% of the target population, retention of 90% of our sample, assessment of the fidelity of the intervention, inclusion of objective outcome measures, and the ability to refine protocols and procedures. Generalizability of the results of this pilot study may be limited due to participant enrolment from a single health service. Consequently, context-specific issues that may be influenced by local, regional or country specific practices are unknown. Furthermore, there were fewer eligible participants than initially anticipated. The exclusion criteria may have resulted in some patients with painful CLUs being overlooked for inclusion in this study.

Bias could have been introduced since the participants, treating nurses and researchers could not be blinded to

the intervention. Additionally, process evaluation was not attended by a neutral party but by the researchers themselves; therefore, there is the potential that further biases may have been introduced. The study protocols and procedures placed considerable demands on the mostly frail aged participants, the treating nurses and health service resources all contributing to missing data. We recognise the need to minimise the difficulties identified in this study that participants and investigators may encounter when designing a protocol for a larger multisite study. Furthermore, regulations for the use of EMLA® on open wounds such as CLUs and its drug schedule status would have to be ascertained prior to an international study as these factors may differ between countries.

## Conclusion

Our goal is to move towards a larger study with wound-related pain as the primary endpoint conducted on individuals with painful chronic leg ulcers. This pilot study provides important feasibility information that can be used to inform a definitive future study. In the interim, this study provides insight into the potential effectiveness of EMLA® on painful chronic leg ulcers, wound healing and health-related quality of life when used as a primary dressing.

## Abbreviations

CI: Chief investigator; CLU: Chronic leg ulcer; DSMB: Data safety monitoring board; EMLA®: Eutectic Mixture of Local Anaesthetics; GCP: Good Clinical Practice; HREC: Human Research Ethics Committee; HRQoL: Health-related quality of life; RA: Research assistant; RCT: Randomised controlled trial; WRP: Wound-related pain

## Acknowledgements

Thank you to the participants and their families for their commitment to this study. Thank you to Jonathan Brinton and Donald Sutherland, Research Assistants, and the CCLHD Community Nursing Service for their contribution to recruitment and data collection. Thank you to Judith Fethney, Biostatistician, University of Sydney, for her assistance with data analysis. Thank you to Wendy White, Director, Wendy White Wound Care, for her role as clinical advisor to the study.

## Funding

This work was supported by the New South Wales (NSW) Health Nursing and Midwifery Innovation Scholarship, Australian Wound Management Research Foundation Grant, Central Coast Health Research Advisory Committee CHARM Research Grant, New South Wales, Australia.

## Availability of data and materials

Ethical approval and associated ethics requirements prevent the sharing of study data.

## Authors' contributions

AP, TB, JK and AM drafted the study protocol and AP analysed the data under the regular supervision of all other co-authors. All authors read and approved the final manuscript.

## Ethics approval and consent to participate

Approved by the Northern Sydney Health (NSH) and the Griffith University Human Research Ethics Committees (HREC). Written informed consent was obtained from all participants.

## Consent for publication

Not applicable.

## Competing interests

The authors declare that they have no competing interests.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## Author details

<sup>1</sup>Menzies Health Institute, Griffith University, Nathan Campus, Building N48, Room 2.10, 170 Kessels Road, Nathan, Queensland 4111, Australia. <sup>2</sup>School of Nursing and Midwifery, Griffith University, Nathan Campus, Building N48, Room 2.06, 170 Kessels Road, Nathan, Queensland 4111, Australia. <sup>3</sup>Central Coast Local Health District, Gosford Hospital, Holden St, Gosford, New South Wales 2250, Australia. <sup>4</sup>Sydney Nursing School, University of Sydney, 88 Mallett St, Camperdown, New South Wales 2050, Australia. <sup>5</sup>Gold Coast University Hospital, Gold Coast Hospital and Health Services, E. 2 015, 1 Hospital Blvd, Southport, Queensland 4215, Australia. <sup>6</sup>National Centre of Research Excellence in Nursing, Room 2.08, G01, Griffith University, Southport, Queensland 4222, Australia.

Received: 16 December 2017 Accepted: 22 June 2018

Published online: 07 July 2018

## References

- Nemeth KA, Harrison MB, Graham ID, Burke S. Pain in pure and mixed aetiology venous leg ulcers: a three-phase point prevalence study. *J Wound Care*. 2003;12(9):336–40.
- Persoon A, Heinen M, Van der Vleuten C, De Rooij M, Van de Kerckhof P, Van Achterberg T. Leg ulcers: a review of their impact on daily life. *J Clin Nurs*. 2004;13:341–54.
- Herber OR, Schnepf W, Rieger MA. A systematic review on the impact of leg ulceration on patients' quality of life. *Health Qual Life Outcomes*. 2007;5:44:1–12
- Price PE, Fagervik-Morton H, Mudge EJ, Beele H, Ruiz JC, Nystrom TH, Lindholm C, Maume S, Melby-Ostergaard B, Peter Y, et al. Dressing-related pain in patients with chronic wounds: an international patient perspective. *Int Wound J*. 2008;5(2):159–71.
- Briggs M, Nelson EA, Martyn-St James M. Topical agents or dressings for pain in venous leg ulcers. *Cochrane Database Syst Rev*. 2012;11:1–50.
- Cuomo R, D'Aniello C, Grimaldi L, Nisi G, Botteri G, Zerini I, Brandi C. EMLA and lidocaine spray: a comparison for surgical debridement in venous leg ulcers. *Adv Wound Care*. 2015;4(6):358–61.
- Fogh K, Andersen MB, Bischoff-Mikkelsen M, Bause R, Zutt M, Schilling S, Schmutz JL, Borbujo J, Jimenez JA, Cartier H, et al. Clinically relevant pain relief with an ibuprofen-releasing foam dressing: results from a randomized, controlled, double-blind clinical trial in exuding, painful venous leg ulcers. *Wound Repair Regen*. 2012;20(6):815–21.
- Purcell A, Marshall A, King J, Buckley T. Eutectic mixture of local anaesthetics (EMLA) 5% cream as a primary dressing on a painful lower leg ulcer: a case report. *J Wound Care*. 2012;21(7):309–14.
- Arnold DM, Burns KE, Adhikari NK, Kho ME, Meade MO, Cook DJ, McMaster Critical Care Interest G. The design and interpretation of pilot trials in clinical research in critical care. *Crit Care Med*. 2009;37(1 Suppl):S69–74.
- Thabane L, Ma J, Chu R, Cheng J, Ismaila A, Rios L, Robson R, Thanbane M, Giangregorio L, Goldsmith C. A tutorial on pilot studies: the what, why and how. *BMC Med Res Methodol*. 2010;10(1): 1–10.
- Purcell A, Buckley T, Fethney J, King J, Moyle W, Marshall AP. The effectiveness of EMLA® as a primary dressing on painful chronic leg ulcers - a pilot randomised controlled trial. *Adv Skin Wound Care*. 2017;30(8):354–63.
- Purcell A, Buckley T, Fethney J, King J, Moyle W, Marshall AP. The effectiveness of EMLA® as a primary dressing on painful chronic leg ulcers: effects on wound healing and health-related quality of life. *Int J Low Extrem Wounds*. 2017;16(3):163–72.
- Schulz KF, Altman DG, Moher D, Group C. CONSORT 2010 statement: updated guidelines for reporting parallel group randomised trials. *BMJ*. 2010;340:c332.
- Therapeutic Goods Administration: The Australia Clinical Trial Handbook: Guidance on conducting clinical trials in Australia using 'unapproved' therapeutic goods. In: Department of Health: Commonwealth of Australia; 2018.

15. Central Coast Health Community Nursing Service: Integrated clinical information system. 2017.
16. Billingham SAM, Whitehead AL, Julious SA. An audit of sample sizes for pilot and feasibility trials being undertaken in the United Kingdom registered in the United Kingdom Clinical Research Network database. *BMC Med Res Methodol*. 2011;13(104):1-6.
17. Teare MD, Dimairo M, Shephard N, Hayman A, Whitehead A, Walters SJ. Sample size requirements to estimate key design parameters from external pilot randomised controlled trials: a simulation study. *Trials*. 2014;15:264.
18. Gottrup F, Apelqvist J, Price P. Outcomes in controlled and comparative studies on non-healing wounds: recommendations to improve the quality of evidence in wound management. *J Wound Care*. 2010;19(6):237-68.
19. Zang D, Yin P, Freemantle N, Jordon R, Zhong N, Cheng K. An assessment of the quality of randomised controlled trials conducted in China. *Trials*. 2008;9(2):1-8.
20. National Heart Lung and Blood Institute. Quality Assessment of Controlled Intervention Studies. USA: National Institutes of Health, Department of Health and Human Services; 2014.
21. Higgins J, Green SE. *Cochrane Handbook for Systematic Reviews of Interventions*. West Sussex: Wiley; 2008.
22. Australian Wound Management Association Inc. *Standards for Wound Management*. 2nd ed. West Leederville WA: Cambridge Publishing; 2010.
23. Flanagan M. Improving accuracy of wound measurement in clinical practice. *Ostomy Wound Manage*. 2003;49(10):28-40.
24. Lavery LA, Barnes SA, Keith MS, Seaman JW Jr, Armstrong DG. Prediction of healing for postoperative diabetic foot wounds based on early wound area progression. *Diabetes Care*. 2008;31(1):26-9.
25. Cardinal M, Eisenbud DE, Phillips T, Harding K. Early healing rates and wound area measurements are reliable predictors of late complete wound closure. *Wound Repair Regen*. 2008;16(1):19-22.
26. Hansson C, Holm J, Lillieborg S, Syren A. Repeated treatment with lidocaine/prilocaine cream (EMLA) as a topical anaesthetic for the cleansing of venous leg ulcers. A controlled study. *Acta Derm Venereol*. 1993;73(3):231-3.
27. Lok C, Paul C, Amblard P, Bessis D, Debure C, Faivre B, Guillot B, Ortonne JP, Huledal G, Kalis B. EMLA cream as a topical anesthetic for the repeated mechanical debridement of venous leg ulcers: a double-blind, placebo-controlled study. *J Am Acad Dermatol*. 1999;40(2 Pt 1):208-13.
28. Aspen Pharmacare Canada Inc.: EMLA cream product monograph. 2017.
29. Paul Hartmann Pty Ltd: Technical data sheet: Zetuvit sterile. Specification no. D61201b. 2003.
30. Elo S, Kyngas H. The qualitative content analysis process. *J Adv Nurs*. 2008; 62(1):107-15.
31. Gardner G, Gardner A, MacLellan L, Osborne S. Reconceptualising the objectives of a pilot study for clinical research. *Int J Nurs Stud*. 2003;40(7):719-24.
32. Van Teijlingen ER, Rennie AM, Hundley V, Graham W. The importance of conducting and reporting pilot studies: the example of the Scottish Births Survey. *J Adv Nurs*. 2001;34(3):289-95.
33. Caldwell PH, Hamilton S, Tan A, Craig JC. Strategies for increasing recruitment to randomised controlled trials: systematic review. *PLoS Med*. 2010;7(11):e1000368.
34. Toerien M, Brookes ST, Metcalfe C, de Salis I, Tomlin Z, Peters TJ, Sterne J, Donovan JL. A review of reporting of participant recruitment and retention in RCTs in six major journals. *Trials*. 2009;10:52.
35. McDonald A, Knight R, Campbell M, Entwistle V, Grant A, Cook J, Elbourne D, Francis D, Garcia J, Roberts I, et al. What influences recruitment to randomised controlled trials? A review of trials funded by two UK funding agencies. *Trials*. 2006;7(9):1-8.
36. Gross D, Fogg L. Clinical trials in the 21st century: the case for participant-centered research. *Res Nurs Health*. 2001;24(6):530-9.
37. Watson JM, Torgerson DJ. Increasing recruitment to randomised trials: a review of randomised controlled trials. *BMC Med Res Methodol*. 2006;6:34.
38. Gul RB, Ali PA. Clinical trials: the challenge of recruitment and retention of participants. *J Clin Nurs*. 2010;19:227-33.
39. Baker SR, Stacey MC. Epidemiology of chronic leg ulcers in Australia. *Aust N Z J Surg*. 1994;64(4):258-61.
40. Graham ID, Harrison MB, Nelson EA. Prevalence of lower-limb ulceration: a systematic review of prevalence studies. *Adv Skin Wound Care*. 2003;16(6):305-16.
41. Moffatt CJ, Franks PJ, Doherty DC, Martin R, Blewett R, Ross F. Prevalence of leg ulceration in a London population. *Q J Med*. 2004;97(7):431-7.
42. Vollenweider D, Ebnetter I, Mayer D, Hafner J, Steurer J, Puhon MA. Dealing with heterogeneous populations in randomized wound trials: challenges and potential solutions. *Wound Repair Regen*. 2012;20(4):466-72.
43. Harris R, Dyson E. Recruitment of frail older people to research: lessons learnt through experience. *J Adv Nurs*. 2001;36(5):643-51.
44. Harrison MB, Graham ID, Lorimer K, Vandenberg E, Buchanan M, Wells PS, Brandys T, Pierscianowski T. Nurse clinic versus home delivery of evidence-based community leg ulcer care: a randomized health services trial. *BMC Health Serv Res*. 2008;8:243.
45. Weller CD, Evans SM, Staples MP, Aldons P, McNeil JJ. Randomized clinical trial of three-layer tubular bandaging system for venous leg ulcers. *Wound Repair Regen*. 2012;20(6):822-9.
46. Nikolovska S, Arsovski A, Damevska K, Gocev G, Pavlova L. Evaluation of two different intermittent pneumatic compression cycle settings in the healing of venous ulcers: a randomized trial. *Med Sci Monit*. 2005;11(7):CR337-43.
47. Lamb KA, Backhouse MR, Adderley UJ. A qualitative study of factors impacting upon the recruitment of participants to research studies in wound care - the community nurses' perspective. *J Tissue Viability*. 2016;25(3):185-8.
48. Bell M, Fiero M, Horton N, Chiu-Hsieh H. Handling missing data in RCTs; a review of the top medical journals. *BMC Med Res Methodol*. 2014;14(118):1-6.
49. Peterson JC, Pirraglia PA, Wells MT, Charlson ME. Attrition in longitudinal randomized controlled trials: home visits make a difference. *BMC Med Res Methodol*. 2012;12:178.
50. Lingler JH, Schmidt KL, Gentry AL, Hu L, Terhorst LA. A new measure of research participant burden: brief report. *J Empir Res Hum Res Ethics*. 2014;9(4):46-9.
51. Ulrich CM, Wallen GR, Feister A, Grady C. Respondent burden in clinical research: when are we asking too much of subjects? *Irb*. 2005;27(4):17-20.
52. Dong Y, Peng CY. *Principled missing data methods for researchers*. SpringerPlus. 2013;2(1):222.
53. Bell ML, Fairclough DL. Practical and statistical issues in missing data for longitudinal patient-reported outcomes. *Stat Methods Med Res*. 2014;23(5):440-59.
54. MacLellan DG, editor. *Wound glossary: terminology for wound practitioners*. Sydney: Health Education & Management Innovations; 2007.
55. AstraZeneca Pty Ltd. *EMLA Cream Product Monograph*. 2016.
56. World Union of Wound Healing Societies. *Principles of best practice: minimising pain at wound dressing-related procedures. A consensus document*. London: MEP Ltd; 2004.
57. Simon DA, Dix FP, McCollum CN. Management of venous leg ulcers. *BMJ*. 2004;328(7452):1358-62.
58. Tanenberg RJ, Donofrio PD. Neuropathic problems of the lower limbs in diabetic patients. In: Bowker JH, Pfeifer M, editors. *Levin and O'Neals the diabetic foot*. 7th ed. Philadelphia: Mosby Elsevier; 2008.
59. Kopecky EA, Jacobson S, Hubley P, Palozzi L, Clarke HM, Koren G. Safety and pharmacokinetics of EMLA in the treatment of postburn pruritis in pediatric patients: a pilot study. *JBurn Care Rehab*. 2001;22(3):235-42.
60. Price P, Harding K. Cardiff wound impact schedule: the development of a condition specific questionnaire to assess health-related quality of life in patients with chronic wounds of the lower limb. *Int Wound J*. 2004;1(1):10-6.
61. Woodbury G, Houghton P, Campbell K, Keast D. Development, validity, reliability, and responsiveness of a new leg ulcer measurement tool. *Adv Skin Wound Care*. 2004;17:187-96.
62. Ferreira-Valente MA, Pais-Ribeiro JL, Jensen MP. Validity of four pain intensity rating scales. *Pain*. 2011;152(10):2399-404.
63. Hjermstad M, Fayers P, Haugen D, et al. Studies comparing numerical rating Scales, Verbal rating scales, and visual analogue, scales for assessment of pain intensity in adults. *J Pain Symptom Manag*. 2011;14(6):1073-93.
64. Hawker GA, Mian S, Kendzerska T, French M. Measures of adult pain. *Arthritis Care Res*. 2011;63(S11):S240-52.
65. Hollinworth H. Pain at wound dressing-related procedures: a template for assessment. *World Wide Wounds*. 2005;
66. Wendelken ME, Berg W, Lichtenstein P, Markowitz L, Comfort C, Alvarez O. Wounds measured from digital photographs using photodigital software: validation and rater reliability. *Wounds*. 2011;23(9):267-75.
67. American Geriatric Society Panel on Persistent Pain in Older Persons. The management of persistent pain in older persons. *American Geriatric Society. J Am Geriatr Soc* 2002;50: June suppl.
68. Reddy M, Kohr R, Queen D, Keast D, Sibbald RG. Practical treatment of wound pain and trauma: a patient-centered approach. An overview. *Ostomy Wound Manage*. 2003;49(4 Suppl):2-15.
69. Woo K, Sibbald G, Fogh K, et al. Assessment and management of persistent (chronic) and total wound pain. *Int Wound J*. 2008;5(2):205-15.