



Inflammatory signaling pathways play a role in SYK inhibitor resistant AML

Sarah Tausch, Christina Villinger, Gabriela Alexe, Daniel J. Urban, Min Shen, Dominique Jahn, Jonas Vischedyk, Sebastian Scheich, Hubert Serve, Matthew D. Hall, Kimberly Stegmaier, Thomas Oellerich, Anjali Cremer

A

**Figure Caption Supplementary Figure 2**

A) Quantification NR3C1 - 3 replicates - Fig. 2A

B) Quantification BIM - 2 replicates - Fig. 2C

C) Quantification NR3C1 - 2 replicates - Fig. 2D

D) Quantification NR3C1- 3 replicates - Fig. 2E


E) Quantification IL3 - 3 replicates - Fig. 3A

F) Quantification IKKB and IKB alpha - 2 replicates - Fig. 4E

For statistical analysis of protein quantification ImageJ was used


(ns, $P \geq 0.05$; *, $P \leq 0.05$; **, $P \leq 0.01$; ***, $P \leq 0.001$;****, $P \leq 0.0001$; t-test)

A




A) Immunoblot Fig. 2A
B) Immunoblot Fig. 2C
C) Immunoblot Fig. 2D
D) Immunoblot Fig. 2E



B



C




D



A) Immunoblot Fig. 3A

B) Immunoblot Fig. 4E

A



B

