

BMJ Open Prevalence and characteristics of hazardous and harmful drinkers receiving general practitioners' brief advice on and support with alcohol consumption in Germany: results of a population survey

Sabrina Kastaun ^{1,2} Claire Garnett ^{3,4} Stefan Wilm ¹ Daniel Kotz ^{2,3}

To cite: Kastaun S, Garnett C, Wilm S, *et al.* Prevalence and characteristics of hazardous and harmful drinkers receiving general practitioners' brief advice on and support with alcohol consumption in Germany: results of a population survey. *BMJ Open* 2022;**12**:e064268. doi:10.1136/bmjopen-2022-064268

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-064268>).

Received 27 April 2022
Accepted 06 September 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Sabrina Kastaun;
sabrina.kastaun@med.uni-duesseldorf.de

ABSTRACT

Objective The German treatment guideline on alcohol-related disorders recommends that general practitioners (GPs) offer brief advice on, and support with, reducing alcohol consumption to hazardous (at risk for health events) and harmful (exhibit health events) drinking patients. We aimed to estimate the implementation of this recommendation using general population data.

Design Cross-sectional analysis of data (2021/2022) of a nationwide, population-based household survey.

Setting Germany.

Participants Population-based sample of 2247 adult respondents who reported hazardous or harmful drinking according to the Alcohol Use Disorders Identification Test-Consumption (AUDIT-C; score women: 4–12 and men: 5–12).

Main outcome measure Ever receipt of 'brief GP advice on, or support with, reducing alcohol consumption'. Differences in the likelihood of ever receiving advice and/or support (yes/no) relative to respondents' sociodemographic, smoking and alcohol consumption characteristics were estimated using logistic regressions.

Results Ever receipt of GP advice on/support with reducing drinking was reported among 6.3% (95% CI=5.3% to 7.4%), and the offer of support among 1.5% (95% CI=1.1% to 2.1%) of the hazardous and harmful drinking respondents. The likelihood of having ever received advice/support was positively associated with being older (OR=1.03 per year, 95% CI=1.01 to 1.04), a current or former (vs never) smoker (OR=2.36, 95% CI=1.46 to 3.80; OR=2.17, 95% CI=1.23 to 3.81) and with increasing alcohol consumption (OR=1.76 per score, 95% CI=1.59 to 1.95). One in two harmful drinking respondents (AUDIT-C score 10–12) reported appropriate advice/support. The likelihood was negatively associated with being woman (eg, OR=0.32, 95% CI=0.21 to 0.48), having a medium or high (vs low) education and with increasing household income.

Conclusions A small proportion of hazardous and harmful drinking people in Germany report having ever received GP advice on, or support with, reducing alcohol consumption. The implementation of advice/support seems to be linked

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ The principal strength of this study is the population-based data collection.
- ⇒ The cross-sectional study design and temporality issues with our measures (alcohol consumption was measured with reference to the present and the outcome as 'ever receipt of GP advice or support') limited our ability to explore causal relationships.
- ⇒ The outcome measure had a complex, not entirely hierarchical structure, which may have led to respondents being unsure of which response option to select.
- ⇒ Data were collected during the COVID-19 pandemic, during which alcohol consumption in Germany seemed to have changed. It is unclear how this might have influenced general practitioners' (GPs) behaviour.
- ⇒ Due to the socially loaded topic, respondents may not have answered truthfully or repressed a previous conversation with their GP on their alcohol consumption. We did not assess the GPs' view on the topic.

to specific socio-demographic characteristics, tobacco smoking and alcohol consumption level. Health policy measures should aim to increase alcohol screening, brief intervention rates and awareness for at-risk populations in primary care.

Trial registration number DRKS00011322, DRKS00017157.

INTRODUCTION

Hazardous drinking is defined as a quantity or a repeated pattern of alcohol consumption that places a person at risk for adverse health events, whereas harmful drinking verifiably results in such events.^{1–3} Alcohol dependence, on the other hand, is seen as a complex drinking pattern, characterised by

persistent consumption despite harmful consequences, craving, the prioritisation of drinking over other activities, tolerance development and withdrawal symptom.¹⁻³ Although this terminology is now commonly used and acknowledges the spectrum of risk that tends to increase with increasing drinking, universal cut-offs with regard to consumption levels and associated risks are missing, making a clear distinction of these theoretical drinking patterns often difficult in practice.¹⁻³

Alcohol misuse contributes to around 3 million deaths each year globally, and is responsible for around 5.1% of global disability-adjusted life-years.⁴ Germany ranks above the average level in the European Union of pure alcohol consumption per capita and year in the adult population (13.4 litres of pure alcohol vs 9.8 litres).⁴ Latest nationally representative prevalence data show that around 20% of the adult population consume alcohol at least at a hazardous level,⁵ including a smaller proportion (approximately 3% per group,⁶) of harmful and alcohol dependent drinkers. There is strong evidence of a dose-response relationship between alcohol consumption and alcohol-related harms.^{7,8} Therefore, people would benefit from reducing their consumption, especially people drinking at a hazardous level.

The implementation of brief interventions in primary healthcare settings is both an effective⁹⁻¹³ and cost-effective⁴ approach to reducing hazardous and harmful drinking. Brief interventions usually include feedback on consumption, brief advice to reduce or quit drinking, motivational enhancement and goal setting and further support such as referral to specialised treatment or the development of a personal reduction plan.^{10,14}

Addressing a patient's alcohol consumption is still perceived as an emotionally difficult and socially loaded issue, and general practitioners (GPs) are thus well placed to address alcohol use disorders and to deliver brief interventions. They commonly see patients of various ages with a broad range of (alcohol-related) health conditions,¹⁴ and the long-lasting patient-GP relationship can help to reduce feelings of stigmatisation and irritation in the patient.² The current German clinical guideline on the treatment of alcohol-related disorders² therefore recommends that brief interventions should be offered to hazardous and harmful drinkers in the primary care setting. According to this guideline, identification of patients at risk can be carried out by means of screening and pragmatic case-finding (ie, when the issue is raised by the patient or when the GP notices alcohol-related conspicuousness). While the latter comes at the costs of missing patients with a risky consumption pattern at an early stage, it seems to be the more common procedure in the GP setting.¹⁵

For individuals with alcohol dependence, evidence on the effectiveness of brief interventions is inconsistent, but according to the German treatment guideline, brief interventions can be a sensible measure among this group for ethical and pragmatic reasons, if other and more intensive interventions (eg, specialised inpatient treatment) are rejected by the patient.²

Despite strong evidence on the effectiveness of brief interventions, international data suggest that its implementation in routine GP practice remains challenging.¹⁶⁻¹⁹ A population survey from England comparing GP advice on drinking with smoking found that only 6.5% of hazardous and harmful drinkers received advice on their alcohol consumption during the past year, compared with half of smokers that received advice on quitting smoking.¹⁸ Although most primary care clinicians report that they ask their patients about alcohol use, far fewer offer advice or recommend treatment.²⁰ A survey which used routine GP data showed that GPs struggled more often at identifying hazardous than dependent drinkers.¹⁹ No representative, national data are available on the implementation of the German clinical guideline on hazardous and harmful drinking in GP settings. Only one study provides initial figures from a single federal state, Bremen, collected in 2016, where 2.9% of all hazardous drinkers were screened by their GPs, and 1.4% received a brief intervention.²¹

Up-to-date and representative national figures on the provision of brief alcohol interventions in GP settings are needed to be able to precisely inform health policy and—if needed—the development of interventions aiming at improving the implementation of treatment recommendations on brief alcohol interventions in primary care. It is also important to report whether the provision of such interventions differs by recipients' characteristics in order to identify potentially underserved groups. This study therefore aimed to explore the following questions using data from a nationwide population-based sample of adults in Germany who self-report hazardous or harmful drinking (operationalised using the Alcohol Use Disorders Identification Test-Consumption (AUDIT-C)).²²

RESEARCH QUESTIONS

Among adults in the population of Germany drinking at hazardous and harmful levels:

1. What proportion reports that a GP ever asked about their alcohol consumption, advised them to drink less, offered help or support with drinking less and offered help with making use of external medical or psychological support because of alcohol consumption?
2. What proportion report ever (vs never) receiving brief GP advice on, or support with, reducing alcohol consumption stratified by person characteristics: age, sex, education, income, smoking status, migration background, region of residence and the overall alcohol consumption?
3. Are there any differences in the likelihood of ever (vs never) receiving brief GP advice on, or support with, reducing alcohol consumption within each measured recipient characteristic?

Since the clinical guideline does not provide a clear recommendation on the exact level of alcohol

consumption at which a brief intervention should take place,² our study will address individuals consuming alcohol at least at a hazardous level. From a preventive medicine perspective, we most likely expect an intervention from GPs among this group of risk.

MATERIALS AND METHODS

Study design

We used data from the cross-sectional German Study on Tobacco Use (DEBRA: ‘Deutsche Befragung zum Rauchverhalten’): an ongoing representative household survey on tobacco and nicotine product use in Germany (www.debra-study.info).²³ This study is conducted by a market research institute and has been registered at the German Clinical Trials Register. Since 2016, the DEBRA study collects data every other month from computer-assisted, face-to-face household interviews of independent samples of people aged >14 years. During the COVID-19 pandemic, interviews were conducted under the applicable COVID-19 protection of the respective federal state where each household was located.

Study population

Data were aggregated from seven survey waves (waves 28–34) collected between February 2021 and February 2022 (N=14 327). Since January 2020, respondents are

selected by using a dual frame design: a composition of random stratified sampling (50% of the sample) and quota sampling (50% of the sample). This sampling design has been described in detail elsewhere: osf.io/s2wxc/. Details on the general sample selection have been published in a study protocol.²³

Alcohol consumption was measured with the AUDIT-C²²; a three-item measure including questions on (1) frequency, (2) quantity and (3) frequency of occasional heavy drinking; full details are given in [table 1](#). The AUDIT-C overall score indicates the level of alcohol consumption and ranges from 0 to 12.²² As recommended in the German treatment guideline on alcohol-related disorders² and in underlying studies,^{24 25} a gender-specific AUDIT-C cut-off score of >5 in men, and of >4 in women was used to operationalise ‘at least hazardous drinking’. Respondents who answered that they never drink alcohol did not receive questions 2 and 3, and were excluded from the analysis. The study population included all adults (aged >18 years, n=14 026) who reported at least hazardous drinking, resulting in a total sample of n=2712 (19.3%) hazardous or harmful drinkers. Respondents aged 14–17 were excluded from the analysis as 16 is the national legal age of sale for beverages containing <15% of alcohol by volume (ABV), and 18 is the legal age of sale for beverages with >15% ABV.

Table 1 Measures of assessing alcohol consumption and ever receiving brief GP advice on/support with reducing consumption

Exposure measure: hazardous and harmful alcohol consumption (AUDIT-C items ²²)	Outcome measure: brief GP advice on alcohol consumption
<p>1. How often do you have a drink containing alcohol?</p> <ul style="list-style-type: none"> ▶ Never (score 0) ▶ At least once a month or less (score 1) ▶ 2–4 times per month (score 2) ▶ 2–3 times per week (score 3) ▶ 4+ times per week (score 4) ▶ No answer <p>2. How many units of alcohol do you drink on a typical day when you are drinking?</p> <ul style="list-style-type: none"> ▶ 1–2 (score 0) ▶ 3–4 (score 1) ▶ 5–6 (score 2) ▶ 7–9 (score 3) ▶ 10+ (score 4) ▶ No answer <p>3. How often have you had six or more units if woman, or eight or more if man, on a single occasion in the last year?</p> <ul style="list-style-type: none"> ▶ Never (score 0) ▶ Less than monthly (score 1) ▶ Monthly (score 2) ▶ Weekly (score 3) ▶ Daily or almost daily (score 4) ▶ No answer 	<p>Has a GP ever asked you about your alcohol consumption and, if so, recommended something to you in this regard?</p> <ol style="list-style-type: none"> 1. No, a GP has never asked me about my alcohol consumption. 2. Yes, a GP has asked me about my alcohol consumption at some point. 3. Yes, a GP has advised me to drink less at some point. 4. Yes, a GP has offered me help or support to drink less at some point. 5. Yes, a GP has advised or helped me to make use of psychological or medical support because of my alcohol consumption at some point. 6. I do not remember if a GP has ever addressed my alcohol consumption. 7. No answer.
AUDIT-C, Alcohol Use Disorders Identification Test-Consumption ; GP, general practitioner.	

Outcome measure

Respondents were asked about 'ever receipt of GP advice on, or support with, reducing alcohol consumption', using a question that was adopted from previous studies on GP advice on smoking cessation in the German,²⁶ and in the Dutch and English population,²⁷ and which was critically reviewed by an experienced GP (see table 1).

For further analyses regarding ever receipt of GP advice on, or support with, reducing alcohol consumption, the response options for this question were dichotomised into 'no' (options 1–2; 0) and 'yes' (options 3–5; 1) (see table 1). Respondents who answered 'I don't remember' (n=443) or refused to answer (n=22) were excluded from the analyses.

Exposure variables

The following socio-demographic characteristics were measured: age; sex (female vs male); region of residence (rural vs urban setting), migration background (yes vs no), current tobacco smoking status (current, former, never) and alcohol consumption as a continuous variable (AUDIT-C score with a possible range of 4–12 among women and of 5–12 among men who drink at least hazardously).

Alcohol consumption varies between regions of residence, with a majority of studies reporting higher drinking rates in rural communities.²⁸ We assumed that this somehow affects the awareness and behaviour of GPs with regard to advice on drinking. This variable was assessed by using the national classification of regions ('BIK Regionsklassifizierung',²⁹), consisting of five categories (eg, metropolitan area or subcentres) which were dichotomised for the analyses: urban versus rural setting.

German population surveys suggest that individuals with migration background relative to those without consume less alcohol.³⁰ Respondents were asked: 'Was one of your parents born abroad?'. Migration background applied if at least one parent did not have German nationality by birth.

Measured socioeconomic status (SES) variables were: educational qualification (low (9 years of education, or no graduation), medium (≥ 10 years), high (≥ 12 years)) and monthly net household income calculated per person (details on the calculation can be found here: <https://osf.io/387fg/>). Income was entered as a continuous variable coded from 0 (€0 income/month) to 7 (\geq €7000/month) in the regression models. For descriptive purposes, income was categorised into: low (<20th percentile), medium (20th–80th percentiles) and high (>80th percentile), approximately reflecting the distribution in the German population.

Statistical analyses

The study protocol and analysis plan was written prior to analysing data and preregistered on the Open Science Framework: osf.io/3fe87.

Data were analysed and reported unweighted as information on true population parameters of the population

of hazardous drinkers were not available. Analyses were performed with IBM SPSS Statistics for Windows, V.27.0 (Armonk, New York, USA: IBM).

To address research question 1, we report descriptive prevalence data on the various levels of GP advice on, or support with, reducing alcohol consumption as percentages together with 95% CIs.

To address research question 2, we present prevalence data including 95% CI of the dichotomous outcome variable 'ever receipt of GP advice on, or support with, reducing alcohol consumption (=yes vs no)' stratified by all categorical exposure variables. For the continuous scaled AUDIT-C score we present a figure showing the prevalence of 'ever receipt of GP advice/support (=yes)' in relation to the range of possible AUDIT-C scores among the sample of hazardous and harmful drinkers.

To address research question 3, a series of univariate logistic regression models were conducted to explore potential differences in the likelihood of ever (yes vs no) receiving brief GP advice on, or support with, reducing alcohol consumption for all exposure variables. Regression models were adjusted for survey wave as a potential confounding variable. As sex differences were found with regard to the likelihood of ever receiving such GP advice/support, it was suggested through the peer-review process to rerun this series of regression models stratified by sex as an exploratory post-hoc analysis.

A considerable number (16.3%) of respondents did not remember if a GP had ever addressed their alcohol consumption and some refused to answer (0.8%). For the analyses of research questions 2 and 3, this group was excluded. We compared whether this group differed systematically from respondents who provided an answer on our outcome question by using the χ^2 test and Mann-Whitney U test (online supplemental table 1).

Dealing with missing data

Missing data were sparse (<0.6% for all variables except for income (2.9%) and migration background (4.8%)), and data were analysed using complete cases. Due to an incorrect questionnaire instruction in four survey waves (28–31), 22.6% of respondents who reported to the first AUDIT-C question with 'at least once a month or less' did not receive the AUDIT-C questions 2 and 3, and were not interviewed on the primary outcome. However, an analysis of AUDIT-C data from earlier waves of the DEBRA study⁵ with the correct questionnaire instruction showed that only 1% of respondents who provided the same answer, were identified as hazardous or harmful drinkers. We thus assume that around 1% of hazardous or harmful drinkers were lost across four of the seven survey waves due to this mistake. These missing data were assumed to be completely at random, and thus excluded from the analyses.

Patient and public involvement statement

Patients or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Table 2 Characteristics of all adult hazardous or harmful drinking respondents (n=2247), and prevalence estimates on the ever receipt of brief GP advice on/support with reducing alcohol consumption (=yes) relative to the respondents' characteristics; including results of univariate regressions models on associations between these characteristics and ever receipt of GP advice

	Total adult sample of hazardous/harmful drinkers†, n=2247, % (n)	Ever GP advice on/support with alcohol consumption, yes vs no (ref), n=142	
		% (n, 95% CI)	OR (95% CI)
Gender			
Male (reference)	55.0 (1235)	9.0 (111, 7.5 to 10.7)	1
Female	45.0 (1012)	3.1 (31, 2.8 to 3.4)	0.32 (0.21 to 0.48)***
Age in years‡§			
18–24	9.9 (222)	1.8 (4, 0.5 to 4.5)	<i>Continuous</i> 1.03 (1.01 to 1.04)***
25–39	21.5 (482)	4.8 (23, 3.0 to 7.1)	
40–64	48.2 (1083)	6.5 (70, 5.1 to 8.1)	
65+	20.5 (460)	9.8 (45, 7.2 to 12.9)	
Educational qualification¶			
Low (ref.)	24.2 (543)	10.7 (58, 8.2 to 13.6)	1
Medium	40.6 (913)	5.5 (50, 4.1 to 7.2)	0.48 (0.33 to 0.72)***
High	34.6 (778)	4.4 (34, 3.0 to 6.1)	0.38 (0.25 to 0.59)***
Household income§††			
Low	9.3 (209)	14.8 (31, 10.3 to 20.4)	<i>Continuous</i> 0.60 (0.48 to 0.76)***
Medium	58.7 (1319)	5.9 (78, 4.7 to 7.3)	
High	29.1 (654)	4.6 (30, 3.1 to 6.5)	
Migration background			
No (ref.)	84.4 (1896)	5.3 (101, 4.4 to 6.4)	1
Yes	10.9 (244)	6.6 (16, 4.0 to 10.4)	1.27 (0.74 to 2.20)
Region of residence			
Urban setting (ref.)	63.7 (1431)	5.9 (85, 4.8 to 7.3)	1
Rural setting	36.3 (816)	7.0 (57, 5.3 to 9.0)	1.19 (0.84 to 1.67)
Tobacco smoking status			
Never smoker (ref.)	29.0 (652)	3.4 (22, 2.1 to 5.1)	1
Former smoker	19.0 (427)	7.0 (30, 4.8 to 9.9)	2.17 (1.23 to 3.81)**
Current smoker	51.9 (1166)	7.6 (89, 6.2 to 9.3)	2.36 (1.46 to 3.80)***
Alcohol consumption per AUDIT-C level††§			1.76 (1.59 to 1.95)***

Data are presented as column percentages (number), row percentages (number, 95% CIs) and as ORs together with 95% CI around OR. ORs were adjusted for the variable 'survey wave' (as design factor); ***p<0.001, **p<0.01, *p<0.05.

Variables with missing data: educational qualification: 0.6%, household income: 2.9%, migration background: 4.8%, smoking status: 0.1%. †According to the Alcohol Use Disorders Identification Test-Consumption (AUDIT-C)²²; an AUDIT-C score of >5 in male respondents, and of >4 in female respondents was used to operationalise hazardous drinking.^{24 25}

‡Gender-specific mean values (including median and SD) for the sample are reported in the results section of this manuscript.

§Entered as continuous variable in regression analysis.

¶German educational qualification levels: low (9 years of education, or no graduation), medium (10 years of education), high (≥12 years of education).

††Monthly net household income per person in the household, based on the Organisation for Economic Co-operation and Development-modified equivalence scale.²⁹ The variable was categorised into three levels: low (<20th income percentile), medium (20th–80th income percentiles) and high (>80th income percentile), approximately reflecting the distribution of income in the German population.^{30–32}

RESULTS

The final analytical sample consisted of all adult respondents who reported at least hazardous drinking, and provided an answer on, or could remember, whether or not a GP had advised them on reducing alcohol consumption

(n=2247). The sample is described in table 2. The mean age of this group was 49.5 years (SD=17.0), and 45.0% (n=1012) of the respondents were women. The mean AUDIT-C score was 5.9 (median: 6, SD=1.3) for men and 4.8 (median: 4, SD=1.1) for women.




Figure 1 Prevalence estimates on the various levels of GP advice on, or support with, reducing alcohol consumption (self-reported) among the total sample of adult hazardous or harmful drinkers (n=2247) reported as percentages together with 95% CIs. GP, general practitioner.

Respondents who did not remember whether or not they received GP advice seem to be more often current smokers and lower educated but did not systematically differ from respondents who did remember (online supplemental table 1).

Proportion of hazardous or harmful drinking adults reporting various levels of GP advice/support

Among hazardous or harmful drinkers, 82.2% (95% CI=80.5% to 83.7%; n=1846) reported that a GP had never asked them about their alcohol consumption, 11.5% (95% CI=10.2% to 12.9%; n=259) reported that they had ever been asked about drinking by a GP and 6.3% (95% CI=5.3% to 7.4%; n=142) said that a GP had ever advised them to drink less, or offered either help or support or advised or helped them to make use of medical or psychological support to drink less (see figure 1). Such support was reported by 1.5% (95% CI=1.1% to 2.1%) of the hazardous and harmful drinking respondents.

Proportion of hazardous or harmful drinking adults reporting ever receipt of GP advice/support stratified by recipients' characteristics

Men (compared with women), respondents of higher age (65+ years), those with low (compared with medium and high) education and household income and current and former smokers (compared with never smokers) reported relatively more often to have ever received brief GP advice on, or support with alcohol consumption (see table 2).

The higher the overall AUDIT-C score, the higher the rate of reporting ever receipt of GP advice on, or support with, reducing alcohol consumption (see figure 2). In persons with an AUDIT-C score >9—which is viewed as




Figure 2 Ever receipt of GP advice on, or support with, reducing alcohol consumption (yes, self-reported) relative to the total AUDIT-C score among the total sample of adult hazardous or harmful drinkers (n=2247) reported as percentages together with 95% CIs (black line; dotted line: trend line, polynomial function, $R^2=0.97$). AUDIT-C, Alcohol Use Disorders Identification Test-Consumption; GP, general practitioner.

harmful or potentially dependent drinking pattern³¹—51.5% (n=17/33) have ever received GP advice on, or support with reducing alcohol consumption.

Likelihood of ever receiving GP advice/support relative to recipients' characteristics

The likelihood of ever receipt of brief GP advice on, or support with, reducing alcohol consumption was positively associated with being older, being a former or a current smoker and reporting a higher alcohol consumption level at the time of the survey (see table 2). The likelihood was negatively associated with being woman, having medium and high (compared with low) educational qualification and with increasing monthly household income.

No significant differences were detected relative to the respondents' migration background or region of residence.

Results of explorative, ancillary post-hoc regression analyses stratified by sex indicate that particularly the negative association between medium and high (compared with low) educational qualification and ever receipt of GP advice/support seem to occur in men rather than in women (online supplemental table 2).

DISCUSSION

In a large population-based sample of the general population in Germany, adults who report hazardous or harmful drinking, about 12% reported having ever been asked by a GP about their alcohol consumption, and about 6% reported having ever received GP advice to drink less, including 1.5% who were also offered support to drink less or to make use of psychological or medical assistance for that purpose. However, in the subgroup of people reporting harmful or potentially dependent drinking

(AUDIT-C score 10–12,³²) around half had ever received brief GP advice on, or support with, reducing alcohol consumption.

To our knowledge, this is the first nationwide study to estimate the implementation of the clinical guideline recommendation on the provision of GPs' brief advice on, or support with alcohol consumption in the population of Germany. Our findings are broadly consistent with a previous study (not methodologically comparable) in a single federal state in Germany,²¹ and a population survey in England (with comparable methodology).¹⁸ Both studies indicate insufficient implementation of brief alcohol interventions in primary care.

In the current study there were significant differences in the likelihood of having ever received GP advice or offer of support by personal characteristics. Hazardous and harmful drinkers of older age, current or former smokers and those with higher alcohol consumption had higher odds of reporting ever receipt of GP advice on, or support with alcohol consumption, whereas women (vs men) and respondents with higher income or medium and high (vs low) educational qualification had substantially lower odds. Higher likelihood of receiving GP advice on alcohol consumption among older respondents and among current or former smokers might be associated with awareness among GPs of existing (alcohol-related) health conditions and of polysubstance use, as well as with higher demand for treatment among these patients. A further possible explanation might be that older patients have had more GP visits during their lifespan and therefore more opportunities of receiving appropriate advice or support. The implementation of alcohol screening and brief intervention in individuals with comorbidities seems to be largely accepted among GPs and higher than in those without comorbidities.^{33–36}

Our findings suggest that hazardous and harmful drinking women and those with higher SES are under-recognised by GPs when it comes to brief alcohol intervention. This is concerning as, although prevalence of hazardous and harmful drinking is higher among men, yet around 1 in 10 women report drinking at least at a hazardous level.⁵ In addition, evidence is good that individuals with higher compared with with lower SES may consume similar or even greater amounts of alcohol and show higher prevalence rates of hazardous drinking.^{5 37} Gender gaps in GP-delivered alcohol interventions have been reported before^{18 38 39} but we can only speculate on underlying reasons. This could be due to greater concerns about stigmatisation or shame, leading women and individuals with higher SES less often admit alcohol use to their GPs (social desirability). On the other hand, implicit cognitive bias and stereotypes might influence the GPs' decision on who to screen for alcohol misuse.⁴⁰ Another possible explanation might be that specific groups of society are less likely to consult a GP, which has been reported for higher SES groups,^{41 42} though is not the case for women who tend to visit their GP more often than men.⁴³

Implications for policy and practise

GPs are a major force to improve the prevention of alcohol-related harm on a population level, and as this study shows, they already intervene in about half of harmful drinking patients. However, from a preventive medicine perspective, this study reveals a need to improve the implementation of guideline recommendations for hazardous drinking in the GP setting. While the clinical guideline does not provide a clear recommendation on the exact level of alcohol consumption at which a brief intervention should take place,² alcohol-related harms are dose-dependent and therefore it is important to provide brief interventions for all individuals at alcohol-related risk. Previous studies showed that education and post-graduate training predicts the GP delivery of brief alcohol interventions to hazardous drinkers,³³ and that training GPs can significantly increase alcohol screening and brief intervention rates in primary care.⁴⁴ This is particularly the case when being tailored to the barriers and facilitators towards the implementation of such interventions in the GP setting and when being developed on the basis of a behaviour change theory.⁴⁵ Further synergistic effects have been shown when financial incentives, training and support were offered together.⁴⁶

In Germany, appropriate training is not offered by default, neither during medical education nor as post-graduate training for physicians. As a consequence, many GPs,⁴⁷ as well as medical students,⁴⁸ in Germany do not feel adequately trained to diagnose and treat patients with alcohol problems. This lack of training had also been identified as a major barrier towards the routine implementation of brief alcohol intervention in primary care in the UK.¹⁶ Other frequently reported barriers comprise the lack of financial reimbursement and the lack of time in primary care settings.⁴⁹ With this regard, alcohol screenings could be, for example, integrated into the regular health check-ups for patients aged 35 and over that already exist in Germany.

Strengths and limitations

A major strength of this study is the large, population-based sample. However, there are also limitations. First, data was self-reported, introducing risk for recall bias that may have affected the prevalence estimates, most likely resulting in an underestimate. Second, the cross-sectional study design and that GP advice and support was measured as 'ever receipt', whereas alcohol consumption was measured with approximate reference to the present, limited our ability to explore causal relationships. Comparable temporality issues might also occur for some of the socio-demographic characteristics such as income or place of residence. Based on this, we did not conduct multivariable regression analyses, and our results are not adjusted for potential confounding through interaction effects between the exposure variables. Future research should look to estimate the potential causal effect of someone's characteristics and the likelihood of

ever receipt of GP advice on, or support with, reducing alcohol consumption.

Third, the question on ‘ever receipt of GP advice and support’ had a complex structure (participants are asked about GP advice on as well as different types of support with drinking less), which could have led to difficulties in understanding. However, the face-to-face interview method and support available from the interviewers may have mitigated this risk. In addition, this measure did not follow an exact hierarchical structure and there was no clear distinction between ‘internal support’ (eg, offered by the GP), and ‘external support’ (eg, psycho-social services). This may have led to respondents being unsure of which response option to select. Fourth, data were collected during the COVID-19 pandemic, during which alcohol consumption in Germany seemed to have decreased on average but increased in specific subgroups of the population.^{50 51} It is unclear how this might have influenced the behaviour of GPs. Fifth, due to the socially loaded topic, respondents may not have answered truthfully or repressed a previous conversation with their GP on the topic. It is therefore important to also assess the GPs’ view on the topic, including in-depth information on barriers to the implementation of the treatment guideline recommendations in their daily practice. Previous surveys among GPs, however, usually only assessed how often, in general, GPs screen for alcohol or provide brief intervention, but not in relation to the number of their hazardous drinking patients.^{33 52} Finally, the approach of the market research institute conducting the survey does not allow for calculation of a response rate or to characterise non-participants.

CONCLUSION

Our findings suggest that hazardous drinking—that places a person at risk for adverse health events—does not seem to be adequately addressed by GPs in Germany, particularly among women and individuals from higher SES groups. From a preventive medicine perspective, this leads to missed opportunities to reduce alcohol-related harm. However, the probability of having ever received brief alcohol intervention by a GP increases with increasing drinking levels. Around every second harmful drinking person seems to have ever received brief GP advice on, or support with, reducing alcohol consumption. Although we did not analyse causal relationships, this study is a call for action in order to further explore underlying reasons why clinical guideline recommendations on brief alcohol interventions are not implemented more often in the German general practice setting, as well as to explore potential approaches for improvement.

Author affiliations

¹Institute of General Practice, Patient–Physician Communication Research Unit, Centre for Health and Society, Medical Faculty of the Heinrich-Heine-University Düsseldorf, Düsseldorf, Germany

²Institute of General Practice, Addiction Research and Clinical Epidemiology Unit, Centre for Health and Society, Medical Faculty of the Heinrich-Heine-University Düsseldorf, Düsseldorf, Germany

³Department of Behavioural Science and Health, University College London, London, UK

⁴SPECTRUM Consortium, London, UK

Acknowledgements The authors thank Erika Baum for her constructive feedback on the outcome measure (the question on general practitioner advice on alcohol consumption) as well as on the first version of the analysis protocol. The authors also thank Constanze Cholmakow-Bodechtel and Franziska Weng from the market research institute ‘Kantar’ for the collection of the data.

Contributors SK coordinates the DEBRA study, conceptualised and drafted the analysis protocol, drafted the manuscript, analysed and interpreted the data. SK also acts as guarantor for this study. CG and SW provided expert advice on the study protocol and critically revised the analysis protocol and the manuscript. DK conceived the DEBRA study, supervised the analyses and critically revised the analysis protocol and the manuscript. All named authors contributed substantially to the manuscript and agreed on its final version.

Funding From 2016 to 2019 (waves 1–18), the DEBRA study was supported by the Ministry of Innovation, Science and Research of the German State of North Rhine–Westphalia (MIWF) in the context of the ‘NRW Rückkehrprogramm’ (the North Rhine–Westphalian postdoc return programme). Since 2019 (wave 19 onwards), the study has been supported by the German Federal Ministry of Health. CG is funded by Cancer Research UK and the National Institute for Health Research.

Competing interests CG is a paid scientific consultant for the behaviour change and lifestyle organisation, One Year No Beer.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and was approved by medical ethics committee of the Heinrich-Heine-University Düsseldorf (HHU 5386R). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. The data underlying this study are third-party data and are available to researchers on reasonable request from the corresponding author (sabrina.kastaun@med.uni-duesseldorf.de). All proposals requesting data access will need to specify how it is planned to use the data, and all proposals will need approval of the DEBRA study team before data release.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Sabrina Kastaun <http://orcid.org/0000-0002-5590-1135>

Claire Garnett <http://orcid.org/0000-0002-6589-299X>

Stefan Wilm <http://orcid.org/0000-0002-1266-5064>

Daniel Kotz <http://orcid.org/0000-0002-9454-023X>

REFERENCES

- 1 Saunders JB, Aasland OG, Babor TF, *et al*. Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO Collaborative Project on Early Detection of Persons with Harmful Alcohol Consumption—II. *Addiction* 1993;88:791–804.
- 2 Association of the Scientific Medical Societies (AWMF) [Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen

- Fachgesellschaften (AWMF)]. S3 Guideline "Screening, diagnostics, and treatment of alcohol use disorders" [S3-Leitlinie "Screening, Diagnostik und Behandlung alkoholbezogener Störungen"]. AWMF-Register Nr. 076-001, 2020. Available: https://www.awmf.org/uploads/tx_szleitlinien/076-001_S3-Screening-Diagnose-Behandlung-alkoholbezogene-Stoerungen_2021-02.pdf [Accessed 13 Feb 2022].
- 3 Alcohol change UK how do we talk about alcohol? Available: <https://alcoholchange.org.uk/policy/policy-insights/how-do-we-talk-about-alcohol> [Accessed 26 Feb 2022].
 - 4 World Health Organization. Global status report on alcohol and health 2018, 2018. Available: <https://apps.who.int/iris/handle/10665/274603> [Accessed 13 Feb 2022].
 - 5 Garnett C, Kastaun S, Brown J, *et al.* Alcohol consumption and associations with sociodemographic and health-related characteristics in Germany: a population survey. *Addict Behav* 2022;125:107159.
 - 6 Atzendorf J, Rauschert C, Seitz N-N. The use of alcohol, tobacco, illegal drugs and medicines. *Dtsch Arztebl Int* 2019;116:577–84.
 - 7 Alcohol and Public Policy Group. Alcohol: no ordinary commodity--a summary of the second edition. *Addiction* 2010;105:769–79.
 - 8 University of Sheffield. Mortality and morbidity risks from alcohol consumption in Australia: analyses using an Australian adaptation of the Sheffield alcohol policy model (v2.7) to inform the development of new alcohol guidelines, 2019. Available: <https://www.nhmrc.gov.au/sites/default/files/documents/attachments/Alcohol/10-mortality-morbidity-risks-from-alcohol-cons.pdf> [Accessed 13 Mar 2022].
 - 9 Kaner EFS, Dickinson HO, Beyer F, *et al.* The effectiveness of brief alcohol interventions in primary care settings: a systematic review. *Drug Alcohol Rev* 2009;28:301–23.
 - 10 Kaner EF, Beyer FR, Muirhead C, *et al.* Effectiveness of brief alcohol interventions in primary care populations. *Cochrane Database Syst Rev* 2018;2:CD004148.
 - 11 Tanner-Smith EE, Parr NJ, Schweer-Collins M, *et al.* Effects of brief substance use interventions delivered in general medical settings: a systematic review and meta-analysis. *Addiction* 2022;117:877–89.
 - 12 Elzerbi C, Donoghue K, Drummond C. A comparison of the efficacy of brief interventions to reduce hazardous and harmful alcohol consumption between European and non-European countries: a systematic review and meta-analysis of randomized controlled trials. *Addiction* 2015;110:1082–91.
 - 13 Ghosh A, Singh P, Das N, *et al.* Efficacy of brief intervention for harmful and hazardous alcohol use: a systematic review and meta-analysis of studies from low middle-income countries. *Addiction* 2022;117:545–58.
 - 14 Whitlock EP, Polen MR, Green CA, *et al.* Behavioral counseling interventions in primary care to reduce risky/harmful alcohol use by adults: a summary of the evidence for the U.S. preventive services Task force. *Ann Intern Med* 2004;140:557–68.
 - 15 Lid TG, Malterud K. General practitioners' strategies to identify alcohol problems: a focus group study. *Scand J Prim Health Care* 2012;30:64–9.
 - 16 Wilson GB, Lock CA, Heather N, *et al.* Intervention against excessive alcohol consumption in primary health care: a survey of GPs' attitudes and practices in England 10 years on. *Alcohol Alcohol* 2011;46:570–7.
 - 17 Geirsson M, Bendtsen P, Spak F. Attitudes of Swedish general practitioners and nurses to working with lifestyle change, with special reference to alcohol consumption. *Alcohol Alcohol* 2005;40:388–93.
 - 18 Brown J, West R, Angus C, *et al.* Comparison of brief interventions in primary care on smoking and excessive alcohol consumption: a population survey in England. *Br J Gen Pract* 2016;66:e1–9.
 - 19 Cheeta S, Drummond C, Oyefeso A, *et al.* Low identification of alcohol use disorders in general practice in England. *Addiction* 2008;103:766–73.
 - 20 Friedmann PD, McCullough D, Chin MH, *et al.* Screening and intervention for alcohol problems. *J Gen Intern Med* 2000;15:84–91.
 - 21 Manthey J, Lindemann C, Verthein U. Care for people with hazardous alcohol consumption and severe alcohol use disorder in Bremen: in line with needs and guidelines? [Versorgung von Personen mit riskantem Alkoholkonsum und schwerer Alkoholkonsumstörung in Bremen: bedarfsgerecht und leitlinienkonform?]. *Bundesgesundheitsblatt - Gesundheitsforschung - Gesundheitsschutz* 2020;63:122–30.
 - 22 Bradley KA, DeBenedetti AF, Volk RJ, *et al.* AUDIT-C as a brief screen for alcohol misuse in primary care. *Alcohol Clin Exp Res* 2007;31:1208–17.
 - 23 Kastaun S, Brown J, Brose LS, *et al.* Study protocol of the German study on tobacco use (DEBRA): a national household survey of smoking behaviour and cessation. *BMC Public Health* 2017;17:378.
 - 24 Ganz T, Braun M, Laging M. Screening for hazardous drinking among German university students: criterion validity and cut-off scores for the alcohol use disorders identification Test–Consumption. *Z Klin Psychol Psychother* 2017;46:187–97.
 - 25 Rumpf H-J, Hapke U, Meyer C, *et al.* Screening for alcohol use disorders and at-risk drinking in the general population: psychometric performance of three questionnaires. *Alcohol Alcohol* 2002;37:261–8.
 - 26 Kastaun S, Kotz D. Brief physician advice for smoking cessation: Results of the DEBRA study [Ärztliche Kurzberatung zur Tabakentwöhnung – Ergebnisse der DEBRA Studie]. *SUCHT* 2019;65:34–41.
 - 27 Kotz D, Willemsen MC, Brown J, *et al.* Light smokers are less likely to receive advice to quit from their GP than moderate-to-heavy smokers: a comparison of national survey data from the Netherlands and England. *Eur J Gen Pract* 2013;19:99–105.
 - 28 Friesen EL, Bailey J, Hyett S, *et al.* Hazardous alcohol use and alcohol-related harm in rural and remote communities: a scoping review. *Lancet Public Health* 2022;7:e177–87.
 - 29 BIK Aschpurwis +Behrens Market, Media and Regional Research [BIK Aschpurwis +Behrens Markt-, Media und Regionalforschung] BIK Regions [BIK Regionen 2020 (780er Systematik)]. Available: https://www.bik-gmbh.de/download/BIK-Regionen_780_Kurzbeschreibung.pdf [Accessed 11 Feb 2022].
 - 30 Rommel A, Saß AC, Born S, *et al.* [Health status of people with a migrant background and impact of socio-economic factors: First results of the German Health Interview and Examination Survey for Adults (DEGS1)]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2015;58:543–52.
 - 31 Hilde P, Ingeborg R, Anne B. Are short audit screeners effective in identifying unhealthy drinking of varying severity? A prison population study. *Drug Alcohol Depend* 2021;229:109153.
 - 32 Khadjesari Z, White IR, McCambridge J, *et al.* Validation of the AUDIT-C in adults seeking help with their drinking online. *Addict Sci Clin Pract* 2017;12:2.
 - 33 Rehm J, Prieto JAA, Beier M, *et al.* The role of alcohol in the management of hypertension in patients in European primary health care practices - a survey in the largest European Union countries. *BMC Fam Pract* 2016;17:130.
 - 34 Hanschmidt F, Manthey J, Kraus L, *et al.* Barriers to alcohol screening among hypertensive patients and the role of stigma: lessons for the implementation of screening and brief interventions in European primary care settings. *Alcohol Alcohol* 2017;52:572–9.
 - 35 Kraus L, Schulte B, Manthey J, *et al.* Alcohol screening and alcohol interventions among patients with hypertension in primary health care: an empirical survey of German general practitioners. *Addict Res Theory* 2017;25:285–92.
 - 36 Rehm J, Allamani A, Elekes Z, *et al.* Alcohol dependence and treatment utilization in Europe - a representative cross-sectional study in primary care. *BMC Fam Pract* 2015;16:90.
 - 37 Pabst A, van der Auwera S, Piontek D, *et al.* Decomposing social inequalities in alcohol consumption in Germany 1995–2015: an age-period-cohort analysis. *Addiction* 2019;114:1359–68.
 - 38 Mauro PM, Askari MS, Han BH. Gender differences in any alcohol screening and discussions with providers among older adults in the United States, 2015 to 2019. *Alcohol Clin Exp Res* 2021;45:1812–20.
 - 39 Delpech R, Bloy G, Panjo H, *et al.* Physicians' preventive practices: more frequently performed for male patients and by female physicians. *BMC Health Serv Res* 2020;20:331.
 - 40 O'Sullivan ED, Schofield SJ. Cognitive bias in clinical medicine. *J R Coll Physicians Edinb* 2018;48:225–32.
 - 41 Krause L, Dini L, Prütz F. Gynaecology and general practitioner services utilisation by women in the age group 50 years and older. *J Health Monit* 2020;5:15–25.
 - 42 Luppma M, Giersdorf J, Riedel-Heller S, *et al.* Frequent attenders in the German healthcare system: determinants of high utilization of primary care services. results from the cross-sectional German health interview and examination survey for adults (degS). *BMC Fam Pract* 2020;21:1–10.
 - 43 Wang Y, Hunt K, Nazareth I, *et al.* Do men consult less than women? An analysis of routinely collected UK general practice data. *BMJ Open* 2013;3:e003320.
 - 44 Friedmann PD, Rose J, Hayaki J, *et al.* Training primary care clinicians in maintenance care for moderated alcohol use. *J Gen Intern Med* 2006;21:1269–75.
 - 45 Rosário F, Vasiljevic M, Pas L, *et al.* Efficacy of a theory-driven program to implement alcohol screening and brief interventions in primary health-care: a cluster randomized controlled trial. *Addiction* 2022;117:1609–21.
 - 46 Anderson P, Bendtsen P, Spak F, *et al.* Improving the delivery of brief interventions for heavy drinking in primary health care:



- outcome results of the optimizing delivery of health care intervention (ODHIN) five-country cluster randomized factorial trial. *Addiction* 2016;111:1935–45.
- 47 Hoffmann T, Voigt K, Kugler J, *et al.* Are German family practitioners and psychiatrists sufficiently trained to diagnose and treat patients with alcohol problems? *BMC Fam Pract* 2019;20:115.
- 48 Hoffmann T, Voigt K, Schließke L. How fit are medical students from Dresden to treat alcohol-related disorders? *ZFA* 2015;91:315–20.
- 49 Hanschmidt F, Manthey J, Kraus L, *et al.* Barriers to alcohol screening among hypertensive patients and the role of stigma: lessons for the implementation of screening and brief interventions in European primary care settings. *Alcohol Alcohol* 2017;52:572–9.
- 50 Manthey J, Kilian C, Schomerus G. Alcohol Use in Germany and Europe during the SARS-CoV-2 Pandemic [Alkoholkonsum in Deutschland und Europa während der SARS-CoV-2 Pandemie]. *SUCHT* 2020;66:247–58.
- 51 Klosterhalfen S, Kotz D, Kastaun S. Did smoking, alcohol consumption, and physical activity change during the COVID-19 restrictions in Germany in spring 2020? *SUCHT* 2022;68:129–38.
- 52 Zaidi Touis L, Bolbrinker J, Riemer TG, *et al.* Moderation of alcohol consumption as a recommendation in European hypertension management guidelines: a survey on awareness, screening and implementation among European physicians. *BMJ Open* 2018;8:e022026.