


Draft Genome Sequences of Two New Zealand *Xanthomonas campestris* pv. *campestris* Isolates, ICMP 4013 and ICMP 21080

Dhairyasheel Desai,^{a,b} Jin-Hua Li,^a Eline van Zijll de Jong,^a Robert Braun,^b Andrew Pitman,^{a,b} Sandra Visnovsky,^b John Hampton,^a  Mary Christey^b

Bio-Protection Research Centre, Lincoln University, Canterbury, New Zealand^a; New Zealand Institute for Plant and Food Research Ltd, Lincoln, New Zealand^b

***Xanthomonas campestris* pv. *campestris* is a necrotrophic bacterial pathogen of crucifers. We report here the draft genome sequences of isolates ICMP 4013 and ICMP 21080 from New Zealand. These sequences will facilitate the identification of race-specific factors in *X. campestris* pv. *campestris*.**

Received 10 September 2015 Accepted 15 September 2015 Published 29 October 2015

Citation Desai D, Li J-H, van Zijll de Jong E, Braun R, Pitman A, Visnovsky S, Hampton J, Christey M. 2015. Draft genome sequences of two New Zealand *Xanthomonas campestris* pv. *campestris* isolates, ICMP 4013 and ICMP 21080. *Genome Announc* 3(5):e01247-15. doi:10.1128/genomeA.01247-15.

Copyright © 2015 Desai et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Mary Christey, mary.christey@plantandfood.co.nz.

Black rot, caused by *Xanthomonas campestris* pv. *campestris*, is one of the most devastating diseases of cruciferous crops. In *X. campestris* pv. *campestris*, nine races have been described (1, 2) and the genome sequences are available for races 1, 3 and 9 (3–5). No sequences of isolates from races 2 and 4 have been published, despite race 4 being considered one of the most prevalent worldwide (6).

X. campestris pv. *campestris* is present on New Zealand crucifer crops. To identify the effectors and other virulence determinants associated with different isolates, the genome sequences of New Zealand isolates ICMP 4013 and ICMP 21080 were obtained by Illumina-based sequencing. Initially, pure cultures of each isolate were incubated overnight in LB media at 28°C. Genomic DNA was then isolated using the QIAamp DNA extraction kit (Qiagen). DNA samples with acceptable quality were then used to generate Illumina HiSeq2000 100-bp paired-end reads (Macrogen).

Approximately 5.6 Gbp and 6.1 Gbp of raw data were generated for ICMP 4013 and ICMP 21080, respectively. A total of 62,928,922 × 2 reads for ICMP 4013 and 69,035,858 × 2 reads for ICMP 21080 were subjected to quality control and trimming using Galaxy (7), and were then *de novo* assembled into contigs using a *k*-mer size of 75 in Velvet version 1.2.08 (8). The draft genome of ICMP 4013 is 4,972,211 bp in length, consisting of 80 contigs with an average 2479-fold coverage and an *N*₅₀ of 29,472. The G+C content of the genome is 64.55 mol% and contains 4,428 predicted coding sequences (CDSs) and at least 60 tRNAs. The draft genome of ICMP 21080 is 4,927,810 bp, consisting of 412 contigs with a 2719-fold coverage and an *N*₅₀ of 36489. The genome has a G+C content of 64.60 mol% and contains 4,371 predicted CDSs and 60 tRNAs.

Genes in both genomes were annotated by searching against the entire collection of FIGfams in the RAST (9) server. A function-based comparison of metabolic reconstruction (9) was then conducted to assess the differential functional roles in the genomes. The comparison revealed 12 functional roles differentially present in ICMP 4013 and eight in ICMP 21080.

The genomes of ICMP 4013 and ICMP 21080 were also com-

pared to other *Xanthomonas* spp. by determining the “closest neighboring genomes” in RAST. Both genomes were closely related to that of the *X. campestris* pv. *campestris* type strain ATCC 33913. Further comparison of ICMP 4013 to ATCC 33913 revealed 61 functional roles differentially present in ATCC 33913 and 94 in ICMP 4013. The comparison of ICMP 21080 to ATCC 33913 revealed 70 functional roles differentially present in ATCC 33913 and 97 differentially present in ICMP 21080.

This is the first report of the genome sequencing of *X. campestris* pv. *campestris* isolates from New Zealand, which adds to the genomic data available and will consequently contribute to our knowledge of genomic diversity among isolates of *X. campestris* pv. *campestris*.

Nucleotide sequence accession numbers. The genome sequences of *Xanthomonas campestris* pv. *campestris* ICMP 4013 and ICMP 21080 have been deposited in GenBank under accession numbers CP012146 and CP012145, respectively.

ACKNOWLEDGMENT

This work was funded by MBIE from contract LINX0702.

REFERENCES

- Fargier E, Manceau C. 2007. Pathogenicity assays restrict the species *Xanthomonas campestris* into three pathovars and reveal nine races within *X. campestris* pv. *campestris*. *Plant Pathol* 56:805–818. <http://dx.doi.org/10.1111/j.1365-3059.2007.01648.x>.
- Vicente JG, Holub EB. 2013. *Xanthomonas campestris* pv. *campestris* (cause of black rot of crucifers) in the genomic era is still a worldwide threat to brassica crops. *Mol Plant Pathol* 14:2–18. <http://dx.doi.org/10.1111/j.1364-3703.2012.00833.x>.
- Bolot S, Guy E, Carrere S, Barbe V, Arlat M, Noël LD. 2013. Genome sequence of *Xanthomonas campestris* pv. *campestris* strain Xca5. *Genome Announc* 1(1):e00032-12. <http://dx.doi.org/10.1128/genomeA.00032-12>.
- Da Silva ACR, Ferro JA, Reinach FC, Farah CS, Furlan LR, Quaggio RB, Monteiro-Vitorello CB, Van Sluys MAV, Almeida NF, Alves LMC, do Amaral AM, Bertolini MC, Camargo LEA, Camarotte G, Cannavan F, Cardozo J, Chambergo F, Ciapina LP, Cicarelli RMB, Coutinho LL, Cursino-Santos JR, El-Dorri H, Faria JB, Ferreira AJS, Ferreira RCC, Ferro MIT, Formighieri EF, Franco MC, Greggio CC, Gruber A, Katsuyama AM, Kishi LT, Leite RP, Lemos EGM, Lemos MVF, Locali EC,

- Machado MA, Madeira AMBN, Martinez-Rossi NM, Martins EC, Meidanis J, Menck CFM, Miyaki CY, Moon DH, Moreira LM, Novo MTM, Okura VK, Oliveira MC, Oliveira VR, Pereira HA, Rossi A, Sena JAD, Silva C, de Souza RF, Spinola LAF, Takita MA, Tamura RE, Teixeira EC, Tezza RID, Trindade dos Santos M, Truffi D, Tsai SM, White FF, Setubal JC, Kitajima JP. 2002. Comparison of the genomes of two *Xanthomonas* pathogens with differing host specificities. *Nature* 417:459–463. <http://dx.doi.org/10.1038/417459a>.
5. Qian W, Jia Y, Ren SX, He YQ, Feng JX, Lu LF, Sun Q, Ying G, Tang DJ, Tang H, Wu W, Hao P, Wang L, Jiang BL, Zeng S, Gu WY, Lu G, Rong L, Tian Y, Yao Z, Fu G, Chen B, Fang R, Qiang B, Chen Z, Zhao GP, Tang JL, He C. 2005. Comparative and functional genomic analyses of the pathogenicity of phytopathogen *Xanthomonas campestris* pv. *campestris*. *Genome Res* 15:757–767. <http://dx.doi.org/10.1101/gr.3378705>.
 6. Vicente JG, Conway J, Roberts SJ, Taylor JD. 2001. Identification and origin of *Xanthomonas campestris* pv. *campestris* races and related pathogens. *Phytopathology* 91:492–499. <http://dx.doi.org/10.1094/PHYTO.2001.91.5.492>.
 7. Goecks J, Nekrutenko A, Taylor J, Galaxy Team T. 2010. Galaxy: a comprehensive approach for supporting accessible, reproducible, and transparent computational research in the life sciences. *Genome Biol* 11:R86. <http://dx.doi.org/10.1186/gb-2010-11-8-r86>.
 8. Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res* 18:821–829. <http://dx.doi.org/10.1101/gr.074492.107>.
 9. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.