

MEETING ABSTRACT

Open Access

Intensive versus conventional insulinotherapy after elective and on-pump myocardial revascularization in the elderly patient: a prospective and randomized study

S Caroleo^{1*}, F Onorati², A Rubino², F Calandese¹, C De Munda¹, E Santangelo³, A Renzulli², B Amantea¹

From de Senectute: Age and Health Forum
Catanzaro, Italy. 5-7 December 2009

Background

In cardiac surgery patients, hyperglycemia was found to be an independent post-operative risk factor for the development of hyperlactatemia and associated with increased morbidity and mortality [1].

Materials and methods

Design: analysis of a prospective and randomized collected database.

Patients: A total of 40 patients (23M; 17F) aged >65 (ASA II-III) submitted to elective on-pump myocardial revascularization from September 2007 to May 2009.

Randomization: on ICU admission, once obtained a preoperative informed consent, the patients were randomly assigned to Group 1 (intensive insulinotherapy during the first 24 hours of ICU stay aimed at glucose levels between 80-110 mg/dl) or Group 2 (conventional insulinotherapy during the first 24 hours of ICU stay aimed at glucose levels between 160-180 mg/dl).

Data collection: Preoperative and each hour (during the first 24 hours in ICU) assessment of Glucose and Lactate blood levels and Body Temperature; Hemodynamic parameters on ICU admission and after 2, 6, 12 and 24 hours. Preoperative and postoperative (12 and 36 hours after ICU admission) assessment of C-Reactive Protein (CRP), White Blood Cells (WBC) and Platelets (PLT) blood levels. Admission, Total Maximum (TMSOFA) and Δ SOFA score (5). Infection Probability Score (IPS) 36 hours after ICU admission.

Statistics: Within-between groups analysis, one-way ANOVA and unpaired t-Test were used when appropriate.

Results

No difference in preoperative and operative variables ($p=NS$ for all measurements). Glucose and Lactate blood levels were lower in Group 1 ($p < 0.0001$). Haemodynamic parameters were comparable between Groups except Indexed Vascular Resistances that were higher in Group 1 ($p < 0.05$). CRP levels were lower in Group 1 ($p < 0.0001$), PLT levels were higher in Group 1 ($p < 0.0001$). Admission and TM SOFA score were lower in Group 1 ($p < 0.0001$), IPS was lower in Group 1 ($p < 0.01$).

Conclusions

Intensive insulinotherapy after elective and on-pump myocardial revascularization modulates the inflammatory pattern and can improve clinical response in this setting.

Author details

¹Department of Anaesthesia and Reanimation, University of Catanzaro, Catanzaro, Italy. ²Department of Cardiac Surgery, University of Catanzaro, Catanzaro, Italy. ³Department of Anaesthesia and Oncological Intensive Care, University of Catanzaro, Catanzaro, Italy.

Published: 19 May 2010

¹Department of Anaesthesia and Reanimation, University of Catanzaro, Catanzaro, Italy

Reference

1. Hoedemaekers CW, Pickkers P, Netea MG, van Deuren M, Van der Hoeven JG: **Intensive insulin therapy does not alter the inflammatory response in patients undergoing coronary artery bypass grafting: a randomized controlled trial [ISRCTN95608630].** *Crit Care* 2005, **9**(6):R790-7.

doi:10.1186/1471-2318-10-S1-A98

Cite this article as: Caroleo *et al.*: Intensive versus conventional insulinotherapy after elective and on-pump myocardial revascularization in the elderly patient: a prospective and randomized study. *BMC Geriatrics* 2010 **10**(Suppl 1):A98.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

