

CORRESPONDENCE

Open Access



# The *pla* gene, encoding plasminogen activator, is not specific to *Yersinia pestis*

Stephanie Hänsch<sup>1</sup>, Elisabetta Cilli<sup>2</sup>, Giulio Catalano<sup>2</sup>, Giorgio Gruppioni<sup>2</sup>, Raffaella Bianucci<sup>1</sup>, Nils C. Stenseth<sup>1</sup>, Barbara Bramanti<sup>1</sup> and Mark J. Pallen<sup>3\*</sup>

## Abstract

Here we present evidence to show that the *pla* gene, previously thought to be specific to *Yersinia pestis*, occurs in some strains of *Citrobacter koseri* and *Escherichia coli*. This means that detection of this gene on its own can no longer be taken as evidence of detection of *Y. pestis*.

## Correspondence

Molecular assays aimed at detecting traces of the etiological agent of plague, *Yersinia pestis*, have focused primarily—and sometimes solely—on the plasminogen activator/coagulase (*pla*) gene [1, 2]. This gene is located on the pPCP1 plasmid and has been considered the target of choice for plague detection due to its assumed specificity to *Y. pestis* and its occurrence in multiple copies [3–5]. However, a recent paper about the amplification of the *pla* gene from tissues from European rats has raised doubts over whether this gene is indeed specific to *Y. pestis* [6]. We can now confirm this suspicion.

We screened archaeological samples from Italy (6th, 14th and 17th centuries CE), amplifying a 70-base-pair fragment from the *pla* gene. Full protocols are described in a previous publication [7], but in brief we performed the work in a dedicated clean laboratory, with physically separated areas for extraction and amplification, following the most stringent criteria for ancient DNA analysis, such as the use of mock extractions and PCR blanks. We used the previously described *pla* primer pair (Forward primer: GACTGGGTTCGGGCACATGC—Reverse primer: CGGATGTCTTCTCACGGA). Cycling conditions started with an initial activation step at 95 °C for 15 min. This was followed by 50 cycles at 94 °C for 30 s, an assay specific annealing temperature at

60 °C for 30 s, and 72 °C for 1 min, ending with a final elongation step at 72 °C for 10 min. Final cooling was carried out at 8 °C until analysis.

Target-fragment amplifications were observed in 17 out of 40 samples. We sequenced one of the fragments and performed a BLASTN search of the NCBI database, which, as expected, revealed full-length identity (70/70 at nucleotide level) with numerous sequences from *Y. pestis*. However, we also found full-length identity with two sequences from outside *Y. pestis*. The first was an annotated contig from a genome assembly of *Citrobacter koseri* (submitted to GenBank in June 2014, with accession number LK931337). This bacterial species has been recognized as a commensal and pathogen in humans and animals [8, 9]. The second was from a contig from an unannotated genome assembly of *Escherichia coli* strain FHI29 (submitted to GenBank in June 2014, with accession number LM995843). This sequence is derived from a human fecal isolate from a case of gastroenteritis in Norway. To conduct a more extensive survey, we then performed BLAST searches with the entire *pla* gene from *Y. pestis* CO92, which confirmed the high level of similarity (>98 %) with sequences from the genome assemblies of *C. koseri* (927/939 identities at nucleotide level) and of the *E. coli* strain FHI29 (925/939 identities) (Fig. 1). The relevant contig from *C. koseri* contained sequences annotated with plasmid-related functions, suggesting that the *pla* gene in this context is also plasmid-encoded.

The presence of *pla* sequences from outside *Y. pestis*, each derived from a distinct geographical or taxonomic

\*Correspondence: m.pallen@warwick.ac.uk

<sup>3</sup> Microbiology and Infection Unit, Warwick Medical School, University of Warwick, Coventry CV4 7AL, UK

Full list of author information is available at the end of the article

```

Yersinia pestis      1 ATGAAGAAAAGTTCTATTGTGGCAACCATTATAACTATTCTGTCCGGGAGTGCTAATGCA
Escherichia coli FHI29 1 .....C.....
Citrobacter koseri   1 .....

Yersinia pestis      61 GCATCATCTCAGTTAATACCAATATATCCCCTGACAGCTTTACAGTTGCAGCCTCCACC
Escherichia coli FHI29 61 .....T.....A
Citrobacter koseri   61 .....T.....

Yersinia pestis      121 GGGATGCTGAGTGAAAGTCTCATGAAATGCTTTATGACGCAGAAACAGGAAGAAAGATC
Escherichia coli FHI29 121 .....G.....
Citrobacter koseri   121 .....G.....A...

Yersinia pestis      181 AGCCAGTTAGACTGGAAGATCAAAAATGTCGCTATCCTGAAAGGTGATATATCCTGGGAT
Escherichia coli FHI29 181 .....A.....T.....
Citrobacter koseri   181 .....

Yersinia pestis      241 CCATACTCATTCTGACCCTGAATGCCAGGGGGTGGACGCTCTGGCTTCCGGGTCAGGT
Escherichia coli FHI29 241 .....A.....
Citrobacter koseri   241 .....A.....A.....

Yersinia pestis      301 AATATGGATGACTACGACTGGATGAATGAAAATCAATCTGAGTGGACAGATCACTCATCT
Escherichia coli FHI29 301 .....A.....
Citrobacter koseri   301 .....

Yersinia pestis      361 CATCCTGCTACAATGTTAATCATGCCAATGAATATGACCTCAATGTGAAAGGCTGGTTA
Escherichia coli FHI29 361 .....
Citrobacter koseri   361 .....

Yersinia pestis      421 CTCCAGGATGAGAATTATAAAGCAGGTATAACAGCAGGATATCAGGAAACACGTTTCAGT
Escherichia coli FHI29 421 .....T..C.....
Citrobacter koseri   421 ..A....C.....

Yersinia pestis      481 TGGACAGCTACAGGTGGTTCATATAGTTATAATAATGGAGCTTATACCGGAAACTCCCCG
Escherichia coli FHI29 481 .....
Citrobacter koseri   481 .....

Yersinia pestis      541 AAAGGAGTGGGGTAATAGTTATAACCAGCGCTTTTCTATGCCATATATTGGACTTGCA
Escherichia coli FHI29 541 .....G.....
Citrobacter koseri   541 .....G.....

Yersinia pestis      601 GGCCAGTATCGCATTAAATGATTTTGGAGTTAAATGCATTATTTAAATTCAGCGACTGGGT
Escherichia coli FHI29 601 .....
Citrobacter koseri   601 .....

Yersinia pestis      661 CGGCACATGATAATGATGAGCACTATATGAGAGATCTTACTTTCCGTGAGAAGACATCC
Escherichia coli FHI29 661 .....
Citrobacter koseri   661 .....

Yersinia pestis      721 GGCTCACGTTATTATGGTACCGTAATTAACGCTGGATATTATGTCACACCTAATGCCAAA
Escherichia coli FHI29 721 .....
Citrobacter koseri   721 .....

Yersinia pestis      781 GTCTTTGCGGAATTTACATACAGTAAATATGATGAGGGCAAAGGAGTACTCAGACCATT
Escherichia coli FHI29 781 .....T.....T.....
Citrobacter koseri   781 .....T.....T.....

Yersinia pestis      841 GATAAGAATAGTGGAGATTCTGTCTCTATTGGCGGAGATGCTGCCGGTATTTCCAATAAA
Escherichia coli FHI29 841 .....T.....
Citrobacter koseri   841 .....A.....

Yersinia pestis      901 AATTATACTGTGACGGCGGGTCTGCAATATCGCTTCTGA
Escherichia coli FHI29 901 .....
Citrobacter koseri   901 .....A.....

```

**Fig. 1** Multiple alignment of *pla* sequences from *Yersinia pestis* and two other species

setting, confirms beyond doubt that this gene can no longer be considered specific to *Y. pestis*. Although there appear to be some potentially informative sequence differences between the *pla* sequences from *Y. pestis* and those from other taxa, these findings call into question any results—whether in contemporary diagnostic microbiology or in an ancient DNA setting—that rely on detection of PCR products from this gene alone. Instead, as many researchers in the field already recognise, it is important to obtain sequences from PCR products and detection or identification of *Y. pestis* should rely on sequences from at least two independent molecular targets. More generally, our observations call into question the wisdom of relying on genes from mobile elements as species-specific markers, given the likelihood that such sequences are able to move from one taxon to another. Interestingly, the roles of the sequence differences between the *pla* genes, some of which are non-synonymous, in the function and evolution of the *pla* gene product remain to be determined.

#### Authors' contributions

EC and GC performed laboratory experiments. SH, GC and MJP performed bioinformatics analyses. RB, GG, NCS and BB conceived of the study, and participated in its design and coordination and helped to draft the manuscript. EC, BB and MJP wrote the manuscript. All authors read and approved the final manuscript.

#### Author details

<sup>1</sup> Centre for Ecological and Evolutionary Synthesis, University of Oslo, Oslo, Norway. <sup>2</sup> Department of Cultural Heritage, University of Bologna, Ravenna Campus, Bologna, Italy. <sup>3</sup> Microbiology and Infection Unit, Warwick Medical School, University of Warwick, Coventry CV4 7AL, UK.

#### Compliance with ethical guidelines

#### Competing interests

The authors declare that they have no competing interests.

Received: 9 July 2015 Accepted: 24 September 2015

Published online: 05 October 2015

#### References

- Martin-Alonso A, Soto M, Foronda P, Aguilar E, Bonnet G, Pacheco R, et al. *Bartonella* spp. and *Yersinia pestis* reservoirs, Cusco, Peru. *Emerg Infect Dis*. 2014;20(6):1069–70. doi:10.3201/eid2006.131194.
- Loiez C, Herwegh S, Wallet F, Armand S, Guinet F, Courcol RJ. Detection of *Yersinia pestis* in sputum by real-time PCR. *J Clin Microbiol*. 2003;41(10):4873–5.
- Sodeinde OA, Goguen JD. Genetic analysis of the 9.5-kilobase virulence plasmid of *Yersinia pestis*. *Infect Immun*. 1988;56:2743–8.
- Norkina OV, Kulichenko AN, Gintsburg AL, Tuchkov IV, Popov YuA, Akse-nov MU, et al. Development of a diagnostic test for *Yersinia pestis* by the polymerase chain reaction. *J Appl Bacteriol*. 1994;76:240–5.
- Parkhill J, Wren BW, Thomson NR, Titball RW, Holden MT, Prentice MB, et al. Genome sequence of *Yersinia pestis*, the causative agent of plague. *Nature*. 2001;413(6855):523–7.
- Janse I, Hamidjaja RA, Reusken C. *Yersinia pestis* plasminogen activator gene homolog in rat tissues. *Emerg Infect Dis*. 2013;19(2):342–4.
- Harbeck M, Seifert L, Hänsch S, Wagner DM, Birdsell D, Parise KL, et al. *Yersinia pestis* DNA from skeletal remains from the 6th Century AD reveals insights into Justinianic plague. *PLoS Pathog*. 2013;9(5):e1003349. doi:10.1371/journal.ppat.1003349.
- Komine M, Massa A, Moon L, Mullaney T. *Citrobacter koseri* Septicaemia in a Holstein Calf. *J Comp Pathol*. 2014;151(4):309–13. doi:10.1016/j.jcpa.2014.07.005.
- Doran TJ. The role of *Citrobacter* in clinical disease of children: review. *Clin Infect Dis*. 1999;28:384–94.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

