




A novel one-pot synthesis of flavones†

 Cite this: *RSC Adv.*, 2021, **11**, 11655

 Meng-Yang Chang,¹ *^{ab} Min-Chen Tsai^{ab} and Chun-Yi Lin^{ab}

 Received 21st January 2021
 Accepted 17th March 2021

DOI: 10.1039/d1ra00534k

rsc.li/rsc-advances

In this paper, a one-pot facile route for the $\text{BiCl}_3/\text{RuCl}_3$ -mediated synthesis of functionalized flavones is described, including: (i) intermolecular *ortho*-acylation of substituted phenols with cinnamoyl chlorides, and (ii) intramolecular cyclodehydrogenation of the resulting *o*-hydroxychalcones. The reaction conditions are discussed herein.

Introduction


For the flavonoid family, flavone (also called 2-arylchromone) with numerous biological activities is attracting many synthetic researchers to develop a number of versatile methodologies.¹ A recent review article^{1a} has revealed the synthetic history of functionalized flavones from the traditional well-known named reactions (*e.g.*, Baker–Venkataraman,² Karl von-Auwers,³ Algar–Flynn–Oyamada,⁴ Allan–Robinson,⁵ Kostanecki,⁶ Mentzer⁷ and Wittig⁸) to modern novel transition-metal promoted protocols (*e.g.*, Suzuki–Miyaura⁹ and Sonogashira¹⁰). By the use of other reaction conditions such as microwave irradiation,¹¹ ionic liquids improvement¹² and photolytic annulation,¹³ the unique methodologies of flavones and their derivatives have been established. For the synthetic chemistry of diversified flavones, the two-step process for (1) *ortho*-acylation of substituted phenols and (2) cyclodehydrogenation of the resulting *o*-hydroxychalcones is a general and direct route to provide access to the core structure, as shown in Scheme 1.

By the involvement of transition metals, various *ortho*-acylations of phenols have been well-developed, including CuCl_2 ,^{14a} FeCl_3 ,^{14b} TiCl_4 ,^{14c} mercury lamp/photolysis,^{14d} MsOH /microwave.^{14e} For the following cyclodehydrogenation step, the uses of $\text{InCl}_3/\text{SiO}_2$,^{15a} $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}/\text{MeOH}$,^{15b} and CuI /ionic liquids^{15c} have been studied. In addition, transition metal-free oxidants-mediated reaction systems, for example, I_2/DMSO ,^{16a} DDQ /dioxane,^{16b} $\text{NaIO}_4/\text{DMSO}$,^{16c} $\text{H}_2\text{O}_2/\text{NaOH}$,^{16d} $\text{SeO}_2/\text{dioxane}$ ^{16e} and Br_2/NaOH ^{16f} have been investigated in the cyclodehydrogenation step.

Among these reported routes with the synthetic sequence of intermolecular *ortho*-acylation followed by intramolecular cyclodehydrogenation, we found that all attempts adopted a two-step stepwise process as the major focused design in the flavone family formation. Despite the above elegant synthetic routes, to date, there are no reports on the one-pot synthesis of substituted flavones on the basis of the formal (3 + 3) annulation. Herein, we present metal chlorides-mediated one-pot synthesis of flavones **4** (Scheme 2) *via* the combination of BiCl_3 and RuCl_3 -mediated intermolecular *ortho*-acylation of substituted phenols **1** with cinnamoyl chlorides **2** (one carbon–carbon bond formation, green), and intramolecular cyclodehydrogenation of the corresponding *o*-hydroxychalcones **3** (one carbon–oxygen bond formation, green). These starting materials, **1** and **2**, were obtained from commercial sources and used without further purification.

Results and discussion

The initial study commenced with the treatment of the model substrate **1a** ($\text{Ar} = \text{Ph}$, 1.0 mmol) with **2a** ($\text{Ar}' = \text{Ph}$, 1.0 mmol) and AlCl_3 (2.0 equiv.) in CCl_4 (20 mL) at reflux (77 °C) for 10 h. Only **3a** was produced at a 54% yield. The results are shown in



Scheme 1 Synthetic route of flavones.



Scheme 2 Our synthetic route towards flavones.

^aDepartment of Medicinal and Applied Chemistry, Kaohsiung Medical University Hospital, Kaohsiung Medical University, Kaohsiung 80708, Taiwan. E-mail: mychang@kmu.edu.tw

^bDepartment of Medical Research, Kaohsiung Medical University Hospital, Kaohsiung 807, Taiwan

† Electronic supplementary information (ESI) available: Scanned photocopies of NMR spectral data for all compounds and X-ray analysis data of **4t** were supported. CCDC 1957347. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/d1ra00534k




Table 1, entry 1. Entries 2–7 show that six trivalent transition-metal chlorides (2.0 equiv., InCl₃, FeCl₃, BiCl₃, CeCl₃, AuCl₃ and RuCl₃) were studied in CCl₄ (20 mL) at reflux for 10 h. However, only InCl₃ and FeCl₃ obtained sole **3a** in 67% and 70% yields, respectively. BiCl₃ provided 65% yield of **3a** along with trace amount of **4a**. In particular, no reactions were observed for CeCl₃ and AuCl₃, while RuCl₃ produced a complex mixture. According to the experimental results, we understood that InCl₃ or FeCl₃-mediated *ortho*-acylation of **1a** proceeded well, but they could not promote the conversion from **3a** to **4a** (entries 2 and 3). In addition to forming **3a**, trace amount of **4a** (3%) could be obtained in the presence of BiCl₃ (entry 4). For CeCl₃ and AuCl₃, the reactivity of Lewis acid was weak to a degree that no desired reaction was initiated (entries 5 and 6). Compared with other MCl₃, RuCl₃ with strong oxidation ability could force the *ortho*-acylation of **1a** to complexation (entry 7). Next, we refocused the synthetic aim to study the equivalents of BiCl₃; however, after increasing the stoichiometric from 2.0 to 3.0 equiv., the conversion efficiency from **3a** to **4a** was similar to that of 2.0 equiv. (5%, entry 8).

In decreasing the equivalence of BiCl₃ from 2.0 to 1.0, no isolation of **4a** was observed (entry 9). On the basis of the above phenomenon, we found that excess amounts of BiCl₃ could not completely drive the conversion of **3a** to **4a**. For this reason, another promoter was required to enhance the reaction condition. Hence, we turned the synthetic focus to study the

combination of BiCl₃ and RuCl₃-mediated synthesis of flavones. In entry 10, we found that the combination of BiCl₃ (1 equiv.) and RuCl₃ (1 equiv.) provided **4a** as the major product (68%) along with a 15% of **3a**. With these results in hand, we envisioned that RuCl₃ could trigger the cyclodehydrogenation step to accomplish the synthesis of **4a**. Three solvents having different boiling points, such as MeNO₂, (CH₂Cl)₂, CH₂Cl₂, and DMF were tested next. Using MeNO₂ (entry 11), complex unknown products were detected due to the high boiling temperature (101 °C). Entry 12 shows that (CH₂Cl)₂ produced a low conversion ratio (2/3) for **3a** and **4a**. For the low boiling point solvent, CH₂Cl₂ showed that only 10% of the amounts of **4a** were isolated, and **3a** was obtained as the major component (78%, entry 13). Among the three chloro-containing solvents, the temperature of boiling CH₂Cl₂ (40 °C) was low; as a result, the cyclodehydrogenation step could not be induced easier. Changing the solvent to DMF (entry 14), however, only unknown and unidentified complex mixture was isolated. Compared with CH₂Cl₂, CCl₄ and (CH₂Cl)₂ with higher boiling points (77 °C and 84 °C) could initiate the occurrence of the cyclodehydrogenation step, besides DMF (153 °C). According to the results, CCl₄ was the preferred solvent to obtain **4a**. To achieve the exhaustive conversion, the reaction time was examined next. Elongating the times to 15 h and 20 h respectively, showed that the transformation from **3a** to **4a** was complete, and the afforded yields of **4a** increased to 81% or 75%, respectively (entries 15 and 16). Even though the reaction time was elongated to 20 h, the yield of **4a** could not be enhanced. Remarkably, adjusting the temperature from reflux to room temperature (25 °C), no reactions were observed by the combination of BiCl₃ and RuCl₃. To increase the yield of **4a**, three combinations were examined next. In entry 17, the combination of BiCl₃ and CuCl₂ provided a 60% yield of **3a** along with trace amount of **4a** (8%). After changing CuCl₂ to FeCl₃, 72% yield of **3a** and 3% yield of **4a** were obtained for the combination of BiCl₃ and CuCl₂ (entry 18). The two results were similar to entries 4 and 8. Under the combination of BiCl₃ and AuCl₃ condition (entry 19), only **3a** was isolated in a 70% yield, and no desired **4a** was detected. Based on the results, we found that CuCl₂, FeCl₃ and AuCl₃ could not trigger the cyclodehydrogenation step easily. From the above screening reaction conditions, we envisioned that the combination of BiCl₃ (1.0 equiv.) and RuCl₃ (1.0 equiv.) could perform better for the formation of **4a** in refluxing CCl₄ for 15 h (entry 15). All the conditions were routinely carried out under an atmosphere of air (open-vessel conditions). The heating mantle was used to provide a stable heat source.


To study the scope and limitations of this one-pot route, substituted phenols **1a–1k** and cinnamoyl chlorides **2a–2o** were examined further. With optimal conditions established (Table 1, entry 14), we found that the one-pot two-step route could allow a direct synthesis of diversified flavones **4a–4y** in moderate to good yields (60–82%), as shown in Table 2, entries 1–26. For the electronic character of different Ar substituents on **1a–1k** and Ar' substituents on **2a–2o**, these various substituents included: (1) electron-donating mono-, di- or trioxxygenated aryl groups, (2) electron-neutral phenyl, biphenyl or naphthyl groups, (3) electron-withdrawing nitroaryl groups, (4) haloaryl

Table 1 Reaction conditions^a



Entry	MCl ₃ (equiv.)	Solvent	Time (h)	3a/4a ^b %
1	AlCl ₃ (2.0)	CCl ₄	10	54/— ^c
2	InCl ₃ (2.0)	CCl ₄	10	67/— ^c
3	FeCl ₃ (2.0)	CCl ₄	10	70/— ^c
4	BiCl ₃ (2.0)	CCl ₄	10	65/3
5	CeCl ₃ (2.0)	CCl ₄	10	— ^d
6	AuCl ₃ (2.0)	CCl ₄	10	— ^d
7	RuCl ₃ (2.0)	CCl ₄	10	— ^e
8	BiCl ₃ (3.0)	CCl ₄	10	68/5
9	BiCl ₃ (1.0)	CCl ₄	10	70/— ^c
10	BiCl ₃ (1.0), RuCl ₃ (1.0)	CCl ₄	10	15/68
11	BiCl ₃ (1.0), RuCl ₃ (1.0)	MeNO ₂	10	— ^e
12	BiCl ₃ (1.0), RuCl ₃ (1.0)	(CH ₂ Cl) ₂	10	35/52
13	BiCl ₃ (1.0), RuCl ₃ (1.0)	CH ₂ Cl ₂	10	78/10
14	BiCl ₃ (1.0), RuCl ₃ (1.0)	DMF	10	— ^e
15	BiCl₃ (1.0), RuCl₃ (1.0)	CCl₄	15	3/81
16	BiCl ₃ (1.0), RuCl ₃ (1.0)	CCl ₄	20	— ^d /75
17	BiCl ₃ (1.0), CuCl ₂ (1.0)	CCl ₄	15	60/8
18	BiCl ₃ (1.0), FeCl ₃ (1.0)	CCl ₄	15	72/3
19	BiCl ₃ (1.0), AuCl ₃ (1.0)	CCl ₄	15	70/— ^c

^a The reactions were run on a 1.0 mmol scale with phenol **1a**, cinnamoyl chloride **2a** (1.0 equiv.), metal chlorides (MCl₃, equiv.), solvent (20 mL), time (h), reflux. ^b Isolated yields. ^c No detection. ^d No reaction. ^e Unknown and unidentified complex mixture was isolated.


Table 2 Synthesis of 4a–4y^a


Entry	1, Ar =	2, Ar' =	4 ^b , (%)
1	1a, Ph	2a, Ph	4a, 81
2	1a, Ph	2b, 4-MeOC ₆ H ₄	4b, 82
3	1a, Ph	2c, 3-MeOC ₆ H ₄	4c, 80
4	1a, Ph	2d, 3,4,5-(MeO) ₃ C ₆ H ₂	4d, 78
5	1a, Ph	2e, 2,3,4-(MeO) ₃ C ₆ H ₂	4e, 72
6	1a, Ph	2f, 3,4-CH ₂ O ₂ C ₆ H ₃	4f, 73
7	1a, Ph	2g, 4-PhC ₆ H ₄	4g, 80
8	1a, Ph	2h, 3,4-(MeO) ₂ C ₆ H ₃	4h, 72
9	1a, Ph	2i, 3,4-Cl ₂ C ₆ H ₃	4i, 78
10	1a, Ph	2a, Ph	4j, 80
11	1b, 4-MeOC ₆ H ₄	2a, Ph	4k, 76
12	1c, 3-MeOC ₆ H ₄	2a, Ph	4l, 68
13	1d, 3-MeO-5-HOC ₆ H ₃	2a, Ph	4m, 64
14	1e, 3,5-(MeO) ₂ C ₆ H ₃	2j, 4-NO ₂ C ₆ H ₄	4n, 76
15	1a, Ph	2k, 2-naphthyl	4o, 70
16	1a, Ph	2l, 2-FC ₆ H ₄	4p, 67
17	1a, Ph	2m, 2-furyl	4q, 60 ^c
18	1a, Ph	2n, 2-thienyl	4r, 62
19	1a, Ph	2o, 1-naphthyl	4s, 75
20	1f, 4-FC ₆ H ₄	2h, 3,4-(MeO) ₂ C ₆ H ₃	4t, 73
21	1g, 4-ClC ₆ H ₄	2h, 3,4-(MeO) ₂ C ₆ H ₃	4u, 72
22	1h, 4-BrC ₆ H ₄	2h, 3,4-(MeO) ₂ C ₆ H ₃	4v, 66
23	1i, 4-MeC ₆ H ₄	2h, 3,4-(MeO) ₂ C ₆ H ₃	4w, 74
24	1c, 3-MeOC ₆ H ₄	2h, 3,4-(MeO) ₂ C ₆ H ₃	4x, 64
25	1j, 3-MeO-2-HOC ₆ H ₃	2a, Ph	4y, 60
26	1k, 2-NO ₂ C ₆ H ₄	2a, Ph	4z, — ^d

^a All reactions were run on a 1.0 mmol scale with phenols **1a–1k**, cinnamoyl chlorides (**2a–2o**, 1.0 equiv.), BiCl₃ (315 mg, 1.0 equiv.), CCl₄ (20 mL), 10 h, reflux (77 °C); then RuCl₃·3H₂O (261 mg, 1.0 equiv.) was added into the reaction mixture, 5 h, reflux (monitored by TLC). ^b Isolated yields. ^c **3q** (10%) was obtained. ^d No detection.

groups and (5) heterocyclic furyl and thienyl groups were highly appropriate. However, when the Ar' substituent was the 2-furyl group, the by-product **3q** was obtained in a 10% yield. Furthermore, with the use of RuCl₃, the conversion from **3q** to **4q** was successful. Therefore, efficient formation of **4a–4y** showed that these substituents (Ar and Ar') did not affect the distribution of the provided yields, besides **4z** (Ar = 4-NO₂C₆H₄). The structures of **4a–4y** could be determined by ¹H NMR analysis. The structure of **4t** was determined by single-crystal X-ray analysis.¹⁷

On the basis of the experimental results, a plausible mechanism for the formation of **4** is illustrated in Scheme 3. Initially, BiCl₃-mediated complexation of **2** forms **A** by one bismuth-chloro (Bi–Cl) bond formation. Then, by the removal of ^oBiCl₄, **B** with a styryl acylium center could be generated. Following the Friedel–Crafts *ortho*-acylation process, **B** reacts with **1** to lead **3** via one carbon–carbon (C–C) bond formation (green mark).^{18a} Dubac *et al.* have reported similar reactions. In addition to BiCl₃, other bismuth salts-mediated Friedel–Crafts reactions have been well-documented.^{18b–18d} Then, by the involvement of




Scheme 3 Plausible mechanism.


RuCl₃, **C** with a ruthenium(III)-chelated complex was produced. Furthermore, the intramolecular oxa-Michael addition provided **D** via one carbon–oxygen (C–O) bond formation (green mark). After the proton exchange, **E** could be formed along with the releasing HCl. Finally, dehydrogenation of **E** obtained **4** by the removal of RuCl and HCl.¹⁹ Under the cyclodehydrogenation step, Laurenczy *et al.* reported the redox conversion between Ru(III) and Ru(I).^{19a}

On the other hand, as an extension of the one-pot two-step synthetic route, changing the combination from BiCl₃/RuCl₃ to BiCl₃/ZnCl₂ was examined. However, the flavanone skeleton **5a** was isolated at a 90% yield, and no flavone **4a** was detected. A possible reason could be that ZnCl₂ lacks sufficient oxidative ability to completely promote the cyclodehydrogenation under the redox condition such that it only served a Lewis acid role in promoting the formation of **5a** via an intramolecular annulation of **3a**. Herein, we also developed the BiCl₃/ZnCl₂-mediated one-pot two-step, novel route for the synthesis of flavanone skeleton (Scheme 4).

With the results in mind, cinnamoyl chlorides **2a–2o** were adjusted next to crotonoyl chloride (**2p**) or acryloyl chloride (**2q**) under one-pot condition (Scheme 5). As shown in eqn (1), the BiCl₃/RuCl₃ mediated reaction of **1a** with **2p** provided **6a** with the chromen-4-one skeleton in only a 31% yield. Compared with **4a** (81%), the isolated yield of **6a** (31%) was low. The possible reason could be that with the hydrogen on **E** (yellow, Scheme 3),



Scheme 4 Synthesis of 5a.



Scheme 5 Synthesis of 6a and 7a.

the low acidity was not easy to abstract by the RuCl_3 for the dehydrogenation step to be triggered efficiently. On the basis of the results, we demonstrated that benzylic hydrogen was acidic and easily eliminated. Furthermore, by the removal of the β -methyl group on **2**, $\text{BiCl}_3/\text{RuCl}_3$ mediated reaction of **1a** with **2q** was shown in eqn (2). In particular, only **7a** was generated in a 67% yield. However, an intramolecular dehydrogenation step was not initiated so that the predicted **7b** was not obtained. The resulting phenomenon meant that cyclodehydrogenation only occurred in cinnamoyl substituents (with benzylic hydrogen). Although substrate **2** was limited to the cinnamoyl substituents, it still provided a novel one-pot synthesis of the flavone skeleton.

Conclusion

In summary, we have developed a concise route for the effective synthesis of functionalized flavones *via* $\text{BiCl}_3/\text{RuCl}_3$ mediating the one-pot, direct intermolecular *ortho*-acylation of substituted phenols with cinnamoyl chlorides followed by intramolecular cyclodehydrogenation of the resulting *o*-hydroxychalcones under refluxing CCl_4 reaction conditions. The process provides a cascade pathway of one carbon–oxygen and one carbon–carbon bond formation. Related plausible mechanisms have been proposed. Further studies regarding the efficient synthetic routes towards flavones will be conducted and published in due course.

Experimental

General

All reagents and solvents were obtained from commercial sources and used without further purification. Reactions were routinely carried out under an atmosphere of air with magnetic stirring. Products in organic solvents were dried with anhydrous magnesium sulfate before concentration *in vacuo*. Melting points were determined with a SMP3 melting apparatus. ^1H and ^{13}C NMR spectra were recorded on a Varian INOVA-400 spectrometer operating at 400 and at 100 MHz, respectively. Chemical shifts (δ) are reported in parts per million (ppm) and the coupling constants (J) are given in Hertz. High resolution mass spectra (HRMS) were measured with a mass spectrometer Finnigan/Thermo Quest MAT 95XL. X-ray crystal structures were obtained with an Enraf-Nonius FR-590 diffractometer (CAD4, Kappa CCD).

A representative synthetic procedure of compounds **4a–4y** and **3q** is as follows

BiCl_3 (315 mg, 1.0 mmol) was added to a solution of phenols **1a–1k** (1.0 mmol) in CCl_4 (20 mL) at 25 °C. The reaction mixture was stirred at 25 °C for 10 min. Cinnamoyl chlorides **2a–2o** (1.0 mmol) was added to the reaction mixture at 25 °C. The reaction mixture was stirred at reflux for 10 h (monitored by TLC). Then, $\text{RuCl}_3 \cdot 3\text{H}_2\text{O}$ (261 mg, 1.0 mmol) was added to the reaction mixture (containing *o*-hydroxychalcones **3**) at reflux. The reaction mixture was stirred at reflux for 5 h (monitored by TLC).

The reaction mixture was cooled to 25 °C and the solvent was concentrated. The residue was diluted with water (10 mL) and the mixture was extracted with CH_2Cl_2 (3 \times 30 mL). The combined organic layers were washed with brine (2 \times 20 mL), dried (MgSO_4), filtered and evaporated to afford crude product under reduced pressure. Purification on silica gel (hexanes/ EtOAc = 20/1–4/1) afforded **4a–4y** and **3q**.

2-Phenylchromen-4-one (4a).²⁰ Yield = 81% (180 mg); white solid; mp = 90–92 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{15}\text{H}_{11}\text{O}_2$ 223.0759, found 223.0768; ^1H NMR (400 MHz, CDCl_3): δ 8.24 (dd, J = 1.6, 8.0 Hz, 1H), 7.94–7.92 (m, 2H), 7.71 (dt, J = 2.0, 8.8 Hz, 1H), 7.57 (dd, J = 0.8, 8.0 Hz, 1H), 7.55–7.50 (m, 3H), 7.42 (dt, J = 0.8, 8.0 Hz, 1H), 6.86 (s, 1H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3): δ 178.4, 163.5, 156.3, 133.8, 131.7, 131.6, 129.0 (2 \times), 126.3 (2 \times), 125.7, 125.2, 123.9, 118.1, 107.5.

2-(4-Methoxyphenyl)chromen-4-one (4b).²¹ Yield = 82% (207 mg); white solid; mp = 172–174 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{16}\text{H}_{13}\text{O}_3$ 253.0865, found 253.0872; ^1H NMR (400 MHz, CDCl_3): δ 8.22 (dd, J = 1.6, 8.0 Hz, 1H), 7.90 (d, J = 9.2 Hz, 2H), 7.69 (dt, J = 1.6, 8.4 Hz, 1H), 7.56 (dd, J = 0.8, 8.4 Hz, 1H), 7.42 (dt, J = 1.2, 8.0 Hz, 1H), 7.03 (d, J = 9.2 Hz, 2H), 6.81 (s, 1H), 3.89 (s, 3H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3): δ 178.4, 163.7, 162.5, 156.2, 133.7, 128.6, 128.1 (2 \times), 125.7, 125.2, 123.9, 117.9, 114.5 (2 \times), 106.0, 55.5.

2-(3-Methoxyphenyl)chromen-4-one (4c).²² Yield = 80% (202 mg); white solid; mp = 92–94 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{16}\text{H}_{13}\text{O}_3$ 253.0865, found 253.0874; ^1H NMR (400 MHz, CDCl_3): δ 8.24 (dd, J = 1.6, 8.0 Hz, 1H), 7.72 (dt, J = 2.0, 8.4 Hz, 1H), 7.59 (dd, J = 0.8, 8.4 Hz, 1H), 7.53 (dt, J = 1.2, 8.0 Hz, 1H), 7.46–7.42 (m, 3H), 7.09 (ddd, J = 0.8, 2.4, 8.8 Hz, 1H), 6.89 (s, 1H), 3.90 (s, 3H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3): δ 178.5, 160.0, 156.3, 134.0, 133.1, 130.2, 125.7 (2 \times), 125.4, 118.9, 118.1 (2 \times), 117.4, 111.8, 107.6, 55.5.

2-(3,4,5-Trimethoxyphenyl)chromen-4-one (4d).²³ Yield = 78% (243 mg); white solid; mp = 168–170 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{18}\text{H}_{17}\text{O}_5$ 313.1076, found 313.1084; ^1H NMR (400 MHz, CDCl_3): δ 8.25 (dd, J = 1.6, 8.0 Hz, 1H), 7.73 (dt, J = 1.6, 8.4 Hz, 1H), 7.61 (d, J = 8.0 Hz, 1H), 7.46 (dt, J = 1.6, 7.6 Hz, 1H), 7.16 (s, 2H), 6.89 (s, 1H), 3.97 (s, 6H), 3.94 (s, 3H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3): δ 178.4, 163.4, 156.3, 153.6 (2 \times), 141.3, 134.0, 126.8, 125.7, 125.5, 123.9, 118.1, 107.1, 103.9 (2 \times), 61.1, 56.4 (2 \times).

2-(2,3,4-Trimethoxyphenyl)chromen-4-one (4e). Yield = 72% (225 mg); white solid; mp = 152–154 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{18}\text{H}_{17}\text{O}_5$ 313.1076, found 313.1067; ^1H NMR (400 MHz, CDCl_3): δ 8.24 (dd, J = 2.0, 8.4 Hz, 1H), 7.68 (dt, J = 2.0, 8.4 Hz, 1H), 7.56 (d, J = 9.2 Hz, 1H), 7.52 (dt, J = 1.6, 8.4 Hz, 1H), 7.41 (dt, J = 1.2, 8.0 Hz, 1H), 7.02 (s, 1H), 6.81 (d, J = 8.8 Hz, 1H), 3.96 (s, 3H), 3.94 (s, 3H), 3.92 (s, 3H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3): δ 178.8, 161.5, 156.5, 156.3, 153.2, 133.5, 128.4, 125.7, 125.0, 124.2, 123.8, 119.1, 117.9, 111.1, 107.4, 61.2, 61.0, 56.1.

2-Benzo[1,3]dioxol-5-ylchromen-4-one (4f).²⁴ Yield = 73% (194 mg); white solid; mp = 200–202 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $C_{16}H_{11}O_4$ 267.0657, found 267.0650; 1H NMR (400 MHz, $CDCl_3$): δ 8.22 (dd, $J = 1.6, 8.0$ Hz, 1H), 7.70 (dt, $J = 1.6, 8.4$ Hz, 1H), 7.55 (d, $J = 8.0$ Hz, 1H), 7.51 (dd, $J = 1.6, 8.0$ Hz, 1H), 7.42 (dt, $J = 0.8, 8.0$ Hz, 1H), 7.38 (d, $J = 1.6$ Hz, 1H), 6.94 (d, $J = 8.4$ Hz, 1H), 6.81 (s, 1H), 6.08 (s, 2H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$): δ 178.3, 163.5, 156.1, 150.8, 148.5, 133.9, 125.7, 125.6, 125.3, 123.6, 121.7, 118.0, 108.8, 106.4, 106.3, 102.0.

2-Biphenyl-4-ylchromen-4-one (4g).²⁵ Yield = 80% (238 mg); white solid; mp = 141–143 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $C_{21}H_{15}O_2$ 299.1072, found 299.1077; 1H NMR (400 MHz, $CDCl_3$): δ 8.26 (dd, $J = 1.6, 8.0$ Hz, 1H), 8.03 (d, $J = 8.4$ Hz, 2H), 7.76 (d, $J = 8.4$ Hz, 2H), 7.74 (dt, $J = 1.6, 8.4$ Hz, 1H), 7.67–7.61 (m, 3H), 7.51–7.40 (m, 4H), 7.00 (s, 1H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$): δ 178.4, 163.6, 156.3, 144.6, 139.7, 134.0, 130.3, 129.0 (2 \times), 128.3, 127.7 (2 \times), 127.2 (2 \times), 126.9 (2 \times), 125.7, 125.4, 123.7, 118.1, 107.1.

2-(3,4-Dimethoxyphenyl)chromen-4-one (4h).²⁶ Yield = 72% (203 mg); white solid; mp = 118–120 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $C_{17}H_{15}O_4$ 283.0970, found 283.0976; 1H NMR (400 MHz, $CDCl_3$): δ 8.23 (dd, $J = 1.6, 8.0$ Hz, 1H), 7.70 (dt, $J = 1.6, 8.4$ Hz, 1H), 7.59 (d, $J = 8.4$ Hz, 1H), 7.58 (d, $J = 8.4$ Hz, 1H), 7.43 (dt, $J = 1.2, 8.0$ Hz, 1H), 7.42 (d, $J = 8.0$ Hz, 1H), 7.00 (d, $J = 8.4$ Hz, 1H), 6.82 (s, 1H), 3.99 (s, 3H), 3.97 (s, 3H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$): δ 178.4, 163.6, 156.2, 152.2, 149.3, 133.7, 125.7, 125.2, 124.2, 123.8, 120.1, 118.0, 111.2, 108.9, 106.4, 56.1 (2 \times).

2-(3,4-Dichlorophenyl)chromen-4-one (4i).²⁶ Yield = 78% (226 mg); white solid; mp = 202–204 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $C_{15}H_9Cl_2O_2$ 290.9980, found 290.9987; 1H NMR (400 MHz, $CDCl_3$): δ 8.23 (dd, $J = 1.6, 8.0$ Hz, 1H), 8.24 (d, $J = 1.6$ Hz, 1H), 7.75–7.71 (m, 2H), 7.60 (dt, $J = 0.8, 8.4$ Hz, 2H), 7.45 (dt, $J = 0.8, 8.0$ Hz, 1H), 6.80 (s, 1H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$): δ 178.1, 160.9, 156.1, 136.0, 134.1, 133.7, 131.7, 131.1, 128.1, 125.8, 125.6, 125.3, 123.9, 118.1, 108.2.

6-Methoxy-2-phenylchromen-4-one (4j).²² Yield = 80% (202 mg); white solid; mp = 165–167 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $C_{16}H_{13}O_3$ 253.0865, found 253.0873; 1H NMR (400 MHz, $CDCl_3$): δ 7.94–7.90 (m, 2H), 7.59 (d, $J = 3.2$ Hz, 1H), 7.54–7.49 (m, 4H), 7.29 (dd, $J = 3.2, 9.2$ Hz, 1H), 6.87 (s, 1H), 3.90 (s, 3H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$): δ 178.3, 163.3, 157.0, 151.1, 131.8, 131.6, 129.0 (2 \times), 126.3 (2 \times), 124.4, 123.9, 119.5, 106.7, 104.8, 55.9.

7-Methoxy-2-phenylchromen-4-one (4k).²² Yield = 76% (192 mg); white solid; mp = 106–108 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $C_{16}H_{13}O_3$ 253.0865, found 253.0860; 1H NMR (400 MHz, $CDCl_3$): δ 8.12 (dd, $J = 0.4, 8.4$ Hz, 1H), 7.91–7.88 (m, 2H), 7.53–7.48 (m, 3H), 6.99–6.96 (m, 2H), 6.80 (s, 1H), 3.93 (s, 3H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$): δ 177.8, 164.3, 163.2, 158.0, 131.7, 131.5, 129.0 (2 \times), 127.0, 126.2 (2 \times), 117.6, 114.5, 107.3, 100.4, 55.8.

5-Hydroxy-7-methoxy-2-phenylchromen-4-one (4l).²⁷ Yield = 68% (182 mg); white solid; mp = 167–169 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $C_{16}H_{13}O_4$ 269.0814, found 269.0821; 1H NMR (400 MHz, $CDCl_3$): δ 10.81 (br s, 1H), 7.87–7.85 (m, 2H), 7.54–7.48 (m, 3H), 6.63 (s, 1H), 6.47 (d, $J = 2.4$ Hz, 1H), 6.35 (d, $J = 2.4$ Hz, 1H), 3.86 (s, 3H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$): δ 182.4, 165.5, 163.9, 162.1, 157.7, 131.8, 131.2, 129.0 (2 \times), 126.2 (2 \times), 105.8, 105.6, 98.1, 92.6, 55.7.

5,7-Dimethoxy-2-phenylchromen-4-one (4m).²² Yield = 64% (181 mg); white solid; mp = 145–147 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $C_{17}H_{15}O_4$ 283.0970, found 283.0973; 1H NMR (400 MHz, $CDCl_3$): δ 7.88–7.83 (m, 2H), 7.54–7.45 (m, 3H), 6.67 (s, 1H), 6.56 (d, $J = 2.4$ Hz, 1H), 6.36 (d, $J = 2.0$ Hz, 1H), 3.95 (s, 3H), 3.90 (s, 3H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$): δ 177.6, 164.0, 160.9, 160.6, 159.9, 131.5, 131.1, 128.9 (2 \times), 125.9 (2 \times), 109.3, 109.0, 96.2, 92.8, 56.4, 55.7.

2-(4-Nitrophenyl)chromen-4-one (4n).²¹ Yield = 76% (203 mg); white solid; mp = 128–130 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $C_{15}H_{10}NO_4$ 268.0610, found 268.0618; 1H NMR (400 MHz, $CDCl_3$): δ 8.39 (d, $J = 9.2$ Hz, 2H), 8.24 (dd, $J = 1.6, 8.0$ Hz, 1H), 8.11 (d, $J = 9.2$ Hz, 2H), 7.76 (dt, $J = 1.6, 8.8$ Hz, 1H), 7.61 (dd, $J = 0.8, 8.4$ Hz, 1H), 7.47 (dt, $J = 1.2, 8.0$ Hz, 1H), 6.92 (s, 1H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$): δ 178.0, 160.6, 156.2, 149.4, 137.6, 134.4, 127.2 (2 \times), 125.9, 125.8, 124.2 (2 \times), 123.6, 118.1, 109.6.

2-Naphthalen-2-ylchromen-4-one (4o).²⁸ Yield = 70% (190 mg); white solid; mp = 141–143 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $C_{19}H_{13}O_2$ 273.0916, found 273.0925; 1H NMR (400 MHz, $CDCl_3$): δ 8.52 (s, 1H), 8.23 (dd, $J = 1.6, 8.0$ Hz, 1H), 8.02–7.90 (m, 4H), 7.76 (dt, $J = 1.6, 8.4$ Hz, 1H), 7.67 (dd, $J = 0.8, 8.4$ Hz, 1H), 7.64–7.57 (m, 2H), 7.47 (dt, $J = 1.2, 7.6$ Hz, 1H), 7.07 (s, 1H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$): δ 178.5, 163.7, 156.4, 134.8, 134.0, 132.9, 129.1, 129.0, 128.8, 128.1, 127.9, 127.1 (2 \times), 125.8, 125.4, 123.8, 122.6, 118.1, 107.7.

2-(2-Fluorophenyl)chromen-4-one (4p).²⁹ Yield = 67% (161 mg); white solid; mp = 100–102 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $C_{15}H_{10}FO_2$ 241.0665, found 241.0669; 1H NMR (400 MHz, $CDCl_3$): δ 8.25 (dd, $J = 1.6, 8.0$ Hz, 1H), 7.94 (dt, $J = 1.6, 8.0$ Hz, 1H), 7.72 (dt, $J = 1.6, 8.0$ Hz, 1H), 7.57–7.50 (m, 2H), 7.44 (dt, $J = 1.2, 7.6$ Hz, 1H), 7.33 (dt, $J = 1.2, 8.0$ Hz, 1H), 7.24 (dd, $J = 1.2, 8.4$ Hz, 1H), 6.96 (s, 1H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$): δ 178.5, 160.6 (d, $J = 254.0$ Hz), 158.9, 156.4, 133.9, 132.9 (d, $J = 9.1$ Hz), 129.1, 125.8, 125.3 (2 \times), 124.6 (d, $J = 3.8$ Hz), 123.8, 118.1, 117.0 (d, $J = 22.7$ Hz), 112.4 (d, $J = 10.6$ Hz).

2-Furan-2-ylchromen-4-one (4q).³⁰ Yield = 60% (127 mg); white solid; mp = 122–124 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $C_{13}H_9O_3$ 213.0552, found 213.0559; 1H NMR (400 MHz, $CDCl_3$): δ 8.22 (dd, $J = 2.0, 8.0$ Hz, 1H), 7.68 (dt, $J = 2.0, 8.4$ Hz, 1H), 7.63 (dd, $J = 0.8, 2.0$ Hz, 1H), 7.50 (dd, $J = 0.8, 8.4$ Hz, 1H), 7.41 (dt, $J = 0.8, 8.0$ Hz, 1H), 7.15 (d, $J = 3.2$ Hz, 1H), 6.76 (s, 1H), 6.62 (dd, $J = 2.0, 3.2$ Hz, 1H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$): δ 177.9, 155.8,

155.2, 146.4, 145.8, 133.8, 125.8, 125.2, 124.1, 117.9, 113.1, 112.5, 105.5.

3-Furan-2-yl-1-(2-hydroxyphenyl)propenone (3q). Yield = 10% (21 mg); white solid; mp = 105–107 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $C_{13}H_{11}O_3$ 215.0708, found 215.0714; 1H NMR (400 MHz, $CDCl_3$): δ 12.89 (s, 1H), 7.92 (dd, $J = 2.0, 8.4$ Hz, 1H), 7.68 (d, $J = 14.8$ Hz, 1H), 7.560 (dd, $J = 0.8, 1.2$ Hz, 1H), 7.558 (d, $J = 14.8$ Hz, 1H), 7.49 (dt, $J = 1.6, 8.8$ Hz, 1H), 7.02 (dd, $J = 0.8, 8.4$ Hz, 1H), 6.94 (dt, $J = 1.2, 8.4$ Hz, 1H), 6.77 (d, $J = 3.2$ Hz, 1H), 6.54 (dd, $J = 1.6, 3.2$ Hz, 1H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$): δ 193.3, 163.5, 151.5, 145.4, 136.3, 131.1, 129.6, 120.0, 118.8, 118.5, 117.6, 117.1, 112.9.

2-Thiophen-2-ylchromen-4-one (4r).³⁰ Yield = 62% (141 mg); white solid; mp = 93–95 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $C_{13}H_9O_2S$ 229.0323, found 229.0332; 1H NMR (400 MHz, $CDCl_3$): δ 8.21 (dd, $J = 1.6, 7.6$ Hz, 1H), 7.34 (dd, $J = 1.2, 3.6$ Hz, 1H), 7.69 (dt, $J = 1.6, 8.4$ Hz, 1H), 7.58 (dd, $J = 1.2, 4.8$ Hz, 1H), 7.54 (dd, $J = 0.8, 8.4$ Hz, 1H), 7.42 (dt, $J = 1.2, 8.4$ Hz, 1H), 7.19 (dd, $J = 3.6, 4.8$ Hz, 1H), 6.72 (s, 1H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$): δ 177.9, 159.1, 155.9, 135.2, 133.8, 130.3, 128.5, 128.5, 125.7, 125.3, 124.0, 117.9, 106.2.

2-Naphthalen-1-ylchromen-4-one (4s).²⁰ Yield = 75% (204 mg); white solid; mp = 139–141 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $C_{19}H_{13}O_2$ 273.0916, found 273.0910; 1H NMR (400 MHz, $CDCl_3$): δ 8.32 (dd, $J = 1.6, 8.0$ Hz, 1H), 8.16–8.12 (m, 1H), 8.04 (d, $J = 8.4$ Hz, 1H), 7.98–7.94 (m, 1H), 7.78 (d, $J = 6.4$ Hz, 1H), 7.74 (dt, $J = 1.6, 8.8$ Hz, 1H), 7.61–7.54 (m, 4H), 7.49 (dt, $J = 1.2, 8.0$ Hz, 1H), 6.73 (s, 1H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$): δ 178.3, 165.6, 156.8, 134.0, 133.8, 131.6, 130.6, 130.4, 128.7, 128.0, 127.5, 126.6, 125.9, 125.4, 125.1, 124.9, 124.0, 118.3, 113.0.

2-(3,4-Dimethoxyphenyl)-6-fluorochromen-4-one (4t). Yield = 73% (219 mg); white solid; mp = 168–170 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $C_{17}H_{14}FO_4$ 301.0876, found 301.0879; 1H NMR (400 MHz, $CDCl_3$): δ 7.86 (dd, $J = 2.8, 8.0$ Hz, 1H), 7.60–7.55 (m, 2H), 7.42 (dt, $J = 3.2, 8.8$ Hz, 1H), 7.38 (br s, 1H), 6.99 (d, $J = 8.0$ Hz, 1H), 6.81 (s, 1H), 3.98 (s, 3H), 3.97 (s, 3H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$): δ 177.5 (d, $J = 2.3$ Hz), 163.9, 160.8, 158.4, 152.4, 149.4, 124.9 (d, $J = 7.6$ Hz), 123.8, 121.8 (d, $J = 25.7$ Hz), 120.2, 120.0 (d, $J = 7.6$ Hz, 1H), 111.2, 110.6 (d, $J = 23.5$ Hz), 108.9, 105.7, 56.13, 56.11. Single-crystal X-ray diagram: crystal of compound **4t** was grown by slow diffusion of EtOAc into a solution of compound **4t** in CH_2Cl_2 to yield colorless prisms. The compound crystallizes in the monoclinic crystal system, space group Pn , $a = 4.0051(3)$ Å, $b = 10.1298(8)$ Å, $c = 16.3327(12)$ Å, $V = 662.13(9)$ Å³, $Z = 2$, $d_{calcd} = 1.506$ g cm⁻³, $F(000) = 312$, 2θ range 2.010–26.380°, R indices (all data) $R1 = 0.0307$, $wR2 = 0.0827$.

6-Chloro-2-(3,4-dimethoxyphenyl)chromen-4-one (4u).³¹ Yield = 72% (228 mg); white solid; mp = 199–201 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $C_{17}H_{14}ClO_4$ 317.0581, found 317.0588; 1H NMR (400 MHz, $CDCl_3$): δ 8.20 (d, $J = 2.4$ Hz, 1H), 7.63 (d, $J = 2.8$ Hz, 1H), 7.56 (dd, $J = 2.4, 8.4$ Hz, 1H), 7.54 (dd, $J = 0.4, 8.8$ Hz, 1H), 7.37 (d, $J = 2.0$ Hz, 1H), 6.99 (d, $J = 8.4$ Hz, 1H), 6.78 (s, 1H), 3.99

(s, 3H), 3.97 (s, 3H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$): δ 177.1, 163.8, 154.5, 152.4, 149.4, 133.8, 131.1, 125.2, 124.9, 123.8, 120.2, 119.7, 111.2, 108.9, 106.3, 56.1 (2 \times).

6-Bromo-2-(3,4-dimethoxyphenyl)chromen-4-one (4v).³² Yield = 66% (238 mg); white solid; mp = 206–208 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $C_{17}H_{14}BrO_4$ 361.0076, found 361.0081; 1H NMR (400 MHz, $CDCl_3$): δ 8.34 (d, $J = 2.4$ Hz, 1H), 7.77 (dd, $J = 2.4, 8.8$ Hz, 1H), 7.55 (dd, $J = 2.0, 8.4$ Hz, 1H), 7.47 (d, $J = 8.8$ Hz, 1H), 7.36 (d, $J = 2.0$ Hz, 1H), 6.98 (d, $J = 8.4$ Hz, 1H), 6.81 (s, 1H), 3.98 (s, 3H), 3.97 (s, 3H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$): δ 176.9, 163.9, 154.9, 152.5, 149.4, 136.6, 128.3, 125.1, 123.7, 120.3, 119.9, 118.7, 111.2, 108.9, 106.2, 56.1 (2 \times).

2-(3,4-Dimethoxyphenyl)-6-methylchromen-4-one (4w).³³ Yield = 74% (219 mg); white solid; mp = 182–184 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $C_{18}H_{17}O_4$ 297.1127, found 297.1138; 1H NMR (400 MHz, $CDCl_3$): δ 8.05 (br s, 1H), 7.64–7.52 (m, 3H), 7.44 (br s, 1H), 7.15 (br s, 1H), 7.00 (s, 1H), 3.98 (s, 6H), 2.50 (s, 3H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$): δ 178.4, 164.8, 154.6, 152.7, 149.4, 135.7, 125.6, 125.0, 123.8, 123.1, 120.8, 117.9, 117.6, 111.3, 109.1, 56.3, 56.2, 21.0.

2-(3,4-Dimethoxyphenyl)-7-methoxychromen-4-one (4x).³⁴ Yield = 64% (200 mg); white solid; mp = 172–174 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $C_{18}H_{17}O_5$ 313.1076, found 313.1085; 1H NMR (400 MHz, $CDCl_3$): δ 8.14 (d, $J = 8.8$ Hz, 1H), 7.57 (dd, $J = 2.0, 8.8$ Hz, 1H), 7.38 (d, $J = 2.0$ Hz, 1H), 7.02–6.97 (m, 3H), 6.89 (s, 1H), 3.99 (s, 3H), 3.97 (s, 3H), 3.95 (s, 3H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$): δ 177.7, 164.4, 163.8, 158.0, 152.3, 149.3, 127.0, 124.0, 120.2, 117.1, 114.7, 111.2, 108.9, 105.8, 100.4, 56.13, 56.10, 55.9.

8-Hydroxy-7-methoxy-2-phenylchromen-4-one (4y). Yield = 60% (161 mg); white solid; mp = 236–238 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $C_{16}H_{13}O_4$ 269.0814, found 269.0816; 1H NMR (400 MHz, $CDCl_3$): δ 8.01–7.99 (m, 2H), 7.80 (d, $J = 9.2$ Hz, 1H), 7.56–7.50 (m, 3H), 7.06 (d, $J = 9.2$ Hz, 1H), 6.96 (s, 1H), 4.05 (s, 3H), 2.80 (br s, 1H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$): δ 178.3, 150.0, 144.9, 134.0, 131.6, 130.8, 129.1 (3 \times), 126.4 (2 \times), 116.5 (2 \times), 108.6, 106.7, 56.7.

2-Phenylchroman-4-one (5a).³⁵ $BiCl_3$ (315 mg, 1.0 mmol) was added to a solution of phenol **1a** (94 mg, 1.0 mmol) in CCl_4 (20 mL) at 25 °C. The reaction mixture was stirred at 25 °C for 10 min. Cinnamoyl chloride **2a** (167 mg, 1.0 mmol) was added to the reaction mixture at 25 °C. The reaction mixture was stirred at reflux for 10 h. Then, $ZnCl_2$ (136 mg, 1.0 mmol) was added to the reaction mixture at reflux. The reaction mixture was stirred at reflux for 5 h. The reaction mixture was cooled to 25 °C and the solvent was concentrated. The residue was diluted with water (10 mL) and the mixture was extracted with CH_2Cl_2 (3 \times 30 mL). The combined organic layers were washed with brine (2 \times 20 mL), dried ($MgSO_4$), filtered and evaporated to afford crude product under reduced pressure. Purification on silica gel (hexanes/EtOAc = 20/1–4/1) afforded **5a**. Yield = 90% (202 mg); colorless solid; mp = 77–79 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $C_{15}H_{13}O_2$ 225.0916, found 225.0924; 1H NMR (400 MHz, $CDCl_3$): δ 7.95

(dd, $J = 2.0, 8.0$ Hz, 1H), 7.54–7.38 (m, 6H), 7.08–7.04 (m, 2H), 5.49 (dd, $J = 2.8, 13.2$ Hz, 1H), 3.10 (dd, $J = 13.2, 16.8$ Hz, 1H), 2.90 (dd, $J = 2.8, 16.8$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3): δ 191.9, 161.5, 138.7, 136.2, 128.8 (2 \times), 128.7, 127.0, 126.1 (2 \times), 121.6, 120.9, 118.1, 79.6, 44.6.

2-Methyl-chromen-4-one (6a).³⁶ and **chroman-4-one (7a)**.³⁷ BiCl_3 (315 mg, 1.0 mmol) was added to a solution of phenol **1a** (94 mg, 1.0 mmol) in CCl_4 (20 mL) at 25 °C. The reaction mixture was stirred at 25 °C for 10 min. Crotonoyl chloride **2p** (104 mg, 1.0 mmol) or acryloyl chloride **2q** (90 mg, 1.0 mmol) was added to the reaction mixture at 25 °C. The reaction mixture was stirred at reflux for 10 h. Then, $\text{RuCl}_3 \cdot 3\text{H}_2\text{O}$ (261 mg, 1.0 mmol) was added to the reaction mixture at reflux. The reaction mixture was stirred at reflux for 5 h. The reaction mixture was cooled to 25 °C and the solvent was concentrated. The residue was diluted with water (10 mL) and the mixture was extracted with CH_2Cl_2 (3 \times 30 mL). The combined organic layers were washed with brine (2 \times 20 mL), dried (MgSO_4), filtered and evaporated to afford crude product under reduced pressure. Purification on silica gel (hexanes/EtOAc = 20/1–4/1) afforded **6a** and **7a**. For **6a**: yield = 31% (50 mg); colorless solid; mp = 62–64 °C (recrystallized from hexanes and EtOAc); HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{10}\text{H}_9\text{O}_2$ 161.0603, found 161.0612; ^1H NMR (400 MHz, CDCl_3): δ 8.17 (dd, $J = 1.6, 8.0$ Hz, 1H), 7.63 (dt, $J = 1.6, 8.8$ Hz, 1H), 7.41 (dd, $J = 0.8, 8.0$ Hz, 1H), 7.35 (dt, $J = 0.8, 8.0$ Hz, 1H), 6.17 (s, 1H), 2.38 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3): δ 178.2, 166.2, 156.5, 133.4, 125.6, 124.9, 123.6, 117.8, 110.6, 20.6. For **7a**: yield = 67% (99 mg); colorless liquid; HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ calcd for $\text{C}_9\text{H}_9\text{O}_2$ 161.0603, found 161.0612; ^1H NMR (400 MHz, CDCl_3): δ 7.86 (dd, $J = 2.0, 8.0$ Hz, 1H), 7.44 (dt, $J = 1.6, 8.4$ Hz, 1H), 6.98 (dt, $J = 1.2, 8.0$ Hz, 1H), 6.94 (dd, $J = 1.2, 8.4$ Hz, 1H), 4.50 (t, $J = 6.4$ Hz, 2H), 2.78 (t, $J = 6.4$ Hz, 2H); ^{13}C NMR (100 MHz, CDCl_3): δ 191.7, 161.8, 135.8, 127.0, 121.2 (2 \times), 117.8, 66.9, 37.7.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The authors would like to thank the Ministry of Science and Technology of the Republic of China (Taiwan) for its financial support (MOST 109-2113-M-037-014-MY3).

Notes and references

- Recent leading review articles with synthesis, see: (a) R. Kshatriya, V. P. Jejurkar and S. Saha, *Tetrahedron*, 2018, **74**, 811–833; (b) C. M. M. Snatos and A. M. S. Silva, *Eur. J. Org. Chem.*, 2017, **2017**, 3115–3133; (c) For recent review articles with the biological activities, see: M. Singh, M. Kaur and O. Silakari, *Eur. J. Med. Chem.*, 2014, **84**, 206–239; (d) J. Reis, A. Gaspar, N. Milhazes and F. Borges, *J. Med. Chem.*, 2017, **60**, 7941–7957.
- (a) W. Baker, *J. Chem. Soc.*, 1933, 1381–1389; (b) D. C. Bhalla, H. S. Mahal and K. Venkataraman, *J. Chem. Soc.*, 1934, 1933–1935.
- K. V. Auwers and K. Muller, *Ber. Dtsch. Chem. Ges.*, 1908, **41**, 4233–4241.
- (a) A. J. Flynn, *Proc. R. Ir. Acad., Sect. B*, 1934, **42**, 1–4; (b) T. J. Oyamada, *Bull. Chem. Soc. Jpn.*, 1935, **10**, 182–186.
- J. Allan and R. Robinson, *J. Chem. Soc., Trans.*, 1924, **125**, 2192–2195.
- (a) S. V. Kostanecki and T. Emilevickz, *Eur. J. Inorg. Chem.*, 1898, **31**, 696–705; (b) S. V. Kostanecki and J. Thamobor, *Eur. J. Inorg. Chem.*, 1895, **28**, 2302–2309.
- (a) C. Mentzer, W. Molho and P. Vercier, *Compt. Rend.*, 1951, **232**, 1488–1490; (b) C. Mentzer and P. Vercier, *Compt. Rents.*, 1952, **232**, 1674–1678.
- (a) P. Kumar and M. S. Bodas, *Org. Lett.*, 2000, **2**, 3821–3823; (b) G. Bose, E. Mondal, A. T. Khan and M. J. Bordoloi, *Tetrahedron Lett.*, 2001, **42**, 8907–8909; (ac) F. Lassagne and F. Pochat, *Tetrahedron Lett.*, 2003, **44**, 9283–9285.
- M. A. Selepe and F. R. van Heerden, *Molecules*, 2013, **18**, 4739–4765.
- (a) E. Awuah and A. Capretta, *Org. Lett.*, 2009, **11**, 3210–3213; (b) H. Miao and Z. Yang, *Org. Lett.*, 2000, **2**, 1765–1768.
- J. A. Seijas, M. P. Vázquez-Tato and R. Carballido-Reboredo, *J. Org. Chem.*, 2005, **70**, 2855–2858.
- Z. Du, H. Ng, K. Zhang, H. Zeng and J. Wang, *Org. Biomol. Chem.*, 2011, **9**, 6930–6935.
- J. Das and S. Ghosh, *Tetrahedron Lett.*, 2011, **52**, 7189–7194.
- (a) Transition metal-mediated *ortho*-acylation of phenols, for Cu(II), see: J. Hu, E. A. Adogla, Y. Ju, D. Fan and Q. Wang, *Chem. Commun.*, 2012, **48**, 11256–11258; (b) For Fe(III), see: H. Naeimi and L. Moradi, *Russ. J. Org. Chem.*, 2007, **43**, 1757–1759; (c) For Ti(IV), see: A. Bensari and N. T. Zaveri, *Synthesis*, 2003, 267–271; (d) For Hg lamp, see: F. Galindo, M. C. Jimenez, M. A. Miranda and R. Tormos, *J. Photochem. Photobiol., A*, 1997, **97**, 151–153; (e) For microwave, see: H. Naeimi, A. Raeisi and M. Moradian, *Arabian J. Chem.*, 2017, **10**, S2723–S2728.
- (a) Transition metal-mediated cyclodehydrogenation methods, for $\text{InCl}_3/\text{SiO}_2$, see: N. Ahmed, H. Ali and J. E. van Lier, *Tetrahedron Lett.*, 2005, **46**, 253–256 and cited references therein; (b) For $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}/\text{MeOH}$, see: K. H. Kumar and P. T. Perumal, *Tetrahedron*, 2007, **63**, 9531–9535; (c) For CuI/ionic liquids, see: Z. Du, H. Ng, K. Zhang, H. Zeng and J. Wang, *Org. Biomol. Chem.*, 2011, **9**, 6930–6933.
- (a) Metal-free-mediated cyclodehydrogenation methods, for I_2/DMSO , see: M. D. L. de la Torre, G. L. Marcorin, G. Pirri, A. C. Tome, A. M. S. Silva and J. A. S. Cavaleira, *Tetrahedron Lett.*, 2002, **43**, 1689–1691; (b) For DDQ/dioxane, see: K. Imafuku, M. Honda and J. F. W. McOmie, *Synthesis*, 1987, 199–201; (c) For $\text{NaIO}_4/\text{DMSO}$, see: N. Hans and S. K. Grover, *Synth. Commun.*, 1993, **23**, 1021–1023; (d) For $\text{H}_2\text{O}_2/\text{NaOH}$, see: S. Gobbi, A. Rampa, A. Bisi, F. Belluti, L. Piazzini, P. Valent, A. Caputo, A. Zampiron and M. Carrara, *J. Med. Chem.*, 2003, **46**, 3662–3669; (e) For $\text{SeO}_2/\text{dioxane}$, see: M. E. Zwaagstra, H. Timmerman,

- A. C. van de Stolpe, F. J. J. de Kanter, M. Tamura, Y. Wada and M.-Q. Zhang, *J. Med. Chem.*, 1998, **41**, 1428–1438; (f) For Br₂/NaOH, see: J. R. Pfister, W. E. Wymann, M. E. Schuler and A. P. Roszkowski, *J. Med. Chem.*, 1980, **23**, 335–338.
- 17 CCDC 1957347 (**4t**) contains the supplementary crystallographic data for this paper.†
- 18 (a) BiCl₃-mediated Friedel-Crafts acylations, see: S. Repichet, C. Le Roux, N. Roques and J. Dubac, *Tetrahedron Lett.*, 2003, **44**, 2037–2040; (b) For recent reviews on bismuth salts-mediated reactions, see: P. Ondet, G. Lemiere and E. Dunach, *Eur. J. Org. Chem.*, 2017, **2017**, 761–780; (c) B. Abnerjee, *ChemistrySelect*, 2017, **2**, 6744–6757; (d) A. Gagnon, J. Dansereau and A. Le Roch, *Synthesis*, 2017, **49**, 1707–1745.
- 19 (a) RuCl₃-mediated dehydrogenations, see: V. Henricks, I. Yuranov, N. Autissier and G. Laurency, *Catalysts*, 2017, **7**, 348–355; (b) Y. Kim, S. Ahn, J. Y. Hwang, D.-H. Ko and K.-Y. Kwon, *Catalysts*, 2017, **7**, 7–17. For review on ruthenium salts-mediated reactions, see: (c) M. Pagliao, S. Campestrini and R. Ciriminna, *Chem. Soc. Rev.*, 2005, **34**, 837–845.
- 20 For **4a** and **4s**, see: M. Golshani, M. Khoobi, N. Jalalimanesh, F. Jafarpour and A. Ariafard, *Chem. Commun.*, 2017, **53**, 10676–10679.
- 21 For **4b** and **4n**, see: R. Charugandla, R. S. Vangala, S. Chidara and R. B. Korupolu, *Tetrahedron Lett.*, 2018, **59**, 3283–3287.
- 22 For **4c**, **4h**, **4j**, **4k**, **4m**, see: J. Lee, J. Yu, S. H. Son, J. Heo, T. Kim, J.-Y. An, K.-S. Inn and N.-J. Kim, *Org. Biomol. Chem.*, 2016, **14**, 777–784.
- 23 For **4d**, see: P. S. Kulkarni, D. D. Kondhare, R. Varala and P. Zubaidha, *J. Serb. Chem. Soc.*, 2013, **78**, 909–916.
- 24 For **4f**, see: G. Priyadarshani, S. Amrutkar, A. Nayak, U. C. Banerjee, C. N. Kundu and S. K. Guchhait, *Eur. J. Med. Chem.*, 2016, **122**, 43–54.
- 25 For **4g**, see: T. Rodrigues, A. S. Ressurreicao, F. P. da Cruz, I. S. Albuquerque, J. Gut, M. P. Carrasco, D. Goncalves, R. C. Guedes, D. J. V. A. dos Santos, M. M. M. Philip, R. Moreira, M. Prudencio and F. Lopes, *Eur. J. Med. Chem.*, 2013, **69**, 872–880.
- 26 For **4i**, see: V. K. Rai, F. Verma, G. P. Sahu, M. Singh and A. Rai, *Eur. J. Org. Chem.*, 2018, **48**, 537–544.
- 27 For **4l**, see: S.-H. Wang, C.-H. Chen, C.-Y. Lo, J.-Z. Feng, H.-J. Lin, P.-Y. Chang, L.-L. Yang, L.-G. Chen, Y.-W. Liu, C.-D. Kuo and J.-Y. Wu, *MedChemComm*, 2015, **6**, 1864–1873.
- 28 For **4o**, see: M. Pérez, D. Ruiz, J. Autino, A. Sathicq and G. Romanelli, *Compt. Rendus Chem.*, 2016, **19**, 551–555.
- 29 For **4p**, see: J. J. Ares, P. E. Outt, S. V. Kakodkar, R. C. Buss and J. C. Geiger, *J. Org. Chem.*, 1993, **58**, 7903–7905.
- 30 For **4q** and **4r**, see: D. Banerjee, U. Kayal and G. Maiti, *Tetrahedron Lett.*, 2016, **57**, 1667–1671.
- 31 For **4u**, see: J. Zhao, Y. Zhao and H. Fu, *Angew. Chem., Int. Ed.*, 2011, **50**, 3769–3773.
- 32 For **4v**, see: T.-L. Shih, C.-E. Chou, W.-Y. Liao and C.-A. Hsiao, *Tetrahedron*, 2014, **70**, 3657–3664.
- 33 For **4w**, see: B. R. Nawghare, S. V. Gaikwad, A. Raheem and P. D. Lokhande, *J. Chil. Chem. Soc.*, 2014, **59**, 2284–2286.
- 34 For **4x**, see: M. Yoshida, Y. Fujino and T. Doi, *Org. Lett.*, 2011, **13**, 4526–4529.
- 35 For **5a**, see: M. Shaikh, K. K. Atyam, M. Sahu and K. V. S. Ranganath, *Chem. Commun.*, 2017, **53**, 6029–6032.
- 36 For **6a**, see: C. Proença, H. M. T. Albuquerque, D. Ribeiro, M. Freitas, C. M. M. Santosb, A. M. S. Silvab and E. Fernandes, *Eur. J. Med. Chem.*, 2016, **122**, 381–392.
- 37 For **7a**, see: W. Li and X.-F. Wu, *Eur. J. Org. Chem.*, 2015, **2015**, 331–335.