

Research article

Open Access

Genome-wide identification of *Xenopus* matrix metalloproteinases: conservation and unique duplications in amphibians

Liezhen Fu, Biswajit Das, Smita Mathew and Yun-Bo Shi*

Address: Section on Molecular Morphogenesis, PCRM, NICHD, NIH, Bldg. 18T, Rm. 106, Bethesda, MD 20892, USA

Email: Liezhen Fu - fuliez@helix.nih.gov; Biswajit Das - dasbiswa@mail.nih.gov; Smita Mathew - mathewsm@mail.nih.gov; Yun-Bo Shi* - shi@helix.nih.gov

* Corresponding author

Published: 17 February 2009

Received: 29 October 2008

BMC Genomics 2009, 10:81 doi:10.1186/1471-2164-10-81

Accepted: 17 February 2009

This article is available from: <http://www.biomedcentral.com/1471-2164/10/81>

© 2009 Fu et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Matrix metalloproteinases (MMPs) are members of the superfamily of Zn²⁺ dependent extracellular or membrane-bound endopeptidases which have been implicated to play critical roles in vertebrate development and human pathogenesis. A number of MMP genes have been found to be upregulated in some or all organs during frog metamorphosis, suggesting that different MMPs may have different functions in various organs/tissues. The recent advances in EST (expressed sequence tag) sequencing and the completion of the genome of *Xenopus* (*X.*) *tropicalis* prompted us to systematically analyze the existence of MMPs in the *Xenopus* genome.

Results: We examined *X. laevis* and *X. tropicalis* ESTs and genomic sequences for MMPs and obtained likely homologs for 20 out of the 25 MMPs known in higher vertebrates. Four of the five missing MMPs, i.e. MMPs 8, 10, 12 and 27, were all encoded on human Chromosome 11 and the other missing MMP, MMP22 (a chicken MMP), was also absent in human genome. In addition, we identified several novel MMPs which appears to be derived from unique duplications over evolution, are present in the genomes of both *Xenopus* species.

Conclusion: We identified the homologs of most of the mammalian MMPs in *Xenopus* and discovered a number of novel MMPs. Our results suggest that MMP genes undergo dynamic changes over evolution. It will be of interest in the future to investigate whether MMP expression and functions during vertebrate development are conserved. The sequence information reported here should facilitate such an endeavor in the near future.

Background

Matrix metalloproteinases (MMPs) are Zn²⁺ dependent extracellular or membrane-bound proteinases with overlapping substrate specificities [1-6]. They are capable of cleaving proteinaceous components of the extracellular matrix (ECM) as well as non-ECM proteins [2-5,7,8], thus affecting cell fate through modifications of cell's microenvironment. MMPs have a similar domain structure that includes a propeptide for secretion, a propeptide to main-

tain latency, and a catalytic domain, featured by the signature sequence HEFGHXXH, for substrate cleavage. The catalytic domain binds to a Zn²⁺ ion through the three-histidine residues within the signature sequence to form the catalytic center [5,9,10]. The propeptide contains a highly conserved sequence, PRCGXP, the so called "cysteine switch", within which the cysteine residue interacts with the catalytic Zn²⁺ to maintain enzyme latency [11]. Most MMPs are secreted as latent enzymes and proc-

essed to the active forms upon the removal of the propeptide domain through various mechanisms. Other MMPs, such as stromelysin 3 (ST3, also known as MMP11), MMP21, MMP23, MMP28, and membrane type MMPs (MT-MMPs) are activated intracellularly through the removal of the propeptide domain by furin, a Golgi enzyme [3,12,13].

MMP expression and distribution have long implicated that MMPs play important roles in many physiological processes including embryonic development, angiogenesis, tissue resorption and remodeling, and pathological events such as tumor invasion and arthritis [8,14-22]. *In vitro* and cell culture studies have provided strong evidence to show that MMPs can regulate cell fate and behavior by remodeling the ECM. On the other hand, increasing evidences indicate that MMPs are capable of cleaving non-ECM extracellular or membrane-bound proteins, suggesting the existence of multiple pathways for MMPs to regulate cells. Despite the extensive *in vitro* and cell culture studies, the *in vivo* functions of MMPs are poorly understood. Surprisingly, with a few exceptions, transgenic overexpression of MMPs and MMP knockouts in mouse have little or weak phenotypes on mouse development [23,24]. This appears to be at least in part due to the redundancy in MMP expression and function. These findings emphasize the need for further *in vivo* studies by using different model systems.

Frog metamorphosis offers a unique opportunity to study MMP function during postembryonic development in vertebrates. This process is totally dependent on the presence of thyroid hormone (TH) and mimics the postembryonic period from a few months before to several months after birth in humans [25-27]. During metamorphosis, dramatic tissue-specific remodeling occurs through TH-regulated cell fate changes. These include complete absorption of the gill and the tail, *de novo* generation of the limbs, and remodeling of most other organs such as the intestine. For example, in the intestine, the larval epithelial cells die through apoptosis and adult epithelial progenitor cells, which may be derived from dedifferentiated larval epithelial cells, proliferate and eventually differentiate to form a multiply folded adult epithelium [28-31]. Numerous studies have shown that the metamorphic effects of TH are mediated by thyroid hormone receptors, which control a gene regulation cascade by regulating the transcription of the so-called direct TH-response genes. These direct response genes in turn affect the expression of indirect TH-response genes to eventually regulate cell fate and behavior during metamorphosis. Initial isolation and characterization of TH-response genes revealed that *Xenopus (X.) laevis* ST3 (MMP11) and collagenase 3 (MMP13), and *Rana catesbeiana* collagenase 1 (MMP1) are regulated by TH during metamorphosis. Sub-

sequent studies have found that essentially all MMPs analyzed so far are regulated by TH in at least some organs/tissues during metamorphosis [32-48]. Among them, ST3 and MMP9-TH in *X. laevis* and collagenase 1 in *Rana catesbeiana* have been shown to be direct response genes with thyroid hormone response elements present in their promoters [43,49,50]. Furthermore, *in vitro* organ culture analysis and *in vivo* analyses have provided strong evidence for the participation of MMPs in metamorphosis [40,41,51-54]. For example, we have demonstrated that ST3 is required for TH-induced ECM remodeling, intestinal larval epithelial apoptosis as well as adult epithelial cell migration in organ cultures and that transgenic overexpression of ST3 alone at premetamorphic stages, e.g., stage 54, can induce larval epithelial apoptosis and ECM remodeling in the intestine in the absence of TH [52,53]. These functional studies directly proved the function of ST3 as first suggested based on expression analyses. Since all MMPs analyzed so far are regulated by TH during metamorphosis, it is pertinent to ask whether the rest of the MMPs are also regulated by TH and whether different MMPs have different functions during metamorphosis in different organs/tissues.

As an initiative to begin to address these important issues, we have carried out a genome-wide analysis of MMP genes in both *X. laevis* and *X. tropicalis* through a bioinformatic approach by making use of the genome sequence information for *X. tropicalis* and cDNA sequences available for *X. laevis* and *tropicalis* genes from the NIH Frog Initiatives Program. We demonstrate that essentially all mammalian MMPs have homologs in *Xenopus*, although the homologs for some MMPs cannot be assigned with certainty. Furthermore, we have discovered a number of novel MMPs and duplications that are uniquely present in the amphibian genome.

Results and discussion

Bioinformatic search for *Xenopus* MMPs

Many *X. laevis* MMPs were previously cloned [32,33,39,41,43,44,46,47,55-58]. These cDNA sequences were used to search for other MMPs in the public EST database at the NCBI <http://www.ncbi.nlm.nih.gov/> and the Gene Index Project in Computational Biology and Function Genomics Laboratory <http://compbio.dfci.harvard.edu/tgi/>. Putative MMP protein sequences that were derived from the retrieved cDNA sequences were pooled and analyzed on a phylogenetic tree. Closely related entries were compared pair-wise by using MacVector (Accelrys Inc., San Diego, CA) and redundant sequences were removed. The resulting *X. laevis* MMPs were listed in Table 1 (see Additional file 1 for their nucleotide sequences). Compared to human MMPs, some *X. laevis* MMPs were missing from the list and some others had highly homologous duplicates, likely due to the pseu-

Table 1: The amino acid identities of XI-MMPs compared to their counterpart Xt-MMPs#.

| <i>Xenopus laevis</i> Name | Amino Acids | Name | <i>Xenopus tropicalis</i> Amino Acids | Scaffold | Identity (%) |
|-------------------------------|-------------|-----------|--|----------|--------------|
| XI-MMP1A | 466 | Xt-MMP1 | 466 | 119 | 87 |
| XI-MMP1B | 466 | | | | 87 |
| XI-MMP2 | 656 | Xt-MMP2 | 655 | 458 | 94 |
| XI-MMP3 | 458 | Xt-MMP3 | 497 | 119 | 84 |
| XI-MMP7A | 252 | Xt-MMP7 | 259 | 119 | 87 |
| XI-MMP7B | 259 | | | | 81 |
| XI-MMP9 | 671 | Xt-MMP9 | 670 | 29 | 87 |
| XI-MMP9TH | 683 | Xt-MMP9TH | 683 | 29 | 91 |
| XI-MMP11 | 477 | Xt-MMP11 | 477 | 12 | 93 |
| XI-MMP13 | 469* | Xt-MMP13 | 472 | 119 | 93 |
| XI-MMP13A | 472 | | | | 93 |
| XI-MMP14A | 575 | Xt-MMP14 | 578 | 792 | 94 |
| XI-MMP14B | 576 | | | | 93 |
| XI-MMP15 | 262* | Xt-MMP15 | 648 | 6 | 97 |
| XI-MMP16 | 592 | Xt-MMP16 | 607 | 452 | 94 |
| XI-MMP17 | 159* | Xt-MMP17 | 588 | 12 | 95 |
| XI-MMP18 | 467 | Xt-MMP18 | 467 | 119 | 86 |
| XI-MMP19 | 123* | Xt-MMP19 | 476 | 101 | 95 |
| XI-MMP20 | 478 | Xt-MMP20 | 458 | 119 | 89 |
| XI-MMP21 | 604 | Xt-MMP21 | 604 | 32 | 92 |
| XI-MMP23 | 381 | Xt-MMP23 | 335* | 414 | 80 |
| XI-MMP24A | 361* | Xt-MMP24 | 603 | 954 | 99 |
| XI-MMP24B | 247* | | | | 97 |
| XI-MMP25 | 546 | Xt-MMP25 | 545 | 1214 | 89 |
| XI-MMP26 | 258 | Xt-MMP26 | 261 | 119 | 88 |
| XI-MMP28A | 496 | Xt-MMP28 | 499 | 72 | 92 |
| XI-MMP28B | 497 | | | | 91 |
| XI-MMP N1 | 562* | Xt-MMP N1 | 573 | 501 | 90 |
| XI-MMP N3 | 519* | Xt-MMP N3 | 627 | 508 | 91 |
| | | Xt-MMP N2 | 260 | 119 | N/A |
| | | Xt-MMP N4 | 455 | 119 | N/A |
| | | Xt-MMP N5 | 422 | 119 | N/A |
| | | Xt-MMP N6 | 364 | 132 | N/A |

#Comparison of *X. laevis* and *X. tropicalis* MMPs. Pair-wise comparisons were done to obtain the percent of identities between the MMPs from the two species (*: incomplete sequences).

dotetraploid *X. laevis* genome. The missing ones could be due to either the absence of the genes in *Xenopus* genome or incomplete sequence data available. Thus, we also searched and analyzed the cDNA sequences for MMPs in the highly related species, *X. tropicalis*. We aligned the cDNA sequences of *X. laevis* or *tropicalis* MMPs to the JGI *X. tropicalis* genomic scaffolds <http://genome.jgi-psf.org/Xentr4/Xentr4.home.html> (Table 1). When needed, we also used human MMP sequences to search the *X. tropicalis* genome database to ensure a complete search of the genome.

Pair-wise sequence comparison (not shown) and phylogenetic analysis (Fig. 1) allowed us to assign most of the *Xenopus* MMPs to the corresponding human homologs (except that chicken MMP22 was also used since MMP22 is absent in human [57]) (where an MMP name had been

previously assigned in the databases, we kept the same name for consistency. As we discussed below, some of these MMPs may not be the true homologs of the human MMPs as currently assigned). As shown in table 1, all *X. laevis* MMPs have a corresponding homolog in *X. tropicalis* and the homologs are highly conserved with over 80% amino acid sequence identities between *X. laevis* and *X. tropicalis*. These include a duplicated gelatinase B (MMP9TH) that is absent in mouse and human genome. As *X. laevis* is a pseudotetraploid organism, it is not surprising that some MMPs (MMP1, MMP7, MMP13, MMP14, MMP24, MMP28) have duplicate copies in *X. laevis* but only one in *X. tropicalis*. Only four *X. tropicalis* MMPs (MMP N2, MMP N4, MMP N5, and MMP N6, see below for more on these MMPs) have no available homologs in *X. laevis* yet, likely due to incomplete cDNA sequence information available for *X. laevis*.

Table 2: Percent homology between *Xenopus tropicalis* and human MMP proteins#.

| MMP | H1 | H2 | H3 | H7 | H8 | H9 | H10 | H11 | H12 | H13 | H14 | H15 | H16 | H17 | H19 | H20 | H21 | C22 | H23 | H24 | H25 | H26 | H27 | H28 |
|------|----|-----------|----|----|----|-----------|-----|-----------|-----|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----|-----------|-----------|-----------|-----|-----|-----------|
| X1 | 51 | 42 | 54 | 48 | 52 | 36 | 51 | 32 | 51 | 53 | 38 | 38 | 40 | 36 | 33 | 46 | 28 | 50 | 25 | 41 | 36 | 39 | 49 | 33 |
| X2 | 44 | 80 | 44 | 49 | 43 | 48 | 44 | 38 | 42 | 48 | 32 | 24 | 32 | 30 | 30 | 40 | 27 | 44 | 27 | 32 | 32 | 39 | 38 | 29 |
| X3 | 48 | 38 | 50 | 50 | 47 | 33 | 47 | 33 | 46 | 46 | 36 | 36 | 37 | 31 | 30 | 43 | 27 | 45 | 26 | 37 | 31 | 41 | 42 | 31 |
| X7 | 54 | 54 | 55 | 51 | 53 | 49 | 53 | 35 | 55 | 55 | 44 | 45 | 46 | 38 | 36 | 51 | 36 | 53 | 29 | 47 | 39 | 43 | 54 | 41 |
| X9 | 37 | 50 | 40 | 41 | 38 | 57 | 40 | 34 | 37 | 41 | 25 | 24 | 26 | 26 | 32 | 34 | 24 | 38 | 32 | 23 | 30 | 37 | 34 | 27 |
| X9TH | 38 | 49 | 40 | 38 | 40 | 59 | 41 | 36 | 37 | 42 | 25 | 25 | 31 | 24 | 32 | 35 | 24 | 40 | 31 | 24 | 30 | 38 | 35 | 30 |
| X11 | 34 | 34 | 37 | 36 | 33 | 34 | 35 | 62 | 31 | 32 | 39 | 38 | 39 | 39 | 33 | 33 | 31 | 33 | 32 | 40 | 40 | 36 | 33 | 30 |
| X13 | 46 | 45 | 49 | 46 | 49 | 38 | 51 | 34 | 43 | 67 | 38 | 42 | 38 | 35 | 33 | 45 | 29 | 48 | 26 | 39 | 34 | 40 | 49 | 31 |
| X14 | 38 | 35 | 38 | 41 | 36 | 33 | 37 | 38 | 39 | 38 | 77 | 60 | 57 | 36 | 34 | 37 | 27 | 38 | 26 | 56 | 38 | 35 | 35 | 34 |
| X15 | 41 | 22 | 42 | 43 | 38 | 25 | 41 | 38 | 40 | 40 | 59 | 69 | 55 | 38 | 37 | 38 | 27 | 38 | 26 | 53 | 42 | 37 | 35 | 36 |
| X16 | 41 | 34 | 38 | 41 | 39 | 31 | 40 | 36 | 39 | 40 | 56 | 57 | 87 | 36 | 35 | 38 | 26 | 39 | 30 | 67 | 38 | 36 | 35 | 32 |
| X17 | 35 | 32 | 37 | 38 | 35 | 24 | 36 | 37 | 36 | 36 | 37 | 38 | 35 | 70 | 34 | 36 | 27 | 34 | 25 | 37 | 47 | 37 | 33 | 31 |
| X18 | 55 | 47 | 53 | 48 | 52 | 36 | 50 | 34 | 48 | 51 | 38 | 40 | 39 | 36 | 35 | 43 | 29 | 50 | 28 | 41 | 34 | 41 | 50 | 33 |
| X19 | 36 | 37 | 36 | 37 | 33 | 34 | 37 | 33 | 35 | 37 | 38 | 38 | 38 | 34 | 60 | 35 | 34 | 35 | 25 | 40 | 35 | 37 | 37 | 35 |
| X20 | 45 | 41 | 47 | 46 | 44 | 36 | 47 | 33 | 44 | 45 | 39 | 39 | 38 | 36 | 32 | 70 | 30 | 44 | 25 | 39 | 36 | 35 | 42 | 35 |
| X21 | 30 | 23 | 30 | 38 | 30 | 19 | 31 | 29 | 32 | 29 | 28 | 26 | 24 | 26 | 28 | 29 | 59 | 29 | 25 | 25 | 27 | 31 | 27 | 28 |
| X23 | 28 | 29 | 24 | 24 | 23 | 33 | 22 | 28 | 24 | 24 | 26 | 27 | 32 | 25 | 22 | 24 | 27 | 25 | 60 | 32 | 26 | 24 | 22 | 25 |
| X24 | 40 | 34 | 41 | 42 | 40 | 27 | 41 | 39 | 41 | 39 | 54 | 56 | 69 | 36 | 35 | 38 | 26 | 38 | 30 | 84 | 38 | 35 | 37 | 34 |
| X25 | 36 | 33 | 34 | 37 | 38 | 31 | 35 | 37 | 36 | 34 | 39 | 42 | 40 | 50 | 35 | 32 | 30 | 34 | 29 | 41 | 49 | 35 | 30 | 31 |
| X26 | 53 | 54 | 55 | 48 | 53 | 51 | 53 | 36 | 53 | 59 | 46 | 44 | 46 | 38 | 38 | 54 | 36 | 53 | 28 | 48 | 38 | 40 | 55 | 39 |
| X28 | 34 | 34 | 34 | 35 | 31 | 30 | 33 | 31 | 33 | 33 | 35 | 37 | 35 | 32 | 35 | 34 | 30 | 31 | 29 | 36 | 32 | 36 | 29 | 56 |
| XN1 | 34 | 29 | 35 | 40 | 34 | 27 | 36 | 31 | 35 | 32 | 30 | 31 | 30 | 26 | 33 | 33 | 24 | 35 | 27 | 29 | 27 | 38 | 31 | 30 |
| XN2 | 56 | 56 | 56 | 54 | 57 | 48 | 56 | 37 | 54 | 57 | 44 | 44 | 47 | 38 | 42 | 54 | 36 | 53 | 33 | 46 | 40 | 41 | 53 | 40 |
| XN3 | 30 | 21 | 30 | 36 | 28 | 18 | 31 | 29 | 29 | 30 | 26 | 27 | 27 | 23 | 28 | 28 | 52 | 29 | 26 | 25 | 27 | 31 | 29 | 30 |
| XN4 | 49 | 40 | 51 | 47 | 48 | 35 | 51 | 34 | 49 | 48 | 38 | 39 | 40 | 35 | 32 | 46 | 28 | 50 | 27 | 42 | 34 | 39 | 49 | 31 |
| XN5 | 47 | 40 | 51 | 44 | 45 | 35 | 50 | 33 | 47 | 48 | 36 | 38 | 39 | 34 | 31 | 45 | 29 | 48 | 27 | 41 | 32 | 40 | 46 | 32 |
| XN6 | 28 | 30 | 28 | 27 | 29 | 34 | 29 | 30 | 28 | 29 | 28 | 31 | 30 | 25 | 30 | 29 | 26 | 28 | 44 | 32 | 29 | 30 | 27 | 32 |

#Comparison of *X. tropicalis* and human MMP protein sequences. Pair-wise comparisons were done to obtain the percent of identities between the MMPs from the two species. H1 ~ 28: human MMP1 ~ 28; C22: chicken MMP22; X1 ~ 28, X9TH, and XN1 ~ N6: *Xenopus tropicalis* MMP1 ~ 28, 9TH, and N1 ~ N6, respectively. The highlighted numbers with bold letters indicate homologies suggesting that *X. tropicalis* and human MMPs are homologs.

Xenopus MMPs with known human homologs

Phylogenetic (Fig. 1) and pair-wise sequence (table 2) analyses suggest that the following human MMPs have true homologs in *Xenopus* genome: MMP2, MMP9, MMP11, MMP13, MMP14, MMP15, MMP16, MMP17, MMP19, MMP20, MMP21, MMP23, MMP24, MMP25, MMP28. The homologous human and *X. tropicalis* MMPs cluster together in the phylogenetic tree (Fig. 1) and share highest sequence identities with each other than with any other MMPs (one exception to this is *X. tropicalis* MMP25, which shares similar homologies with human MMP17 and MMP25. Since the structurally related *X. tropicalis* MMP17 share 70% homology with the human MMP17 but only 47% homology with the human MMP25, we assigned this MMP as *X. tropicalis* MMP25) (table 2). In addition, the homologous MMPs have similar lengths and domain organizations (data not shown).

Likely Xenopus homologs of human MMPs

The homologs of the rest of human MMPs could not be easily identified based on sequence comparison and phylogenetic analysis. These MMPs may have corresponding homologs in *Xenopus* but their sequences have diverged significantly that it is difficult to match the human and *Xenopus* counterparts. For these MMPs, we kept the putative names for any *Xenopus* MMPs with previously assigned names in the public databases or assigned the names as described below.

MMP1

There were two entries for *X. laevis* MMP1 (GenBank accession # [BC054233](#) and [BC084836](#)), encoding two closely related MMPs of 466 amino acids (aa) that are 90% identical (data not shown). Alignment of these *X. laevis* MMP1s (MMP1A and MMP1B) to the *X. tropicalis* genomic scaffolds showed significant homology at three different loci on the Scaffold_119 (Note that the *X. tropicalis* genomic sequence is not complete and the individual sequences are assembled into scaffolds instead of individual chromosomes). The putative cDNA sequences were derived from these loci and used to deduce the protein sequences of three related MMPs. Among these three putative MMP genes, the best-matched one has 87% identities with *X. laevis* MMP1s and has the same length; it was therefore named as *X. tropicalis* MMP1 (Table 1). The other two were tentatively named as *X. tropicalis* MMP N4 and N5 (Table 1).

Phylogenetic analysis revealed that *X. tropicalis* MMP1, MMP N4 and N5 cluster together with MMP N4 and N5 more closely related to each other than to MMP1 (Fig. 1). These MMPs are related to several human MMP sub-families including collagenases (MMP1 and 8) and stromelysins (MMP3 and 10), etc. The MMP that is most closely related to these three MMPs is *X. tropicalis* MMP18,

a homolog of *X. laevis* MMP18 (Table 1). *X. laevis* MMP18 is a known collagenase [42], suggesting that these three MMPs are collagenases. Apart from the typical MMP domains (i.e., the signal peptide, the conserved zinc binding motif characteristic of MMPs, and the conserved cysteine-switch domain within the propeptide), *X. tropicalis* MMP1, as well as *X. laevis* MMP1A and 1B, contains a 16 aa proline-rich motif after the catalytic domain that distinguishes collagenases from stromelysins, although the *X. tropicalis* MMP N4 and N5 have deletions within the region (*X. tropicalis* MMP N5 lacks the entire hinge domain but has the intact catalytic and hemopexin domain, a characteristic similar to that of the MMP21s) (Fig. 2). These MMPs share less than 60% identity with *X. laevis* collagenases MMP13 and MMP18. In addition, they are also three amino acids shorter than the human MMP1 at the C-terminus (Fig. 2), just like the *X. laevis* MMP13 and MMP18. Taken together, these MMP genes are likely *X. tropicalis* collagenases, but it is possible that the *X. tropicalis* MMP1 is not the true homolog of human MMP1, especially considering that the MMP1 from another amphibian species, *Rana catesbeiana*, is much more homologous to human MMP1, although much shorter, compared to *Xenopus* MMP1 (Fig. 1) [42].

MMP3

MMP3 and MMP10, also referred to as stromelysin 1 and 2, respectively, are MMPs that have quite broad substrate specificities and were originally described as proteoglycanases [59-62]. Neither *Xenopus* MMP3 nor MMP10 has been characterized. A putative MMP deduced from a cDNA entry for each *Xenopus* species (GenBank accession number: [BC077966](#) of *X. laevis* clone and [NM_001030331](#) of *X. tropicalis* clone) structurally resembles MMP3 and MMP10 (Fig. 3). Similar to human MMP3 and MMP10, the *Xenopus* MMP has an insertion (8 aa for the *X. tropicalis* MMP and 14 aa for the *X. laevis* MMP) in the 16 aa proline-rich motif after the catalytic domain whose integrity is important for collagenase activity (Fig. 3). This suggests that the *Xenopus* MMP is likely a stromelysin. The two *Xenopus* homologs share 84% identity, although the *X. laevis* one is 45 aa shorter than *X. tropicalis* MMP3 at the N-terminus, likely due to incomplete 5'-end cDNA sequence. Thus, the *Xenopus* MMPs are homologs of each other and are tentatively named as MMP3 since they are slightly more similar to human MMP3 than MMP10 (Table 2).

MMP23

MMP23 is characterized by the presence of a furin activation site, a type II transmembrane domain at the N-terminus, and a unique truncated C-terminal domain unrelated to the hemopexin domain found in other MMPs, and it lacks a typical prepeptide [63-66]. There are two reported MMP23, MMP23A and MMP23B, in human that are




Figure 2
Sequence comparison of MMP1 with MMP N4 and MMP N5. *X. tropicalis* (Xt) MMP1, N4 and N5, and *X. laevis* (Xl) MMP1A and 1B were aligned with human (h) MMP1 for comparison. The sequences of the putative signal peptide are underlined. The predicted cleavage site between the signal peptide and the propeptide is indicated by an arrow, and the predicted cleavage site between the propeptide and the catalytic domain is indicated by solid arrowhead. The conserved sequence in the propeptide involved in the "cysteine-switch" is boxed, and the zinc-binding motif within the bracketed catalytic domain is indicated by a solid line on top. The three conserved histidine residues in the zinc binding motif and the conserved methionine residue of the nearby "Met-turn" are indicated by stars below. The 16 aa sequence (shaded) at the end of the catalytic domain (bracketed) indicates the region whose integrity is important for collagenase specificity for collagen. An insertion of 8 or more aa within this region at the site indicated by an arrowhead is characteristics of stromelysins. The arrow marked "C" shows the beginning of the C-terminal hemopexin-like domain. A dot indicates an identical amino acid as the corresponding one in Xt-MMP1. Gaps (dashes) are introduced to optimize the alignment among proteins. Note that MMP N4 and N5 contain internal deletions in the linker region between the catalytic domain and C-terminal hemopexin-like domain.

Scaffold_132. *X. tropicalis* MMP23 and MMP N6 differ from each other at both the cDNA and protein sequence levels, unlike the two human MMP23 genes (note that there is only one MMP23 in the mouse genome). Sequence comparison showed that *X. laevis* MMP23, *X. tropicalis* MMP23 and MMP N6 have the same features of human MMP23, although there is a sequence gap in *X. tropicalis* MMP23 (Fig. 4), possibly due to incomplete genomic sequence information (see Additional file 1). The *Xenopus* MMP 23, MMP N6, and human MMP23 cluster together and are away from all other MMPs (Fig. 1) with the MMP23 sharing 60% identity between *Xenopus* and human, similar to other homologous MMPs (table 2).

encoded by two genes, likely due to a very recent, partial duplication at Chromosome 1p36.3 [65]. Human MMP23A and B are identical in amino acid sequences and thus both are referred to as MMP23 here. Two overlapping EST entries (CD302225 and CD302813) encode a putative *X. laevis* MMP23 of 381aa. Sequence search of the *X. tropicalis* genome identified a putative *X. tropicalis* MMP23 on Scaffold_414. There was no EST entry representing the *X. tropicalis* MMP23. However, two *X. tropicalis* EST entries (CX344815 and CX344816) composed of cDNA sequences that together encode another putative MMP related to MMP23 (tentatively named as MMP N6). These cDNA sequences aligned on to *X. tropicalis* genomic

MMP21

MMP21 was first cloned in *X. laevis* [67]. It has since been found to be present in other vertebrates including human. The common features of this MMP across different species, in addition to those characteristics of MMPs, are a putative furin cleavage site between the propetide and the catalytic domain, a relative long insertion (20~44 aa) between the PRCGXPD cysteine switch motif and the furin cleavage site (RXKR), and a unique cysteine residue in the catalytic domain (Fig. 5). The putative *X. tropicalis* MMP21 composes of 604 aa and shares 92% and 59% identities with the *X. laevis* and human MMP21, respectively (Tables 1 and 2).




Figure 3
Comparison of *Xenopus* MMP3 with human MMPI, 3, and 10. Note that the shadowed region at the end of the catalytic domain corresponds to the same region in Fig. 2 except the insertion of 8–14 aa in stromelysins (MMP3, 10) compared to collagenases (e.g., the MMPI shown here). See Fig. 2 for other information.

| | | | |
|-----------|-----|---|-----|
| Xt-MMP23 | 1 | <u>MGDIQEIEHWQK</u> ----RYIWTFLAIFAGTVLVAGIFTASNSVSLDSKVDF | 46 |
| Xt-MMP N6 | 1 | <u>MDWGLADR</u> ----S.G.ICVSAAVTFL.LSNW.CYQOKSLIFP.FQN | 42 |
| Xl MMP23 | 1 | .DGT.D...RR.----.I.A...L.....L.....V.K..P.ET.... | 46 |
| hMMP23 | 1 | .RGARVPSEAPGAGVERR.LGA.LV.LCL.P.LVLLARLGAPAVPAWSA | 50 |
| Xt-MMP23 | 47 | VVAPTPALQL----PLQLPRHL-- RNKRY TLTPGLLKWDHYNLTyrIVSF | 90 |
| Xt-MMP N6 | 43 | E.VECGTNCS-----FRSSILT-- RIN.LGY....L....K..Q. | 85 |
| Xl MMP23 | 47 |Q.....-----.....Y..... | 90 |
| hMMP23 | 51 | AQGDVA..G.SAVP.TRV.GP.AP. RRAR.R...F.....L. | 100 |
| Xt-MMP23 | 91 | PRNLINESDTKKGMAQAFQMWSEVSPFHFKEVPADQPSDLVIGFYGINHT | 140 |
| Xt-MMP N6 | 86 | -.TL.KD..ERAL.L..R...K..SLT.QR.QSH.V...R...TF..S | 134 |
| Xl MMP23 | 91 |R.....D.....K.....H...E..... | 140 |
| hMMP23 | 101 | ...LSPRE.RRAL.A..R...D...S.R..APE...R...P.... | 150 |
| Xt-MMP23 | 141 | DCLESYIHYCFDGTGELAHAYFPKTGEIHFDDSEFWILGNTRFSWKKGV | 190 |
| Xt-MMP N6 | 135 | .WG.PL.P...LN.....FL.PR.....NH...V..PS.....Q.. | 184 |
| Xl MMP23 | 141 |Y.....W..... | 190 |
| hMMP23 | 151 | ...V.AL.H...PT.....F..PH.G.....Y.V..P..Y..... | 200 |
| Xt-MMP23 | 191 | WLTDLVHVAAH----- | 203 |
| Xt-MMP N6 | 185 | .YN...Q...EIGHALGLWHSSNVTALMHPNATYTRIRHVTKDDIMAIQ | 234 |
| Xl MMP23 | 191 |EIGHALGLMHSLSNALMHINATLTGKKLISQDEIWGIF | 240 |
| hMMP23 | 201 |EIGHALGLMHSQHGRALMHLNATLRGWKALSQDELWGLH | 250 |
| | | * * * * | |
| Xt-MMP23 | 204 | ----}KDKYLVCASWAHKG Y CD S RKRIMKK Y CP Y SCDF C YDFPFPTKLPT | 251 |
| Xt-MMP N6 | 235 | SLY G CPSSGSR.Y.LEPS.S.G-----Q.HLK..S.KESLD--QR.K | 275 |
| Xl MMP23 | 241 | RLY G C.....R.L.....Q.. | 290 |
| hMMP23 | 251 | RLY G CL.RLF.....RR.F.DA.R.L..RL..S.....E.....VAT. | 300 |
| | | # # # # # # | |
| Xt-MMP23 | 252 | PPPPRTKVRLVPEGRNVTLRCGKKIMHKKGKVFYKDKELLEYSYGYLS | 301 |
| Xt-MMP N6 | 276 | QYRIKI.N.Y.SQ..P..FH.SH.VSQASKR.S....GAR.SS.TP.LVN | 325 |
| Xl MMP23 | 291 |T.....Y.....L.....S.... | 340 |
| hMMP23 | 301 |TRL.....F...Q..L..... <u>Y...Q.P..F..P...A</u> | 350 |
| Xt-MMP23 | 302 | LDDDHMSIIANAINEGLYTCIVKKRDRILTTYSW | 335 |
| Xt-MMP N6 | 326 | .SLSSLVLK.EEETQ.R...VIRHGKV.VGGK..NLHIT | 364 |
| Xl MMP23 | 341 | .N.....Q.....KHPTANV | 381 |
| hMMP23 | 351 | .GEA.L.II...V...T...V.RRQQRVL.....RVRVRG | 390 |

Figure 4
Comparison of frog and human MMP23 with MMP N6. The predicted signal anchor (transmembrane domain) sequences are underlined and the putative furin recognition sequences are in bold. The cysteine residues in the "cysteine-array" unique to MMP23 are in bold and indicated with # below. The amino acid residues characteristic of an Ig (immunoglobulin)-fold are indicated with rectangle boxes below. See Fig. 2 for other information.

| | | | |
|-----------|-----|---|-----|
| Xt-MMP N3 | 1 | <u>MESRI</u> <u>RPSLDRMEQFAHRGKMPVTVAIFLMLIAGW</u> | 36 |
| Xl-MMP N3 | 1 | | 0 |
| Xt-MMP21 | 1 | <u>MPSIKLLI</u> <u>WCSLCLISSRL</u> | 19 |
| Xl-MMP21 | 1 | <u>MPSIKLL</u> <u>WCCLCVLSPLRL</u> | 19 |
| hMMP21 | 1 | <u>MLAASIFRP</u> <u>PLLLCLVAAPWP</u> | 21 |
| Xt-MMP N3 | 37 | <u>SDSEKIF</u> <u>HNHRDHSDFDVQNI</u> <u>PHAGAILTPEAAQ</u> <u>YLVKYGWVAPVNWEEQ</u> | 86 |
| Xl-MMP N3 | 1 | | 0 |
| Xt-MMP21 | 20 | <u>CH</u> <u>.L.S.R.LQPSG</u> <u>.QQ.ELVKD</u> <u>LLS.A.T</u> | 61 |
| Xl-MMP21 | 20 | <u>CH</u> <u>.L.S.R.LQPSA</u> <u>.EQ.ELVKD</u> <u>MLS.F.A.T</u> | 61 |
| hMMP21 | 22 | <u>TQP</u> <u>.SL.S.R.LEPSP</u> <u>LRQ.KP.ADLH</u> <u>.RF.SR.SG</u> | 64 |
| Xt-MMP N3 | 87 | <u>AFTDFP</u> <u>VHDLAPQDVSLISEG</u> <u>EEHPRVSNSENETE</u> <u>INPFSIDSLKK</u> | 136 |
| Xl-MMP N3 | 1 | | 30 |
| Xt-MMP21 | 62 | <u>EPVI</u> <u>NDPSSANENEP</u> <u>.KDFSL</u> <u>MQ.GVSNARQ</u> <u>EVA.PTKS.Q.A</u> | 109 |
| Xl-MMP21 | 62 | <u>QVI</u> <u>NDPSSTNENEP</u> <u>.KDFSL</u> <u>MQ.GVSNARQ</u> <u>EVA.PTKS.Q.A</u> | 109 |
| hMMP21 | 65 | <u>VVA</u> <u>AWGSPSEGP</u> | 91 |
| Xt-MMP N3 | 137 | <u>FQDANGL</u> <u>TATGVLDSATKI</u> <u>AMNRC</u> <u>RGVDFK</u> <u>VASRRK</u> <u>NATLSFTD</u> <u>ADSN</u> | 186 |
| Xl-MMP N3 | 31 | <u>E</u> <u>.N.I.SOM</u> | 80 |
| Xt-MMP21 | 110 | <u>KL</u> <u>.N.P.I.D.TN</u> | 159 |
| Xl-MMP21 | 110 | <u>KL</u> <u>.N.P.V.T.D.TN</u> | 159 |
| hMMP21 | 92 | <u>R</u> <u>.A.P.S.E.A.LA</u> <u>.R</u> | 141 |
| Xt-MMP N3 | 187 | <u>SHSSQA</u> <u>----GAT</u> <u>RRKRS</u> <u>F</u> <u>LSKMVEH</u> <u>YRQREAK</u> <u>DTMGSS</u> <u>NKRFS</u> | 230 |
| Xl-MMP N3 | 81 | <u>HN</u> <u>----SOM</u> | 122 |
| Xt-MMP21 | 160 | <u>TKD</u> <u>.ENV</u> <u>TQONP</u> <u>DPKPV</u> <u>....FLD</u> <u>ML--SNKY</u> <u>.EQE</u> <u>LQK</u> <u>.TG</u> <u>.V</u> <u>.T</u> | 207 |
| Xl-MMP21 | 160 | <u>TKD</u> <u>.ENV</u> <u>TQONP</u> <u>DPKPI</u> <u>....FLD</u> <u>ML--YSNKY</u> <u>.EQE</u> <u>LQK</u> <u>.TG</u> <u>.V</u> <u>.T</u> | 207 |
| hMMP21 | 142 | <u>PPRA</u> <u>.SR</u> <u>.PDAPL</u> <u>-SL</u> <u>SRG</u> <u>WQPR</u> <u>GRG</u> <u>YDP</u> <u>GGAAQ</u> <u>LR</u> | 172 |
| Xt-MMP N3 | 231 | <u>RNTV</u> <u>KWRLMGE</u> <u>GYSMQ</u> <u>LTIQQ</u> <u>ORAIL</u> <u>LALAF</u> <u>RMWSE</u> <u>VVLLF</u> <u>VED</u> <u>LTG--D</u> | 278 |
| Xl-MMP N3 | 123 | <u>L</u> <u>.....R</u> <u>.....Q</u> <u>.....S</u> <u>.....N</u> <u>.....G</u> <u>.....Q</u> | 170 |
| Xt-MMP21 | 208 | <u>KKLL</u> <u>..MI</u> <u>.....N</u> <u>.S</u> <u>.NE</u> <u>.YVFR</u> <u>.....IM</u> <u>.D</u> <u>.E</u> <u>.N</u> <u>.SPL</u> <u>S</u> | 257 |
| Xl-MMP21 | 208 | <u>KKLL</u> <u>..MI</u> <u>.....N</u> <u>.S</u> <u>.NE</u> <u>.YVFR</u> <u>.....M</u> <u>.D</u> <u>.E</u> <u>.N</u> <u>.SPL</u> <u>S</u> | 257 |
| hMMP21 | 173 | <u>KR</u> <u>.LS</u> <u>.L</u> <u>.AL</u> <u>.S</u> <u>.SVAD</u> <u>.R</u> <u>.V</u> <u>.....T</u> <u>.D</u> <u>.R</u> <u>.LA</u> <u>APGA</u> | 222 |
| Xt-MMP N3 | 279 | <u>DIDIRI</u> <u>GFGTG</u> <u>QHLGCSQ</u> <u>AFD</u> <u>GVGQ</u> <u>QFAHAW</u> <u>YLGDI</u> <u>HFD</u> <u>DEH</u> <u>FVGP</u> <u>SS</u> | 328 |
| Xl-MMP N3 | 171 | | 220 |
| Xt-MMP21 | 258 | <u>Q</u> <u>..KL</u> <u>..R</u> <u>.R</u> <u>.....R</u> <u>.....S</u> <u>.E</u> <u>.....F</u> <u>.....TA</u> <u>....</u> | 307 |
| Xl-MMP21 | 258 | <u>Q</u> <u>..KL</u> <u>..R</u> <u>.R</u> <u>.....R</u> <u>.....S</u> <u>.E</u> <u>.....F</u> <u>.....TA</u> <u>....</u> | 307 |
| hMMP21 | 223 | <u>AV</u> <u>..KL</u> <u>..R</u> <u>.R</u> <u>.....PR</u> <u>.....S</u> <u>.E</u> <u>.....R</u> <u>.....TP</u> <u>.T</u> <u>.D</u> | 272 |
| Xt-MMP N3 | 329 | <u>HGISL</u> <u>LKVAV</u> <u>HEIGHAL</u> <u>GLSHIN</u> <u>REGSV</u> <u>MQPN</u> <u>YI</u> <u>QERH</u> <u>FEL</u> <u>DWED</u> <u>RKAV</u> | 378 |
| Xl-MMP N3 | 221 | | 270 |
| Xt-MMP21 | 308 | <u>T</u> <u>.....V</u> <u>.....H</u> <u>.V</u> <u>.....I</u> <u>.....DSG</u> <u>....RS</u> <u>..R</u> <u>.I</u> | 357 |
| Xl-MMP21 | 308 | <u>A</u> <u>.....V</u> <u>.....H</u> <u>.V</u> <u>.....I</u> <u>.....DSG</u> <u>....RS</u> <u>..R</u> <u>.I</u> | 357 |
| hMMP21 | 273 | <u>T</u> <u>.....V</u> <u>.....P</u> <u>.TY</u> <u>.T</u> <u>.....I</u> <u>.....PA</u> <u>.....S</u> <u>.....I</u> | 322 |
| Xt-MMP N3 | 379 | <u>QEKY</u> <u>GACD</u> <u>GSF</u> <u>STV</u> <u>FDW</u> <u>IRK</u> <u>ETPR</u> <u>GETV</u> <u>YR</u> <u>FNTY</u> <u>FFK</u> <u>TN</u> <u>WY</u> <u>MY</u> <u>EN</u> <u>KS</u> <u>N</u> | 428 |
| Xl-MMP N3 | 271 | <u>R</u> <u>.....R</u> <u>.....D</u> <u>.G</u> <u>F</u> <u>X</u> <u>.....S</u> <u>.....G</u> <u>.....S</u> | 320 |
| Xt-MMP21 | 358 | <u>NL</u> <u>.S</u> <u>.E</u> <u>.P</u> <u>.D</u> <u>.....Y</u> <u>.KNQY</u> <u>.L</u> <u>.V</u> <u>.Y</u> <u>.....R</u> <u>N</u> <u>S</u> <u>.....R</u> <u>.R</u> <u>....</u> | 407 |
| Xl-MMP21 | 358 | <u>NL</u> <u>.S</u> <u>.E</u> <u>.P</u> <u>.D</u> <u>.....Y</u> <u>.KNQY</u> <u>.L</u> <u>.V</u> <u>.Y</u> <u>.....R</u> <u>N</u> <u>S</u> <u>.....R</u> <u>.R</u> <u>....</u> | 407 |
| hMMP21 | 323 | <u>KL</u> <u>.S</u> <u>.E</u> <u>.....D</u> <u>.A</u> <u>.....NQY</u> <u>.VMV</u> <u>.S</u> <u>.....R</u> <u>N</u> <u>S</u> <u>.....L</u> <u>.L</u> <u>.....R</u> | 372 |
| Xt-MMP N3 | 429 | <u>RTRF</u> <u>GDPL</u> <u>PIKGG</u> <u>WNGI</u> <u>PKDK</u> <u>IDAY</u> <u>VHV</u> <u>WTW</u> <u>NIDA</u> <u>QY</u> <u>FF</u> <u>KG</u> <u>TQ</u> <u>VW</u> <u>RY</u> <u>DP</u> | 478 |
| Xl-MMP N3 | 321 | <u>L</u> <u>.....I</u> <u>.....S</u> <u>.....L</u> <u>.....L</u> <u>.....L</u> <u>.....L</u> <u>.....L</u> | 370 |
| Xt-MMP21 | 408 | <u>Y</u> <u>.....A</u> <u>.AT</u> <u>.S</u> <u>.AQN</u> <u>.....F</u> <u>.....TR</u> <u>.T</u> <u>.....Y</u> <u>.H</u> <u>.SE</u> | 457 |
| Xl-MMP21 | 408 | <u>Y</u> <u>.....A</u> <u>.AN</u> <u>.H</u> <u>.VQN</u> <u>.....F</u> <u>.....TR</u> <u>.S</u> <u>.....Y</u> <u>.SE</u> | 457 |
| hMMP21 | 373 | <u>Y</u> <u>.....I</u> <u>.Q</u> <u>.LT</u> <u>.P</u> <u>.....THN</u> <u>.....F</u> <u>.....I</u> <u>.....KR</u> <u>.ER</u> <u>.....Q</u> <u>.N</u> <u>.Y</u> <u>.....S</u> | 422 |
| Xt-MMP N3 | 479 | <u>KDRA</u> <u>FTED</u> <u>WRK</u> <u>VY</u> <u>QPI</u> <u>AD</u> <u>VFP</u> <u>GP</u> <u>IS</u> <u>PV</u> <u>DA</u> <u>AY</u> <u>FN</u> <u>KK</u> <u>ER</u> <u>FI</u> <u>Y</u> <u>FR</u> <u>GN</u> <u>V</u> <u>T</u> | 528 |
| Xl-MMP N3 | 371 | <u>N</u> <u>.....K</u> <u>.....V</u> <u>.....VEI</u> <u>.....V</u> <u>.....F</u> <u>.....I</u> <u>.....I</u> | 420 |
| Xt-MMP21 | 458 | <u>N</u> <u>.K</u> <u>.YA</u> <u>..TQ</u> <u>GKS</u> <u>..RL</u> <u>.SEG</u> <u>.....IN</u> <u>.....DRR</u> <u>QY</u> <u>.....DSQ</u> <u>.F</u> | 507 |
| Xl-MMP21 | 458 | <u>N</u> <u>.K</u> <u>.YA</u> <u>..AQ</u> <u>GKS</u> <u>..RL</u> <u>.SEG</u> <u>.....IN</u> <u>.....DRR</u> <u>QY</u> <u>.....DSQ</u> <u>.F</u> | 507 |
| hMMP21 | 423 | <u>Q</u> <u>.L</u> <u>.....EQ</u> <u>GKS</u> <u>..KL</u> <u>.SEG</u> <u>.....L</u> <u>.T</u> <u>.....DRR</u> <u>Q</u> <u>L</u> <u>.....KES</u> <u>L</u> <u>.F</u> | 472 |
| Xt-MMP N3 | 529 | <u>AF</u> <u>SV</u> <u>DK</u> <u>NQ</u> <u>K</u> <u>VE</u> <u>K</u> <u>PK</u> <u>SI</u> <u>ID</u> <u>V</u> <u>FP</u> <u>VP</u> <u>DN</u> <u>DH</u> <u>PR</u> <u>GN</u> <u>IDA</u> <u>V</u> <u>F</u> <u>S</u> <u>Y</u> <u>Q</u> <u>T</u> <u>FF</u> <u>I</u> <u>K</u> | 578 |
| Xl-MMP N3 | 421 | <u>P</u> <u>.....P</u> <u>.....I</u> <u>.....I</u> <u>.....I</u> <u>.....I</u> <u>.....I</u> <u>.....I</u> | 470 |
| Xt-MMP21 | 508 | <u>DIN</u> <u>.RV</u> <u>.PD</u> <u>.....R</u> <u>.F</u> <u>.....A</u> <u>.AA</u> <u>.N</u> <u>.K</u> <u>.....A</u> <u>.Y</u> <u>.T</u> <u>SS</u> <u>L</u> <u>.LF</u> | 557 |
| Xl-MMP21 | 508 | <u>DIN</u> <u>.RV</u> <u>.APD</u> <u>.....R</u> <u>.L</u> <u>.F</u> <u>.....A</u> <u>.AA</u> <u>.N</u> <u>.K</u> <u>.....VA</u> <u>.Y</u> <u>.T</u> <u>SS</u> <u>L</u> <u>.LF</u> | 557 |
| hMMP21 | 473 | <u>D</u> <u>.NR</u> <u>.R</u> <u>.V</u> <u>L</u> <u>NS</u> <u>.Y</u> <u>.R</u> <u>.TE</u> <u>.....A</u> <u>.I</u> <u>.QN</u> <u>.FR</u> <u>.....SA</u> <u>.Y</u> <u>.A</u> <u>.NS</u> <u>L</u> <u>.F</u> | 522 |
| Xt-MMP N3 | 579 | <u>DR</u> <u>Y</u> <u>W</u> <u>K</u> <u>M</u> <u>I</u> <u>D</u> <u>Q</u> <u>ER</u> <u>Q</u> <u>T</u> <u>NS</u> <u>SL</u> <u>PL</u> <u>NG</u> <u>LW</u> <u>PR</u> <u>K</u> <u>I</u> <u>N</u> <u>S</u> <u>Q</u> <u>W</u> <u>F</u> <u>I</u> <u>D</u> <u>I</u> <u>C</u> <u>D</u> <u>V</u> <u>H</u> <u>P</u> <u>S</u> <u>V</u> <u>L</u> <u>L</u> <u>S</u> <u>T</u> | 627 |
| Xl-MMP N3 | 471 | <u>F</u> <u>.....V</u> <u>.S</u> <u>.R</u> <u>.....S</u> <u>.....V</u> <u>.....V</u> <u>.....V</u> <u>.....V</u> <u>.....V</u> | 519 |
| Xt-MMP21 | 558 | <u>GR</u> <u>E</u> <u>F</u> <u>..V</u> <u>.S</u> <u>.K</u> <u>D</u> <u>.R</u> <u>Q</u> <u>.P</u> <u>.....Y</u> <u>.....F</u> <u>.....RT</u> <u>.SQ</u> <u>.....N</u> <u>.....L</u> <u>.KM</u> | 604 |
| Xl-MMP21 | 558 | <u>G</u> <u>.E</u> <u>F</u> <u>..V</u> <u>.S</u> <u>.K</u> <u>D</u> <u>.R</u> <u>Q</u> <u>.P</u> <u>.....Y</u> <u>.....F</u> <u>.....RA</u> <u>.SQ</u> <u>.....N</u> <u>.....L</u> <u>.KI</u> | 604 |
| hMMP21 | 523 | <u>GNA</u> <u>..V</u> <u>.N</u> <u>.K</u> <u>D</u> <u>K</u> <u>.Q</u> <u>.....W</u> <u>.....A</u> <u>.....F</u> <u>.....K</u> <u>.F</u> <u>.SEK</u> <u>.....V</u> <u>.....I</u> <u>.T</u> <u>.NM</u> | 569 |

Figure 5 Comparison of MMP21 with MMP N3. The predicted signal peptide is underlined and the putative furin recognition sequences within the propeptide are in bold. The sequence in white on dark background indicates the unique insertion in the propeptide in MMP21. A unique cysteine residue in the catalytic domain is in bold and indicated with a black diamond below. Note that the sequence for XI-MMP N3 is incomplete at the N-terminus. See Fig. 2 for other information.

Interestingly, sequence search also revealed a cDNA entry (TC84199), composed of *X. tropicalis* ESTs DR836290 and CX386748, that encodes a putative MMP of 524 aa. This MMP is highly homologous to *X. tropicalis* MMP21 and is tentatively named as *X. tropicalis* MMP N3 (Fig. 5). In addition, 4 highly homologous *X. laevis* ESTs, BG234242, BU905338, CB558404, and CF547511, were also found to constitute a putative cDNA sequence encoding a homolog of *X. tropicalis* MMP N3 (see Additional file 1, Table 1, and Fig. 5). The *X. tropicalis* MMP N3 is located on the *X. tropicalis* genomic Scaffold_508, from which the rest 5'-end cDNA sequence was predicted. The putative *X. tropicalis* MMP N3 encodes a predicted protein of 627 aa. The N-terminal sequence of *X. laevis* MMP N3 is still missing in the databases. Sequence comparison clearly showed that MMP N3 is highly homologous to MMP21 and is likely derived from a gene duplication event (Fig. 5). Furthermore, phylogenetic analysis indicated that *Xenopus* MMP21 clusters with human MMP21 and is closely related to MMP N3.

MMP7 and 26

MMP7 and MMP26 are the smallest MMPs known and have no hemopexin domain at their C-terminus [62,68-70]. *X. laevis* MMP7 was previously reported (GenBank accession # [AF573380](#)) and found to be expressed specifically in tissue resident macrophages [56]. Bioinformatic search revealed a closely related *X. laevis* cDNA clone (GenBank accession # [BC056040](#)) encoding a protein of 259 aa vs. 252 aa for *X. laevis* MMP7. These two *X. laevis* MMPs shared 85% identities in amino acid sequences and 82% identities in nucleotide sequences, and therefore, are likely duplicated MMP7 genes in the pseudotetraploid genome. We designated them as MMP7A (GenBank accession # [AF573380](#)) and 7B (GenBank accession # [BC056040](#)), respectively. Both *X. laevis* MMP7A and 7B aligned to *X. tropicalis* Scaffold_119 at the location that encodes a *X. tropicalis* cDNA (GenBank accession # [NM_001005043](#)). Thus, this *X. tropicalis* cDNA represents *X. tropicalis* MMP7. *X. tropicalis* MMP7 is 259 aa in length and highly homologous to *X. laevis* MMP7A/B (Tables 1 and Fig. 6).

In addition to the MMP7 genes, the *X. laevis* cDNA clone MGC69070 (GenBank accession # [BC056080](#)) and *X. tropicalis* cDNA clone MGC108008 (GenBank accession # [NM_001032335](#)) also encode a small MMP each (258 aa and 261 aa, respectively). Like MMP7, both proteins lacked the hemopexin domain (Fig. 6). They share 88% identities with each other and the *X. tropicalis* gene is also located on *X. tropicalis* genomic Scaffold_119. These two proteins are closely related to *X. laevis* MMP7A/B and *X. tropicalis* MMP7 on the phylogenetic tree and thus are tentatively named as *X. laevis* MMP26 and *X. tropicalis*




Figure 6

Comparison of MMP N2 with MMP7 and MMP26. Note that like human and *Xenopus* MMP7 and MMP26, MMP N2 lacks the linker peptide and hemopexin-like domain at the C-terminal. See Fig. 2 for other information.

MMP26, respectively. (It should be pointed out that it is difficult to assign with certainty which of the *Xenopus* gene is the homolog of human MMP7 and which is that of human MMP26. For consistency, we kept *Xenopus* MMP7 for the previously published sequence [56]).

Surprisingly, an additional *X. tropicalis* clone (IMAGE7719439, EST# CX982585 and CX982586) was also found to encode a similar MMP lacking the hemopexin domain. The cDNA sequence had two in-frame stop codons after the 3'-end of the coding sequence (see Additional file 1) and another independent EST sequence (EST #CX979196) overlapped with this region with 100% identity (data not shown). Thus, this gene represents a novel MMP that is structurally similar to MMP7 and

MMP26. It is tentatively named as *X. tropicalis* MMP N2 (Fig. 6). *X. tropicalis* MMP N2 is also located on *X. tropicalis* Scaffold_119 in between MMP7 and MMP26, suggesting that it was derived from a gene duplication event.

Novel *Xenopus* MMPs

Gelatinases

Sequence analysis revealed the existence of an alternatively spliced form of *X. laevis* gelatinase A (MMP2) (MMP2asv, Fig. 7). This alternatively spliced MMP2 transcript encodes a MMP that lacks most of the catalytic domain, including the zinc binding motif, and part of the C-terminal hemopexin domain. To date, no such spliced form of MMP2 has been reported for other vertebrate species, including *X. tropicalis*.

A

```

1   ATTCATACAAAGAACTGCAACACAGACATTAACCTCAAAGCTGTTTACCTGGAATGAGCAGTCCATTTATCTATTAGAACAGTTCATAAAGGTTTTCGG 100
101 CAGAAAGTCAGGCTTTAGCATGCGGACAATAAAATTATTAACGTTTTAGTACTCATTTCATAGTTTATAGTATATTTTATTATGTTTCCCTGCTCCA 200
      M R T I K I I N V L V L I F N S F S I F Y Y V S P A P
201 TCTCCTATCATAAAGTTTCCAGGAGACAAATCTCAAAGACAGACGTGGAACCTGCTGCGCAATACCTGAATCAGTTTTATGGCTGCCCTAAGGAAAAAT 300
      S P I I K F P G D K S P K T D V E L A A Q Y L N Q F Y G C P K E K
301 GTCACCTGATGGTCTTAAGGATGCCCTAAAGAAAATGCAAAGTTTTTTGGTCTTCCTGAAACTGGGGAGTTTGATCAAAATACCATTGAAACCATGAA 400
      C H L M V L K D A L K K M Q S F F G L P E T G E F D Q N T I E T M K
401 AAAGCCAAGATGTGGGAACCCCGATGTAGCCAACATAATTCTTTCCAGGAAACCGAAATGGGACAAAAACCATCTGACATACAGGATTCTTGGTTAT 500
      K P R C G N P D V A N Y N F F P R K P K W D K N H L T Y R I L G Y
      ↑
501 ACAACAGATCTGGATTCTGAAACAGTTGATGATGCTTTTTCACGTGCTTTTAAAGTTGGAGTGATGTCACACCATTGGAATTTAACAGAATTCATGATG 600
      T T D L D S E T V D D A F A R A F K V W S D V T P L E F N R I H D
601 GAGAAGCAGATATCATGATCAATTTGGACGATGGGAACATGGTGTACCAGATAATTTAGATGCTGTTCTAGATCAAACCGGAAGTGGCTACAGCTACTT 700
      G E A D I M I N F G R W E // H G V P D N L D A V L D Q T G S G Y S Y F
701 TTTCAAAGACTGGTATTACTTTCAAGTTGAAGACAAAAGTGTGAAGATTGTAAGGTTGGCAATGTCAAATGACTGGCTACGCTGCTGAAATATATTC 800
      F K D W Y Y F Q V E D K S V K I V K V G N V K N D W L R C *
801 ATTTAATTACTCTGTATTGCTTAACATGAACCTTTCATACGTTGATGGCATTTTTATAGCATAATAAATATAAGGGACCAGTCTGGTACTGGTATTT 900
901 GGGGAGCATACCAAGATATACATACAGTAACACTCAAACACTGCATTTTTAAGAATTTGTTTCATCCATTAGTTTTTTTCACTTGCTTGCTTAACTATAGC 1000
1001 TTTTAACTTTCATCTTAAATGTTTTTATTATTGTAAGCAGTGTGACAATGAGTTAACTCCTGCTACTTTTTTATATTGACCTTAATAAATCAAATCAAAT 1100
1101 TFACTAACCCATTTGTACATAGGTACCGCTGCCTATACGTGACACTTAAATAATGCTATGGATTTTTGAGAGAAAACACTGATAAGACCCCATTTCTGT 1200
1201 AAGTGCTTTCAATTTTATTTACCAAACAGTGCAGTGAGAGACTACAGTGCAAAATTAATCAACTATTGTGTGCTGTTATTTTACACAGCTGTATGTTATCC 1300
1301 ATATTTCAAAGAAAATCTGTACATCATCAAAGAACGTATATAATTATAACATAAAAGATAAACATTCAAAGTGTCTAATTTGTGCGCAAAAAGTAAAA 1400
1401 TAAATGGCAATTGTAGTTATGCATTGACATTACTTTTGGAAAGCTTTTAAAGGTTATTTAAAGTGTACCTGTCAACCCAGACACAAAAGCTGTATAATA 1500
1501 AAAGTCCTTTTCAAATTTGAAAAAATAAAAAAAAAAAAAAAAAAAAAAAAAA 1549
    
```

B





Figure 7
Putative alternative splicing variant of *X. laevis* MMP2 (MMP2asv). A) Nucleotide and deduced amino acid sequences of MMP2asv. The protein contains, from the N-terminus to C-terminus, a signal peptide (underlined), the conserved sequence in the propeptide involved in the "cysteine-switch" (in bold letters), a truncated catalytic domain linked to a truncated hemopexin domain (separated by double slash lines). The predicted cleavage site between the propeptide and the catalytic domain is indicated by an arrow. B) Comparison of the full length and alternatively spliced *X. laevis* MMP2 exon/intron organization. Solid blocks stand for exons present in the mRNAs and lines are introns.

Xenopus tropicalis Scaffold_119



Homo sapiens Chromosome 11




Figure 9

Comparison of the MMPs cluster on *X. tropicalis* Scaffold_119 to that on human Chromosome 11. *X. tropicalis* MMP cDNA sequences were used to do BLAST search against the *X. tropicalis* genomic sequences to locate the genes on the assembly scaffolds. MMP 1, 3, 7, 13, 18, 20, 26, as well as the novel ones MMP N2, N4 and N5 are found on Scaffold_119. They were arranged on the scaffold according to their location and orientation. The human MMPs on Chromosome 11 were arranged according to the annotations for their locations and orientations in Human Genome Build 36.3 on the NCBI website. The MMPs shown above the line for the chromosome/scaffold are MMPs specific to *X. tropicalis* or human while those shown below are the MMPs present in both species. Mb, mega base pair; kb, kilo base pairs. Note the gene size was not drawn to scale for clarity.

MMP N3

MMP N3 is one of the two genes in both *X. laevis* and *X. tropicalis* that have similar levels of homology to human MMP21 (Figs. 1 and 5). Although MMP N3 and MMP21 are located in different scaffolds in *X. tropicalis* genomic sequence, it is possible that they are located adjacent to each other in a chromosome as the genomic sequence annotation is incomplete in *X. tropicalis*. Thus, these two genes might have derived from a unique gene duplication event in amphibians.

Collagenases

Based on sequence features and its enzymatic activity, *X. laevis* MMP18 was proposed to be a novel collagenase [42]. Indeed, potential homologs of mammalian collagenases MMP1 and MMP13 have been reported for *Xenopus*. Interestingly, while the amphibian MMP13 and

human MMP13 are 67% identical, the corresponding MMP1s are only 51% identical (Table 2), similar to the levels of homology that MMP18 has with human collagenases (MMP1, 8, and 13). Furthermore, human MMP1 and MMP8 have 3 extra amino acids (RKN) at the C-terminus that are lacking in *Xenopus* MMP1 and MMP18 as well as the putative, novel *Xenopus* collagenases (MMP N4 and N5) (Fig. 2). In addition, a most likely homolog of human MMP1 has been reported for another amphibian species, *Rana catesbeiana* [40]. The *Rana* MMP1 is much shorter but much more homologous to human MMP1 (about 80%), compared to these *Xenopus* collagenases (data not shown and [42]). Furthermore, unlike the *Xenopus* MMP1, the *Rana* MMP1 clusters with human MMP1 on the phylogenetic tree (Fig. 1). Thus, it is possible that MMP1 has diverged extensively between amphibians and mammals, leading to a very different size in *Rana* or its

loss in *Xenopus*. On the other hand, all known and putative collagenases in *Xenopus* are located consecutively in a single chromosome in the order of MMP1, MMP N4, MMP N5, MMP18, and MMP13 (Fig. 9). This suggests that multiple duplication events might be responsible for the generation of these MMPs. It is interesting to note that human MMP 1, 3, 7, 8, 10, 12, 13, 20, 26 and 27 are all encoded by Chromosome 11 (Fig. 9). With the exception of four MMPs (MMP8, 10, 12 and 27) that have no apparent homologs in *X. tropicalis*, the other six MMPs found on human Chromosome 11 are clustered on the Scaffold_119 together with four frog-specific ones (MMP18, MMP N2, N4, and N5) in *X. tropicalis* (Fig. 9). Thus, *X. tropicalis* Scaffold_119 appears to contain a large region syntenic to human Chromosome 11 [71]. On the other hand, with the exception of MMP13 and MMP20, it is difficult to determine which MMP in this cluster of 10 *X. tropicalis* MMPs is the homolog of an MMP in the human cluster based on sequence homology and syntenic analyses. *X. tropicalis* MMP13 and MMP20 share high degrees of homology (about 70%) with their human counterparts, supporting that they are true homologs of the human MMPs. Consistently, phylogenetic analysis of *X. tropicalis* and human MMPs showed that these two MMPs evolved earlier than the other MMPs in the cluster (see Additional file 2). In addition, the drastic differences in the distances between MMP7 and MMP26 on the chromosome (100 mega bp in human and 100 kb in *Xenopus*) (Fig. 9) and on the phylogenetic trees (Supplemental Fig. 2), suggest that this duplication occurred after the separation of amphibians from mammals. The other six MMPs in the cluster appear to have diverged rapidly during evolution and/or evolved through duplications and/or losses independently in amphibians and mammals.

Conclusion

Through a bioinformatic approach, we have identified *Xenopus* homologs for most human MMPs. By comparing the MMPs in the two highly related species, *X. tropicalis* and *X. laevis*, we have been able to discover several unique duplications of MMPs genes in amphibians that are absent in mammals. On the other hand, several human MMPs have no apparent homologs in *Xenopus* and were possibly evolved *de novo* in mammals. Among the likely duplicated genes, genes in the following two pairs, MMP9 and MMP9TH, MMP21 and MMP N3, MMP 7 (or MMP26) and MMP N2, are more homologous to each other than to their human homologs (Fig. 1), suggesting that the duplications occurred after the separation of amphibians from mammals. On the other hand, MMP23 clusters closer to its human homolog than to the putative duplicate MMP N6 (Fig. 1), suggesting the possibility that MMP23 might have duplicated before the separation of amphibians and mammals and one copy was lost subsequently in mammals. Duplications and loss in MMP

genes are also evident when comparing the largest MMP cluster located on human Chromosome 11 with the MMP cluster on *X. tropicalis* Scaffold_119, where a number of novel MMPs were found and several MMPs were lost in *X. tropicalis*. Our findings thus demonstrate a dynamic process for MMP gene evolution. It will be of interest in the future to investigate whether MMP expression and function are conserved during vertebrate development. The sequence information reported here and the advantage of the amphibian metamorphosis for functional studies *in vivo* should facilitate such an endeavor in the near future.

Methods

We first searched the public EST database at the National Center for Biotechnology Information (NCBI, <http://www.ncbi.nlm.nih.gov/>) and the Gene Index Project in Computational Biology and Function Genomics Laboratory <http://compbio.dfci.harvard.edu/tgi/> with known *Xenopus* MMP genes for other possible MMP sequences based on sequence similarities. The identities of putative *Xenopus* MMPs were tentatively determined by building a phylogenetic tree with human MMPs. This was done through Multiple Sequence Alignment by CLUSTALW <http://align.genome.jp/>. Human MMPs used were: MMP1 (NP_002412), MMP2 (NP_004521), MMP3 (NP_002413), MMP7 (NP_002414), MMP8 (NP_002415), MMP9 (NP_004985), MMP10 (NP_002416), MMP11 (NP_005931), MMP12 (NP_002417), MMP13 (NP_002418), MMP14 (NP_004986), MMP15 (NP_002419), MMP16 (NP_005932), MMP17 (NP_057239), MMP19 (NP_002420), MMP20 (NP_004762), MMP21 (NP_671724), MMP22 (NP_990331), MMP23 (NP_008914), MMP24 (NP_006681), MMP25 (NP_071913), MMP26 (NP_068573), MMP27 (NP_071405), and MMP28 (NP_077278). The *X. laevis* MMPs were: MMP1A (BC054233), MMP1B (BC084836), MMP2 (AY037943), MMP3 (BC077966), MMP7A (AY573380), MMP7B (BC056040), MMP9 (AF072455), MMP9TH (AB288054), MMP11 (Z27093), MMP13 (L49412), MMP13A (U41824), MMP14A (AY633953), MMP14B (BC077870), MMP15 (AY573378), MMP16 (AY310397), MMP17 (CK806816), MMP18 (L76275), MMP19 (BX847184), MMP20 (DQ885892), MMP21 (U82541), MMP23 (CD302225 & CD302813), MMP24A (CA791076 & EB480268), MMP24B (EB483310), MMP25 (BC078136), MMP26 (BC056080), MMP28A (EF187277), MMP28B (BC061659), MMP N1 (Assembly of BI032306, BE509380, EC276067, BX852582, BI047339 and BG578455) and MMP N3 (Assembly of BG234242, BU905338, CB558404, and CF547511) (See Supplemental Fig. 1). Pair-wise comparison of protein sequences was conducted by using MacVector (Accelrys Inc., San Diego, CA) to further confirm the identity assignment. *X. laevis* or *tropicalis* cDNA sequences were used to

do BLAST search against the *X. tropicalis* genome assembly 4.1 <http://genome.jgi-psf.org/Xentr4/Xentr4.home.html> to determine the corresponding gene structures and predict cDNA sequences if necessary. The *X. tropicalis* MMPs thus obtained from the GenBank database were: MMP2 (NM_001015789), MMP3 (NM_001030331), MMP7 (NM_001005043), MMP9 (NM_001006842), MMP14 (NM_001030388), MMP15 (NM_001015921), MMP16 (NM_001015992), MMP17 (NM_001102999), MMP18 (NM_001030330), MMP19 (BC153750), MMP25 (CU075461), MMP26 (NM_001032335), MMP N1 (BC155487, DN028798, DN034177 and DN076875), MMP N2 (CX982585, CX982586 and CX979196), MMP N6 (CX344816 and CX344815). Other *X. tropicalis* MMP sequences were derived from predicted exons of the genomic sequences (see Additional file 1 and Table 1 for their sequences and locations in the genome).

Abbreviations

MMP: matrix metalloproteinase; ECM: extracellular matrix; TH: thyroid hormone; ST3: stromelysin 3; EST: expressed sequence tags; aa: amino acid; bp: base pair.

Authors' contributions

LF collected sequence information, performed bioinformatic analysis and wrote the first draft; BD and SM performed bioinformatic analysis and edited the manuscript; and YS supervised the research and finalized the paper. All the authors critically revised and approved the final version of the paper.

Additional material

Additional file 1

The nucleotide sequences of the X. tropicalis and X. laevis MMPs. The data presented all the all the nucleotide sequences of the X. tropicalis and X. laevis MMPs that were used for deducing Xenopus MMPs in the study. GenBank accession numbers or the scaffold of the X. tropicalis genome on which the Xenopus MMP locates were included if applicable.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-10-81-S1.doc>]

Additional file 2

Phylogenetic trees of X. tropicalis and human MMPs. X. tropicalis MMPs along with Rana catesbeiana MMP1 (RcMMP1) or human MMPs along with chicken MMP22 (CMMP22) were analyzed using the multiple sequence alignment program CLUSTALW to generate the corresponding phylogenetic trees with defined ancestral nodes marked by purple square. The MMPs located on human Chromosome 11 and those located on X. tropicalis Scaffold_119 are in red.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-10-81-S2.tiff>]

Acknowledgements

This research was supported by the Intramural Research Program of NICHD, NIH.

References

- Overall CM: **Molecular determinants of metalloproteinase substrate specificity.** *Molecular Biotechnology* 2002, **22**:51-86.
- Alexander CM, Werb Z: **Extracellular matrix degradation.** In *Cell Biology of Extracellular Matrix* 2nd edition. Edited by: Hay ED. New York: Plenum Press; 1991:255-302.
- McCawley LJ, Matrisian LM: **Matrix metalloproteinases: they're not just for matrix anymore!** *Current Opinion in Cell Biology* 2001, **13**:534-540.
- Parks WC, Mecham RP: **Matrix metalloproteinases.** New York: Academic Press; 1998.
- Nagase H: **Cell surface activation of progelatinase A (proMMP-2) and cell migration.** *Cell Res* 1998, **8**(3):179-186.
- Pei D: **Leukolysin/MMP25/MT6-MMP: a novel matrix metalloproteinase specifically expressed in the leukocyte lineage.** *Cell Res* 1999, **9**(4):291-303.
- Mott JD, Werb Z: **Regulation of matrix biology by matrix metalloproteinases.** *Current Opinion in Cell Biology* 2004, **16**:558-564.
- Uria JA, Werb Z: **Matrix metalloproteinases and their expression in mammary gland.** *Cell Res* 1998, **8**(3):187-194.
- Murphy G, Stanton H, Cowell S, Butler G, Knauper V, Atkinson S, Gavrilovic J: **Mechanisms for pro matrix metalloproteinase activation.** *Apmis* 1999, **107**(1):38-44.
- Nagase J, Suzuki K, Morodomi T, Englund JJ, Salvesen G: **Activation Mechanisms of the Precursors of Matrix Metalloproteinases 1, 2, and 3.** *Matrix Suppl* 1992, **1**:237-244.
- van Wart HE, Birkedal-Hansen H: **The cysteine switch: A principle of regulation of metalloproteinase activity with potential applicability to the entire matrix metalloproteinase gene family.** *Proc Natl Acad Sci USA* 1990, **87**:5578-5582.
- Pei D, Weiss SJ: **Furin-dependent intracellular activation of the human stromelysin-3 zymogen.** *Nature* 1995, **375**(6528):244-247.
- Sato H, Seiki M: **Membrane-Type Matrix Metalloproteinases (MT-MMPs) in Tumor Metastasis.** *J Biochem* 1996, **119**:209-215.
- Basset P, Bellocq JP, Lefebvre O, Noel A, Chenard MP, Wolf C, Anglard P, Rio MC: **Stromelysin-3: a paradigm for stroma-derived factors implicated in carcinoma progression.** *Crit Rev Oncol Hematol* 1997, **26**(1):43-53.
- Lochter A, Bissell MJ: **An odyssey from breast to bone: multi-step control of mammary metastases and osteolysis by matrix metalloproteinases.** *APMIS* 1999, **107**(1):128-136.
- MacDaougall JR, Matrisian LM: **Contributions of tumor and stromal matrix metalloproteinases to tumor progression, invasion and metastasis.** *Cancer Metastasis Rev* 1995, **14**:351-362.
- Tryggvason K, Hoyhtya M, Salo T: **Proteolytic degradation of extracellular matrix in tumor invasion.** *Biochim Biophys Acta* 1987, **907**(3):191-217.
- Matrisian LM, Hogan BL: **Growth factor-regulated proteases and extracellular matrix remodeling during mammalian development.** *Curr Top Dev Biol* 1990, **24**:219-259.
- Shi Y-B, Fu L, Hasebe T, Ishizuya-Oka A: **Regulation of ECM remodeling and cell fate determination by matrix metalloproteinase stromelysin-3 during thyroid hormone-dependent postembryonic development.** *Pharmacology & Therapeutics* 2007, **116**:391-400.
- Fu L, Hasebe T, Ishizuya-Oka A, Shi Y-B: **Roles of matrix metalloproteinases and ECM remodeling during thyroid hormone-dependent intestinal metamorphosis in Xenopus laevis.** *Organogenesis* 2007, **3**:14-19.
- Sarras MPJ, Yan L, Leontovich A, Zhang JS: **Structure, expression, and developmental function of early divergent forms of metalloproteinases in hydra.** *Cell Research* 2002, **12**:163-176.
- Sang QX: **Complex role of matrix metalloproteinases in angiogenesis.** *Cell Res* 1998, **8**(3):171-177.
- Shapiro SD: **Matrix metalloproteinase degradation of extracellular matrix: biological consequences.** *Curr Opin Cell Biol* 1998, **10**(5):602-608.

24. Page-McCaw A, Ewald AJ, Werb Z: **Matrix metalloproteinases and the regulation of tissue remodelling.** *Nat Rev Mol Cell Biol* 2007, **8(3)**:221-233.
25. Yen PM: **Physiological and molecular basis of thyroid hormone action.** *Physiol Rev* 2001, **81(3)**:1097-1142.
26. Tata JR: **Gene expression during metamorphosis: an ideal model for post-embryonic development.** *Bioessays* 1993, **15(4)**:239-248.
27. Shi Y-B: **Amphibian Metamorphosis: From morphology to molecular biology.** New York: John Wiley & Sons, Inc; 1999.
28. Amano T, Noro N, Kawabata H, Kobayashi Y, Yoshizato K: **Metamorphosis-associated and region-specific expression of calbindin gene in the posterior intestinal epithelium of *Xenopus laevis* larva.** *Dev Growth Differ* 1998, **40**:177-188.
29. Schreiber AM, Cai L, Brown DD: **Remodeling of the intestine during metamorphosis of *Xenopus laevis*.** *Proc Natl Acad Sci USA* 2005, **102(10)**:3720-3725.
30. Shi Y-B, Ishizuya-Oka A: **Thyroid hormone regulation of apoptotic tissue remodeling: Implications from molecular analysis of amphibian metamorphosis.** *Prog Nucleic Acid Res Mol Biol* 2001, **65**:53-100.
31. Shi Y-B, Ishizuya-Oka A: **Biphasic intestinal development in amphibians: Embryogenesis and remodeling during metamorphosis.** *Curr Top Dev Biol* 1996, **32**:205-235.
32. Shi Y-B, Brown DD: **The earliest changes in gene expression in tadpole intestine induced by thyroid hormone.** *J Biol Chem* 1993, **268**:20312-20317.
33. Patterson D, Hayes W, Shi YB: **Transcriptional activation of the matrix metalloproteinase gene stromelysin-3 coincides with thyroid hormone-induced cell death during frog metamorphosis.** *Dev Biol* 1995, **167(1)**:252-262.
34. Ishizuya-Oka A, Ueda S, Shi Y-B: **Transient expression of stromelysin-3 mRNA in the amphibian small intestine during metamorphosis.** *Cell Tissue Res* 1996, **283(2)**:325-329.
35. Damjanovski S, Ishizuya-Oka A, Shi YB: **Spatial and temporal regulation of collagenases-3, -4, and stromelysin - 3 implicates distinct functions in apoptosis and tissue remodeling during frog metamorphosis.** *Cell Res* 1999, **9(2)**:91-105.
36. Berry DL, Schwartzman RA, Brown DD: **The expression pattern of thyroid hormone response genes in the tadpole tail identifies multiple resorption programs.** *Dev Biol* 1998, **203**:12-23.
37. Berry DL, Rose CS, Remo BF, Brown DD: **The expression pattern of thyroid hormone response genes in remodeling tadpole tissues defines distinct growth and resorption gene expression programs.** *Dev Biol* 1998, **203**:24-35.
38. Shi YB, Fu L, Hsia SC, Tomita A, Buchholz D: **Thyroid hormone regulation of apoptotic tissue remodeling during anuran metamorphosis.** *Cell Res* 2001, **11(4)**:245-252.
39. Wang Z, Brown DD: **Thyroid hormone-induced gene expression program for amphibian tail resorption.** *J Biol Chem* 1993, **268**:16270-16278.
40. Oofusa K, Yomori S, Yoshizato K: **Regionally and hormonally regulated expression of genes of collagen and collagenase in the anuran larval skin.** *Int J Dev Biol* 1994, **38(2)**:345-350.
41. Jung J-C, Leco KJ, Edwards DR, Fini ME: **Matrix metalloproteinase mediate the dismantling of mesenchymal structures in the tadpole tail during thyroid hormone-induced tail resorption.** *Dev Dyn* 2002, **223**:402-413.
42. Stelow MA, Bauzon DD, Li J, Sedgwick T, Liang VC, Sang QA, Shi YB: **Identification and characterization of a novel collagenase in *Xenopus laevis*: possible roles during frog development.** *Mol Biol Cell* 1996, **7(10)**:1471-1483.
43. Fujimoto K, Nakajima K, Yaoita Y: **One of the duplicated matrix metalloproteinase-9 genes is expressed in regressing tail during anuran metamorphosis.** *Dev Growth Differ* 2006, **48(4)**:223-241.
44. Fujimoto K, Nakajima K, Yaoita Y: **Expression of matrix metalloproteinase genes in regressing or remodeling organs during amphibian metamorphosis.** *Dev Growth Differ* 2007, **49(2)**:131-143.
45. Hasebe T, Kajita M, Fujimoto K, Yaoita Y, Ishizuya-Oka A: **Expression profiles of the duplicated matrix metalloproteinase-9 genes suggest their different roles in apoptosis of larval intestinal epithelial cells during *Xenopus laevis* metamorphosis.** *Dev Dyn* 2007, **236(8)**:2338-2345.
46. Hasebe T, Hartman R, Matsuda H, Shi YB: **Spatial and temporal expression profiles suggest the involvement of gelatinase A and membrane type I matrix metalloproteinase in amphibian metamorphosis.** *Cell Tissue Res* 2006, **324**:105-116.
47. Hammoud L, Walsh LA, Damjanovski S: **Cloning and developmental characterization of *Xenopus laevis* membrane type-3 matrix metalloproteinase (MT3-MMP).** *Biochem Cell Biol* 2006, **84**:167-177.
48. Walsh LA, Carere DA, Cooper CA, Damjanovski S: **Membrane type-I matrix metalloproteinases and tissue inhibitor of metalloproteinases-2 RNA levels mimic each other during *Xenopus laevis* metamorphosis.** *PLoS ONE* 2007, **2(10)**:e1000.
49. Fu L, Tomita A, Wang H, Buchholz DR, Shi Y-B: **Transcriptional regulation of the *Xenopus laevis* stromelysin-3 gene by thyroid hormone is mediated by a DNA element in the first intron.** *J Biol Chem* 2006, **281**:16870-16878.
50. Oofusa K, Yoshizato K: **Thyroid hormone-dependent expression of bullfrog tadpole collagenase gene.** *Roux's Arch Dev Biol* 1996, **205**:241-251.
51. Damjanovski S, Amano T, Li Q, Pei D, Shi Y-B: **Overexpression of Matrix Metalloproteinases Leads to Lethality in Transgenic *Xenopus Laevis*: Implications for Tissue-Dependent Functions of Matrix Metalloproteinases during Late Embryonic development.** *Dev Dyn* 2001, **221(1)**:37-47.
52. Ishizuya-Oka A, Li Q, Amano T, Damjanovski S, Ueda S, Shi Y-B: **Requirement for matrix metalloproteinase stromelysin-3 in cell migration and apoptosis during tissue remodeling in *Xenopus laevis*.** *J Cell Biol* 2000, **150(5)**:1177-1188.
53. Fu L, Ishizuya-Oka A, Buchholz DR, Amano T, Matsuda H, Shi YB: **A causative role of stromelysin-3 in extracellular matrix remodeling and epithelial apoptosis during intestinal metamorphosis in *Xenopus laevis*.** *J Biol Chem* 2005, **280(30)**:27856-27865.
54. Hasebe T, Fu L, Amano T, Shi Y-B: **Evidence for a cooperative role of gelatinase A and membrane type-I matrix metalloproteinase during *Xenopus laevis* development.** *Mechanisms of Development* 2007, **124**:11-22.
55. Shintani S, Kobata M, Kamakura N, Toyosawaa S, Ooshima T: **Identification and characterization of matrix metalloproteinase-20 (MMP20; enamelysin) genes in reptile and amphibian.** *Gene* 2007, **392**:89-97.
56. Harrison M, Abu-Elmagd M, Grocott T, Yates C, Gavrilovic J, Wheeler GN: **Matrix metalloproteinase genes in *Xenopus* development.** *Dev Dyn* 2004, **231(1)**:214-220.
57. Yang M, Kurkinen M: **Cloning and characterization of a novel matrix metalloproteinase (MMP), CMMP, from chicken embryo fibroblasts.** *J Biol Chem* 1998, **273**:17893-17900.
58. Tomlinson ML, Garcia-Morales C, Abu-Elmagd M, Wheeler GN: **Three matrix metalloproteinases are required in vivo for macrophage migration during embryonic development.** *Mech Dev* 2008, **125**:1059-1070.
59. Saus J, Quinones S, Otani Y, Nagase H, Harris ED Jr, Kurkinen M: **The complete primary structure of human matrix metalloproteinase-3. Identity with stromelysin.** *J Biol Chem* 1988, **263(14)**:6742-6745.
60. Whitham SE, Murphy G, Angel P, Rahmsdorf H-J, Smith BJ, Lyons A, Harris TJR, Reynolds JJ, Herrlich P, Docherty AJP: **Comparison of human stromelysin and collagenase by cloning and sequence analysis.** *Biochem J* 1986, **240(3)**:913-916.
61. Murphy GJP, Murphy G, Reynolds JJ: **The origin of matrix metalloproteinases and their familial relations.** *FEBS Letters* 1991, **289**:4-7.
62. Muller D, Quantin B, Gesnel MC, Millon-Collard R, Abecassis J, Breathnach R: **The collagenase gene family in humans consists of at least four members.** *Biochem J* 1988, **253(1)**:187-192.
63. Pei D: **CA-MMP: a matrix metalloproteinase with a novel cysteine array, but without the classic cysteine switch.** *FEBS Letters* 1999, **457**:262-270.
64. Velasco G, Pendas AM, Fueyo A, Knäuper V, Murphy G, López-Otín C: **Cloning and characterization of human MMP23, a new matrix metalloproteinase predominantly expressed in reproductive tissues and lacking conserved domains in other family members.** *J Biol Chem* 1999, **274**:4570-4576.
65. Gururajan R, Grenet J, Lahti JM, Kidd VJ: **Isolation and characterization of two novel metalloproteinase genes linked to the**

- Cdc2L locus on human chromosome 1p36.3.** *Genomics* 1998, **52**:101-106.
66. Ohnishi J, Onishi E, Jin M, Hirano W, Nakane D, Matsui H, Kimura A, Sawa H, Nakayama K, Shibuya H, et al.: **Cloning and characterization of a rat ortholog of MMP23 (Matrix metalloproteinase-23), a unique type of membrane-anchored matrix metalloproteinase and conditioned switching of its expression during the ovarian follicular development.** *Mol Endocrinol* 2001, **15**:747-764.
 67. Yang M, Murray MT, Kurkinen M: **A novel matrix metalloproteinase gene (XMMP) encoding vitronectin-like motif is transiently expressed in Xenopus laevis early embryo development.** *J Biol Chem* 1997, **272**:13527-13533.
 68. Park HI, Ni J, Gerkema FE, Liu D, Belozero V, Sang Q-XA: **Identification and characterization of human endometase (Matrix Metalloproteinase-26) from endometrial tumor.** *J Biol Chem* 2000, **275**:20540-20544.
 69. de Coignac AB, Elson G, Delneste Y, Magistrelli G, Jeannin P, Aubry J-P, Berthier O, Schmitt D, Bonnefoy J-Y, Gauchat J-F: **Cloning of MMP-26: A novel matrilysin-like proteinase.** *Eur J Biochem* 2000, **267**:3323-3329.
 70. Marchenko GN, Ratnikov BI, Rozanov DV, Godzik A, Deryugina EI, Strongin AY: **Characterization of matrix metalloproteinase-26, a novel metalloproteinase widely expressed in cancer cells of epithelial origin.** *Biochem J* 2001, **356**:705-718.
 71. Loots GG, Ovcharenko I: **ECRbase: Database of Evolutionary Conserved Regions, Promoters, and Transcription Factor Binding Sites in Vertebrate Genomes.** *Bioinformatics* 2007, **23**:122-124.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

