

4-Ethoxyanilinium hexafluorophosphate monohydrate

Xue-qun Fu

Ordered Matter Science Research Center, Southeast University, Nanjing 210096, People's Republic of China
Correspondence e-mail: fuxuequn222@163.com


Received 12 May 2010; accepted 18 May 2010

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; disorder in main residue; R factor = 0.059; wR factor = 0.164; data-to-parameter ratio = 12.5.

In the crystal of the title compound, $\text{C}_8\text{H}_{12}\text{NO}^+\cdot\text{PF}_6^-\cdot\text{H}_2\text{O}$, intermolecular $\text{N}-\text{H}\cdots\text{F}$, $\text{N}-\text{H}\cdots\text{O}$ and $\text{O}-\text{H}\cdots\text{F}$ hydrogen bonds link the molecules into chains along the c axis and $\text{C}-\text{H}\cdots\pi$ contacts further stabilize the structure. The F atoms of one of the hexafluorophosphate anions are disordered over two sets of sites with site-occupancy factors of 0.27 (3) and 0.73 (3).

Related literature

For related structures, see: Fu (2009a,b). The title compound was studied as part of our search for ferroelectric compounds, which usually have a phase transition. For background to phase transition materials, see: Li *et al.* (2008); Zhang *et al.* (2009).



Experimental

Crystal data

$\text{C}_8\text{H}_{12}\text{NO}^+\cdot\text{PF}_6^-\cdot\text{H}_2\text{O}$	$V = 1232.4(4)\text{ \AA}^3$
$M_r = 301.17$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 17.498(4)\text{ \AA}$	$\mu = 0.29\text{ mm}^{-1}$
$b = 5.1236(10)\text{ \AA}$	$T = 298\text{ K}$
$c = 14.793(3)\text{ \AA}$	$0.4 \times 0.3 \times 0.2\text{ mm}$
$\beta = 111.68(3)^\circ$	

Data collection

Rigaku SCXmini diffractometer
Absorption correction: multi-scan
(*CrystalClear*; Rigaku, 2005)
 $T_{\min} = 0.900$, $T_{\max} = 0.943$

12124 measured reflections
2820 independent reflections
1890 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.054$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.059$
 $wR(F^2) = 0.164$
 $S = 1.05$
2820 reflections
226 parameters
235 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.45\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.48\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$Cg1$ is the centroid of the C1–C6 ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1A \cdots F2'	0.89	2.16	3.03 (2)	167
N1—H1A \cdots F2	0.89	2.17	3.049 (10)	169
N1—H1B \cdots F1 ⁱ	0.89	2.22	3.066 (8)	158
N1—H1B \cdots F1 ⁱⁱ	0.89	2.28	3.09 (3)	152
N1—H1C \cdots O1W ⁱⁱ	0.89	2.22	2.883 (3)	131
N1—H1C \cdots F3 ⁱⁱ	0.89	2.46	3.137 (12)	133
N1—H1C \cdots F3 ⁱⁱⁱ	0.89	2.40	3.01 (3)	126
N1—H1C \cdots F6 ⁱⁱ	0.89	2.51	2.96 (2)	112
O1W—H1WB \cdots F3 ⁱⁱⁱ	0.82 (1)	2.20 (3)	2.97 (3)	156 (4)
O1W—H1WB \cdots F3 ⁱⁱ	0.82 (1)	2.28 (3)	3.030 (12)	152 (4)
O1W—H1WA \cdots F4 ⁱ	0.82 (3)	2.24 (2)	3.040 (12)	164 (4)
O1W—H1WA \cdots F4 ⁱⁱ	0.82 (3)	2.11 (3)	2.88 (2)	155 (4)
O1W—H1WA \cdots F5 ⁱ	0.82 (3)	2.48 (4)	2.951 (16)	117 (3)
C8—H8C \cdots Cg1 ^{iv}	0.96	3.16	4.023 (5)	150

Symmetry codes: (i) $x, -y + \frac{3}{2}, z + \frac{1}{2}$; (ii) $x, y - 1, z$; (iii) $-x, y - \frac{1}{2}, -z + \frac{1}{2}$; (iv) $-x + 1, -y, -z + 1$.

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *PRPKAPPA* (Ferguson, 1999).

The author is grateful to the starter fund of Southeast University for financial support to buy the X-ray diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZQ2040).

References

- Ferguson, G. (1999). *PRPKAPPA*. University of Guelph, Canada.
- Fu, X. (2009a). *Acta Cryst. E65*, o2345.
- Fu, X. (2009b). *Acta Cryst. E65*, o2520.
- Li, X. Z., Qu, Z. R. & Xiong, R. G. (2008). *Chin. J. Chem.*, **11**, 1959–1962.
- Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Zhang, W., Chen, L. Z., Xiong, R. G., Nakamura, T. & Huang, S. D. (2009). *J. Am. Chem. Soc.*, **131**, 12544–12545.

supplementary materials

Acta Cryst. (2010). E66, o1461 [doi:10.1107/S1600536810018404]

4-Ethoxyanilinium hexafluorophosphate monohydrate

X. Fu

Comment

As a continuation of our study of dielectric-ferroelectric materials, including organic ligands (Li *et al.*, 2008), metal-organic coordination compounds (Zhang *et al.*, 2009), and organic-inorganic hybrids, we studied the dielectric properties of the title compound. Unfortunately, there was no distinct anomaly observed from 93 to 350 K. In this article, the crystal structure of this compound is presented. The crystal structures of 4-ethoxyanilinium together with other anions are known (Fu, 2009a,b).

The asymmetric unit of the crystal structure consists of one almost coplanar protonated 4-ethoxyanilinium cation with the C2—C1—O1—C7 and C1—O1—C7—C8 torsion angles of 172.9 (3) and 177.4 (3) $^{\circ}$, respectively, one hexafluorophosphate anion for which the F atoms are disordered over two sets of positions with site-occupancy factors of 0.27 (3) and 0.73 (3), and one water molecule (Fig. 1). In the crystal structure, several intermolecular N—H \cdots F and O—H \cdots F hydrogen bonds link all species to chains along the c axis (Fig. 2). In addition, C—H \cdots π interactions further stabilize the crystal structure.

Experimental

1.37 g (10 mmol) of 4-Ethoxybenzenamine was firstly dissolved in 50 ml ethanol, to which hexafluorophosphoric acid (70%, w/w) was then added until the solution becomes acidic under stirring. Single crystals of the title compound were prepared by slow evaporation at room temperature of the acidic solution after 3 days.

Refinement

The water H atoms were found in Fourier difference maps and were refined freely. Positional parameters of all other H atoms were calculated geometrically and allowed to ride on the C and N atoms to which they are bonded, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$. The F atoms of the hexafluorophosphate anion are disordered in two orientations with site-occupancy factors of 0.73 (3) and 0.27 (3).

Figures




Fig. 1. The molecular structure of the title compound with the atomic numbering scheme. Displacement ellipsoids are drawn at the 30% probability level, and all H atoms have been omitted for clarity.

supplementary materials




Fig. 2. A view of the packing of the title compound, stacking along the c axis. Dashed lines indicate hydrogen bonds.

4-Ethoxyanilinium hexafluorophosphate monohydrate

Crystal data

$C_8H_{12}NO^+\cdot PF_6^- \cdot H_2O$	$F(000) = 616$
$M_r = 301.17$	$D_x = 1.623 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 4774 reflections
$a = 17.498 (4) \text{ \AA}$	$\theta = 3.1\text{--}27.7^\circ$
$b = 5.1236 (10) \text{ \AA}$	$\mu = 0.29 \text{ mm}^{-1}$
$c = 14.793 (3) \text{ \AA}$	$T = 298 \text{ K}$
$\beta = 111.68 (3)^\circ$	Prism, colourless
$V = 1232.4 (4) \text{ \AA}^3$	$0.4 \times 0.3 \times 0.2 \text{ mm}$
$Z = 4$	

Data collection

Rigaku SCXmini diffractometer	2820 independent reflections
Radiation source: fine-focus sealed tube graphite	1890 reflections with $I > 2\sigma(I)$
Detector resolution: 13.6612 pixels mm^{-1}	$R_{\text{int}} = 0.054$
ω scans	$\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 3.1^\circ$
Absorption correction: multi-scan (<i>CrystalClear</i> ; Rigaku, 2005)	$h = -22 \rightarrow 22$
$T_{\text{min}} = 0.900, T_{\text{max}} = 0.943$	$k = -6 \rightarrow 6$
12124 measured reflections	$l = -19 \rightarrow 19$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.059$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.164$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.05$	$w = 1/[\sigma^2(F_o^2) + (0.0764P)^2 + 0.6523P]$
2820 reflections	where $P = (F_o^2 + 2F_c^2)/3$
	$(\Delta/\sigma)_{\text{max}} = 0.003$

226 parameters	$\Delta\rho_{\max} = 0.45 \text{ e } \text{\AA}^{-3}$
235 restraints	$\Delta\rho_{\min} = -0.48 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
N1	0.15275 (14)	0.3564 (5)	0.42609 (17)	0.0414 (6)	
H1A	0.1427	0.4860	0.3832	0.050*	
H1B	0.1304	0.3936	0.4697	0.050*	
H1C	0.1311	0.2092	0.3953	0.050*	
O1	0.49349 (12)	0.2376 (5)	0.60339 (18)	0.0624 (7)	
C1	0.40946 (18)	0.2556 (6)	0.5644 (2)	0.0458 (7)	
C4	0.24232 (16)	0.3242 (5)	0.47587 (19)	0.0366 (6)	
C3	0.29478 (19)	0.4801 (6)	0.4497 (2)	0.0515 (8)	
H3A	0.2739	0.6074	0.4021	0.062*	
C5	0.27196 (18)	0.1368 (7)	0.5450 (2)	0.0497 (8)	
H5A	0.2360	0.0333	0.5625	0.060*	
C6	0.35635 (19)	0.1007 (7)	0.5895 (2)	0.0558 (9)	
H6A	0.3769	-0.0286	0.6364	0.067*	
C2	0.37808 (19)	0.4465 (7)	0.4943 (3)	0.0561 (9)	
H2A	0.4137	0.5527	0.4773	0.067*	
C8	0.6217 (2)	0.0478 (10)	0.6954 (3)	0.0783 (12)	
H8A	0.6483	-0.0935	0.7380	0.117*	
H8B	0.6405	0.2108	0.7279	0.117*	
H8C	0.6347	0.0396	0.6378	0.117*	
C7	0.5304 (2)	0.0266 (8)	0.6678 (3)	0.0632 (10)	
H7A	0.5107	-0.1390	0.6359	0.076*	
H7B	0.5164	0.0372	0.7254	0.076*	
O1W	0.08195 (12)	0.8613 (5)	0.44539 (16)	0.0473 (5)	
H1WB	0.0467 (16)	0.843 (9)	0.3908 (11)	0.086 (15)*	
H1WA	0.071 (2)	0.856 (9)	0.4948 (14)	0.086 (15)*	
P1	0.11668 (4)	0.96937 (15)	0.18642 (5)	0.0396 (3)	
F3	0.0519 (7)	1.103 (3)	0.2240 (8)	0.064 (2)	0.73 (3)
F5	0.1765 (9)	0.833 (3)	0.1430 (10)	0.065 (2)	0.73 (3)
F4	0.0451 (7)	0.782 (2)	0.1246 (9)	0.072 (2)	0.73 (3)
F1	0.0911 (6)	1.1714 (17)	0.0948 (6)	0.0618 (15)	0.73 (3)

supplementary materials

F2	0.1410 (8)	0.783 (2)	0.2777 (7)	0.086 (2)	0.73 (3)
F6	0.1869 (6)	1.169 (2)	0.2477 (7)	0.0704 (18)	0.73 (3)
F6'	0.1908 (13)	1.081 (6)	0.2715 (16)	0.066 (4)	0.27 (3)
F4'	0.0396 (15)	0.830 (6)	0.1040 (17)	0.053 (4)	0.27 (3)
F2'	0.1153 (17)	0.730 (5)	0.256 (2)	0.070 (4)	0.27 (3)
F3'	0.0594 (18)	1.160 (7)	0.2229 (19)	0.056 (4)	0.27 (3)
F5'	0.184 (3)	0.796 (8)	0.164 (3)	0.062 (4)	0.27 (3)
F1'	0.1166 (18)	1.168 (5)	0.1154 (19)	0.075 (4)	0.27 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0383 (13)	0.0429 (13)	0.0428 (13)	-0.0011 (11)	0.0147 (11)	-0.0042 (11)
O1	0.0333 (11)	0.0710 (15)	0.0758 (16)	0.0024 (11)	0.0119 (11)	0.0209 (13)
C1	0.0335 (14)	0.0500 (17)	0.0516 (17)	0.0022 (13)	0.0128 (13)	0.0029 (14)
C4	0.0337 (14)	0.0378 (15)	0.0383 (14)	-0.0002 (12)	0.0132 (12)	-0.0041 (12)
C3	0.0425 (16)	0.0509 (18)	0.0603 (19)	0.0031 (14)	0.0181 (15)	0.0191 (16)
C5	0.0406 (17)	0.0564 (19)	0.0508 (17)	-0.0050 (15)	0.0153 (14)	0.0124 (15)
C6	0.0443 (18)	0.060 (2)	0.0577 (19)	0.0035 (16)	0.0128 (15)	0.0216 (17)
C2	0.0425 (17)	0.0553 (19)	0.073 (2)	-0.0024 (15)	0.0237 (16)	0.0172 (17)
C8	0.045 (2)	0.108 (3)	0.071 (3)	0.018 (2)	0.0099 (19)	0.020 (2)
C7	0.0450 (18)	0.073 (2)	0.064 (2)	0.0094 (18)	0.0113 (16)	0.0147 (19)
O1W	0.0395 (12)	0.0532 (13)	0.0448 (13)	-0.0011 (10)	0.0105 (11)	0.0014 (11)
P1	0.0366 (4)	0.0479 (5)	0.0323 (4)	0.0065 (3)	0.0103 (3)	-0.0011 (3)
F3	0.055 (3)	0.082 (6)	0.065 (3)	0.018 (3)	0.034 (2)	-0.005 (3)
F5	0.055 (3)	0.086 (4)	0.054 (5)	0.022 (2)	0.019 (4)	-0.012 (3)
F4	0.068 (3)	0.074 (4)	0.071 (5)	-0.032 (3)	0.022 (3)	-0.011 (3)
F1	0.054 (4)	0.073 (2)	0.054 (3)	0.014 (3)	0.014 (2)	0.026 (2)
F2	0.092 (6)	0.101 (5)	0.054 (3)	0.027 (4)	0.015 (3)	0.036 (3)
F6	0.057 (2)	0.078 (4)	0.066 (4)	-0.013 (3)	0.011 (2)	-0.025 (3)
F6'	0.042 (4)	0.089 (9)	0.054 (7)	0.011 (6)	0.004 (5)	-0.032 (6)
F4'	0.044 (5)	0.075 (8)	0.040 (6)	-0.004 (5)	0.015 (4)	-0.013 (6)
F2'	0.074 (10)	0.075 (7)	0.060 (8)	0.017 (6)	0.025 (6)	0.029 (6)
F3'	0.048 (6)	0.066 (9)	0.042 (6)	0.022 (6)	0.003 (5)	-0.007 (5)
F5'	0.050 (6)	0.087 (10)	0.049 (10)	0.018 (6)	0.018 (7)	-0.017 (7)
F1'	0.065 (10)	0.081 (6)	0.066 (7)	-0.007 (7)	0.010 (6)	0.035 (6)

Geometric parameters (\AA , $^\circ$)

N1—C4	1.474 (4)	C8—H8B	0.9600
N1—H1A	0.8900	C8—H8C	0.9600
N1—H1B	0.8900	C7—H7A	0.9700
N1—H1C	0.8900	C7—H7B	0.9700
O1—C1	1.370 (4)	O1W—H1WB	0.821 (10)
O1—C7	1.428 (4)	O1W—H1WA	0.82 (3)
C1—C6	1.373 (4)	P1—F1'	1.46 (2)
C1—C2	1.384 (4)	P1—F6'	1.546 (18)
C4—C5	1.358 (4)	P1—F4	1.575 (10)
C4—C3	1.376 (4)	P1—F5	1.578 (13)

C3—C2	1.371 (4)	P1—F2	1.580 (8)
C3—H3A	0.9300	P1—F3	1.590 (11)
C5—C6	1.389 (4)	P1—F6	1.598 (7)
C5—H5A	0.9300	P1—F5'	1.60 (4)
C6—H6A	0.9300	P1—F2'	1.61 (2)
C2—H2A	0.9300	P1—F4'	1.61 (3)
C8—C7	1.500 (5)	P1—F3'	1.63 (3)
C8—H8A	0.9600	P1—F1	1.632 (7)
C4—N1—H1A	109.5	F6'—P1—F3	92.8 (9)
C4—N1—H1B	109.4	F4—P1—F3	87.1 (6)
H1A—N1—H1B	109.5	F5—P1—F3	176.5 (7)
C4—N1—H1C	109.5	F2—P1—F3	87.9 (6)
H1A—N1—H1C	109.5	F1'—P1—F6	76.1 (9)
H1B—N1—H1C	109.5	F4—P1—F6	177.8 (5)
C1—O1—C7	118.8 (3)	F5—P1—F6	91.6 (8)
O1—C1—C6	125.2 (3)	F2—P1—F6	89.7 (3)
O1—C1—C2	115.5 (3)	F3—P1—F6	91.4 (6)
C6—C1—C2	119.4 (3)	F1'—P1—F5'	92.5 (15)
C5—C4—C3	120.9 (3)	F6'—P1—F5'	85.2 (18)
C5—C4—N1	119.7 (3)	F4—P1—F5'	91.5 (18)
C3—C4—N1	119.4 (3)	F2—P1—F5'	81.6 (13)
C2—C3—C4	119.5 (3)	F3—P1—F5'	169.4 (13)
C2—C3—H3A	120.2	F6—P1—F5'	90.3 (18)
C4—C3—H3A	120.2	F1'—P1—F2'	174.5 (15)
C4—C5—C6	119.6 (3)	F6'—P1—F2'	88.1 (9)
C4—C5—H5A	120.2	F4—P1—F2'	73.3 (9)
C6—C5—H5A	120.2	F5—P1—F2'	95.5 (12)
C1—C6—C5	120.1 (3)	F3—P1—F2'	85.4 (12)
C1—C6—H6A	119.9	F6—P1—F2'	108.2 (8)
C5—C6—H6A	119.9	F5'—P1—F2'	84.1 (17)
C3—C2—C1	120.4 (3)	F1'—P1—F4'	89.3 (11)
C3—C2—H2A	119.8	F6'—P1—F4'	174.7 (12)
C1—C2—H2A	119.8	F5—P1—F4'	89.5 (10)
C7—C8—H8A	109.5	F2—P1—F4'	105.3 (9)
C7—C8—H8B	109.5	F3—P1—F4'	87.1 (10)
H8A—C8—H8B	109.5	F6—P1—F4'	164.9 (9)
C7—C8—H8C	109.5	F5'—P1—F4'	94.0 (19)
H8A—C8—H8C	109.5	F2'—P1—F4'	86.7 (10)
H8B—C8—H8C	109.5	F1'—P1—F3'	89.0 (15)
O1—C7—C8	107.4 (3)	F6'—P1—F3'	86.7 (13)
O1—C7—H7A	110.2	F4—P1—F3'	96.0 (12)
C8—C7—H7A	110.2	F5—P1—F3'	169.1 (14)
O1—C7—H7B	110.2	F2—P1—F3'	94.9 (12)
C8—C7—H7B	110.2	F6—P1—F3'	82.3 (12)
H7A—C7—H7B	108.5	F5'—P1—F3'	171.8 (19)
H1WB—O1W—H1WA	122 (2)	F2'—P1—F3'	94.9 (17)
F1'—P1—F6'	96.0 (11)	F4'—P1—F3'	94.1 (14)
F1'—P1—F4	102.6 (9)	F6'—P1—F1	109.3 (11)
F6'—P1—F4	161.3 (10)	F4—P1—F1	89.4 (4)

supplementary materials

F1'—P1—F5	80.8 (12)	F5—P1—F1	87.8 (5)
F6'—P1—F5	90.7 (10)	F2—P1—F1	177.8 (5)
F4—P1—F5	89.9 (8)	F3—P1—F1	90.3 (5)
F1'—P1—F2	164.6 (8)	F6—P1—F1	89.0 (3)
F6'—P1—F2	69.4 (10)	F5'—P1—F1	100.2 (13)
F4—P1—F2	91.9 (4)	F2'—P1—F1	162.3 (8)
F5—P1—F2	94.0 (6)	F4'—P1—F1	75.9 (10)
F1'—P1—F3	98.1 (12)	F3'—P1—F1	83.1 (11)
C7—O1—C1—C6	6.9 (5)	C2—C1—C6—C5	-0.3 (5)
C7—O1—C1—C2	-172.9 (3)	C4—C5—C6—C1	0.7 (5)
C5—C4—C3—C2	-0.2 (5)	C4—C3—C2—C1	0.7 (5)
N1—C4—C3—C2	-178.5 (3)	O1—C1—C2—C3	179.3 (3)
C3—C4—C5—C6	-0.5 (5)	C6—C1—C2—C3	-0.4 (5)
N1—C4—C5—C6	177.8 (3)	C1—O1—C7—C8	177.4 (3)
O1—C1—C6—C5	180.0 (3)		


Hydrogen-bond geometry (\AA , $^\circ$)

Cg1 is the centroid of the C1—C6 ring.

$D\cdots H$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
N1—H1A…F2'	0.89	2.16	3.03 (2)	167.
N1—H1A…F2	0.89	2.17	3.049 (10)	169.
N1—H1B…F1 ⁱ	0.89	2.22	3.066 (8)	158.
N1—H1B…F1 ⁱ	0.89	2.28	3.09 (3)	152.
N1—H1C…O1W ⁱⁱ	0.89	2.22	2.883 (3)	131.
N1—H1C…F3 ⁱⁱ	0.89	2.46	3.137 (12)	133.
N1—H1C…F3 ⁱⁱⁱ	0.89	2.40	3.01 (3)	126.
N1—H1C…F6 ⁱⁱⁱ	0.89	2.51	2.96 (2)	112.
O1W—H1WB…F3 ⁱⁱⁱ	0.82 (1)	2.20 (3)	2.97 (3)	156 (4)
O1W—H1WB…F3 ⁱⁱⁱ	0.82 (1)	2.28 (3)	3.030 (12)	152 (4)
O1W—H1WA…F4 ⁱ	0.82 (3)	2.24 (2)	3.040 (12)	164 (4)
O1W—H1WA…F4 ⁱ	0.82 (3)	2.11 (3)	2.88 (2)	155 (4)
O1W—H1WA…F5 ⁱ	0.82 (3)	2.48 (4)	2.951 (16)	117 (3)
C8—H8C…Cg1 ^{iv}	0.96	3.16	4.023 (5)	150.

Symmetry codes: (i) $x, -y+3/2, z+1/2$; (ii) $x, y-1, z$; (iii) $-x, y-1/2, -z+1/2$; (iv) $-x+1, -y, -z+1$.

Fig. 1



supplementary materials

Fig. 2

