

RESEARCH ARTICLE

Environmental DNA sampling reveals high occupancy rates of invasive Burmese pythons at wading bird breeding aggregations in the central Everglades

Sophia C. M. Orzechowski^{1*}, Peter C. Frederick¹, Robert M. Dorazio^{2,3}, Margaret E. Hunter²

1 Department of Wildlife Ecology and Conservation, University of Florida, Gainesville, Florida, United States of America, **2** U.S. Geological Survey, Wetland and Aquatic Research Center, Gainesville, Florida, United States of America, **3** Department of Biology, San Francisco State University, San Francisco, California, United States of America

* sorzechowski@ufl.edu



OPEN ACCESS

Citation: Orzechowski SCM, Frederick PC, Dorazio RM, Hunter ME (2019) Environmental DNA sampling reveals high occupancy rates of invasive Burmese pythons at wading bird breeding aggregations in the central Everglades. PLoS ONE 14(4): e0213943. <https://doi.org/10.1371/journal.pone.0213943>

Editor: Christopher M. Somers, University of Regina, CANADA

Received: December 19, 2018

Accepted: March 4, 2019

Published: April 10, 2019

Copyright: This is an open access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the [Creative Commons CC0](https://creativecommons.org/licenses/by/4.0/) public domain dedication.

Data Availability Statement: All ddPCR and covariate data files needed to replicate our analyses will be available in the public USGS-hosted ScienceBase website ([sciencebase.gov](https://www.sciencebase.gov)). The accession numbers and DOI will be available after acceptance.

Funding: The authors received funding from the Florida Fish and Wildlife Conservation Commission (myfwc.com; award # 13416-TA-3020). Publication of this article was funded in part by the

Abstract

The Burmese python (*Python bivittatus*) is now established as a breeding population throughout south Florida, USA. However, the extent of the invasion, and the ecological impacts of this novel apex predator on animal communities are incompletely known, in large part because Burmese pythons (hereafter “pythons”) are extremely cryptic and there has been no efficient way to detect them. Pythons are recently confirmed nest predators of long-legged wading bird breeding colonies (orders Ciconiiformes and Pelecaniformes). Pythons can consume large quantities of prey and may not be recognized as predators by wading birds, therefore they could be a particular threat to colonies. To quantify python occupancy rates at tree islands where wading birds breed, we utilized environmental DNA (eDNA) analysis—a genetic tool which detects shed DNA in water samples and provides high detection probabilities. We fitted multi-scale Bayesian occupancy models to test the prediction that pythons occupy islands with wading bird colonies at higher rates compared to representative control islands containing no breeding birds. Our results suggest that pythons are widely distributed across the central Everglades in proximity to active wading bird colonies. In support of our prediction that pythons are attracted to colonies, site-level python eDNA occupancy rates were higher at wading bird colonies ($\psi = 0.88$, 95% credible interval [0.59–1.00]) than at the control islands ($\psi = 0.42$ [0.16–0.80]) in April through June ($n = 15$ colony-control pairs). We found our water temperature proxy (time of day) to be informative of detection probability, in accordance with other studies demonstrating an effect of temperature on eDNA degradation in occupied samples. Individual sample concentrations ranged from 0.26 to 38.29 copies/ μL and we generally detected higher concentrations of python eDNA in colony sites. Continued monitoring of wading bird colonies is warranted to determine the effect pythons are having on populations and investigate putative management activities.

University of Florida Open Access Publishing Fund. The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: I have read the journal's policy and the authors of this manuscript have the following competing interests: Coauthor MEH is an Academic Editor at PLOS ONE. We confirm that this does not alter our adherence to PLOS ONE policies on sharing data and materials.

Introduction

Highly secretive or cryptic invasive species present unique monitoring challenges. Effective detection methodologies are necessary to successfully manage elusive invasive species, whether populations are at low densities or well-established [1, 2]. These detection methods must also allow researchers to broadly survey the landscape because of the possibility of multiple and often unknown introduction points [3, 4]. Traditional survey methods, such as traps or visual and acoustic signs, can be inadequate and result in false negatives and ill-informed management decisions [5, 6].

Environmental DNA (eDNA) monitoring has emerged as an effective genetic tool to monitor biodiversity and detect cryptic or low-density species when traditional methods fail [7–12]. The use of eDNA as an indicator of species presence is predicated on the fact that organisms shed DNA into the environments they inhabit. Environmental DNA monitoring has been used to determine range limits, track invasion fronts, detect invasive species prior to confirmed visual sightings, and in some cases to obtain relative estimates of biomass and abundance [13–16].

The Greater Everglades Ecosystem (GEE) in Florida is a vast, shallow marsh with slow laminar sheet flow, and is amenable to the application of eDNA monitoring to detect invasive aquatic or semi-aquatic species. Ecosystems in southern Florida have high invasibility as well as significant inputs from the exotic pet trade, and effective detection methods are therefore important for identifying and managing the frequent introduction of exotic species [17–21]. The Burmese python (*Python bivittatus*) is a highly cryptic and well-established invasive apex predator in southern Florida. While detection rates using traditional methods have been very low (<1%), high detection rates of Burmese pythons using eDNA occupancy analysis (>90%) have enabled occupancy rates to be quantified in logistically difficult-to-survey habitats like sawgrass marsh and tree islands far from roads and canals [5, 6, 8, 22].

As novel superpredators (*sensu* Terborgh [23]), Burmese pythons (hereafter “pythons”) are exerting negative effects on multiple trophic levels in the GEE. In Everglades National Park, pythons are linked to dramatic declines in small mammal populations (>90%) over the past decade(s) [24–26]. Pythons have also negatively impacted mammalian species richness throughout the GEE [27]. Over 25 species of birds have been found in gut contents of pythons, but the extent to which pythons are impacting bird populations is unclear [28]. Pythons are likely to pose a particular threat to long-legged wading bird species that breed colonially on tree islands throughout the greater Everglades and represent dense, accessible prey. A python telemetry study in Everglades National Park found that tree islands are common-use areas for pythons [29]. Predation of wading bird nest contents by pythons in the Everglades was five times the rate of predation by native nest predators in wading bird colonies in 2017 [30]. Pythons are semi-aquatic, utilize arboreal habitat, and may engage in wide-ranging foraging for prey—all traits which would make colonies throughout the Everglades accessible [29, 31–33]. We used eDNA in water samples as a tool to test the prediction that pythons are attracted to tree islands with bird colonies compared to representative control islands without breeding birds in the central Everglades. We found the site-level occupancy estimate to be higher in colony islands (0.88, 95% credible interval (CRI) [0.59–1.00]) than control islands (0.42, 95% CRI [0.16–0.80]), supporting our prediction, and we generally detected higher concentrations of python eDNA in colony sites, as well.

Materials and methods

Sampling design


We compared python occupancy estimates and eDNA concentrations in 15 wading bird colonies and 15 islands that had no wading birds nesting (= “control” islands). All samples were

taken at sites within a 1,000 km² area of Water Conservation Area 3 in the central Everglades (25.989444°, -80.654444°; Fig 1). The central Everglades is characterized by vast expanses of sawgrass marsh (*Cladium jamaicense*) punctuated by tree islands. Tree islands that are dominated by willow (*Salix caroliniana*) support wading bird colonies during the dry season (January–June). Water moves slowly as laminar sheet flow in a roughly north-south direction across the nearly flat floodplain [34]. We targeted willow islands (N = 20) associated with pools made or enlarged by alligators excavating sediment to create refugia in the dry season, known as ‘alligator hole islands’ [35, 36]. Willow trees grow up on the banks of sediment surrounding the pool. Alligator holes provided a constant, mostly isolated source of water to sample eDNA in direct proximity to breeding species. These species included Little blue herons (*Egretta caerulea*), Tricolored herons (*Egretta tricolor*), Snowy egrets (*Egretta thula*), and Anhingas (*Anhinga anhinga*), which nest in the trees around alligator holes. Colony breeding aggregations ranged between 8–84 pairs in our sampled alligator hole islands (229–3803 m²). The alligator hole at the northern end of each island ranged in area from 10–134 m², with an average area of 47.4 m². In the tree islands not associated with alligator holes, 25–94 breeding pairs of mostly Great egrets (*Ardea alba*) were present in four colonies (2319–16,197 m²) and 11,352 pairs of mostly White ibises (*Eudocimus albus*) were present in one outlier (colony B3; 86,295 m²). For each of the 15 colony islands, we selected a paired control island of similar size, habitat type and geographic location that was known to be unoccupied by wading bird colonies for a minimum of 4 years using long-term survey data. The five colony-control pairs that did not contain alligator holes were willow strand islands that were surrounded by water or partially inundated.

To reduce disturbance during times of day with high thermal stress to chicks and eggs, we generally visited colonies between 8–11AM to take eDNA samples. Most controls were also sampled before 11AM, except for four control sites visited between 11AM and 1PM. We measured water temperature for the majority of samples and imputed values for the remaining 10.4% of samples (out of 265 in total) without temperature measurements because of thermometer malfunction (S1 Appendix). We found a positive relationship between sample collection time and sample temperature (samples with imputed values excluded: Pearson correlation coefficient = 0.34, $t_{(238)} = 5.5$, $P = < 0.0001$, S1 Fig). To reduce potential temperature-related bias when comparing eDNA concentrations in paired colonies and controls, we excluded three colony-control pairs in which the median temperature of control samples was at least 8°C higher than the median temperature of the paired colony sample. This left 12 acceptable colony-control pairs for comparing eDNA concentrations. We retained all 15 pairs in the occupancy analysis because we did not expect water temperature at a single point in time to determine sample or site eDNA presence, although it could influence detection probability when eDNA is present [42–44]. Our recorded water temperatures ranged from 19 to 35°C. In a related camera-trapping study on python predation rates in colonies, the majority of python detections were nocturnal, and ambient temperatures recorded during detections ranged from 13.9 to 30°C [30]. Therefore, although eDNA studies involving other ectotherms, like fish, have found that water temperature can affect eDNA shedding rates (via changes in life-history states or by increasing metabolic and physical activities) [45–48], we expect the range of temperatures that might impact python activity/behavior (and therefore sample or site presence) to be more extreme than the ranges we observed at the temporal scale of a day (and which recurred daily) during our sampling window.

Field collection protocol

We collected 265 water samples (mean 9.2 ± 2.1 [SD] samples at colony sites; mean 8.5 ± 2.3 [SD] samples at control sites) in 1-liter DNase and RNase free Nalgene bottles and added 34



colony islands (yellow) and control islands (blue). Negative locations, where no python eDNA was detected, are marked with crosses. The boundary of Everglades National Park (ENP) is marked in green. The blue rectangle in the upper inset marks the study area in WCA-3. The black inset inside the main map details a cluster of colony and control sites located in the western half of WCA-3. The geospatial data layers used in this figure were obtained from the Florida Geographic Data Library [37–39], the South Florida Water Management District [40], and the National Park Service [41].

<https://doi.org/10.1371/journal.pone.0213943.g001>

mL of DNA preservative (33 mL 95% ethanol, 1 mL sodium acetate [3M]), and changed nitrile gloves at each sampling point. We stored each sample in a separate sealed bag. In alligator hole islands, we took samples at regular intervals of 2–5 m depending on the circumference of the hole. For islands without alligator holes, we approached each location to within 2 m and then reached horizontally to collect undisturbed water 5–25 cm below the surface, depending on the total water depth. At the five colony-control island pairs (1–3,5,8) that did not contain alligator holes, we sampled water at regular intervals (20–30 m) around the island perimeter ($n = 2$ non-inundated pairs; 2,8) and along 100–150 m transects downstream or within partially inundated islands, oriented normal (90°) to the direction of flow ($n = 3$ pairs; 1,3,5). We collected water temperature and depth measurements at each point except on a handful of occasions when the thermometer malfunctioned. We established a negative field control at the end of each sampling excursion by adding distilled water and DNA preservative to a sterile bottle. All samples were stored in a -20°C chest freezer during the field season, then transported to the USGS lab facility and stored immediately in a -20°C freezer.

Research activities were carried out in Water Conservation Area 3. Although this is a public access recreational area, we entered breeding colonies under the authority of Florida Fish and Wildlife Conservation Commission scientific collecting permit LSSC-15-0004. The field work did not involve endangered or protected species. The research was conducted under the approved University of Florida Institutional Animal Care and Use Committee (IACUC) protocol #201708305, which specifically covered all aspects of this study pertaining to animals. No field permit was required to collect water samples in WCA-3 at colonies or control sites.

Laboratory analysis

We vacuum-filtered water samples through polyethersulfone filters following Hunter et al. [49]. All filtration equipment was soaked in a 20% bleach solution (mixed from 6% sodium hypochlorite stock solution) for 10–15 minutes between usage, rinsed with UltraPure H_2O and dried in a fume hood. After 6–8 uses of the filtration apparatus, we ran a negative filtering control, consisting of 1 L RNase-free water plus DNA preservative, to check for contamination. We incubated the filters in cetyl-trimethylammonium-bromide (CTAB) buffer for 4–7 days at 4°C following Hunter et al. [49] and performed a phenol-chloroform-isoamyl (PCI) DNA isolation following Renshaw et al. [50]. We rehydrated the extracted DNA pellet in 100 μL 1X Tris-EDTA (TE) Buffer and stored the samples in a -80°C freezer. Every 6–8 isolations, we performed a negative isolation control by adding three sterile filters to CTAB buffer and isolating alongside samples.

We performed 1–5 rounds of Zymo Research OneStep Inhibitor Removal kits (IRK) to remove polymerase chain reaction (PCR) inhibitors such as tannins, phenols, and other acids. We used sample color to help determine the number of IRK treatments. If a sample was still inhibited after the first PCR run (determined by the internal positive control, see below), we administered more IRKs and assessed using droplet digital PCR (ddPCR) again (described below). When administering IRKs, a trade-off exists between removal of inhibitors and the possible loss of target eDNA. Hunter et al. [49] found no measurable difference in eDNA concentration after one IRK treatment, but because we often had to administer multiple IRKs, we

use the term ‘minimum occurrence’ to reflect the possible loss of eDNA during this step. For samples with multiple ddPCR runs due to additional IRK treatments, we selected the analysis that contained the highest concentration of python eDNA. We used an exogenous internal positive control assay (YWHAZ Rhesus Monkey template and PreAmp from Bio-Rad) to test for the presence of inhibition and differentiate between a true negative or inhibited sample. It was still possible to detect python DNA molecules in inhibited samples because we measured inhibition based on known concentrations of the internal positive control (IPC) template present in each sample well. Our python assay appeared less susceptible to inhibition than the IPC assay, possibly due to assay efficiency. Inhibition after applying IRKs was especially prevalent in colony samples (S2 and S3 Figs).

We performed ddPCR amplification using the QX200 Bio-Rad platform, following Hunter et al. [51]. ddPCR is thought to be the best tool to cope with high levels of inhibitors found in our system because each sample is partitioned into 15,000–20,000 nanofluidic droplets in which PCR occurs independently and endpoint amplification is detected with a fluorescent probe [52, 53]. In addition, ddPCR enables absolute quantification of molecules and allows detection down to a single molecule. Our sample PCR mixture included 12.5 μL of ddPCR supermix from Bio-Rad, 4 μL sample, 150 nM VIC TaqMan probe, 800 nM of each primer, and the IPC assay (0.20 μL template; 1.0 μL PreAmp) for a total volume of 25 μL . We added 20 μL of the reaction mixture and 70 μL of droplet generation oil to form droplets using the Bio-Rad QX200 Droplet Generator. We transferred 40 μL of suspended droplets to a 96-well plate and applied the PCR protocol used by Hunter et al. [51]. Following amplification, each well was scanned for the presence of amplified target using the QX200 Droplet Reader. The ddPCR output data were analyzed with Quantasoft v1.7.4.0917. We set thresholds manually according to recommended specifications from Bio-Rad.

We subdivided each sample into PCR replicates ($n = 5$) to account for potential PCR error and to estimate the detection probability of eDNA given sample presence. On each plate we ran four negative no template control (NTC) replicates (RNase-free H_2O) and two positive control replicates of DNA (0.00085 ng/ μL) extracted from python tissue and purified. To guard against potential contamination and/or false positives [51, 54, 55] we applied a universal limit of blank (LOB) threshold to the concentration and presence-absence data following Hunter et al. [56]. We arrived at the LOB value by inspecting all negative controls (field, filtration, extraction, and NTC controls) for target amplification. The maximum average concentration of python eDNA we observed in any control was an NTC with a concentration of 0.258 copies/ μL . We therefore zeroed all samples at or below this concentration (see S1 Appendix for analysis of plate-specific standard curves as an alternative). We calculated python eDNA concentration estimates and confidence intervals following Dorazio and Hunter [57].

Statistical analysis

We fitted occupancy models using the hierarchical Bayesian Monte Carlo Markov Chain (MCMC) algorithm developed by Dorazio and Erickson [58], which quantified the observational error of our multi-scale eDNA sampling design in which we 1) sampled multiple sites within a region, 2) took multiple water samples within each site and 3) subdivided each sample into multiple PCR replicates to estimate detection probabilities. The hierarchical model estimated a latent site-level probability of eDNA occupancy (ψ), an average conditional probability of eDNA occurrence in a single sample ($\bar{\theta}$), and the conditional probability of detecting eDNA in each PCR replicate (p), given sample presence. Sites were defined as tree islands.

We ran 100,000 iterations of the MCMC algorithm per model and set burn-in at 5000 to discard the initial transient region of the chain and obtain precise parameter estimates,

Table 1. *A priori* predictions of covariates included in multi-scale occupancy models to refine estimates of eDNA detection probabilities, sample occupancy, and site occupancy.

Covariate	<i>A priori</i> predictions:	Empirical support
Island type (colony or control)	Pythons are attracted to colony islands because they contain high densities of avian prey. This would lead to higher eDNA site occupancy rates and concentrations at colonies compared to representative control islands. We assumed environmental factors impacting eDNA detection probability did not vary by island type.	Orzechowski et al. [30]: Pythons are novel predators in colonies
Time of day (temperature proxy)	The time of sampling negatively covaries with eDNA detection probability due to molecular degradation at higher temperatures later in the day. Water temperature increased daily across all sites in conjunction with daily ambient temperature increases. We did not expect the proxy for water temperature at a single point in time to predict site or sample occupancy.	Tsuji et al. [42] Strickler et al. [43] Eichmiller et al. [44] Barnes et al. [60]
Water depth	Water depth served as an indirect measure of long-term differences in UV-B exposure and water temperature. Shallow sites heat faster, and UV-B rays can penetrate more of the water column—both causing eDNA degradation. We predicted water depth should positively covary with detection probability and sample occupancy. Since water depth was correlated with collection date (see next), we included collection date as a sole predictor of sample occupancy.	Jane et al. [61] Kiesecker et al. [62]: UV radiation attenuated within 20 cm water depth
Collection date	Daily water temperatures in the marsh gradually increase in conjunction with increasing air temperatures and water recession in April-May in South Florida. The net temperature increase is not extreme (approx. 5–10°C) but could affect sample occupancy or detection probability. We included depth instead of collection date as a predictor of eDNA detection probability since both were correlated.	Duever et al. [63]; Romigh et al. [64], Schaffranek and Jenter [65], S4 Fig

<https://doi.org/10.1371/journal.pone.0213943.t001>

following Hunter et al. [8]. We inspected the Markov chain of each parameter for visual confirmation that it was mixing well in the sample space and likely converging to the posterior distribution ([59]; S2 Appendix). We assessed autocorrelation values to determine whether we were over-parameterizing the model and checked the Monte Carlo standard errors for each parameter estimate (S2 Appendix). We ran the occupancy analyses in software R v3.4.0, using the R package eDNAoccupancy, which fit the MCMC algorithm developed by Dorazio and Erickson [58] for eDNA sampling scenarios.

Model comparison and selection. We conducted an exploratory graphical analysis of the relationship between eDNA presence at each scale and the covariates described in the supplementary materials (S1 Appendix). The lack of support for most covariates considered in the exploratory analysis prompted us to include only variables for which we had *a priori* predictions based on previous empirical studies (Table 1).

In our models we included the categorical variable ‘island type’ as a predictor of site occupancy to test our prediction that pythons are attracted to active wading bird colonies compared to the control islands. We included collection date to account for potential differences in sample occupancy across our sampling window. We included time of day (a proxy for sample temperature), and water depth (accounting for site-level differences in temperature and UV-B exposure) as predictors of eDNA detection probabilities. See Table 1 for explicit predictions regarding each covariate. In addition to models containing these covariates, we ran a null intercept-only model.

All numeric variables were scaled and centered. We performed model selection using the Widely Applicable Information Criterion (WAIC) of a multiscale occupancy model. WAIC is a Bayesian criterion and uses the full posterior distribution to compute predictive variance and goodness of fit. Like AIC, models with more parameters are penalized with WAIC’s predictive variance term and lower values of WAIC indicate models that better fit the data. In the best model, we compared the posterior probability density curves of site occupancy in colonies versus controls to determine how much the distributions overlapped, and whether they were skewed.

Results

We detected python eDNA at 10 of 15 colony sites and 4 of 15 control sites (Table 2). Of 137 water samples taken at colonies, 19 (13.9%) contained python eDNA; in 128 control samples, 5 (3.9%) contained python eDNA (Table 2).

Of the seven models we compared (Table 3), the model with the lowest WAIC score included island type (β typeControl = -0.19, 95% CRI [-0.99–0.84]; β typeColony = 1.17, 95% CRI [0.21–5.58]) as a predictor of site occupancy, and collection time (δ = -0.33, 95% CRI [-0.64– -0.04]) as a predictor of detection probability. The estimated posterior distributions of python eDNA site occupancy clearly differed between control sites and colony sites (Fig 2). The posterior for colony sites was left-skewed with a peak at 1, whereas the posterior for control sites was more symmetrical and centered around an occupancy estimate of 0.42 (Fig 2). Estimates of detection probability declined with sample collection time, ranging from 0.82 to 0.40 (Fig 3).

The estimated median minimum site eDNA occupancy rate at the 15 colony islands was 0.88 (95% CRI [0.59–1.00]), while the estimate for the 15 control islands was 0.42 (95% CRI [0.16–0.80]). The differences in python eDNA occupancy between colony and control sites supported our prediction that pythons are attracted to colonies. Occupied sites were scattered throughout the 1000 km² sampling region, although an apparent cluster of occupied colonies existed in the western part of WCA-3 (Fig 1). We discovered a universally low minimum sample occupancy rate ($\bar{\theta}$ = 0.14, 95% CRI [0.09–0.20]) suggesting patchiness of naturally occurring python eDNA at all occupied sites [8, 66–68].

eDNA Concentration

The total concentration of python eDNA at each bird colony (mean 3.20 ± sd 9.80 copies/μL, n = 15) was an order of magnitude higher than the total concentration at each control site (mean 0.11 ± sd 0.21 copies/μL, n = 15). At 8 of 9 acceptable pairs where we detected python eDNA, colonies contained higher eDNA concentrations than controls (Fig 4). The mean difference in concentration at colonies compared to paired local controls (mean 3.77 ± sd 6.99 copies/μL) was not significant (paired t-test, $t(11) = -1.19$, $p = 0.26$). The total amount of python eDNA detected at any site ranged from 0–38.6 copies/μL (Fig 4). Python eDNA concentrations were within a similar range at all island types (alligator hole or non-alligator hole islands [pairs 1–3, 5, 8]), except for one sample at colony B6. This was an alligator hole site,

Table 2. Naïve and model estimates of minimum site and sample occupancy, average detection probability, and eDNA concentration estimates at colony and control sites.

Site	Positive/ total samples (% positive)	Positive/ total sites	Average ^a total [eDNA] copies/μL	Min ^b /max sample [eDNA] copies/μL	Ψ (95% CRI)	$\bar{\theta}$ (95% CRI)	\bar{p} (95% CRI)
Colonies	19/137 (13.9%)	10/15	3.20	0.26/ 38.29	0.88 (0.59– 1.00)	0.14 (0.09– 0.20)	0.61 (0.48– 0.73)
Controls	5/128 (3.9%)	4/15	0.11	0.26/ 0.39	0.42 (0.16– 0.80)		

Environmental DNA concentration abbreviated [eDNA] and measured in copies per μL. Estimates of Ψ (site-level python eDNA occupancy probability in colonies and controls), $\bar{\theta}$ (average conditional probability of eDNA occurrence in a single sample) and \bar{p} (eDNA detection probability, averaged over all sites) are from the top ranked model in Table 3. 95% credible intervals (CRI) are reported for each estimate. Colonies were defined as active wading bird breeding sites and controls were empty islands of similar size and geographic location.

^aIncluding zeros

^bExcluding zeros

<https://doi.org/10.1371/journal.pone.0213943.t002>

Table 3. Model comparison using the Widely Applicable Information Criterion (WAIC).

	Model	WAIC	Predictive variance	Lack of fit
(1)	$\psi(\text{type})\theta(\cdot)p(\text{time})$	41.85	6.96	34.89
(2)	$\psi(\text{type})\theta(\cdot)p(\cdot)$	42.14	6.00	36.14
(3)	$\psi(\cdot)\theta(\cdot)p(\cdot)$	42.17	6.03	36.14
(4)	$\psi(\text{type})\theta(\text{date})p(\cdot)$	42.22	6.08	36.14
(5)	$\psi(\text{type})\theta(\text{date})p(\text{time})$	42.37	7.34	35.03
(6)	$\psi(\text{type})\theta(\cdot)p(\text{time} + \text{depth})$	42.70	8.04	34.66
(7)	$\psi(\text{type})\theta(\cdot)p(\text{depth})$	42.94	7.44	35.50

Models are listed in order of increasing WAIC score. The components of each WAIC score—predictive variance and lack of fit—are also reported.

<https://doi.org/10.1371/journal.pone.0213943.t003>


which contained an outlying concentration (38.29 copies/ μL) compared to the rest of the samples. We found that eDNA concentration declined with sample water temperature, as expected (S5 Fig).

Discussion

We quantified occupancy and detection rates of python eDNA at tree islands in the central Everglades, a critical habitat for breeding wading birds. The estimate of python eDNA site occupancy was higher in colonies than control islands, suggesting that pythons are attracted to wading bird colonies during the wading bird breeding season. A low estimate of sample occupancy indicated spatial heterogeneity of eDNA and could also in part be reduced by high levels of PCR inhibitors throughout our sampling locations. We found a strong relationship between our eDNA detection probability and temperature proxy in occupied samples, which corroborates the results of other studies demonstrating the effect of temperature on eDNA degradation [42–44].

The high python eDNA occupancy rates we observed in colonies were notable because the Everglades is not thought to be a hospitable environment for eDNA. eDNA likely degrades quickly in the Everglades because the water is shallow, warm, acidic, and contains abundant microbial fauna, all of which degrades eDNA [42, 43]. In addition, sun exposure provides constant UV radiation during the day. Acidic pH in conjunction with water at 35°C and UV exposure has been demonstrated to reduce longevity of eDNA to 8 days or less [43]. The rapid break-down of eDNA in harsh conditions, such as those found in the Everglades, therefore enhances the temporal resolution of a positive eDNA detection. Based on research in other systems, the typical lifespan of eDNA is likely 1–27 days [69–71].

The spatial resolution of a positive eDNA detection in the Everglades in part depends on water flow velocity along with eDNA lifespan. Average flow velocities in slough (open-water marsh) habitat have been estimated at 173 m day⁻¹ in mid-January, when bird colonies start to form during the dry season [34]. The majority of our sites were alligator hole islands, which consisted of a sun-exposed pool of stagnant water with no outflow at the time of sampling. Inflow from the slough was mostly cut off, except for an alligator path typically at the north apex of the island. The spatial resolution for detecting recently shed python eDNA should be especially high in these locations. In addition, water flow was strongly curtailed across our sampling region as a result of the rapid recession in water levels in March and April. Harvey et al. [34] found a strong reduction in mean flow velocity in conjunction with receding water levels in January onward. For colony-control pairs sampled around the island perimeter (n = 2



Site-level eDNA occurrence probability (ψ)

Fig 2. Estimated posterior distributions of site occupancy (ψ) in colony versus control sites.

<https://doi.org/10.1371/journal.pone.0213943.g002>

pairs) or those that were partially inundated willow strand islands ($n = 3$ pairs), we therefore assumed upstream contamination of eDNA was negligible in April and May.

Given the likely brief lifespan of eDNA and its short travel distance during our sampling period, the high site occupancy estimates in this study appear conservative. In addition to presence, the concentration of eDNA (hereafter [eDNA]) may also allow some inference regarding site usage. DNA molecules are shed into the environment in a clumped state, and as time elapses, they disperse, become fragmented and destroyed, or filtered out of the water. Longer eDNA fragments, that are more likely to contain the assay barcode, degrade faster than short fragments of eDNA [72]. Pilliod et al. [15] found that removing animals from a flowing stream drastically reduced [eDNA] detected within an hour. Similarly, in other studies involving closed systems, [eDNA] steadily declined over time after removing study organisms from experimental tanks [69, 73]. A relationship between [eDNA] and relative animal abundance has also been demonstrated in other systems where abundance was known using traditional methods or calibrated with experimental eDNA trials in tanks [13, 74]. Given these findings, it appears reasonable that a sample with high [eDNA] may indicate higher abundance or more recent presence of an animal (barring the complication of dead animal eDNA sources) with greater spatiotemporal precision than a sample containing one or two detected molecules. [eDNA] can be highly heterogeneous, however. The eDNA in the sample containing the outlying [eDNA] from site B6 could have come from a highly concentrated and localized release of




Fig 3. Estimated relationship between detection probability and sample collection time (temperature proxy).

<https://doi.org/10.1371/journal.pone.0213943.g003>

gametes or feces from a single animal, for example, or alternatively from a dead python in the area. We suggest the need for field-based trials with live pythons in a natural system to further explore the strength of this relationship between heterogeneous [eDNA] and abundance or recent presence in an open system.

Our detection assay appeared to be robust in the presence of inhibitors. Overall, PCR inhibition as measured with the exogenous IPC was quite high in our samples. This meant our ability to detect python eDNA molecules was diminished since inhibitors impede or prevent the replication of target eDNA, which is essential for detection. High levels of bird guano were apparent from the algal blooms frequently observed in colony alligator holes, and never observed in control islands [75]. Bird guano contains calcium, an inhibitor of PCR [76]. The fact that we detected the highest concentrations of python eDNA in colonies, despite the possibility of generally higher levels of inhibition in colonies, suggests our estimates of eDNA concentration were conservative relative to those in controls.

We conclude eDNA monitoring is useful for detection of python presence in the Everglades ecosystem and is the most powerful and quantitatively accurate technique currently available, especially in habitats where inaccessibility bias is high and the crypticity of pythons is augmented by impenetrable tree island vegetation. The inference we draw from our data is that Burmese pythons are widely distributed across the central Everglades in proximity to active bird colonies. Taken in conjunction with evidence of repeated python predation of bird nests by multiple individual snakes at some colonies [30] and observations of a telemetered python performing directed movement towards a colony (B. Smith, USGS Biologist, 25 May 2018, personal communication), it seems likely that pythons are preferentially attracted to wading bird colonies as sources of prey. Continued monitoring is necessary to determine whether python management in wading bird colonies will be needed to protect these avian species.




Fig 4. Total concentration of Burmese python eDNA at colony (dark grey bars) and control (light grey bars) pairs with 95% confidence intervals. Pair 6 is plotted on a separate axis because the colony concentration is much higher than any other site. At pairs 1, 3 and 8, (§) any differences in concentration were likely subject to temperature-related bias.

<https://doi.org/10.1371/journal.pone.0213943.g004>

Supporting information

S1 Appendix. Supplementary methods information.

(DOCX)

S2 Appendix. Trace and ACF plots of Markov chains from all occupancy models and MCMC standard errors of parameter estimates.

(DOCX)

S1 Fig. Relationship between time collected and sample water temperature.

(TIFF)

S2 Fig. Sample inhibition rates as a function of the number of Zymo inhibitor removal kits (IRKs) applied to each sample. Zero to five Zymo kits were administered to colony and control samples. “MB” refers to the Mo Bio inhibitor removal kit, which was administered to two colony samples. Sample size is noted underneath each boxplot. Inhibition rates (as measured with the IPC) were often high for both colony and control samples, even when more IRKs were applied. Of samples given three to five IRKs, colony samples were still more inhibited than control samples.

(TIFF)

S3 Fig. Violin plot of inhibition rates in colony and control samples. The amount of inhibition was quantified by dividing the total concentration of the internal positive control (IPC) in each sample by the IPC concentration of the standards on each plate, which served as a completely uninhibited reference point. The median inhibition rate in colony samples was close to 100% whereas it was less than 25% in control samples (solid black lines denote the median). Points are slightly jittered to reduce overlapping.

(TIFF)

S4 Fig. Relationship between water temperature and collection date. We separated the data into early versus late morning groups (before or after 11AM) because of the influence daily increases in air temperature had on water temperature. We took 70.8% (97/137) of colony samples before 11AM and 71.1% (91/128) of control samples before 11 AM.

(TIFF)

S5 Fig. Relationship between eDNA concentration and sample water temperature. Samples with zero concentration are excluded from the regression line. The dashed vertical lines mark the temperature range represented by both colony and control samples. The majority of samples (218/265 or 82.3%) were taken within this range (20.3–27.8 °C). The size of the triangles flanking the x axis indicate the number of negative colony and control samples taken at each temperature. One outlier in sample concentration from colony site B6 (38.29 copies/μL taken at 23.2 °C) was omitted to enhance readability.

(TIFF)

Acknowledgments

We thank L. Garner, J. Zabala Albizua, V. Dobrinova Vitanova, A. Bacher, D. LaFlamme, S. Carvey, and D. MacRae for assistance with sample collection in 2017. We are grateful to G. Meigs-Friend in particular, and J. Ferrante, B. Caton, and M. Jubair for guidance on laboratory protocols and analyses. We thank our lab technicians A. Ulmer, M. Giles, and S. Gamez for processing and preparing samples for ddPCR. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Author Contributions

Conceptualization: Sophia C. M. Orzechowski, Peter C. Frederick, Margaret E. Hunter.

Data curation: Sophia C. M. Orzechowski.

Formal analysis: Sophia C. M. Orzechowski, Robert M. Dorazio, Margaret E. Hunter.

Funding acquisition: Sophia C. M. Orzechowski, Peter C. Frederick, Margaret E. Hunter.

Investigation: Sophia C. M. Orzechowski, Margaret E. Hunter.

Methodology: Sophia C. M. Orzechowski, Peter C. Frederick, Robert M. Dorazio, Margaret E. Hunter.

Project administration: Sophia C. M. Orzechowski, Peter C. Frederick, Margaret E. Hunter.

Resources: Peter C. Frederick, Margaret E. Hunter.

Software: Robert M. Dorazio.

Supervision: Sophia C. M. Orzechowski, Peter C. Frederick, Robert M. Dorazio, Margaret E. Hunter.

Validation: Sophia C. M. Orzechowski, Robert M. Dorazio, Margaret E. Hunter.

Visualization: Sophia C. M. Orzechowski, Robert M. Dorazio.

Writing – original draft: Sophia C. M. Orzechowski.

Writing – review & editing: Sophia C. M. Orzechowski, Peter C. Frederick, Robert M. Dorazio, Margaret E. Hunter.

References

1. McCarthy MA, Moore JL, Morris WK, Parris KM, Garrard GE, Vesk PA, et al. The influence of abundance on detectability. *Oikos*. 2013; 122(5):717–26.
2. Trebitz AS, Hoffman JC, Darling JA, Pilgrim EM, Kelly JR, Brown EA, et al. Early detection monitoring for aquatic non-indigenous species: Optimizing surveillance, incorporating advanced technologies, and identifying research needs. *Journal of Environmental Management*. 2017; 202:299–310. <https://doi.org/10.1016/j.jenvman.2017.07.045> PMID: 28738203
3. Novak SJ. The role of evolution in the invasion process. *Proceedings of the National Academy of Sciences of the United States of America*. 2007; 104(10):3671–2. <https://doi.org/10.1073/pnas.0700224104> PMID: 17360409
4. Dlugosch KM, Parker IM. Founding events in species invasions: genetic variation, adaptive evolution, and the role of multiple introductions. *Molecular Ecology*. 2008; 17(1):431–49. <https://doi.org/10.1111/j.1365-294X.2007.03538.x> PMID: 17908213
5. Reed RN, Hart KM, Rodda GH, Mazzotti FJ, Snow RW, Cherkiss M, et al. A field test of attractant traps for invasive Burmese pythons (*Python molurus bivittatus*) in southern Florida. *Wildlife Research*. 2011; 38(2):114–21. <https://doi.org/10.1071/wr10202>
6. Dorcas ME, Willson JD. Hidden giants: problems associated with studying secretive invasive pythons. In: Lutterschmidt W, editor. *Reptiles in Research: Investigations of Ecology, Physiology, and Behavior from Desert to Sea*. Hauppauge, NY, USA: Nova Science Publ. Inc.; 2013. p. 367–85.
7. Dejean T, Valentini A, Miquel C, Taberlet P, Bellemain E, Miaud C. Improved detection of an alien invasive species through environmental DNA barcoding: the example of the American bullfrog *Lithobates catesbeianus*. *Journal of Applied Ecology*. 2012; 49(4):953–9. <https://doi.org/10.1111/j.1365-2664.2012.02171.x>
8. Hunter ME, Oyler-McCance SJ, Dorazio RM, Fike JA, Smith BJ, Hunter CT, et al. Environmental DNA (eDNA) Sampling Improves Occurrence and Detection Estimates of Invasive Burmese Pythons. *Plos One*. 2015; 10(4). <https://doi.org/10.1371/journal.pone.0121655> PMID: 25874630
9. Thomsen PF, Willerslev E. Environmental DNA—An emerging tool in conservation for monitoring past and present biodiversity. *Biological Conservation*. 2015; 183:4–18. <https://doi.org/10.1016/j.biocon.2014.11.019>

10. Bohmann K, Evans A, Gilbert MTP, Carvalho GR, Creer S, Knapp M, et al. Environmental DNA for wildlife biology and biodiversity monitoring. *Trends in Ecology & Evolution*. 2014; 29(6):358–67. <https://doi.org/10.1016/j.tree.2014.04.003> PMID: 24821515
11. Smart AS, Tingley R, Weeks AR, van Rooyen AR, McCarthy MA. Environmental DNA sampling is more sensitive than a traditional survey technique for detecting an aquatic invader. *Ecological Applications*. 2015; 25(7):1944–52. <https://doi.org/10.1890/14-1751.1> PMID: 26591459
12. Biggs J, Ewald N, Valentini A, Gaboriaud C, Dejean T, Griffiths RA, et al. Using eDNA to develop a national citizen science-based monitoring programme for the great crested newt (*Triturus cristatus*). *Biological Conservation*. 2015; 183:19–28. <https://doi.org/10.1016/j.biocon.2014.11.029>
13. Doi H, Inui R, Akamatsu Y, Kanno K, Yamanaka H, Takahara T, et al. Environmental DNA analysis for estimating the abundance and biomass of stream fish. *Freshwater Biology*. 2017; 62(1):30–9. <https://doi.org/10.1111/fwb.12846>
14. Jerde CL, Chadderton WL, Mahon AR, Renshaw MA, Corush J, Budny ML, et al. Detection of Asian carp DNA as part of a Great Lakes basin-wide surveillance program. *Canadian Journal of Fisheries and Aquatic Sciences*. 2013; 70(4):522–6. <https://doi.org/10.1139/cjfas-2012-0478>
15. Pilliod DS, Goldberg CS, Arkle RS, Waits LP. Estimating occupancy and abundance of stream amphibians using environmental DNA from filtered water samples. *Canadian Journal of Fisheries and Aquatic Sciences*. 2013; 70(8):1123–30. <https://doi.org/10.1139/cjfas-2013-0047>
16. Takahara T, Minamoto T, Doi H. Using Environmental DNA to Estimate the Distribution of an Invasive Fish Species in Ponds. *Plos One*. 2013; 8(2). <https://doi.org/10.1371/journal.pone.0056584> PMID: 23437177
17. Ewel J. Invasibility: lessons from south Florida. In: Mooney HA, Drake JA, editors. *Ecology of biological invasions of North America and Hawaii*. Ecological Studies (Analysis and Synthesis), vol 58. New York, NY: Springer; 1986. p. 214–30.
18. Fujisaki I, Hart KM, Mazzotti FJ, Rice KG, Snow S, Rochford M. Risk assessment of potential invasiveness of exotic reptiles imported to south Florida. *Biological Invasions*. 2010; 12(8):2585–96. <https://doi.org/10.1007/s10530-009-9667-1>
19. Simberloff D, Schmitz DC, Brown TC. *Strangers in paradise: impact and management of nonindigenous species in Florida*. Washington, DC: Island press; 1997.
20. Blackburn TM, Cassey P. Patterns of non-randomness in the exotic avifauna of Florida. *Diversity and Distributions*. 2007; 13(5):519–26. <https://doi.org/10.1111/j.1472-4642.2007.00358.x>
21. Romagosa CM, Guyer C, Wooten MC. Contribution of the Live-Vertebrate Trade toward Taxonomic Homogenization. *Conservation Biology*. 2009; 23(4):1001–7. <https://doi.org/10.1111/j.1523-1739.2009.01194.x> PMID: 19627323
22. Piaggio AJ, Engeman RM, Hopken MW, Humphrey JS, Keacher KL, Bruce WE, et al. Detecting an elusive invasive species: a diagnostic PCR to detect Burmese python in Florida waters and an assessment of persistence of environmental DNA. *Molecular Ecology Resources*. 2014; 14(2):374–80. <https://doi.org/10.1111/1755-0998.12180> PMID: 24119154
23. Terborgh JW. Toward a trophic theory of species diversity. *Proceedings of the National Academy of Sciences of the United States of America*. 2015; 112(37):11415–22. <https://doi.org/10.1073/pnas.1501070112> PMID: 26374788
24. Dorcas ME, Willson JD, Reed RN, Snow RW, Rochford MR, Miller MA, et al. Severe mammal declines coincide with proliferation of invasive Burmese pythons in Everglades National Park. *Proceedings of the National Academy of Sciences of the United States of America*. 2012; 109(7):2418–22. <https://doi.org/10.1073/pnas.1115226109> PMID: 22308381
25. Sovie AR, McCleery RA, Fletcher RJ Jr., Hart KM. Invasive pythons, not anthropogenic stressors, explain the distribution of a keystone species. *Biological Invasions*. 2016; 18(11):3309–18. <https://doi.org/10.1007/s10530-016-1221-3>
26. McCleery RA, Sovie A, Reed RN, Cunningham MW, Hunter ME, Hart KM. Marsh rabbit mortalities tie pythons to the precipitous decline of mammals in the Everglades. *Proceedings Biological sciences / The Royal Society*. 2015; 282(1805). <https://doi.org/10.1098/rspb.2015.0120> PMID: 25788598
27. Reichert BE, Sovie AR, Udell BJ, Hart KM, Borkhataria RR, Bonneau M, et al. Urbanization may limit impacts of an invasive predator on native mammal diversity. *Diversity and Distributions*. 2017; 23(4):355–67. <https://doi.org/10.1111/ddi.12531>
28. Dove CJ, Snow RW, Rochford MR, Mazzotti FJ. Birds consumed by the invasive Burmese python (*Python molurus bivittatus*) in Everglades National Park, Florida, USA. *Wilson Journal of Ornithology*. 2011; 123(1):126–31. <https://doi.org/10.1676/10-092.1>

29. Hart KM, Cherkiss MS, Smith BJ, Mazzotti FJ, Fujisaki I, Snow RW, et al. Home range, habitat use, and movement patterns of non-native Burmese pythons in Everglades National Park, Florida, USA. *Animal Biotelemetry*. 2015; 3(8).
30. Orzechowski SCM, Romagosa CM, Frederick PC. Invasive Burmese pythons (*Python bivittatus*) are novel nest predators in wading bird colonies of the Florida Everglades. *Biological Invasions*. 2019; *Forthcoming*. <https://doi.org/10.1007/s10530-019-01979-x>
31. Hart KM, Schofield PJ, Gregoire DR. Experimentally derived salinity tolerance of hatchling Burmese pythons (*Python molurus bivittatus*) from the Everglades, Florida (USA). *Journal of Experimental Marine Biology and Ecology*. 2012; 413:56–9. <https://doi.org/10.1016/j.jembe.2011.11.021>
32. Dorcas ME, Willson JD. *Invasive pythons in the United States: ecology of an introduced predator*. Athens, GA: University of Georgia Press; 2011.
33. Greene HW. *Snakes: the evolution of mystery in nature*. Berkeley, CA: Univ of California Press; 1997.
34. Harvey JW, Schaffranek RW, Noe GB, Larsen LG, Nowacki DJ, O'Connor BL. Hydroecological factors governing surface water flow on a low-gradient floodplain. *Water Resources Research*. 2009; 45. <https://doi.org/10.1029/2008wr007129>
35. Campbell MR, Mazzotti FJ. Characterization of natural and artificial alligator holes. *Southeastern Naturalist*. 2004; 3(4):583–94. [https://doi.org/10.1656/1528-7092\(2004\)003\[0583:conaaa\]2.0.co;2](https://doi.org/10.1656/1528-7092(2004)003[0583:conaaa]2.0.co;2)
36. Palmer ML, Mazzotti FJ. Structure of Everglades alligator holes. *Wetlands*. 2004; 24(1):115–22. [https://doi.org/10.1672/0277-5212\(2004\)024\[0115:soeah\]2.0.co;2](https://doi.org/10.1672/0277-5212(2004)024[0115:soeah]2.0.co;2)
37. Florida Natural Areas Inventory. Water Conservation Areas—Line, UTM [Data file]. 1999. Available from: <http://www.fgdl.org> (Accessed 23 June 2012). Florida Geographic Data Library, University of Florida, Gainesville, FL.
38. Florida Fish and Wildlife Conservation Commission—Fish and Wildlife Research Institute. Florida coastline [Data file]. 2004. Available from: <http://www.fgdl.org> (Accessed 26 June 2008). Florida Geographic Data Library, University of Florida, Gainesville, FL.
39. Florida Department of Transportation, Transportation Statistics Office. Florida Department of Transportation RCI Roads—February 2011 [Data file]. 2011. Available from: <http://www.fgdl.org> (Accessed 14 March 2011). Florida Geographic Data Library, University of Florida, Gainesville, FL.
40. South Florida Water Management District. SFWMD Canals [Data file]. 1997. Available from: <http://fcelter.fiu.edu/data/GIS/> (Accessed 23 January 2019).
41. National Park Service. Everglades National Park Boundary [Data file]. 2002. Available from: <http://fcelter.fiu.edu/data/GIS/> (Accessed 8 August 2003).
42. Tsuji S, Ushio M, Sakurai S, Minamoto T, Yamanaka H. Water temperature-dependent degradation of environmental DNA and its relation to bacterial abundance. *Plos One*. 2017; 12(4). <https://doi.org/10.1371/journal.pone.0176608> PMID: 28448613
43. Strickler KM, Fremier AK, Goldberg CS. Quantifying effects of UV-B, temperature, and pH on eDNA degradation in aquatic microcosms. *Biological Conservation*. 2015; 183:85–92. <https://doi.org/10.1016/j.biocon.2014.11.038>
44. Eichmiller JJ, Best SE, Sorensen PW. Effects of Temperature and Trophic State on Degradation of Environmental DNA in Lake Water. *Environmental Science & Technology*. 2016; 50(4):1859–67. <https://doi.org/10.1021/acs.est.5b05672> PMID: 26771292
45. Buxton AS, Groombridge JJ, Zakaria NB, Griffiths RA. Seasonal variation in environmental DNA in relation to population size and environmental factors. *Scientific Reports*. 2017; 7. <https://doi.org/10.1038/s41598-017-00035-9>
46. Bylemans J, Furlan EM, Hardy CM, McGuffie P, Lintermans M, Gleeson DM. An environmental DNA-based method for monitoring spawning activity: a case study, using the endangered Macquarie perch (*Macquaria australasica*). *Methods in Ecology and Evolution*. 2017; 8(5):646–55. <https://doi.org/10.1111/2041-210x.12709>
47. Lacoursiere-Roussel A, Rosabal M, Bernatchez L. Estimating fish abundance and biomass from eDNA concentrations: variability among capture methods and environmental conditions. *Molecular Ecology Resources*. 2016; 16(6):1401–14. <https://doi.org/10.1111/1755-0998.12522> PMID: 26946353
48. Robson HLA, Noble TH, Saunders RJ, Robson SKA, Burrows DW, Jerry DR. Fine-tuning for the tropics: application of eDNA technology for invasive fish detection in tropical freshwater ecosystems. *Molecular Ecology Resources*. 2016; 16(4):922–32. <https://doi.org/10.1111/1755-0998.12505> PMID: 26849294
49. Hunter ME, Ferrante JA, Meigs-Friend G, Ulmer A. Improving eDNA yield and inhibitor reduction through increased water volumes and multi-filter isolation techniques. *Scientific Reports*. 2019; 9:5259. <https://doi.org/10.1038/s41598-019-40977-w> PMID: 30918268
50. Renshaw MA, Olds BP, Jerde CL, McVeigh MM, Lodge DM. The room temperature preservation of filtered environmental DNA samples and assimilation into a phenol-chloroform-isoamyl alcohol DNA

- extraction. *Molecular Ecology Resources*. 2015; 15(1):168–76. <https://doi.org/10.1111/1755-0998.12281> PMID: 24834966
51. Hunter ME, Dorazio RM, Butterfield JSS, Meigs-Friend G, Nico LG, Ferrante JA. Detection limits of quantitative and digital PCR assays and their influence in presence-absence surveys of environmental DNA. *Molecular Ecology Resources*. 2017; 17(2):221–9. <https://doi.org/10.1111/1755-0998.12619> PMID: 27768244
 52. Doi H, Takahara T, Minamoto T, Matsuhashi S, Uchii K, Yamanaka H. Droplet Digital Polymerase Chain Reaction (PCR) Outperforms Real-Time PCR in the Detection of Environmental DNA from an Invasive Fish Species. *Environmental Science & Technology*. 2015; 49(9):5601–8. <https://doi.org/10.1021/acs.est.5b00253> PMID: 25850372
 53. Dingle TC, Sedlak RH, Cook L, Jerome KR. Tolerance of Droplet-Digital PCR vs Real-Time Quantitative PCR to Inhibitory Substances. *Clinical Chemistry*. 2013; 59(11):1670–2. <https://doi.org/10.1373/clinchem.2013.211045> PMID: 24003063
 54. Strain MC, Lada SM, Luong T, Rought SE, Gianella S, Terry VH, et al. Highly Precise Measurement of HIV DNA by Droplet Digital PCR. *Plos One*. 2013; 8(4). <https://doi.org/10.1371/journal.pone.0055943> PMID: 23573183
 55. Kiselinova M, Pasternak AO, De Spiegelaere W, Vogelaers D, Berkhout B, Vandekerckhove L. Comparison of Droplet Digital PCR and Seminested Real-Time PCR for Quantification of Cell-Associated HIV-1 RNA. *Plos One*. 2014; 9(1). <https://doi.org/10.1371/journal.pone.0085999> PMID: 24465831
 56. Hunter ME, Meigs-Friend G, Ferrante JA, Kamla AT, Dorazio RM, Diagne LK, et al. Surveys of environmental DNA (eDNA): a new approach to estimate occurrence in vulnerable manatee populations. *Endangered Species Research*. 2018; 35:101–11. <https://doi.org/10.3354/esr00880>
 57. Dorazio RM, Hunter ME. Statistical Models for the Analysis and Design of Digital Polymerase Chain Reaction (dPCR) Experiments. *Analytical Chemistry*. 2015; 87(21):10886–93. <https://doi.org/10.1021/acs.analchem.5b02429> PMID: 26436653
 58. Dorazio RM, Erickson RA. ednaoccupancy: An R package for multiscale occupancy modelling of environmental DNA data. *Molecular ecology resources*. 2017.
 59. Gilks WR, Richardson S, Spiegelhalter D. Markov chain Monte Carlo in practice. Boca Raton, FL: CRC press; 1995.
 60. Barnes MA, Turner CR, Jerde CL, Renshaw MA, Chadderton WL, Lodge DM. Environmental Conditions Influence eDNA Persistence in Aquatic Systems. *Environmental Science & Technology*. 2014; 48(3):1819–27. <https://doi.org/10.1021/es404734p> PMID: 24422450
 61. Jane SF, Wilcox TM, McKelvey KS, Young MK, Schwartz MK, Lowe WH, et al. Distance, flow and PCR inhibition: eDNA dynamics in two headwater streams. *Molecular Ecology Resources*. 2015; 15(1):216–27. <https://doi.org/10.1111/1755-0998.12285> PMID: 24890199
 62. Kiesecker JM, Blaustein AR, Belden LK. Complex causes of amphibian population declines. *Nature*. 2001; 410(6829):681–4. <https://doi.org/10.1038/35070552> PMID: 11287952
 63. Duever M, Meeder J, Meeder L, McCollom J. The climate of south Florida and its role in shaping the Everglades ecosystem. In: Davis S, Ogden J, editors. *Everglades: The ecosystem and its restoration*. Boca Raton, FL: St. Lucie Press; 1994. p. 225–48.
 64. Romigh MM, Davis SE, Rivera-Monroy VH, Twilley RR. Flux of organic carbon in a riverine mangrove wetland in the Florida Coastal Everglades. *Hydrobiologia*. 2006; 569:505–16. <https://doi.org/10.1007/s10750-006-0152-x>
 65. Schaffranek RW, Jenter HL. Observations of daily temperature patterns in the southern Florida Everglades. *Wetlands Engineering & River Restoration*. 2001: 1–4.
 66. Eichmiller JJ, Bajer PG, Sorensen PW. The Relationship between the Distribution of Common Carp and Their Environmental DNA in a Small Lake. *Plos One*. 2014;9(11). <https://doi.org/10.1371/journal.pone.0112611> PMID: 25383965
 67. Davison PI, Copp GH, Creach V, Vilizzi L, Britton JR. Application of environmental DNA analysis to inform invasive fish eradication operations. *Science of Nature*. 2017; 104(3–4). <https://doi.org/10.1007/s00114-017-1453-9> PMID: 28357478
 68. Chestnut T, Anderson C, Popa R, Blaustein AR, Voytek M, Olson DH, et al. Heterogeneous Occupancy and Density Estimates of the Pathogenic Fungus *Batrachochytrium dendrobatidis* in Waters of North America. *Plos One*. 2014; 9(9). <https://doi.org/10.1371/journal.pone.0106790> PMID: 25222122
 69. Dejean T, Valentini A, Duparc A, Pellier-Cuit S, Pompanon F, Taberlet P, et al. Persistence of Environmental DNA in Freshwater Ecosystems. *Plos One*. 2011; 6(8). <https://doi.org/10.1371/journal.pone.0023398> PMID: 21858099

70. Thomsen PF, Kielgast J, Iversen LL, Wiuf C, Rasmussen M, Gilbert MTP, et al. Monitoring endangered freshwater biodiversity using environmental DNA. *Molecular Ecology*. 2012; 21(11):2565–73. <https://doi.org/10.1111/j.1365-294X.2011.05418.x> PMID: 22151771
71. Pilliod DS, Goldberg CS, Arkle RS, Waits LP. Factors influencing detection of eDNA from a stream-dwelling amphibian. *Molecular Ecology Resources*. 2014; 14(1):109–16. <https://doi.org/10.1111/1755-0998.12159> PMID: 24034561
72. Jo T, Murakami H, Masuda R, Sakata MK, Yamamoto S, Minamoto T. Rapid degradation of longer DNA fragments enables the improved estimation of distribution and biomass using environmental DNA. *Molecular Ecology Resources*. 2017; 17(6):e25–e33. <https://doi.org/10.1111/1755-0998.12685> PMID: 28449215
73. Sassoubre LM, Yamahara KM, Gardner LD, Block BA, Boehm AB. Quantification of Environmental DNA (eDNA) Shedding and Decay Rates for Three Marine Fish. *Environmental Science & Technology*. 2016; 50(19):10456–64. <https://doi.org/10.1021/acs.est.6b03114> PMID: 27580258
74. Lodge DM, Turner CR, Jerde CL, Barnes MA, Chadderton L, Egan SP, et al. Conservation in a cup of water: estimating biodiversity and population abundance from environmental DNA. *Molecular Ecology*. 2012; 21(11):2555–8. <https://doi.org/10.1111/j.1365-294X.2012.05600.x> PMID: 22624944
75. Irick DL, Gu BH, Li YCC, Inglett PW, Frederick PC, Ross MS, et al. Wading bird guano enrichment of soil nutrients in tree islands of the Florida Everglades. *Science of the Total Environment*. 2015; 532:40–7. <https://doi.org/10.1016/j.scitotenv.2015.05.097> PMID: 26057723
76. Schrader C, Schielke A, Ellerbroek L, Johne R. PCR inhibitors—occurrence, properties and removal. *Journal of Applied Microbiology*. 2012; 113(5):1014–26. <https://doi.org/10.1111/j.1365-2672.2012.05384.x> PMID: 22747964