

2-(4-Bromobenzenesulfonamido)benzoic acid

Muhammad Nadeem Arshad,^a Islam Ullah Khan,^a Mehmet Akkurt,^{b*} Muhammad Shafiq^a and Ghulam Mustafa^a

^aMaterials Chemistry Laboratory, Department of Chemistry, Government College University, Lahore, Pakistan, and ^bDepartment of Physics, Faculty of Arts and Sciences, Erciyes University, 38039 Kayseri, Turkey

Correspondence e-mail: akkurt@erciyes.edu.tr, mnachemist@hotmail.com


Received 8 June 2009; accepted 12 June 2009

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.038; wR factor = 0.093; data-to-parameter ratio = 18.8.

In the title compound, $\text{C}_{13}\text{H}_{10}\text{BrNO}_4\text{S}$, the dihedral angle between the benzene rings is $82.75(15)^\circ$. An intramolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond generates an $S(6)$ ring motif. In the crystal structure, two molecules form an $R_2^2(8)$ centrosymmetric dimer through a pair of $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds. Intra- and intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds are also observed.

Related literature

For background to sulfonamide derivatives, see: Allison *et al.* (2006); Sheppard *et al.* (2006). For related structures, see: Arshad *et al.* (2009); Shafiq *et al.* (2009); Asiri *et al.* (2009). For hydrogen-bond graph-set terminology, see: Bernstein *et al.* (1995); Etter (1990).



Experimental

Crystal data

$\text{C}_{13}\text{H}_{10}\text{BrNO}_4\text{S}$

$M_r = 356.19$

Monoclinic, $C2/c$

$a = 27.8316(11)\text{ \AA}$

$b = 8.5684(4)\text{ \AA}$

$c = 11.6632(5)\text{ \AA}$

$\beta = 98.196(2)^\circ$

$V = 2752.9(2)\text{ \AA}^3$

$Z = 8$

Mo $K\alpha$ radiation

$\mu = 3.15\text{ mm}^{-1}$

$T = 296\text{ K}$

$0.23 \times 0.19 \times 0.11\text{ mm}$

Data collection

Bruker Kappa-APEXII CCD area-detector diffractometer

Absorption correction: refined from

ΔF

[Cubic fit to $\sin(\theta)/\lambda - 24$

parameters; Parkin *et al.*, 1995]

$T_{\min} = 0.497$, $T_{\max} = 0.707$

14822 measured reflections

3416 independent reflections

1764 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.042$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$

$wR(F^2) = 0.093$

$S = 0.98$

3416 reflections

182 parameters

H-atom parameters constrained

$\Delta\rho_{\max} = 0.27\text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.29\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1 \cdots O1	0.86	2.13	2.670 (3)	121
O2—H2A \cdots O1 ⁱ	0.82	1.89	2.703 (3)	173
C6—H6 \cdots O4 ⁱⁱ	0.93	2.56	3.185 (4)	125
C11—H11 \cdots O3 ⁱⁱⁱ	0.93	2.47	3.369 (3)	164
C12—H12 \cdots O4	0.93	2.31	2.987 (3)	130

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $x, -y + 1, z + \frac{1}{2}$; (iii) $x, y - 1, z$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON* (Spek, 2009).

The authors MNA, IUK, MS and GM acknowledge the Higher Education Commission, Pakistan, for support under the indigenous PhD 5000 Fellowship Programme.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS2431).

References

- Allison, B. D., Phuong, V. K., McAtee, L. C., Rosen, M., Magda Morton, M., Prendergast, C., Barrett, T., Lagaud, G., Jamie Freedman, J., Lina Li, L., Xiaodong Wu, X., Venkatesan, H., Pippel, M., Woods, C., Rizzolio, M. C., Hack, M., Hoey, K., Deng, X., King, C., Shankley, N. P. & Rabinowitz, M. H. (2006). *J. Med. Chem.* **49**, 6371–6390.
- Altomare, A., Burla, M. C., Camalli, M., Carrozzini, B., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Rizzi, R. (1999). *J. Appl. Cryst.* **32**, 339–340.
- Arshad, M. N., Khan, I. U., Shafiq, M. & Mukhtar, A. (2009). *Acta Cryst. E65*, o549.
- Asiri, A. M., Akkurt, M., Khan, S. A., Arshad, M. N., Khan, I. U. & Sharif, H. M. A. (2009). *Acta Cryst. E65*, o1246–o1247.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Etter, M. C. (1990). *Acc. Chem. Res.* **23**, 120–126.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Parkin, S., Moezzi, B. & Hope, H. (1995). *J. Appl. Cryst.* **28**, 53–56.
- Shafiq, M., Tahir, M. N., Khan, I. U., Arshad, M. N. & Asghar, M. N. (2009). *Acta Cryst. E65*, o1182.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.

- Sheppard, G. S., Wang, J., Kawai, M., Fidanze, S. D., Bamaung, N. Y., Erickson, S. A., Barnes, D. M., Tedrow, J. S., Kolaczkowski, L., Vasudevan, A., Park, D. C., Wang, G. T., Sanders, W. J., Mantei, R. A., Palazzo, F., Tucker-Garcia, L., Lou, P. P., Zhang, Q., Park, C. H., Kim, K. H., Petros, A., Olejniczak, E., Nettesheim, D., Hajduk, P., Henkin, J., Lesniewski, R., Davidsen, S. K. & Bell, R. L. (2006). *J. Med. Chem.* **49**, 3832–3849.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supplementary materials

Acta Cryst. (2009). E65, o1610-o1611 [doi:10.1107/S1600536809022545]

2-(4-Bromobenzenesulfonamido)benzoic acid

M. N. Arshad, I. U. Khan, M. Akkurt, M. Shafiq and G. Mustafa

Comment

Sulfonamides are biologically active organic compounds. The anthranilic sulfonamide derivative has been reported as inhibitors of Methionine aminopeptidase-2 (MetAP2) (Sheppard *et al.*, 2006) and halogenated anthranilic sulfonamide derivatives have been identified as novel, selective Cholecystokinin-2 Receptor Antagonists (Allison *et al.*, 2006). The title compound is halogenated sulfonamide in continuation of our studies on the synthesis of sulfonamide derivatives (Arshad *et al.*, 2009) and benzothiazines (Shafiq *et al.*, 2009).

In the title compound, (I), (Fig. 1), the values of the geometric parameters are normal, and they are comparable with those in the reported structure of the isomorf compound 2-benzenesulfonamidobenzoic acid (Asiri *et al.*, 2009). The angle between the benzene rings is 82.75 (15)°.

The crystal packing is stabilized by C—H···O and O—H···O hydrogen bonds (Table 1). The intramolecular N—H···O hydrogen bond generates a graph set motif S(6). The O—H···O hydrogen bond forms a cyclic dimer, with a $R^2_2(8)$ motif (Bernstein *et al.*, 1995; Etter, 1990), about a inversion centre (Fig. 2). Figure 3 shows the molecular packing for (I) viewed down the *b* axis, generating a zigzag layer running along the *a* axis.

Experimental

Anthranilic acid (2 g, 14.6 mmol) was dissolved in distilled water (10 ml) in a round bottom flask (25 ml). The pH of the solution was maintained at 8–9 using 1*M*, Na₂CO₃. 4- Bromobenzene sulfonylchloride (3.72 g, 14.6 mmol) was suspended to the above solution and stirred at room temperature until all the 4-bromobenzene sulfonyl chloride was consumed. Progress of the reaction was observed by disappearing of suspension to clear solution. On completion of the reaction the pH was adjusted 1–2, using 1 *N* HCl. The precipitate obtained was filtered, washed with distilled water, dried and recrystallized in methanol to yield dark brown crystals.

Refinement

H atoms were fixed geometrically and treated as riding, with C—H = 0.93 Å, N—H = 0.86 Å and O—H = 0.82 Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$ and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$.

supplementary materials

Figures




Fig. 1. View of the title molecule showing the atom labelling scheme and displacement ellipsoids at the 30% probability level.




Fig. 2. View of the dimeric structure of (I). Hydrogen bonds are indicated by dashed lines. The atom labelled with the suffix a is generated by the symmetry operator $(3/2 - x, 1/2 - y, 1 - z)$.




Fig. 3. The molecular packing and hydrogen bonding for (I), viewed down the b axis. Hydrogen atoms not involved in the showed interactions have been omitted for clarity.

2-(4-Bromobenzenesulfonamido)benzoic acid

Crystal data

$C_{13}H_{10}BrNO_4S$	$F_{000} = 1424$
$M_r = 356.19$	$D_x = 1.719 \text{ Mg m}^{-3}$
Monoclinic, $C2/c$	Mo $K\alpha$ radiation
Hall symbol: -C 2yc	$\lambda = 0.71073 \text{ \AA}$
$a = 27.8316 (11) \text{ \AA}$	Cell parameters from 3121 reflections
$b = 8.5684 (4) \text{ \AA}$	$\theta = 2.5\text{--}22.0^\circ$
$c = 11.6632 (5) \text{ \AA}$	$\mu = 3.15 \text{ mm}^{-1}$
$\beta = 98.196 (2)^\circ$	$T = 296 \text{ K}$
$V = 2752.9 (2) \text{ \AA}^3$	Block, dark brown
	$0.23 \times 0.19 \times 0.11 \text{ mm}$

$Z = 8$

Data collection

Bruker Kappa-APEXII CCD area-detector diffractometer	3416 independent reflections
Radiation source: sealed tube	1764 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.042$
$T = 296$ K	$\theta_{\text{max}} = 28.3^\circ$
φ and ω scans	$\theta_{\text{min}} = 2.5^\circ$
Absorption correction: part of the refinement model (ΔF) [Cubic fit to $\sin(\theta)/\lambda$ - 24 parameters; Parkin <i>et al.</i> , 1995]	$h = -37 \rightarrow 36$
$T_{\text{min}} = 0.497$, $T_{\text{max}} = 0.707$	$k = -11 \rightarrow 11$
14822 measured reflections	$l = -15 \rightarrow 15$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.038$	H-atom parameters constrained
$wR(F^2) = 0.093$	$w = 1/[\sigma^2(F_o^2) + (0.033P)^2 + 1.5317P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 0.98$	$(\Delta/\sigma)_{\text{max}} = 0.001$
3416 reflections	$\Delta\rho_{\text{max}} = 0.27 \text{ e } \text{\AA}^{-3}$
182 parameters	$\Delta\rho_{\text{min}} = -0.29 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement on $F^{2\wedge}$ for ALL reflections except those flagged by the user for potential systematic errors. Weighted R -factors wR and all goodnesses of fit S are based on $F^{2\wedge}$, conventional R -factors R are based on F , with F set to zero for negative $F^{2\wedge}$. The observed criterion of $F^{2\wedge} > \sigma(F^{2\wedge})$ is used only for calculating - R -factor-obs *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on $F^{2\wedge}$ are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.81917 (1)	0.43825 (6)	0.51473 (4)	0.1153 (2)
S1	0.62425 (3)	0.44674 (8)	0.15544 (6)	0.0547 (2)
O1	0.53281 (7)	0.4722 (2)	0.39261 (16)	0.0628 (7)

supplementary materials

O2	0.50891 (7)	0.2889 (2)	0.50674 (17)	0.0721 (8)
O3	0.61345 (8)	0.6071 (2)	0.13563 (17)	0.0713 (8)
O4	0.63014 (7)	0.3473 (2)	0.06120 (15)	0.0665 (7)
N1	0.58022 (7)	0.3824 (2)	0.22029 (18)	0.0555 (8)
C1	0.67716 (9)	0.4336 (3)	0.2571 (2)	0.0514 (9)
C2	0.71403 (11)	0.3319 (3)	0.2413 (3)	0.0659 (11)
C3	0.75605 (11)	0.3324 (4)	0.3185 (3)	0.0788 (14)
C4	0.76082 (11)	0.4310 (4)	0.4111 (3)	0.0705 (11)
C5	0.72375 (12)	0.5304 (4)	0.4281 (3)	0.0782 (14)
C6	0.68182 (12)	0.5303 (4)	0.3517 (3)	0.0714 (12)
C7	0.57612 (8)	0.2291 (3)	0.2636 (2)	0.0488 (9)
C8	0.55269 (8)	0.2054 (3)	0.3611 (2)	0.0505 (9)
C9	0.54903 (10)	0.0538 (4)	0.4003 (3)	0.0685 (11)
C10	0.56676 (11)	-0.0710 (4)	0.3463 (3)	0.0788 (14)
C11	0.58873 (11)	-0.0459 (3)	0.2499 (3)	0.0726 (11)
C12	0.59331 (10)	0.1022 (3)	0.2086 (3)	0.0606 (10)
C13	0.53137 (9)	0.3347 (4)	0.4206 (2)	0.0562 (10)
H1	0.55740	0.44700	0.22910	0.0670*
H2	0.71040	0.26350	0.17880	0.0790*
H2A	0.49850	0.36530	0.53760	0.1080*
H3	0.78130	0.26550	0.30760	0.0940*
H5	0.72720	0.59740	0.49150	0.0940*
H6	0.65640	0.59580	0.36380	0.0860*
H9	0.53410	0.03640	0.46550	0.0820*
H10	0.56390	-0.17150	0.37460	0.0940*
H11	0.60060	-0.13020	0.21220	0.0870*
H12	0.60810	0.11760	0.14300	0.0730*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0724 (2)	0.1642 (5)	0.1042 (4)	-0.0277 (2)	-0.0048 (2)	0.0332 (3)
S1	0.0662 (4)	0.0525 (4)	0.0493 (4)	0.0040 (3)	0.0221 (3)	0.0044 (3)
O1	0.0667 (12)	0.0678 (13)	0.0586 (12)	0.0035 (10)	0.0251 (10)	0.0005 (10)
O2	0.0768 (13)	0.0831 (13)	0.0641 (13)	-0.0003 (11)	0.0362 (11)	0.0049 (11)
O3	0.0920 (14)	0.0549 (12)	0.0700 (14)	0.0115 (10)	0.0218 (11)	0.0149 (10)
O4	0.0860 (13)	0.0710 (12)	0.0478 (11)	-0.0033 (10)	0.0273 (10)	-0.0051 (10)
N1	0.0548 (12)	0.0553 (13)	0.0600 (15)	0.0105 (10)	0.0206 (11)	0.0067 (11)
C1	0.0595 (15)	0.0492 (15)	0.0499 (16)	-0.0006 (12)	0.0226 (13)	0.0021 (13)
C2	0.0743 (19)	0.0599 (18)	0.066 (2)	0.0077 (15)	0.0182 (16)	-0.0043 (15)
C3	0.069 (2)	0.084 (2)	0.086 (3)	0.0129 (17)	0.0198 (19)	0.006 (2)
C4	0.0598 (17)	0.089 (2)	0.064 (2)	-0.0170 (17)	0.0137 (15)	0.0162 (18)
C5	0.076 (2)	0.100 (3)	0.061 (2)	-0.0116 (19)	0.0177 (18)	-0.0176 (18)
C6	0.074 (2)	0.079 (2)	0.065 (2)	0.0075 (16)	0.0227 (17)	-0.0166 (17)
C7	0.0403 (13)	0.0554 (16)	0.0510 (16)	-0.0008 (11)	0.0079 (12)	-0.0010 (13)
C8	0.0388 (12)	0.0612 (17)	0.0529 (16)	-0.0032 (12)	0.0112 (11)	0.0012 (14)
C9	0.0602 (17)	0.073 (2)	0.077 (2)	-0.0094 (15)	0.0259 (16)	0.0083 (18)
C10	0.074 (2)	0.0595 (19)	0.106 (3)	-0.0088 (16)	0.024 (2)	0.0123 (19)

C11	0.076 (2)	0.0557 (19)	0.090 (2)	-0.0043 (15)	0.0251 (18)	-0.0065 (17)
C12	0.0607 (17)	0.0576 (18)	0.0665 (19)	-0.0056 (13)	0.0196 (14)	-0.0032 (15)
C13	0.0402 (13)	0.080 (2)	0.0490 (16)	-0.0046 (13)	0.0080 (12)	0.0015 (15)

Geometric parameters (\AA , $^{\circ}$)

Br1—C4	1.882 (3)	C7—C12	1.382 (4)
S1—O3	1.4184 (19)	C7—C8	1.404 (3)
S1—O4	1.4186 (19)	C8—C13	1.476 (4)
S1—N1	1.625 (2)	C8—C9	1.386 (4)
S1—C1	1.758 (3)	C9—C10	1.368 (5)
O1—C13	1.225 (4)	C10—C11	1.371 (5)
O2—C13	1.317 (3)	C11—C12	1.370 (4)
O2—H2A	0.8200	C2—H2	0.9300
N1—C7	1.418 (3)	C3—H3	0.9300
N1—H1	0.8600	C5—H5	0.9300
C1—C6	1.371 (4)	C6—H6	0.9300
C1—C2	1.379 (4)	C9—H9	0.9300
C2—C3	1.371 (5)	C10—H10	0.9300
C3—C4	1.363 (5)	C11—H11	0.9300
C4—C5	1.374 (5)	C12—H12	0.9300
C5—C6	1.364 (5)		
Br1···C9 ⁱ	3.662 (3)	C12···O4	2.987 (3)
Br1···C10 ⁱ	3.540 (3)	C13···N1 ^{vi}	3.325 (3)
Br1···C11 ⁱ	3.599 (3)	C13···O3 ⁱⁱⁱ	3.182 (3)
S1···H12	2.8600	C13···C7 ^{vi}	3.543 (3)
O1···N1	2.670 (3)	C2···H5 ^v	3.0500
O1···O2 ⁱⁱ	2.703 (3)	C5···H2 ^{ix}	3.0900
O2···O3 ⁱⁱⁱ	3.205 (3)	C6···H3 ^{ix}	3.0200
O2···O1 ⁱⁱ	2.703 (3)	C9···H9 ^{vii}	3.0700
O3···C11 ^{iv}	3.369 (3)	C13···H1	2.6300
O3···C13 ^v	3.182 (3)	C13···H1 ^{vi}	2.9700
O3···O2 ^v	3.205 (3)	C13···H2A ⁱⁱ	2.7700
O4···C6 ^v	3.185 (4)	H1···O1	2.1300
O4···C5 ^v	3.384 (4)	H1···C13	2.6300
O4···C12	2.987 (3)	H1···O1 ^{vi}	2.7100
O1···H1	2.1300	H1···C13 ^{vi}	2.9700
O1···H2A ⁱⁱ	1.8900	H2···O4	2.5500
O1···H1 ^{vi}	2.7100	H2···H5 ^v	2.5900
O2···H9	2.3500	H2···C5 ^x	3.0900
O2···H10 ^{vii}	2.8000	H2A···O1 ⁱⁱ	1.8900
O3···H11 ^{iv}	2.4700	H2A···C13 ⁱⁱ	2.7700
O3···H6	2.7600	H2A···H2A ⁱⁱ	2.4700
O4···H12	2.3100	H3···C6 ^x	3.0200

supplementary materials

O4···H2	2.5500	H5···C2 ⁱⁱⁱ	3.0500
O4···H6 ^v	2.5600	H5···H2 ⁱⁱⁱ	2.5900
N1···O1	2.670 (3)	H6···O3	2.7600
N1···C13 ^{vi}	3.325 (3)	H6···O4 ⁱⁱⁱ	2.5600
C5···O4 ⁱⁱⁱ	3.384 (4)	H9···O2	2.3500
C6···O4 ⁱⁱⁱ	3.185 (4)	H9···C9 ^{vii}	3.0700
C7···C13 ^{vi}	3.543 (3)	H9···H9 ^{vii}	2.2500
C9···Br1 ⁱ	3.661 (3)	H10···O2 ^{vii}	2.8000
C10···Br1 ⁱ	3.540 (3)	H11···O3 ^{viii}	2.4700
C11···Br1 ⁱ	3.599 (3)	H12···S1	2.8600
C11···O3 ^{viii}	3.369 (3)	H12···O4	2.3100
O3—S1—O4	120.06 (12)	C9—C8—C13	119.7 (2)
O3—S1—N1	104.36 (12)	C8—C9—C10	122.2 (3)
O3—S1—C1	108.04 (13)	C9—C10—C11	119.2 (3)
O4—S1—N1	109.57 (11)	C10—C11—C12	120.6 (3)
O4—S1—C1	107.82 (12)	C7—C12—C11	120.7 (3)
N1—S1—C1	106.21 (11)	O1—C13—O2	121.8 (3)
C13—O2—H2A	109.00	O1—C13—C8	124.5 (2)
S1—N1—C7	125.76 (16)	O2—C13—C8	113.7 (3)
C7—N1—H1	117.00	C1—C2—H2	120.00
S1—N1—H1	117.00	C3—C2—H2	120.00
S1—C1—C2	121.2 (2)	C2—C3—H3	120.00
S1—C1—C6	118.6 (2)	C4—C3—H3	120.00
C2—C1—C6	120.2 (3)	C4—C5—H5	120.00
C1—C2—C3	119.5 (3)	C6—C5—H5	120.00
C2—C3—C4	120.0 (3)	C1—C6—H6	120.00
Br1—C4—C3	120.5 (2)	C5—C6—H6	120.00
Br1—C4—C5	118.9 (3)	C8—C9—H9	119.00
C3—C4—C5	120.6 (3)	C10—C9—H9	119.00
C4—C5—C6	119.7 (3)	C9—C10—H10	120.00
C1—C6—C5	120.0 (3)	C11—C10—H10	120.00
C8—C7—C12	119.6 (2)	C10—C11—H11	120.00
N1—C7—C12	120.9 (2)	C12—C11—H11	120.00
N1—C7—C8	119.5 (2)	C7—C12—H12	120.00
C7—C8—C9	117.9 (2)	C11—C12—H12	120.00
C7—C8—C13	122.5 (2)		
O3—S1—N1—C7	-176.5 (2)	Br1—C4—C5—C6	177.9 (3)
O4—S1—N1—C7	53.8 (2)	C3—C4—C5—C6	0.0 (5)
C1—S1—N1—C7	-62.4 (2)	C4—C5—C6—C1	-1.3 (5)
O3—S1—C1—C2	-135.0 (2)	N1—C7—C8—C9	179.5 (2)
O4—S1—C1—C2	-3.8 (3)	N1—C7—C8—C13	0.7 (3)
N1—S1—C1—C2	113.6 (2)	C12—C7—C8—C9	1.8 (4)
O3—S1—C1—C6	43.2 (3)	C12—C7—C8—C13	-177.0 (2)
O4—S1—C1—C6	174.3 (2)	N1—C7—C12—C11	-179.2 (3)
N1—S1—C1—C6	-68.3 (3)	C8—C7—C12—C11	-1.6 (4)
S1—N1—C7—C8	149.12 (19)	C7—C8—C9—C10	-0.9 (4)

S1—N1—C7—C12	−33.3 (3)	C13—C8—C9—C10	177.9 (3)
S1—C1—C6—C5	−175.6 (3)	C7—C8—C13—O1	−1.4 (4)
S1—C1—C2—C3	175.7 (2)	C7—C8—C13—O2	176.9 (2)
C6—C1—C2—C3	−2.4 (4)	C9—C8—C13—O1	179.9 (3)
C2—C1—C6—C5	2.5 (5)	C9—C8—C13—O2	−1.9 (3)
C1—C2—C3—C4	1.1 (5)	C8—C9—C10—C11	−0.3 (5)
C2—C3—C4—C5	0.1 (5)	C9—C10—C11—C12	0.6 (5)
C2—C3—C4—Br1	−177.7 (2)	C10—C11—C12—C7	0.3 (5)

Symmetry codes: (i) $-x+3/2, -y+1/2, -z+1$; (ii) $-x+1, -y+1, -z+1$; (iii) $x, -y+1, z+1/2$; (iv) $x, y+1, z$; (v) $x, -y+1, z-1/2$; (vi) $-x+1, y, -z+1/2$; (vii) $-x+1, -y, -z+1$; (viii) $x, y-1, z$; (ix) $-x+3/2, y+1/2, -z+1/2$; (x) $-x+3/2, y-1/2, -z+1/2$.

Hydrogen-bond geometry (\AA , °)

$D—H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
N1—H1 \cdots O1	0.86	2.13	2.670 (3)	121
O2—H2A \cdots O1 ⁱⁱ	0.82	1.89	2.703 (3)	173
C6—H6 \cdots O4 ⁱⁱⁱ	0.93	2.56	3.185 (4)	125
C11—H11 \cdots O3 ^{viii}	0.93	2.47	3.369 (3)	164
C12—H12 \cdots O4	0.93	2.31	2.987 (3)	130

Symmetry codes: (ii) $-x+1, -y+1, -z+1$; (iii) $x, -y+1, z+1/2$; (viii) $x, y-1, z$.

supplementary materials

Fig. 1





Fig. 2



supplementary materials

Fig. 3

