

Nuclear Factor κB Signaling and Its Related Non-coding RNAs in Cancer Therapy

Xiaomin Liu,^{1,2,3} Yang Shao,^{2,3} Jinbao Zhou,² Guangren Qian,¹ and Zhongliang Ma²

¹School of Environmental and Chemical Engineering, Shanghai University, Shanghai 200444, China; ²Lab for Noncoding RNA & Cancer, School of Life Sciences, Shanghai University, Shanghai 200444, China

Nuclear factor κB (NF-κB) acts as a nuclear factor that is composed of five main subunits. It is a pluripotent and crucial dimer transcription factor that has a close relationship with many serious illnesses, especially its influences on cell proliferation, inflammation, and cancer initiation and progression. NF-κB acts as part of the signaling pathway and determines its effect on the expression of several other genes, such as epidermal growth factor receptor (EGFR), p53, signal transducer and activator of transcription 3 (STAT3), and non-coding RNA (ncRNA). Continuous activation of the NF-κB signaling pathway has been seen in many cancer types. While the NF-κB signaling pathway is tightly regulated in physiological settings, quite frequently it is constitutively activated in cancer, and the molecular biology mechanism underlying the deregulated activation of NF-κB signaling remains unclear. In this review, we discuss the regulatory role and possible clinical significance of ncRNA (microRNA [miRNA] and long non-coding RNA [lncRNA]) in NF-κB signaling in cancer, including in the conversion of inflammation to carcinogenesis. Non-coding RNA plays an essential and complex role in the NF-κB signaling pathway. NF-κB activation can also induce the ncRNA status. Targeting NF-κB signaling by ncRNA is becoming a promising strategy of drug development and cancer treatment.

INTRODUCTION

Cancer is a malignancy disease, which results in aberrant and amplification function of several oncogenes.¹ Many studies have shown that the occurrence and development process of tumors are related to inflammation, and activation of the inflammation-related signaling pathway plays vital roles in the tumor microenvironment. It is worth noting that the tumor cells also have abnormal activation of the inflammatory signaling pathway independent of external stimuli. Nuclear factor κB (NF-κB) signaling pathway activation promotes the transcription of inflammatory factors, chemokines, adhesion molecules, growth factors, and other related genes, which in turn leads to tumor development.²

The transcription factor NF-κB, a nuclear factor that binds to the enhancer of the immunoglobulin kappa light-chain of activated B cell tumors, was discovered in 1986.^{3,4} NF-κB has also been presented as being closely connected with cancer development and progression. It induces the expression of many genes involved in the develop-

mental process, the inflammatory response, immunoreaction, as well as cellular proliferation apoptosis and angiogenesis.^{5–7}

At present, non-coding RNA (ncRNA) and NF-κB act as crucial factors in the development of cancer, both of which can be applied as tumor markers and drug targets in clinical therapeutics.⁸ ncRNAs mainly contain microRNA (miRNA) and long non-coding RNA (lncRNA). In addition, there are other ncRNAs, including such circular RNA (circRNA) and piwi-interacting RNA (piRNA), among others.

miRNAs are a class of endogenous regulatory ncRNAs involved in cell carcinogenesis that are about 20–25 nt in length.⁹ They can bind to the 3' UTR of protein-coding mRNAs, typically leading to mRNA degradation. Alternatively, they can also cause repression of target mRNA translation.¹⁰ miRNA cleavage is related to cancer progression and clinical features. Thus, as an important regulator of life processes, miRNAs have a close relationship with tumors.¹¹

lncRNA is a type of ncRNA with a length >200 nt. It has been shown that lncRNAs participate in a variety of biological processes, such as regulation of gene, epigenetic, cell cycle and cell differentiation.¹² Therefore, a more in-depth study of the functions and mechanisms of lncRNA will help reveal their role in tumorigenesis as a primary tool for diagnosis and cancer treatment.

In this review, we provide a highlighted overview of the impact of miRNAs and lncRNAs on the inflammatory regulation of the NF-κB signaling pathway in cancer.

NF-κB Signaling Pathway and Its Activation

The NF-κB transcription factor family is composed of five main subunits, including Rel (c-Rel), RelB, p65 (RelA, NF-κB3), p50 (NF-κB1), and p52 (NF-κB2), all of which have an N-terminal segment containing about 300 aa residues called Rel homology domain (RHD)

<https://doi.org/10.1016/j.omtn.2019.11.007>.

³These authors contributed equally to this work

Correspondence: Zhongliang Ma, Lab for Noncoding RNA & Cancer, School of Life Sciences, Shanghai University, Shanghai 200444, China.

E-mail: zlma@shu.edu.cn

Correspondence: Guangren Qian, School of Environmental and Chemical Engineering, Shanghai University, Shanghai 200444, China.

E-mail: grqian@shu.edu.cn






Figure 1. Differences in the Structures of NF-κB Family Members

(Figure 1). RHD contains the binding domain of DNA, dimerization domain, and the inhibitor of κB (IkB) binding domain besides the nuclear localization signal (NLS). Among them, p65, c-Rel, and RelB possess C-terminal transactivation domains (TADs) that endow the ability to activate NF-κB, thereby positively modulating its expression. Although p50 and p52 lack the transcription activation domain, their homodimers can inhibit transcription.¹³ These family members can form many homodimers or heterodimers, among which the p50/p65 heterodimers play a major role in physical function.^{14,15} Different NF-κB/Rel dimers have varied binding sites, providing them with diverse properties. As a consequence, different target genes are distinctively induced by different NF-κB dimers and play different functions.

Activation of NF-κB has three forms: release from the IkB molecules, cleavage of the inhibitory ankyrin repeat domains of p100 and p105, and activation of inhibitor of NF-κB kinase (IKK) through the CBM (CARMA1, BCL10, and MALT1) complex. In general, the NF-κB signaling pathway is split into the canonical and the non-canonical NF-κB pathway (Figure 2).

Canonical NF-κB Pathway

(1) In the cytoplasm, NF-κB will bind to IkB as a trimer when in the inert state. (2) Upon recognition of external stimulation such as bacteria, viruses, parasites, inflammatory cytokines, and fibroblast growth factors, the tumor necrosis factor receptor (TNFR), Toll-like receptor (TLR), and interleukin-1 receptor (IL-1R) will be activated, which myeloid differentiation factor 88 (MyD88) and IL-1R-associated kinase (IRAK) will bind and pass the activated signal to downstream proteins as important adaptors. The signal passes through the downstream proteins TNFR-associated factor (TRAF)2, NF-κB inducing kinase (NIK), and receptor interacting protein (RIP) and consequently activates IKK, leading to ubiquitination and phosphorylation of IkB, which further causes the release of NF-κB dimers, and the dissociated IkB molecules are degraded by proteasomes. (3) The NF-κB dimers enter the nucleus and bind to the κB domain of the target genes, making a difference in gene expression.

Non-canonical NF-κB Pathway

The non-canonical NF-κB pathway is composed of IkB family members, the main parts of which are p100 and p105.

Upon stimulation, the B cell-activating factor receptor (BAFR), lymphotxin β receptor (LTβR), receptor activator of NF-κB (RANK), and CD40 will be activated, a consequence of which is that p100 will be phosphorylated by the downstream NIK, eventually leading to the ubiquitination of p100 and its degradation into p52. The free p52-RelB dimers will enter the nucleus and bind to the target genes of the κB domain.

NF-κB Signaling and miRNA in Cancer

In tumorigenesis, abnormal expression of miRNA is usually accompanied by activation of the NF-κB signaling pathway. miRNAs can regulate the NF-κB pathway through translocating themselves to some upstream signal molecules or NF-κB members. Meanwhile, the NF-κB pathway can also regulate the expression level of miRNAs by synthesis of proteins.¹⁶


p50/p65 and miRNAs

Studies have shown that the synthesis of miR-9 is under the control of pro-inflammatory cytokine TNF-α and IL-1β regulation, whereas the precursor of p50, p105, is actually a target protein of miR-9. They also suggest that miR-9 will function as a trimming agent in the anti-inflammatory responses to hinder the negative control by p50 homodimers in some monocytes. In addition, miR-9 can inhibit the development of ovarian cancer and gastric cancer through regulation of the NF-κB pathway.¹⁷⁻¹⁹ Moreover, researchers have discovered that miR-143 can function as the transcription target of NF-κB and promote metastasis in lung cancer.²⁰ Meanwhile, overexpression of miR-143 will decrease the viability of the cells and have an effect on the expression of p65 in the NF-κB pathway.²¹ Besides, there are plenty of miRNAs found to have a direct or indirect regulatory function on p50/p65. This can explain that NF-κB does not work alone, but rather that it forms part of a network with a variety of miRNAs.

One study testified that miR-145-5p is a tumor inhibitor,²² during regulation of the NF-κB/YY1 pathway, whose aberrant expression will lead to the synthesis of rhabdomyosarcoma.²³ In addition, a study by Mott et al.²⁴ also verified that the inhibitory effect NF-κB has on miR-29b exists in the biliary duct and cholangiocarcinoma. Certainly, a series of miRNAs such as miR-125b-1, miR-125a-5p, miR-494, and miR-130b are under the regulation of NF-κB²⁵⁻²⁹ (Table 1), which determines the mode of its effects on the expression of several other genes and thus its function (Figure 3).

TNF-α and miRNAs

TNF-α is a secreted proinflammatory cytokine. Chen et al.³⁰ found that miR-27a can protect human mitral valve interstitial cells from

**Figure 2. NF-κB Signaling Pathway**

(Left) Canonical pathway of NF-κB activation. This pathway is mediated by the IκB kinase (IKK) complex, leading to phosphorylation and degradation of IκB. (Right) Non-canonical pathway of NF-κB activation. This pathway involves IKK α activation via NF-κB-inducing kinase (NIK) and induces the phosphorylation of p100, leading to the generation and translocation of p52.

TNF- α induced through the upregulation of NELL-1 in patients with heart valve disease. Overexpression of miR-27a protected human mitral valve interstitial cells (hMVICs) from TNF- α -induced cellular injury, indicating that miR-27a has a vital role in the regulation of TNF- α release. To date, miR-27a is considered as an important negative regulatory factor in human inflammations, which has a close connection with TNF- α .

Additionally, studies have shown that miR-146a has a connection with TNF- α in gastric diseases and psoriasis patients.^{31,32} miR-146a can increase the secretion of TNF- α , which is confirmed by experiments. Compared with the control group, the overexpression of miR-146a significantly upregulated the expression level of TNF- α in a gastritis group, while that of the gastric cancer group was significantly reduced.

Thus, TNF- α and NF-κB have a close interaction in which miRNAs have an essential role. Some miRNAs reacted with TNF- α in inflammation, while TNF- α has a vital joint role in NF-κB pathway regulation.^{33,34} p50 and p65 regulate the expression and secretion of TNF- α indirectly through interacting with important members of NF-κB pathway.³⁵

IκB and miRNAs

The regulation of IκB by miRNAs is not direct but instead requires the activation of IKK. The IKK family members mainly include IKK α , IKK β , IKK γ , and IKK ϵ . First, miRNAs bind with the IKK complex to activate IKK, causing phosphorylation of IκB, after which IκB separates from the p50/p65/IκB trimers and is ubiquitinated. Then, NF-κB switches from an inert state to an activated state, activating a series of subsequent reactions.³⁶

Studies have shown that during the differentiation of human macrophages, the expression levels of miR-223, miR-23a, miR-15a, and

miR-16 decrease rapidly, which directly leads to the increase of IKK α expression and causes an elevated expression level of p52.

Overexpression of p52 will remarkably inhibit the foundation level of target genes in the NF-κB pathway. When miRNA expression decreases, the activation of macrophages is apparently inhibited.^{37,38} In tumors with NF-κB activity, miR-199a targets IKK β and inhibits its expression in endometrial stromal cells (ESCs).³⁹ As the IKK β expression decreases, miR-199a suppresses activation of the NF-κB pathway and the synthesis of IL-8, which means that miR-199a can reduce the invasiveness of ESCs through inhibiting the NF-κB pathway.^{39,40} IKK ϵ also had a contradictory role in activation of the NF-κB pathway. Studies have verified that miR-155 is involved in the negative-feedback loop through down-regulation of IKK ϵ , IKK β and other transcription factor of NF-κB.^[41] Additionally, building the regulatory feedback loop of IKK ϵ /YAP1/miR-Let-7b/I, IKK ϵ promotes glioblastoma progression.⁴²

MyD88 and miRNAs

d'Adhemar et al.⁴³ found that miR-146a and miR-21 bind to the mRNA of MyD88 in ovarian cancer, influencing the regulatory function of MyD88 in the TLR4 pathway, whose activation does not directly promote the activation of downstream NF-κB. Meanwhile, they found that the regulatory function of miR-146a and miR-21 on MyD88 is not only limited in activating the TLR4 and NF-κB pathways, but also in regulating the expression level of MyD88 to directly influence the sensitivity of ovarian cancer cells to pharmacotherapy. miR-200b and miR-200c also play a significant role in activation of the TLR4 and NF-κB pathway, which is to bind with MyD88.⁴⁴ They observed that after transfecting miR-200b and miR-200c into HEK293-TLR4 cells, the expression of MyD88 clearly decrease, while other target genes such as IRAK-1 and TRAF6 have no effect. This shows that miR-200b and miR-200c can influence the TLR4 pathway through regulation of MyD88, which has vital physiological significance in the defense of microbial pathogens.^{44,45}

In short, miRNAs can adjust NF-κB efficiently through MyD88, which has a crucial role in the biogenesis and development of inflammations and tumors.

**Table 1. miRNAs and the NF-κB Signaling Pathway**

miRNA	Target Gene	Disease/Cancer	Function	Reference
miR-302e	RelA	allergic inflammation	anti-inflammatory	106
miR-194, miR-195	TRAF6, IKK α	intervertebral disc degeneration, Wilms tumor	anti-inflammatory proliferation, apoptosis	107,108
miR-199a-3p	IKK β	cystic fibrosis	chronic pulmonary inflammation	109
miR-23a	IKK β	articular chondrocytes	anti-inflammatory	38
miR-429	IKK β	cervical cancer	proliferation, apoptosis	110
miR-7	Myd88	<i>Eriocheir sinensis</i>	enhance white spot syndrome	111
let-7b	IKK ϵ	human glioma	invasion, migration	112
miR-342-3p	IKK γ	hepatocellular carcinoma	proliferation	113
miR-503	IKK β	NSCLC	metastasis	114
miR-451	IKK β	hepatocellular carcinoma	proliferation	115
miR-199a	IKK β	endometrial stromal cell	cell adhesion, migration, invasion	39
miR-194, miR-502-5p	TRAF6, TRAF2	nucleus pulposus cells of the intervertebral disc, chondrocyte injury	anti-inflammatory proliferation, apoptosis	107,116
miR-18a-5p	IRF2	NSCLC	proliferation, metastasis, apoptosis	61
miR-708-3p	STAT3	idiopathic pulmonary fibrosis	inhibit fibrosis	75
miR-411	STAT3	cervical cancer	cell proliferation and invasion	117
miR-29a	STAT3	human retinoblastoma	cell proliferation, migration, invasion, apoptosis	118
miR-124	STAT3	NSCLC	proliferation, apoptosis	119
miR-181b	STAT3	atherosclerosis	proliferation, apoptosis, cell cycle	120
miR-26b	TAK1	hepatocellular carcinoma	apoptosis	49
miR-20a	TAK1	osteosarcoma	proliferation	121
miR-23b	TRAF1	sepsis	proliferation, immunosuppression	122

NSCLC, non-small cell lung cancer.

CBM Complex and miRNAs

The CBM complex plays an essential role mainly by modulating nuclear transcription factor activation, such as NF-κB, which is initiated by antigen receptor activation. CARMA1 is a cytoskeletal protein that changes its conformation after antigen receptor activation and recruits downstream signaling pathway components BCL10 and MALT1. The mutual recognition of CARMA1 and the linker protein BCL10 is mediated by the CARD domain, whereas BCL10 and MALT1 are cytoplasmic. The constitutive complex forms a complex, and the interaction between these three proteins stabilizes the CBM complex. The CBM complex also has a regulatory effect on miRNA. Recent studies have shown that miR-181d acts as a tumor suppressor by targeting MALT1 to suppress glioblastoma through NF-κB pathways.⁴⁶ Additionally, miR-26 restrains TNF- α /NF-κB signaling and IL-6 expression by silencing MALT1.⁴⁷ Gu and Sun⁴⁸ found that miR-539 can regulate migration and invasion through targeting CARMA1 in human thyroid cancer.

TAK1 and miRNAs

Zhao et al.⁴⁹ found that miR-26b can inhibit activation of the NF-κB pathway by targeting transforming growth factor β (TGF- β)-activated kinase-1 (TAK1) and TAK1 binding protein (TAB3). They observed that in the hepatocellular carcinoma (HCC) cell lines QGY-7703 and MHCC-97H, miR-26b can bind directly to the 3'


UTR of TAK1 and TAB3 to suppress their expression, inhibiting the TNF- α -induced NF-κB pathway and eventually enhancing the sensitivity of lung cancer cells to pharmacotherapy. TAK1 has a special and important position in the NF-κB pathway, which can bind to TAB1 to activate NF-κB-induced kinase and IκB kinase.⁵⁰

Other NF-κB-Associated Components and miRNAs Involved

There is increasing evidence that miRNAs play a crucial part in the regulation of NF-κB signaling pathways in a variety of cancer types. In addition to the genes mentioned above, other NF-κB-associated genes and miRNAs are involved in the regulation of NF-κB signaling pathways in cancer.

Epidermal Growth Factor Receptor

As a vital regulator, epidermal growth factor receptor (EGFR) has a pivotal effect on the NF-κB signaling pathway and cancer. In some ways, EGFR and NF-κB are partners in cancer.^{51,52} For example, EGFR can indirectly or directly activate signal transducer and activator of transcription (STAT) members.⁵³ NF-κB mediated extracellular signal-regulated kinase (ERK) reactivation through gefitinib resistance in EGFR mutant non-small cell lung cancer (NSCLC) cells.⁵⁴ Studies have elucidated that NF-κB is activated by EGFR activation.⁵⁵⁻⁵⁷ In addition, environmental factors can also affect EGFR expression through regulating NF-κB.⁵⁸ Our laboratory has proved that miR-34a

**Figure 3. miRNA and NF-κB Signaling Pathway**

Relationship between miRNAs and the NF-κB signaling pathway.

ally, through suppressing the STAT3 signaling pathway activation, miR-125a-5p heightens the sensitivity of cisplatin in esophageal squamous carcinoma.⁷⁸ Meanwhile, miR-26a-5p regulates ITGB8-JAK2/STAT3 to potentiate metastasis of lung cancer.⁷⁹ STAT3 and NF-κB co-regulate a series of target genes, including cell cycle and anti-apoptotic genes. Research has shown that STAT3 regulates post-transcriptional translation of p65 by regulating the expression of acetyltransferase p300, thereby promoting pro-inflammatory cytokines in tumor microenvironments.⁸⁰

Other ncRNAs and NF-κB Signal Transduction

lncRNA has been found to play a vital part in pathological and physiological processes, containing tumor formation and metastasis.

Different lncRNAs have different molecular mechanisms that play different biological functions.⁸¹ The activation of NF-κB is also related with lncRNA.

Tumor-associated NF-κB activation and lncRNA overexpression are most directly related to the inhibition of IκB, which acts as a negative regulator of NF-κB signaling. Liu et al.⁸² found that NKILA (NF-κB interacting lncRNA) binds to NF-κB, which is upregulated by inflammatory factors in breast cancer. NKILA inhibits activation of the NF-κB signaling pathway by masking the location of phosphorylation of IκB for IKK phosphorylation suppression. Further studies revealed that there are two hairpin structures, A and B, at 300–500 nt of NKILA. Hairpin A binds to the DNA-binding region of NF-κB, and hairpin B binds to S51-R73 of p65, preventing IκBα detachment that forming a stable NKILA/NF-κB/IκBα complex. Yang et al.⁸³ found that FTH1P3 (long non-protein coding RNA ferritin heavy chain 1 pseudogene 3) regulated metastasis and invasion through SP1 (specificity protein 1)/NF-κB (p65) in esophageal squamous cell carcinoma (ESCC). The mechanism of lncRNAs has been studied. A deeper understanding of the molecular mechanisms and biological functions of lncRNA will help to find new effective anticancer strategies in tumorigenesis.

and miR-107-5p could directly target EGFR to reduce cell proliferation, metastasis, cell cycle, and promote apoptosis *in vitro* and *vivo*.^{59,60} Moreover, our laboratory has also verified that miR-18a-5p can downregulate the expression of interferon regulatory factor 2 (IRF2), which is related to the NF-κB signaling pathway.⁶¹

There is a mutual inhibitory relationship between NF-κB and p53. Studies have shown that p65 can inhibit p53-related transcriptional activation, and p53 can also inhibit NF-κB transcriptional activity.⁶² Mutant p53 can increase p52 expression through acetylation by regulating the amount of histone acetyltransferase CBP.⁶³ In addition, the interaction of NF-κB, such as TNF-α, p53, and NF-κB, with certain specific stimuli also plays a vital role. The miR-34 family is the first miRNA associated with p53.⁶⁴ It was shown that upregulated expression of the p53-binding region will increase all members of miR-34 family, which also implies an important function of the miR-34 family in the p53 signaling pathway. The miRNA family is also significant in the anti-proliferative and pro-apoptotic processes regulated by p53 through binding cell cycle genes and proto-oncogenes.^{65–68} In other studies, miR-142-3p, miR-155-5p, miR-518, miR-211, miR-1307, and miR-30a can all directly or indirectly stimulate the p53 signaling pathway, which shows the close relationship between miRNA and p53.^{69–74}

STAT3

STAT3 activation is responsible for a variety of genes that promote diagnostic markers and therapeutic targets for many diseases.^{75,76} A study has shown that miR-124 inhibits the growth of colon cancer through targeting STAT3.⁷⁷ This shows that the tumor-suppressing effect of miR-124 is based on the combination with STAT3. Addition-

miRNAs and lncRNAs have been demonstrated to play an irreplaceable role in the NF-κB signaling pathway. In addition, other types of ncRNAs, such as circRNA and piRNA, are also involved in the regulation of tumor-associated signaling pathways. Studies have shown that miR-7 sponge circRNA (ciRS-7) has more than 70 conserved



miR-7 binding sites, which can effectively sponge miR-7 and suppress the inhibition of miR-7 target genes, promoting the correlation of gene expression and activating NF-κB signaling pathways.^{84,85} There are few reports on piRNA in cancer. Leng et al.⁸⁶ found that piR-DQ590027, miR-377, and miR-153 in glioma-conditioned endothelial cells (GECs) had lower expression. piR-DQ590027 bound to MIR17HG. FOXR2 was a downstream target of miR-377 and miR-153. This study proved that the piR-DQ590027/MIR17HG/miR-153(miR-377)/FOXR2 pathway plays a crucial role in regulating the permeability of glioma-conditioned normal blood-brain barrier (BBB). Another study has shown that piR-021285 can induce the methylation of genes in breast cancer.⁸⁷ The roles of these ncRNAs in tumors remain to be further explored.

NF-κB Signaling in Chronic Inflammation to Cancer

Since the discovery of NF-κB more than 30 years ago, researchers have demonstrated that NF-κB is the “master switch” of gene expression in various pro-inflammatory mediators.⁸⁸ It is expressed in many cancers, including HCC, colon cancer, breast cancer, prostate cancer, cervical cancer, and chronic lymphocytic leukemia.^{89–92} NF-κB has an important function in inflammation by regulating the expression of pro-inflammatory cytokines such as TNF, IL-1, and monocyte chemoattractant protein 1 (MCP-1).⁹³ Drugs such as aminosalicylates and lipopolysaccharide, which are used clinically to combat inflammation work by inhibiting NF-κB. At the same time, many studies have verified that these drugs can inhibit the activation of some cancers.^{94,95} Although NF-κB activation can prevent self-activated apoptosis, it acts as a transcription factor that regulates the production of cytotoxic drugs such as nitric oxide, which in turn may indirectly lead to apoptosis in other cells.⁹⁶ Environmental factors affect NF-κB activation and inflammation secretion, which induce to promote lung tumor growth cancer.⁹⁷ Particularly, in glioblastoma, it produces neurotoxic reactive oxygen species, which induces by cytokines through NF-κB activation. Moreover, NF-κB and STAT3 synergistically regulate a number of target genes affecting growth, angiogenesis, and immunosuppression.⁹⁸ In addition, they also collaboratively control a common set of genes encoding for cytokines.⁹⁹ This means the mechanism of NF-κB activity regulated by STAT3 influences and plays a significant role in cancer.

Studies have shown that the occurrence and development of cancers are closely related to inflammation, and activation of the inflammation-related signaling pathway plays a crucial role in formation of the tumor microenvironment. It is worth noting that cancers also have abnormal activation of the inflammatory signaling pathway independent of external stimuli. Activation of the NF-κB signaling pathway promotes the transcription of inflammatory factors, chemokines, adhesion molecules, growth factors, and other related genes, which in turn leads to tumor development.²

Abnormal activation of inflammatory signaling pathways in tumors can promote the expression of cellular inflammatory factors or activate genes associated with tumor progression. At the same time, inflammatory factors can accelerate the development of malignant cells

through paracrine effects. For instance, low expression of miR-892b leads to aberrant activation of the NF-κB signaling pathway, increasing tumor cell proliferation, metastasis, and angiogenesis in breast cancer.¹⁰⁰ In addition, lncRNA-SRLR binds to the IL-6 promoter region, activating the STAT3 signaling pathway in tumor cells by autocrine or paracrine secretion.¹⁰¹ Tumor cell inflammation is similar to a series of phenotypes in traditional inflammatory cells that participate in immune responses. It has the characteristics of rapid proliferation, anti-apoptosis, invasion and migration, and secretion of inflammatory factors. At present, most of the evidence indicates that inflammation in tumors is one of the vital causes of tumor deterioration.

Conclusion and Future Perspectives

In cancer, activation of the NF-κB signaling pathway is ubiquitous. Its activation is not caused by stimulation of external pathogens; instead, it is caused by the abnormal regulation of cells. Previous studies have found that ncRNA may be involved in this regulation process. Compared with protein, the production of RNA products in cells is faster. When cells are stimulated by some factors, DNA rapidly transcribes RNA and undergoes splice processing, without the need for protein modification to function. Different RNA products appear in different splicing modes and perform different functions. Meanwhile, RNA has more modifications than proteins, which have an effect on the subcellular localization, activity, and structure of RNA that makes RNA more diverse in functioning. These properties of RNA determine their status in regulating the signaling pathway. RNA participates in regulation of the NF-κB signaling pathway, which requires more in-depth research and is of great significance for finding new cancer therapies.

Activation of NF-κB signaling can cause cells to be resistant to chemotherapeutic drugs and inhibit cell apoptosis.¹⁰² Combined with the effects of NF-κB signaling on tumor formation and drug resistance, targeting NF-κB can effectively inhibit the development of tumors and alleviate the tolerance to chemotherapeutic drugs caused by the activation of NF-κB signaling.

As a vital transcriptional regulator, miRNA, a non-coding protein single-chain small molecule RNA, has been widely demonstrated to be closely related to the development of various tumors and potential therapeutic approach.¹⁰³ miRNA has great clinical significance for research in tumors. Among those pathways in which miRNAs are involved, the NF-κB pathway is one of the most important ones. It is found to perform an important regulatory function in immune reactions, inflammations, and carcinogenesis.^{16,49} The main connection of miRNAs and NF-κB lies in miRNAs directly or indirectly regulating the expression level of target genes in the NF-κB pathway. Currently, the most commonly used method for inhibiting the function of miRNA is using anti-miR oligonucleotides. After the endogenous mature miRNA is inhibited by the anti-miR oligonucleotide, the downstream gene that is inhibited by the miRNA is released. Therefore, anti-miR oligonucleotides are effective in inhibiting miRNAs that promote the NF-κB signaling pathway in cancer.¹⁰⁴ In general,



the clinical application of miRNA-related products requires more in-depth experiments.¹⁰⁵ In addition, lncRNA is a potential therapeutic target, and it is essential to find an effective way to inhibit it. With the study of lncRNA in the NF-κB signaling pathway, it is thought that there will be drugs or treatments designed for lncRNA.

Furthermore, research is increasing on the relationship between non-coding RNA and the NF-κB signaling pathways in tumors. Undoubtedly, the study of this relationship has extremely important clinical significance related to the mechanism of tumor development and for the provision of new drug target sites for tumor treatment. Accordingly, further research is needed to elucidate whether members of the pathway are clinically significant as biomarkers or therapeutic targets.

AUTHOR CONTRIBUTIONS

Conception and Design, Z.M. and G.R.; Writing, Review, and/or Revision of the Manuscript, Z.M., X.L., Y.S., J.B., and G.R. All authors read and approved the contents of the manuscript and its publication.

CONFLICTS OF INTEREST

The authors declare no competing interests.

ACKNOWLEDGMENTS

This work was supported by National Natural Science Foundation of China grant 381570056 and Training Program of the Major Research Plan of the National Natural Science Foundation of China (grant 91543123).

REFERENCES

- Antoni, S., Ferlay, J., Soerjomataram, I., Znaor, A., Jemal, A., and Bray, F. (2017). Bladder cancer incidence and mortality: a global overview and recent trends. *Eur. Urol.* **71**, 96–108.
- Pikarsky, E., Porat, R.M., Stein, I., Abramovitch, R., Amit, S., Kasem, S., Gutkovich-Pyest, E., Urieli-Shoval, S., Galun, E., and Ben-Neriah, Y. (2004). NF-κB functions as a tumour promoter in inflammation-associated cancer. *Nature* **431**, 461–466.
- Sen, R., and Baltimore, D. (1986). Multiple nuclear factors interact with the immunoglobulin enhancer sequences. *Cell* **46**, 705–716.
- Sen, R., and Baltimore, D. (2006). Multiple nuclear factors interact with the immunoglobulin enhancer sequences. *Cell* **1986**, 46: 705–716. *J. Immunol.* **177**, 7485–7496.
- Tian, B., Liu, Z., Yang, J., Sun, H., Zhao, Y., Wakamiya, M., Chen, H., Ryttig, E., Zhou, J., and Brasier, A.R. (2018). Selective antagonists of the bronchiolar epithelial nf-κb-bromodomain-containing protein 4 pathway in viral-induced airway inflammation. *Cell Rep.* **23**, 1138–1151.
- Wu, M.Y., Li, C.J., Yang, G.T., Cheng, Y.L., Tsai, A.P., Hou, Y.T., Ho, Y.C., Hou, M.F., and Chu, P.Y. (2018). Molecular regulation of bone metastasis pathogenesis. *Cell. Physiol. Biochem.* **46**, 1423–1438.
- Liu, Y.D., Ji, C.B., Li, S.B., Yan, F., Gu, Q.S., Balic, J.J., Yu, L., and Li, J.K. (2018). Toll-like receptor 2 stimulation promotes colorectal cancer cell growth via PI3K/Akt and NF-κB signaling pathways. *Int. Immunopharmacol.* **59**, 375–383.
- Fang, J., Zhang, H., and Jin, S. (2014). Epigenetics and cervical cancer: from pathogenesis to therapy. *Tumour Biol.* **35**, 5083–5093.
- Croce, C.M. (2009). Causes and consequences of microRNA dysregulation in cancer. *Nat. Rev. Genet.* **10**, 704–714.
- Bartel, D.P. (2004). MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* **116**, 281–297.
- Hayes, J., Peruzzi, P.P., and Lawler, S. (2014). MicroRNAs in cancer: biomarkers, functions and therapy. *Trends Mol. Med.* **20**, 460–469.
- Bhan, A., Soleimani, M., and Mandal, S.S. (2017). Long noncoding RNA and cancer: a new paradigm. *Cancer Res.* **77**, 3965–3981.
- Hayden, M.S., and Ghosh, S. (2012). NF-κB, the first quarter-century: remarkable progress and outstanding questions. *Genes Dev.* **26**, 203–234.
- Kim, J.W., Jang, S.M., Kim, C.H., An, J.H., Kang, E.J., and Choi, K.H. (2012). New molecular bridge between RelA/p65 and NF-κB target genes via histone acetyltransferase TIP60 cofactor. *J. Biol. Chem.* **287**, 7780–7791.
- Caamaño, J., and Hunter, C.A. (2002). NF-κB family of transcription factors: central regulators of innate and adaptive immune functions. *Clin. Microbiol. Rev.* **15**, 414–429.
- Olarerin-George, A.O., Anton, L., Hwang, Y.C., Elovitz, M.A., and Hogenesch, J.B. (2013). A functional genomics screen for microRNA regulators of NF-κB signaling. *BMC Biol.* **11**, 19.
- Guo, L.M., Pu, Y., Han, Z., Liu, T., Li, Y.X., Liu, M., Li, X., and Tang, H. (2009). MicroRNA-9 inhibits ovarian cancer cell growth through regulation of NF-κB1. *FEBS J.* **276**, 5537–5546.
- Wan, H.Y., Guo, L.M., Liu, T., Liu, M., Li, X., and Tang, H. (2010). Regulation of the transcription factor NF-κB1 by microRNA-9 in human gastric adenocarcinoma. *Mol. Cancer* **9**, 16.
- Wang, Q., Sun, Z.X., Allgayer, H., and Yang, H.S. (2010). Downregulation of E-cadherin is an essential event in activating β-catenin/Tcf-dependent transcription and expression of its target genes in Pcd4 knockdown cells. *Oncogene* **29**, 128–138.
- Zhang, X., Liu, S., Hu, T., Liu, S., He, Y., and Sun, S. (2009). Up-regulated microRNA-143 transcribed by nuclear factor kappa B enhances hepatocarcinoma metastasis by repressing fibronectin expression. *Hepatology* **50**, 490–499.
- Borrallo, P.M., Kren, B.T., Castro, R.E., da Silva, I.B., Steer, C.J., and Rodrigues, C.M. (2009). MicroRNA-143 reduces viability and increases sensitivity to 5-fluorouracil in HCT116 human colorectal cancer cells. *FEBS J.* **276**, 6689–6700.
- Mei, L.L., Wang, W.J., Qiu, Y.T., Xie, X.F., Bai, J., and Shi, Z.Z. (2017). miR-145-5p suppresses tumor cell migration, invasion and epithelial to mesenchymal transition by regulating the Sp1/NF-κB signaling pathway in esophageal squamous cell carcinoma. *Int. J. Mol. Sci.* **18**, E1833.
- Wang, H., Garzon, R., Sun, H., Ladner, K.J., Singh, R., Dahlman, J., Cheng, A., Hall, B.M., Qualman, S.J., Chandler, D.S., et al. (2008). NF-κB-YY1-miR-29 regulatory circuitry in skeletal myogenesis and rhabdomyosarcoma. *Cancer Cell* **14**, 369–381.
- Mott, J.L., Kurita, S., Cazanave, S.C., Bronk, S.F., Werneburg, N.W., and Fernandez-Zapico, M.E. (2010). Transcriptional suppression of mir-29b-1/mir-29a promoter by c-Myc, hedgehog, and NF-κB. *J. Cell. Biochem.* **110**, 1155–1164.
- Zhou, R., Hu, G., Liu, J., Gong, A.Y., Drescher, K.M., and Chen, X.M. (2009). NF-κB p65-dependent transactivation of miRNA genes following *Cryptosporidium parvum* infection stimulates epithelial cell immune responses. *PLoS Pathog.* **5**, e1000681.
- Zhou, R., Hu, G., Gong, A.Y., and Chen, X.M. (2010). Binding of NF-κB p65 subunit to the promoter elements is involved in LPS-induced transactivation of miRNA genes in human biliary epithelial cells. *Nucleic Acids Res.* **38**, 3222–3232.
- Tang, Z.P., Zhao, W., Du, J.K., Ni, X., Zhu, X.Y., and Lu, J.Q. (2018). miR-494 contributes to estrogen protection of cardiomyocytes against oxidative stress via targeting (NF-κB) repressing factor. *Front. Endocrinol. (Lausanne)* **9**, 215.
- Fan, X., Liang, Y., Liu, Y., Bai, Y., Yang, C., and Xu, S. (2018). The upregulation of TMPRSS4, partly ascribed to the downregulation of miR-125a-5p, promotes the growth of human lung adenocarcinoma via the NF-κB signaling pathway. *Int. J. Oncol.* **53**, 148–158.
- Datta, C., Subuddhi, A., Kumar, M., Lepcha, T.T., Chakraborty, S., Jana, K., Ghosh, Z., Mukhopadhyay, A.K., Basu, J., and Kundu, M. (2018). Genome-wide mRNA-miRNA profiling uncovers a role of the microRNA miR-29b-1-5p/PHLPP1 signalling pathway in *Helicobacter pylori*-driven matrix metalloproteinase production in gastric epithelial cells. *Cell. Microbiol.* **20**, e12859.
- Chen, H., Zhang, Z., Zhang, L., Wang, J., Zhang, M., and Zhu, B. (2018). miR-27a protects human mitral valve interstitial cell from TNF-α-induced inflammatory injury via up-regulation of NELL-1. *Braz. J. Med. Biol. Res.* **51**, e6997.



31. Zabaglia, L.M., Sallas, M.L., Santos, M.P.D., Orcini, W.A., Peruquetti, R.L., Constantino, D.H., Chen, E., Smith, M.A.C., Payão, S.M., and Rasmussen, L.T. (2018). Expression of miRNA-146a, miRNA-155, IL-2, and TNF- α in inflammatory response to *Helicobacter pylori* infection associated with cancer progression. *Ann. Hum. Genet.* **82**, 135–142.
32. Mensà, E., Recchioni, R., Marcheselli, F., Giuliodori, K., Consales, V., Molinelli, E., Prattichizzo, F., Rippo, M.R., Campanati, A., Procopio, A.D., et al. (2018). miR-146a-5p correlates with clinical efficacy in psoriasis patients treated with the tumor necrosis factor-alpha inhibitor adalimumab. *Br. J. Dermatol.* **179**, 787–789.
33. Zhang, C., Qian, D., Zhao, H., Lv, N., Yu, P., and Sun, Z. (2018). miR17 improves insulin sensitivity through inhibiting expression of ASK1 and anti-inflammation of macrophages. *Biomed. Pharmacother.* **100**, 448–454.
34. Feng, Y., Chen, L., Luo, Q., Wu, M., Chen, Y., and Shi, X. (2018). Involvement of microRNA-146a in diabetic peripheral neuropathy through the regulation of inflammation. *Drug Des. Devel. Ther.* **12**, 171–177.
35. Lorente-Cebrián, S., Mejhert, N., Kulyté, A., Laurencikiene, J., Åström, G., Hedén, P., Rydén, M., and Arner, P. (2014). MicroRNAs regulate human adipocyte lipolysis: effects of miR-145 are linked to TNF- α . *PLoS ONE* **9**, e86800.
36. Neumann, M., and Naumann, M. (2007). Beyond IκBs: alternative regulation of NF-κB activity. *FASEB J.* **21**, 2642–2654.
37. Li, T., Morgan, M.J., Choksi, S., Zhang, Y., Kim, Y.S., and Liu, Z.G. (2010). MicroRNAs modulate the noncanonical transcription factor NF-κB pathway by regulating expression of the kinase IKK α during macrophage differentiation. *Nat. Immunol.* **11**, 799–805.
38. Hu, J., Zhai, C., Hu, J., Li, Z., Fei, H., Wang, Z., and Fan, W. (2017). miR-23a inhibited IL-17-mediated proinflammatory mediators expression via targeting IKK α in articular chondrocytes. *Int. Immunopharmacol.* **43**, 1–6.
39. Dai, L., Gu, L., and Di, W. (2012). miR-199a attenuates endometrial stromal cell invasiveness through suppression of the IKK β /NF-κB pathway and reduced interleukin-8 expression. *Mol. Hum. Reprod.* **18**, 136–145.
40. Zhou, H.J., Wang, L.Q., Xu, Q.S., Fan, Z.X., Zhu, Y., Jiang, H., Zheng, X.J., Ma, Y.H., and Zhan, R.Y. (2016). Downregulation of miR-199b promotes the acute spinal cord injury through IKK β -NF-κB signaling pathway activating microglial cells. *Exp. Cell Res.* **349**, 60–67.
41. Tili, E., Michaille, J.J., Cimino, A., Costinean, S., Dumitru, C.D., Adair, B., Fabbri, M., Alder, H., Liu, C.G., Calin, G.A., and Croce, C.M. (2007). Modulation of miR-155 and miR-125b levels following lipopolysaccharide/TNF- α stimulation and their possible roles in regulating the response to endotoxin shock. *J. Immunol.* **179**, 5082–5089.
42. Zhang, Z., Lu, J., Guo, G., Yang, Y., Dong, S., Liu, Y., Nan, Y., Zhong, Y., Yu, K., and Huang, Q. (2017). IKBKE promotes glioblastoma progression by establishing the regulatory feedback loop of IKBKE/YAP1/miR-Let-7b/i. *Tumour Biol.* **39**, 1010428317705575.
43. d'Adhemar, C.J., Spillane, C.D., Gallagher, M.F., Bates, M., Costello, K.M., Barry-O'Crowley, J., Haley, K., Kernan, N., Murphy, C., Smyth, P.C., et al. (2014). The MyD88 $^{+}$ phenotype is an adverse prognostic factor in epithelial ovarian cancer. *PLoS ONE* **9**, e100816.
44. Wendlandt, E.B., Graff, J.W., Gioannini, T.L., McCaffrey, A.P., and Wilson, M.E. (2012). The role of microRNAs miR-200b and miR-200c in TLR4 signaling and NF-κB activation. *Innate Immun.* **18**, 846–855.
45. Feng, Y., Zou, L., Yan, D., Chen, H., Xu, G., Jian, W., Cui, P., and Chao, W. (2017). Extracellular microRNAs induce potent innate immune responses via TLR7/MyD88-dependent mechanisms. *J. Immunol.* **199**, 2106–2117.
46. Yang, F., Liu, X., Liu, Y., Liu, Y., Zhang, C., Wang, Z., Jiang, T., and Wang, Y. (2017). miR-181d/MALT1 regulatory axis attenuates mesenchymal phenotype through NF-κB pathways in glioblastoma. *Cancer Lett.* **396**, 1–9.
47. Chen, C.Y., Chang, J.T., Ho, Y.F., and Shyu, A.B. (2016). miR-26 down-regulates TNF- α /NF-κB signalling and IL-6 expression by silencing HMGA1 and MALT1. *Nucleic Acids Res.* **44**, 3772–3787.
48. Gu, L., and Sun, W. (2015). miR-539 inhibits thyroid cancer cell migration and invasion by directly targeting CARMA1. *Biochem. Biophys. Res. Commun.* **464**, 1128–1133.
49. Zhao, N., Wang, R., Zhou, L., Zhu, Y., Gong, J., and Zhuang, S.M. (2014). MicroRNA-26b suppresses the NF-κB signaling and enhances the chemosensitivity of hepatocellular carcinoma cells by targeting TAK1 and TAB3. *Mol. Cancer* **13**, 35.
50. Morioka, S., Broglie, P., Omori, E., Ikeda, Y., Takaesu, G., Matsumoto, K., and Ninomiya-Tsuji, J. (2014). TAK1 kinase switches cell fate from apoptosis to necrosis following TNF stimulation. *J. Cell Biol.* **204**, 607–623.
51. Sirinian, C., Papanastasiou, A.D., Schizas, M., Spella, M., Stathopoulos, G.T., Repanti, M., Zarkadis, I.K., King, T.A., and Kalofonos, H.P. (2018). RANK-c attenuates aggressive properties of ER-negative breast cancer by inhibiting NF-κB activation and EGFR signaling. *Oncogene* **37**, 5101–5114.
52. Shostak, K., and Chariot, A. (2015). EGFR and NF-κB: partners in cancer. *Trends Mol. Med.* **21**, 385–393.
53. Clay, T.D., Russell, P.A., Do, H., Sundararajan, V., Conron, M., Wright, G.M., Solomon, B., Dobrovic, A., McLachlan, S.A., and Moore, M.M. (2017). EGFR and KRAS mutations do not enrich for the activation of IL-6, JAK1 or phosphorylated STAT3 in resected lung adenocarcinoma. *Med. Oncol.* **34**, 175.
54. Cho, J.H., You, Y.M., Yeom, Y.I., Lee, D.C., Kim, B.K., Won, M., Cho, B.C., Kang, M., Park, S., Yang, S.J., et al. (2018). RNF25 promotes gefitinib resistance in EGFR-mutant NSCLC cells by inducing NF-κB-mediated ERK reactivation. *Cell Death Dis.* **9**, 587.
55. Ma, J., Lyu, H., Huang, J., and Liu, B. (2014). Targeting of erbB3 receptor to overcome resistance in cancer treatment. *Mol. Cancer* **13**, 105.
56. Jiang, C., Zhu, Y., Zhou, Z., Gumin, J., Bengtsson, L., Wu, W., Songyang, Z., Lang, F.F., and Lin, X. (2017). TMEM43/LUMA is a key signaling component mediating EGFR-induced NF-κB activation and tumor progression. *Oncogene* **36**, 2813–2823.
57. Chen, Z.D., Xu, L., Tang, K.K., Gong, F.X., Liu, J.Q., Ni, Y., Jiang, L.Z., Hong, J., Han, F., Li, Q., et al. (2016). NF-κB-dependent transcriptional upregulation of cyclin D1 exerts cytoprotection against hypoxic injury upon EGFR activation. *Exp. Cell Res.* **347**, 52–59.
58. Wang, J., Zhu, M., Wang, L., Chen, C., and Song, Y. (2019). Amphiregulin potentiates airway inflammation and mucus hypersecretion induced by urban particulate matter via the EGFR-PI3K α -AKT/ERK pathway. *Cell. Signal.* **53**, 122–131.
59. Li, Y.L., Liu, X.M., Zhang, C.Y., Zhou, J.B., Shao, Y., Liang, C., Wang, H.M., Hua, Z.Y., Lu, S.D., and Ma, Z.L. (2017). MicroRNA-34a/EGFR axis plays pivotal roles in lung tumorigenesis. *Oncogenesis* **6**, e372.
60. Wang, P., Liu, X., Shao, Y., Wang, H., Liang, C., Han, B., and Ma, Z. (2017). MicroRNA-107-5p suppresses non-small cell lung cancer by directly targeting oncogene epidermal growth factor receptor. *Oncotarget* **8**, 57012–57023.
61. Liang, C., Zhang, X., Wang, H.M., Liu, X.M., Zhang, X.J., Zheng, B., Qian, G.R., and Ma, Z.L. (2017). MicroRNA-18a-5p functions as an oncogene by directly targeting IRF2 in lung cancer. *Cell Death Dis.* **8**, e2764.
62. Heyne, K., Winter, C., Gerten, F., Schmidt, C., and Roemer, K. (2013). A novel mechanism of crosstalk between the p53 and NFκB pathways: MDM2 binds and inhibits p65RelA. *Cell Cycle* **12**, 2479–2492.
63. Vaughan, C.A., Singh, S., Windle, B., Sankala, H.M., Graves, P.R., Andrew Yeadall, W., Deb, S.P., and Deb, S. (2012). p53 mutants induce transcription of NF-κB2 in H1299 cells through CBP and STAT binding on the NF-κB2 promoter and gain of function activity. *Arch. Biochem. Biophys.* **518**, 79–88.
64. Bommer, G.T., Gerin, I., Feng, Y., Kaczorowski, A.J., Kuick, R., Love, R.E., Zhai, Y., Giordano, T.J., Qin, Z.S., Moore, B.B., et al. (2007). p53-mediated activation of miRNA34 candidate tumor-suppressor genes. *Curr. Biol.* **17**, 1298–1307.
65. Chang, T.C., Wentzel, E.A., Kent, O.A., Ramachandran, K., Mullendore, M., Lee, K.H., Feldmann, G., Yamakuchi, M., Ferlito, M., Lowenstein, C.J., et al. (2007). Transactivation of miR-34a by p53 broadly influences gene expression and promotes apoptosis. *Mol. Cell* **26**, 745–752.
66. Raver-Shapira, N., Marciano, E., Meiri, E., Spector, Y., Rosenfeld, N., Moskovits, N., Bentwich, Z., and Oren, M. (2007). Transcriptional activation of miR-34a contributes to p53-mediated apoptosis. *Mol. Cell* **26**, 731–743.
67. Yamakuchi, M., Ferlito, M., and Lowenstein, C.J. (2008). miR-34a repression of SIRT1 regulates apoptosis. *Proc. Natl. Acad. Sci. USA* **105**, 13421–13426.
68. Bayraktar, R., Ivan, C., Bayraktar, E., Kanlikilicer, P., Kabil, N., Kahraman, N., Mokhlis, H.A., Karakas, D., Rodriguez-Aguayo, C., Arslan, A., et al. (2018). Dual



- suppressive effect of microRNA-34a on the FOXM1/eEF2-kinase axis regulates triple-negative breast cancer growth and invasion. *Clin. Cancer Res.* 24, 4225–4241.
69. Wang, Y., Zheng, Z.J., Jia, Y.J., Yang, Y.L., and Xue, Y.M. (2018). Role of p53/miR-155-5p/sirt1 loop in renal tubular injury of diabetic kidney disease. *J. Transl. Med.* 16, 146.
 70. Godfrey, J.D., Morton, J.P., Wilczynska, A., Sansom, O.J., and Bushell, M.D. (2018). miR-142-3p is downregulated in aggressive p53 mutant mouse models of pancreatic ductal adenocarcinoma by hypermethylation of its locus. *Cell Death Dis.* 9, 644.
 71. Feng, C., Xian, Q., and Liu, S. (2018). Micro RNA-518 inhibits gastric cancer cell growth by inducing apoptosis via targeting MDM2. *Biomed. Pharmacother.* 97, 1595–1602.
 72. Chang, C.C., Lin, C.C., Wang, C.H., Huang, C.C., Ke, T.W., Wei, P.L., Yeh, K.T., Hsu, K.C., Hsu, N.Y., and Cheng, Y.W. (2018). miR-211 regulates the expression of *RRM2* in tumoral metastasis and recurrence in colorectal cancer patients with a *k-ras* gene mutation. *Oncol. Lett.* 15, 8107–8117.
 73. Wang, X., and Zhu, J. (2018). mir-1307 regulates cisplatin resistance by targeting Mdm4 in breast cancer expressing wild type P53. *Thorac. Cancer* 9, 676–683.
 74. di Gennaro, A., Damiano, V., Brisotto, G., Armellin, M., Perin, T., Zucchetto, A., Guardascione, M., Spaink, H.P., Doglioni, C., Snaar-Jagalska, B.E., et al. (2018). A p53/miR-30a/ZEB2 axis controls triple negative breast cancer aggressiveness. *Cell Death Differ.* 25, 2165–2180.
 75. Liu, B., Li, R., Zhang, J., Meng, C., Zhang, J., Song, X., and Lv, C. (2018). MicroRNA-708-3p as a potential therapeutic target via the ADAM17-GATA/STAT3 axis in idiopathic pulmonary fibrosis. *Exp. Mol. Med.* 50, e465.
 76. Wang, Z., Yuan, W., Li, B., Chen, X., Zhang, Y., Chen, C., et al. (2018). PEITC promotes neurite growth in primary sensory neurons via the miR-17-5p/STAT3/GAP-43 axis. *J. Drug Target.* 27, 82–93.
 77. Zhang, J., Lu, Y., Yue, X., Li, H., Luo, X., Wang, Y., Wang, K., and Wan, J. (2013). miR-124 suppresses growth of human colorectal cancer by inhibiting STAT3. *PLoS ONE* 8, e70300.
 78. Zhao, Y., Ma, K., Yang, S., Zhang, X., Wang, F., Zhang, X., Liu, H., and Fan, Q. (2018). MicroRNA-125a-5p enhances the sensitivity of esophageal squamous cell carcinoma cells to cisplatin by suppressing the activation of the STAT3 signaling pathway. *Int. J. Oncol.* 53, 644–658.
 79. Song, Q., Liu, B., Li, X., Zhang, Q., Cao, L., Xu, M., Meng, Z., Wu, X., and Xu, K. (2018). miR-26a-5p potentiates metastasis of human lung cancer cells by regulating ITGB8–JAK2/STAT3 axis. *Biochem. Biophys. Res. Commun.* 501, 494–500.
 80. Lee, H., Herrmann, A., Deng, J.H., Kujawski, M., Niu, G., Li, Z., Forman, S., Jove, R., Pardoll, D.M., and Yu, H. (2009). Persistently activated Stat3 maintains constitutive NF-κB activity in tumors. *Cancer Cell* 15, 283–293.
 81. Quinodoz, S., and Guttmann, M. (2014). Long noncoding RNAs: an emerging link between gene regulation and nuclear organization. *Trends Cell Biol.* 24, 651–663.
 82. Liu, B., Sun, L., Liu, Q., Gong, C., Yao, Y., Lv, X., Lin, L., Yao, H., Su, F., Li, D., et al. (2015). A cytoplasmic NF-κB interacting long noncoding RNA blocks IκB phosphorylation and suppresses breast cancer metastasis. *Cancer Cell* 27, 370–381.
 83. Yang, L., Sun, K., Chu, J., Qu, Y., Zhao, X., Yin, H., Ming, L., Wan, J., and He, F. (2018). Long non-coding RNA FTH1P3 regulated metastasis and invasion of esophageal squamous cell carcinoma through SP1/NF-κB pathway. *Biomed. Pharmacother.* 106, 1570–1577.
 84. Hansen, T.B., Jensen, T.I., Clausen, B.H., Bramsen, J.B., Finsen, B., Damgaard, C.K., and Kjems, J. (2013). Natural RNA circles function as efficient microRNA sponges. *Nature* 495, 384–388.
 85. Hansen, T.B., Kjems, J., and Damgaard, C.K. (2013). Circular RNA and miR-7 in cancer. *Cancer Res.* 73, 5609–5612.
 86. Leng, X., Ma, J., Liu, Y., Shen, S., Yu, H., Zheng, J., Liu, X., Liu, L., Chen, J., Zhao, L., et al. (2018). Mechanism of piR-DQ590027/MIR17HG regulating the permeability of glioma conditioned normal BBB. *J. Exp. Clin. Cancer Res.* 37, 246.
 87. Fu, A., Jacobs, D.I., Hoffman, A.E., Zheng, T., and Zhu, Y. (2015). PIWI-interacting RNA 021285 is involved in breast tumorigenesis possibly by remodeling the cancer epigenome. *Carcinogenesis* 36, 1094–1102.
 88. Tsoulfas, G., and Geller, D.A. (2001). NF-κB in transplantation: friend or foe? *Transpl. Infect. Dis.* 3, 212–219.
 89. Ehedego, H., Mohs, A., Jansen, B., Hiththetiya, K., Sicinski, P., Liedtke, C., and Trautwein, C. (2018). Loss of Cyclin E1 attenuates hepatitis and hepatocarcinogenesis in a mouse model of chronic liver injury. *Oncogene* 37, 3329–3339.
 90. Liu, H., Li, G., Zhang, B., Sun, D., Wu, J., Chen, F., Kong, F., Luan, Y., Jiang, W., Wang, R., and Xue, X. (2018). Suppression of the NF-κB signaling pathway in colon cancer cells by the natural compound Riccardin D from Dumortierahirsute. *Mol. Med. Rep.* 17, 5837–5843.
 91. Romagny, S., Bouauouiche, S., Lucchi, G., Ducoroy, P., Bertoldo, J.B., Terenzi, H., Bettaieb, A., and Plencharre, S. (2018). S-nitrosylation of cIAP1 switches cancer cell fate from TNFα/TNFR1-mediated cell survival to cell death. *Cancer Res.* 78, 1948–1957.
 92. Deswaerte, V., Nguyen, P., West, A., Browning, A.F., Yu, L., Ruwanpura, S.M., Balic, J., Livis, T., Girard, C., Preaudet, A., et al. (2018). Inflammasome adaptor ASC suppresses apoptosis of gastric cancer cells by an IL18-mediated inflammation-independent mechanism. *Cancer Res.* 78, 1293–1307.
 93. Mattson, M.P., and Camandola, S. (2001). NF-κB in neuronal plasticity and neurodegenerative disorders. *J. Clin. Invest.* 107, 247–254.
 94. Roebuck, K.A., Carpenter, L.R., Lakshminarayanan, V., Page, S.M., Moy, J.N., and Thomas, L.L. (1999). Stimulus-specific regulation of chemokine expression involves differential activation of the redox-responsive transcription factors AP-1 and NF-κB. *J. Leukoc. Biol.* 65, 291–298.
 95. Qureshi, A.A., Zuvanich, E.G., Khan, D.A., Mushtaq, S., Silswal, N., and Qureshi, N. (2018). Proteasome inhibitors modulate anticancer and anti-proliferative properties via NF-κB signaling, and ubiquitin-proteasome pathways in cancer cell lines of different organs. *Lipids Health Dis.* 17, 62.
 96. Ahmadian, E., Khosrourshahi, A.Y., Eftekhar, A., Farajnia, S., Babaei, H., and Eghbal, M.A. (2018). Novel angiotensin receptor blocker, azilsartan induces oxidative stress and NFκB-mediated apoptosis in hepatocellular carcinoma cell line HepG2. *Biomed. Pharmacother.* 99, 939–946.
 97. Poźniak, J., Nsengimana, J., Laye, J.P., O'Shea, S.J., Diaz, J.M.S., Droop, A.P., Filia, A., Harland, M., Davies, J.R., Mell, T., et al. (2019). Genetic and environmental determinants of immune response to cutaneous melanoma. *Cancer Res.* 79, 2684–2696.
 98. Silva, T.C.C., de Jesus Menezes-Filho, N., de Faria Lopes, G.P., de Oliveira, D.M., Pereira, E., Pitanga, B.P.S., Dos Santos Costa, R., da Silva Velozo, E., Freire, S.M., Clarêncio, J., Borges, H.L., et al. (2018). Specific cytostatic and cytotoxic effect of dihydrochelerythrine in glioblastoma cells: role of NF-κB/β-catenin and STAT3/IL-6 pathways. *Anticancer Agents Med. Chem.* 18, 1386–1393.
 99. Yang, J., Liao, X., Agarwal, M.K., Barnes, L., Auron, P.E., and Stark, G.R. (2007). Unphosphorylated STAT3 accumulates in response to IL-6 and activates transcription by binding to NFκB. *Genes Dev.* 21, 1396–1408.
 100. Jiang, L., Yu, L., Zhang, X., Lei, F., Wang, L., Liu, X., Wu, S., Zhu, J., Wu, G., Cao, L., et al. (2016). miR-892b silencing activates NF-κB and promotes aggressiveness in breast cancer. *Cancer Res.* 76, 1101–1111.
 101. Xu, Z., Yang, F., Wei, D., Liu, B., Chen, C., Bao, Y., Wu, Z., Wu, D., Tan, H., Li, J., et al. (2017). Long noncoding RNA-SRLR elicits intrinsic sorafenib resistance via evoking IL-6/STAT3 axis in renal cell carcinoma. *Oncogene* 36, 1965–1977.
 102. Yu, L., Li, L., Medeiros, L.J., and Young, K.H. (2017). NF-κB signaling pathway and its potential as a target for therapy in lymphoid neoplasms. *Blood Rev.* 31, 77–92.
 103. Rupaimooole, R., and Slack, F.J. (2017). MicroRNA therapeutics: towards a new era for the management of cancer and other diseases. *Nat. Rev. Drug Discov.* 16, 203–222.
 104. Soldevilla, M.M., Meraviglia-Crivelli de Caso, D., Menon, A.P., and Pastor, F. (2018). Aptamer-irRNAs as therapeutics for cancer treatment. *Pharmaceuticals (Basel)* 11, E108.
 105. Catela Ivkovic, T., Voss, G., Cornellà, H., and Ceder, Y. (2017). microRNAs as cancer therapeutics: A step closer to clinical application. *Cancer Lett.* 407, 113–122.
 106. Xiao, L., Jiang, L., Hu, Q., and Li, Y. (2018). miR-302e attenuates allergic inflammation *in vitro* model by targeting RelA. *Biosci. Rep.* 38, BSR20180025.
 107. Kong, L., Sun, M., Jiang, Z., Li, L., and Lu, B. (2018). MicroRNA-194 inhibits lipopolysaccharide-induced inflammatory response in nucleus pulposus cells of the



- intervertebral disc by targeting tnf receptor-associated factor 6 (TRAF6). *Med. Sci. Monit.* **24**, 3056–3067.
108. Zhu, S., Fu, W., Zhang, L., Fu, K., Hu, J., Jia, W., and Liu, G. (2018). LINC00473 antagonizes the tumour suppressor miR-195 to mediate the pathogenesis of Wilms tumour via IKK α . *Cell Prolif.* **51**, e12416.
 109. Bardin, P., Marchal-Duval, E., Sonneville, F., Blouquit-Laye, S., Rousselet, N., Le Rouzic, P., Corvol, H., and Tabary, O. (2018). Small RNA and transcriptome sequencing reveal the role of miR-199a-3p in inflammatory processes in cystic fibrosis airways. *J. Pathol.* **245**, 410–420.
 110. Fan, J.Y., Fan, Y.J., Wang, X.L., Xie, H., Gao, H.J., Zhang, Y., Liu, M., and Tang, H. (2017). miR-429 is involved in regulation of NF- κ B activity by targeting IKK β and suppresses oncogenic activity in cervical cancer cells. *FEBS Lett.* **591**, 118–128.
 111. Huang, Y., Wang, W., Xu, Z., Pan, J., Zhao, Z., and Ren, Q. (2018). Eriocheir sinensis microRNA-7 targets crab Myd88 to enhance white spot syndrome virus replication. *Fish Shellfish Immunol.* **79**, 274–283.
 112. Tian, Y., Hao, S., Ye, M., Zhang, A., Nan, Y., Wang, G., Jia, Z., Yu, K., Guo, L., Pu, P., et al. (2015). MicroRNAs let-7b/i suppress human glioma cell invasion and migration by targeting IKBKE directly. *Biochem. Biophys. Res. Commun.* **458**, 307–312.
 113. Zhao, L., and Zhang, Y. (2015). miR-342-3p affects hepatocellular carcinoma cell proliferation via regulating NF- κ B pathway. *Biochem. Biophys. Res. Commun.* **457**, 370–377.
 114. Yang, Y., Liu, L., Zhang, Y., Guan, H., Wu, J., Zhu, X., Yuan, J., and Li, M. (2014). miR-503 targets PI3K p85 and IKK- β and suppresses progression of non-small cell lung cancer. *Int. J. Cancer* **135**, 1531–1542.
 115. Li, H.P., Zeng, X.C., Zhang, B., Long, J.T., Zhou, B., Tan, G.S., Zeng, W.X., Chen, W., and Yang, J.Y. (2013). miR-451 inhibits cell proliferation in human hepatocellular carcinoma through direct suppression of IKK- β . *Carcinogenesis* **34**, 2443–2451.
 116. Zhang, G., Sun, Y., Wang, Y., Liu, R., Bao, Y., and Li, Q. (2016). miR-502-5p inhibits IL-1 β -induced chondrocyte injury by targeting TRAF2. *Cell. Immunol.* **302**, 50–57.
 117. Shan, D., Shang, Y., and Hu, T. (2019). MicroRNA-411 inhibits cervical cancer progression by directly targeting STAT3. *Oncol. Res.* **27**, 349–358.
 118. Liu, S., Zhang, X., Hu, C., Wang, Y., and Xu, C. (2018). miR-29a inhibits human retinoblastoma progression by targeting STAT3. *Oncol. Rep.* **39**, 739–746.
 119. Wang, M., Meng, B., Liu, Y., Yu, J., Chen, Q., and Liu, Y. (2017). miR-124 inhibits growth and enhances radiation-induced apoptosis in non-small cell lung cancer by inhibiting STAT3. *Cell. Physiol. Biochem.* **44**, 2017–2028.
 120. Zhong, X., Ma, X., Zhang, L., Li, Y., Li, Y., and He, R. (2018). MIAT promotes proliferation and hinders apoptosis by modulating miR-181b/STAT3 axis in ox-LDL-induced atherosclerosis cell models. *Biomed. Pharmacother.* **97**, 1078–1085.
 121. Yuan, G., Zhao, Y., Wu, D., Gao, C., and Jiao, Z. (2018). miRNA-20a upregulates TAK1 and increases proliferation in osteosarcoma cells. *Future Oncol.* **14**, 461–469.
 122. Zhang, H., Li, H., Shaikh, A., Caudle, Y., Yao, B., and Yin, D. (2018). Inhibition of microRNA-23b attenuates immunosuppression during late sepsis through NIK, TRAF1, and XIAP. *J. Infect. Dis.* **218**, 300–311.