

TECHNICAL NOTES

Open Access

# An agmatine-inducible system for the expression of recombinant proteins in *Enterococcus faecalis*

Daniel M Linares, Marta Perez, Victor Ladero, Beatriz del Rio, Begoña Redruello, M<sup>a</sup> Cruz Martin, María Fernandez\* and Miguel A Alvarez

## Abstract

**Background:** Scientific interest in *Enterococcus faecalis* has increased greatly over recent decades. Some strains are involved in food fermentation and offer health benefits, whereas others are vancomycin-resistant and cause infections that are difficult to treat. The limited availability of vectors able to express cloned genes efficiently in *E. faecalis* has hindered biotechnological studies on the bacterium's regulatory and pathogenicity-related genes. The agmatine deiminase (AGDI) pathway of *E. faecalis*, involved in the conversion of agmatine into putrescine, is driven by a response inducer gene *aguR*.

**Results:** This study describes that the exposure to the induction factor (agmatine) results in the transcription of genes under the control of the *aguB* promoter, including the *aguBDAC* operon. A novel *E. faecalis* expression vector, named pAGEnt, combining the *aguR* inducer gene and the *aguB* promoter followed by a cloning site and a stop codon was constructed. pAGEnt was designed for the overexpression and purification of a protein fused to a 10-amino-acid His-tag at the C-terminus. The use of GFP as a reporter of gene expression in *E. faecalis* revealed that under induction with 60 mM agmatine, fluorescence reached 40 arbitrary units compared to 0 in uninduced cells.

**Conclusion:** pAGEnt vector can be used for the overexpression of recombinant proteins under the induction of agmatine in *E. faecalis*, with a close correlation between agmatine concentration and fluorescence when GFP was used as reporter.

**Keywords:** *Enterococcus*, Overexpression, Expression vector, Agmatine

## Background

Members of the genus *Enterococcus* are found throughout the normal gut of vertebrates and insects and are commonly associated with fermented foods. In artisanal Mediterranean cheeses they are believed important in the development of flavour and aroma [1-3]. *Enterococcus* has been the subject of much research given its members' potential as biopreservatives. Some strains isolated from cheese produce enterocins (bacteriocins); their bactericidal activity may be effective against food-borne pathogens such as *Listeria monocytogenes*, *Vibrio cholerae*, *Staphylococcus aureus*, *Clostridium botulinum*, *Salmonella enterica* and *Bacillus cereus*, as well as against spoilage microorganisms [4-8]. Indeed, non-pathogenic enterococci have promising commercial potential [9].

They are already used as starter cultures in certain artisanal food fermentations and are commercially available as probiotics [10-12]. For example, *Enterococcus faecalis* Symbioflor 1 clone DSM 16431 is included in a commercial probiotic product with potential benefits for human health - it has been used for over 50 years with no reports of infection [13]. Given their ability to improve the microbial equilibrium of the intestine, some strains of *Enterococcus faecium* have also been used as probiotic adjunct cultures in the making of Cheddar cheese [7]. In addition, enterococci have been used as animal feed additives in the European Union [14], and as probiotics for farm animals destined for human consumption [15-17]. The well-studied probiotic strain *E. faecium* SF68, produced in Switzerland under the name Cylactin® (DSM Nutritional Products Ltd.), is currently authorised by the European Food Safety Authority (EFSA) for use in goat and kid feed for the treatment of diarrhoea [18,19].

\* Correspondence: mfernandez@ipla.csic.es  
Instituto de Productos Lácteos de Asturias, IPLA-CSIC, Paseo Río Linares s/n,  
33300 Villaviciosa, Asturias, Spain

Enterococci can, however, also be harmful. Some strains have been studied because of their production of biogenic amines (mostly tyramine and putrescine) in fermented foods [20-23]. Other strains have virulence factors and are recognised as emerging pathogens responsible for serious clinical infections [24,25]. Enterococci are also among the most important multidrug-resistant organisms affecting immunocompromised patients [26]. This bacterial group can acquire genetic determinants that confer resistance to antibiotics and it is particularly concerning the acquisition of vancomycin resistance [27]. Indeed, the emergence of vancomycin-resistant enterococci (VRE) has alarmed the global infectious diseases community given the few options left for the management of their associated diseases. The transfer of resistance genes from enterococci to other strains, the presence of selection pressures for VRE proliferation, and the rapid expansion of resistant populations are matters of great concern. The prevalence of VRE has been reported for Europe, Asia, Australia, South America and some African countries [26].

Our knowledge of the regulation of virulence factors in *E. faecalis*, and of the environmental signals that contribute towards pathogenicity, remains poor [28]; a pressing need exists to understand this infectious agent at the genetic and physiological level. Studies in these areas, however, require molecular and genetic tools be available. For many investigations, e.g., on genetic regulation, knock-out complementation, the heterologous expression of bacteriocins in bacteriocin-negative hosts, or research on membrane permeases, the overexpression of genes is often required [29-31]. For *Enterococcus*, some expression vectors based on bacteriocin-inducible promoters have been developed, but these require the genes encoding the kinase and regulator proteins supplied *in trans* [32]. Beside this, the widely used *Lactococcus lactis* NICE system is suitable for use in other Gram-positive species including *Enterococcus*, either by using two simultaneous compatible plasmids - one harbouring the expression cassette and the other carrying the *nisRK* genes- [33], or as a single-plasmid expression vector including both cassettes [34]. In addition, other *E. faecalis* expression systems based on rhamnose as inducer [35] or under control of the pheromone cCF10 [36] have been described. Yet, the molecular genetic tools available for *E. faecalis* are more limited than for some other microorganisms. In recent years, research has therefore focused on the development of alternative Gram-positive bacterial expression systems for the production of industrially important proteins [36]. The development of controlled gene expression systems for homologous and heterologous gene expression therefore remains an important goal.

The agmatine deiminase (AGDI) genes cluster of *E. faecalis* contains the genes necessary for the biosynthesis

of putrescine from agmatine. *aguD* encodes for the agmatine/putrescine antiporter, *aguA* encodes for the agmatine deiminase, *aguB* encodes for the putrescine transcarbamylase, and *aguC* encodes for a specific carbamate kinase [22]. Overall, the locus is organized in an operon constituted by the metabolic genes *aguB*, *aguD*, *aguA* and *aguC*, which are cotranscribed in a single mRNA from the *aguB* promoter ( $P_{aguB}$ ), in a divergent orientation with the putative regulator gene (*aguR*) that has its own promoter ( $P_{aguR}$ ) [37] (Figure 1A). Our working hypothesis is that *AguR* would activate the transcription from  $P_{aguB}$  and therefore, we could construct a genetic expression system inducible by the addition of agmatine in the culture medium.

In the present work, by combining the *aguR* inducer gene and the  $P_{aguB}$  promoter (with its natural ribosome binding site), a new agmatine-inducible expression system - the "agmatine controlled expression (ACE)" system - was developed. This system was successfully used in the production of green fluorescent protein (GFP) in *E. faecalis* V583, and offers a practical and straightforward method for the heterologous expression of other recombinant proteins in *E. faecalis*.


## Results

### Influence of agmatine on putrescine production

Since agmatine is the substrate for putrescine biosynthesis in the AGDI pathway, the potential of *E. faecalis* V583 to produce putrescine in GM17 supplemented with a range of different concentrations of agmatine (0, 0.05, 0.1, 0.25, 0.5, 1, 5, 10 or 20 mM) was investigated. Liquid cultures were inoculated (2% v/v) with overnight cultures and incubated at 37°C for 24 h. Accumulation of putrescine in the supernatants was analysed by UPLC™ (Figure 1B). As expected, a positive correlation was seen between agmatine supplementation and putrescine accumulation in the extracellular medium. Putrescine was first detectable in the supernatant after treatment with 0.1 mM agmatine. Above this concentration, putrescine accumulation was gradually increased with increasing agmatine concentration.

### Transcriptional regulation of the AGDI operon by agmatine


The expression profile of the genes of the AGDI operon was analysed by RT-qPCR in GM17 media with supplementation of 0, 0.05, 0.1, 0.25, 0.5, 1, 5, 10 or 20 mM agmatine. RNA samples were taken at the end of the exponential phase of growth ( $t = 6$  h; i.e., once putrescine production had started) and mRNA levels compared. Since *aguB*, *aguD*, *aguA* and *aguC* are coexpressed in a single transcript, the expression of *aguA* is shown as representative of all *aguBDAC* mRNA.



**Figure 1 Genetic organization of the AGDI cluster of *E. faecalis* V583 and putrescine production.** **A.** The AGDI cluster of *E. faecalis* is composed of five genes, *aguR* encoding a putative regulator, followed by *aguB*, *aguD*, *aguA* and *aguC* which encode the putrescine biosynthesis pathway (accession number NC\_004668). The  $P_{aguR}$  and  $P_{aguB}$  promoters are shaded and their predicted transcripts are indicated below (dotted arrows). **B.** Putrescine production by *E. faecalis* V583 in 24 h cultures grown with 0, 0.05, 0.1, 0.25, 0.5, 1, 5, 10 or 20 mM agmatine. Values are the means of three independent assays and error bars denote standard deviation.

The expression level of *aguR* remained almost constant under all conditions of agmatine supplementation (Figure 2). However, the expression of *aguBDAC* was strictly dependent on agmatine, and a significant increase in *aguBDAC* mRNA was observed under agmatine

supplementation. In fact, agmatine induction of *aguR* transcription (approximately 2-fold increase in relative transcript levels) was much lower than that of the *aguBDAC* operon (approximately 10,000-fold increase in relative transcript levels) (Figure 2).



**Figure 2 Relative gene expression of *aguR* (A) and *aguA* (B) in cultures grown with 0, 0.05, 0.1, 0.25, 0.5, 1, 5, 10 or 20 mM agmatine.** The expression of each gene at 0 mM was independently normalised to 1, and used as the reference condition. Values shown are the means of three independent assays and error bars denote standard deviation. Asterisks indicate statistically significant differences in relative expression levels with respect to the no-agmatine condition (\* $P < 0.05$ ; \*\* $P < 0.01$ ; \*\*\* $P < 0.005$ ; Student *t* test).

### Construction of the expression vector pAGent

The core of the lactococcal vector pNZ8048 [38,39], which includes the replication cassette and the chloramphenicol resistance marker, was used as a starting point for the construction of the pAGent vector. First, a fragment of the chromosomal AGDI cluster from *E. faecalis* V583 including the *aguR* promoter ( $P_{aguR}$ ), the *aguR* gene, and the *aguB* promoter ( $P_{aguB}$ ), was amplified by PCR using the primers indicated in Table 1, and cloned into the *Bgl*II-*Nco*I sites of pNZ8048. A fragment including the multicloning site and the histidine tag (His-tag) sequence was then PCR-amplified using plasmid pNZErmC [40] as a template, employing the primers Expvfor1 and Expvrev1. The products were digested and cloned into the *Nco*I-*Hind*III sites of the previous vector, thus providing plasmid pAGent (Figure 3).

The resulting pAGent vector offers the possibility of traditional cloning without the His-tag, i.e., the insertion of the gene of interest into the extensive range of sites provided by the multiple cloning site, or the fusion of an optional C-terminal 10x His-Tag by cloning the target gene specifically into the *Nco*I-*Pst*I sites. The latter alternative enables the insertion of a promoterless gene into the correct frame, fused to the His-tag and ready for expression. Subsequent protein purification is therefore possible. Thus, this vector can be used for gene overexpression, protein immunodetection, and purification purposes.

### Controlled heterologous expression of *gfp*

To test the usefulness of pAGent in the expression of a foreign gene, tests were performed to determine whether the *aguR*/ $P_{aguB}$  cassette was able to drive the expression of heterologous genes. The promoterless gene *gfp* encoding green fluorescent protein was cloned (using the

primers indicated in Table 1) as a reporter into the *Nco*I-*Pst*I sites of pAGent, generating the vector pAGent-GFP. Fluorescence measurements revealed that, upon the addition of the induction factor agmatine (20 mM), AguR significantly enhanced the expression of *gfp* by up to 10.13 arbitrary units. Parallel cultures harbouring the empty vector pAGent, or the vector pAGent-GFP with no agmatine present, provided negative controls; expression of the gene of interest was undetectable (<0.5 arbitrary units).

### Sensitivity of the pAGent system to the inducer: dose-response curve

The control afforded by the agmatine-induced expression system in the production of recombinant proteins was tested. The production of GFP in cultures, induced with a range of agmatine concentrations (between 0 and 60 mM), was analysed by whole-cell fluorescence. Figure 4 shows an increase in fluorescence with even low concentrations of agmatine (1 mM). Induction levels increased with the agmatine concentration; indeed, a close correlation was seen between agmatine concentration and fluorescence ( $R^2 = 0.979$ ). In the presence of 60 mM agmatine, fluorescence reached 40 arbitrary units compared to 0 in uninduced cells. The virtual lack of fluorescence with 0 mM agmatine demonstrates the absence of any leaky activity of the promoter  $P_{aguB}$ .


### Toxicological assessment of agmatine in *Enterococcus faecalis*

Agmatine has been reported to act as an anti-proliferative agent in several non-intestinal mammalian cell models [41]. Since high induction concentrations were tested in the present work, assays were performed to see whether these affected bacterial viability.


**Table 1 Oligonucleotides used**

Primer	Function	Sequence (5' to 3')
<i>aguAq</i> F	<i>aguA</i> expression analysis (F)	TTGTCCCGCTTCATAAAATGG
<i>aguAq</i> R	<i>aguA</i> expression analysis (R)	CACCTGGTGAAGTGGCTTGATT
<i>aguRq</i> F	<i>aguR</i> expression analysis (F)	CGGGTTCATCTGATTGATTTCTTC
<i>aguRq</i> R	<i>aguR</i> expression analysis (R)	CGTGATTTCTCTGTCGGTTCCT
<i>recA</i> F	<i>recA</i> internal control (F)	CAAGGCTTAGAGATTGCCGATG
<i>recA</i> R	<i>recA</i> internal control (R)	ACGAGGAACTAACGCAGCAAC
<i>EFV583-tufF</i>	<i>tuf</i> internal control (F)	CAGGACATGCGGACTACGTAA
<i>EFV583-tufR</i>	<i>tuf</i> internal control (R)	TAGGACCATCAGCAGCAGAAAC
<i>AguR-EntBglII</i>	Cloning $P_{aguR}$ -AguR- $P_{aguB}$ cassette (F)	CCCCAGATCT TAAAAAGAAACAAGGTGGTGGCCG
<i>AguR EntNco</i>	Cloning $P_{aguR}$ -AguR- $P_{aguB}$ cassette (R)	CCCCATGG TGTGTTCTCCTAAAAGTTGTTTTG
<i>Expvfor1</i>	Insertion of His-tag (F)	CACACACCCCATGGCTAATCGACTGCAGGAAAATTATACTTCCAAGGTC
<i>Expvrev1</i>	Insertion of His-tag (R)	CTATCAATCAAAGCAACACGTG
<i>Gff1</i>	Cloning of GFP (F)	CACACACCCCATGGAATTCAGTAAGGGAGAAGAACTTTTC
<i>Gfr1</i>	Cloning of GFP (R)	CACACACCTGCAG ACTAGTTTTGTAGAGCTCATCCATGC

F, forward; R, reverse;  $P_{aguR}$ , *aguR* promoter;  $P_{aguB}$ , *aguB* promoter; *recA*, recombinase A; *tuf*, elongation factor Tu.



**Figure 3 Genetic map of the expression vector pAGent and zoom-in of the expression cassette.** *repC* and *repA*, replication genes; *cm<sup>r</sup>*, chloramphenicol resistance gene; *aguR*, gene encoding the regulatory protein Agur;  $P_{aguR}$ , *aguR* promoter; RBS, ribosome binding site; T, transcription terminator;  $P_{aguB}$ , *aguB* agmatine-inducible promoter; TGene, open reading frame for target gene of interest; His-tag; C-terminal histidine tag. Representative restriction sites are indicated.



**Figure 4 Sensitivity of the *aguR*/ $P_{aguB}$  expression system to the inducer.** **A.** Induction strength of *gfp* in *E. faecalis* V583 cultures containing pAGent-GFP under induction with a range of agmatine concentrations. Cultures harbouring the empty vector pAGent, or the vector pAGent-GFP with no agmatine present, were carried in parallel as negative controls. Error bars: mean standard deviation of three independent experiments. **B.** Correlation between inducer concentration (mM) and reporter fluorescence (arbitrary units).



*E. faecalis* cultures were grown in liquid GM17 supplemented with, 20, 40 or 60 mM agmatine. After 24 h incubation, serial dilutions of each suspension were prepared and immediately plated to determine the number of viable bacteria (colony forming units per millilitre [cfu/ml]). Under all conditions, the average viable count ranged from 8.85 to 9.39  $\log_{10}$ [cfu/ml] units; no significant differences were seen between the different agmatine-treated and untreated cultures (Figure 5). Thus, agmatine has no toxic effect on *E. faecalis* V583 cell viability after 24 hours of exposure, even at the high concentrations required for maximal induction.

## Discussion


Some strains of *E. faecalis* have commercial uses, e.g., as probiotics that promote a beneficial gut environment [42] and in food fermentation processes [1,43-45]. Other strains, however, can pose food safety problems and yet others are a leading cause of opportunistic, hospital-acquired infections (including urinary tract infections, septicaemia, bacteraemia and endocarditis) [2]. The extensive use of antibiotics has resulted in the rise of multiresistant *E. faecalis* strains, making the treatment of infections difficult. The identification of traits that contribute to their pathogenicity is important in understanding the dual nature of this organism [24,46,47].

Despite scientific interest in this bacterium in recent years, the genetic toolbox and methodologies available

for overexpressing genes of interest are more limited than for some other microorganisms [48]. Although heterologous gene expression in an *E. coli* background is the most common method of producing recombinant proteins from other bacteria, autologous gene expression is an alternative when heterologous expression has failed [49]. Availability of appropriate expression vectors for *E. faecalis* becomes essential to express cloned genes efficiently, to study regulatory proteins, or to carry genetic complementation studies [50]. A number of expression systems for *Enterococcus* are available. Probably the best known controllable expression system for Gram-positive bacteria is based on the autoregulatory properties of nisin biosynthesis in *L. lactis* [38,39,51,52]. The system can be used in a wide range of genera including *Lactococcus*, *Streptococcus*, *Enterococcus*, *Leuconostoc* and *Lactobacillus* [53]. However, when used in *Enterococcus*, the regulatory genes *nisR* and *nisK* need to be supplied *in trans*. Similarly, the enterocin regulatory system for controlling the expression of heterologous genes in *Enterococcus* requires the regulator- and kinase-encoding genes *entR* and *entK* be expressed *in trans* [32]. In order to tackle this disadvantage, the NICE system was optimised to be used in *Enterococcus* as a single-plasmid expression vector [34]. Other inducible expression vectors for *E. faecalis* are available: the rhamnose-inducible system [35] and the recently described one that is under control of the pheromone cCF10 [54]. The present work describes an alternative system for controlled gene expression in *E. faecalis* using a single-plasmid expression system.

The present results show putrescine accumulation by *E. faecalis* V583 to be strictly agmatine-dependent. They also show that *aguR* gene expression is not modified by an increase in agmatine concentration of the culture medium, whereas *aguBDAC* gene expression is significantly upregulated by agmatine concentrations of over 0.25 mM. The suitability of the *aguR/P<sub>aguB</sub>* system (the regulatory part of the AGDI system of *E. faecalis* V583) as an agmatine-induced gene expression system was therefore explored. An expression vector combining the *aguR* activator gene and the *aguB* promoter, followed by convenient cloning sites for introducing the gene of interest, was constructed. An important feature of the developed pAGent system is the addition of a histidine tag, which makes the vector an option for protein purification purposes. By cloning the target gene into the *NcoI-PstI* sites, the gene is inserted into the correct frame and fused to the His-tag. This His-tag was obtained from expression vectors previously shown to perform efficiently when used in protein overproduction, immunodetection and purification settings [40,55].

This expression system was assessed by expressing the reporter gene *gfp* (which codes for green fluorescent protein).



**Figure 5 Susceptibility of *E. faecalis* V583 to agmatine.** *E. faecalis* cultures were grown in liquid GM17 supplemented with 0 mM, 20 mM, 40 mM or 60 mM agmatine. After 24 h of incubation, serial dilutions of the suspension were prepared and immediately plated to determine the number of viable bacteria (cfu/ml). The number of bacteria observed was within the range of 30 to 300 cfu per plate. Error bars represent standard deviation of three independent experiments.

Very strong fluorescence induction (40 arbitrary units compared to 0 in uninduced cells) was seen in the presence of 60 mM agmatine. It should be noted that agmatine concentrations of 60 mM are within the range associated with no toxic effect in *Enterococcus*. A potential benefit of having an expression system based on *aguR/P<sub>aguB</sub>* is that gene expression can be finely controlled upon the addition of appropriate concentrations of agmatine. Agmatine is an intermediate in polyamine biosynthesis, and ubiquitous in living cells. However, it is normally present only in trace amounts - tissue concentrations are usually below 1  $\mu$ M [56]. It should also be noted that this system requires the expression of no additional proteins supplied *in trans*.

The present results clearly show that this system can effectively control the expression of genes in response to non-toxic agmatine in *E. faecalis*. This could be a useful tool for the overexpression of proteins in this species, and expands the toolbox available to use with it. Moreover, this system may be active in other *Enterococcus* species and perhaps in other Gram-positive hosts (as the NICE system may be used), although this needs to be confirmed.

## Conclusions

The described agmatine-inducible system represents an attractive means for the overproduction and purification of recombinant proteins in *E. faecalis*. The present work describes the construction of an *E. faecalis aguR/P<sub>aguB</sub>* controlled expression system and demonstrates its potential as a means of overproducing recombinant proteins. This system was assessed by expressing the reporter gene *gfp*, and very strong fluorescence was induced in the presence of 60 mM agmatine (40 arbitrary units compared to 0 in uninduced cells). The potential benefit of this system is that gene expression can be finely controlled by the addition of appropriate concentrations of agmatine. The addition of a histidine tag to the pAGent vector renders the system suitable for protein purification purposes.

## Methods

### Bacterial strains and growth conditions

*L. lactis* NZ9000 was grown at 30°C in M17 medium (Oxoid, Basingstoke, United Kingdom) supplemented with 30 mM glucose. *E. faecalis* V583 was grown at 37°C in M17 medium (Oxoid, Basingstoke, United Kingdom) supplemented with 30 mM glucose (GM17). When required, the indicated concentration of agmatine (Sigma-Aldrich, St. Louis, MO) was added to the medium. Chloramphenicol (5  $\mu$ g ml<sup>-1</sup>) was added as required.

### Analytical chromatography methods

Cultures were centrifuged at 8000g for 10 min and the resulting supernatants filtered through a 0.2  $\mu$ m Supor membrane (Pall, NY). Putrescine and agmatine concentrations

were analysed by ultra-performance liquid chromatography (UPLC™) using a Waters H-Class ACQUITY UPLC™ apparatus controlled by Empower 2.0 software and employing a UV-detection method based on derivatization with diethylethoxymethylene malonate (Sigma-Aldrich), as previously described [57].

### DNA manipulation procedures

The procedures used for DNA manipulation and recombination were essentially those described by Sambrook et al. [58]. Table 1 lists the primer sequences used. Genetic constructs for *Enterococcus* were achieved using *L. lactis* NZ9000 as an intermediate host. Plasmid and total DNA of *L. lactis* and *Enterococcus* were isolated and transformed as previously described [59]. All plasmid constructs were verified by nucleotide sequencing at Macrogen Inc. (Seoul, Republic of Korea). All enzymes for DNA technology were used according to the manufacturer's specifications.

### RNA extraction

Total RNA was extracted at 6 h of incubation using the TRI Reagent (Sigma) as previously described [21]. *E. faecalis* was grown in M17 or in M17 supplemented with 30 mM glucose and 0, 0.05, 0.1, 0.25, 0.5, 1, 5, 10 or 20 mM agmatine. The cells were then harvested by centrifugation and disrupted using glass beads (diameter up to 50  $\mu$ m) in a Fast-Prep FP120 apparatus (Thermo Savant-BIO101/Q-Biogen) at 4°C for 6  $\times$  30 s (power setting 6). The resulting samples were treated as recommended by the manufacturer. Purified RNAs were resuspended in RNase-free water. After extraction, RNA samples were treated with DNase (Fermentas, Vilnius, Lithuania), as described by the manufacturer, to eliminate any genomic contamination. Total RNA concentrations were determined by UV spectrophotometry by measuring absorbance at 260 nm in a BioPhotometer (Eppendorf, Germany).

### Gene expression quantification by RT-qPCR

Gene expression analysis was performed by reverse transcription-quantitative PCR (RT-qPCR) in a 7500 Fast Real-Time PCR System (Applied Biosystems, Carlsbad, CA) using SYBR® Green PCR Master Mix (Applied Biosystems). After 2-fold dilution of the cDNA, 5  $\mu$ l were added to 20  $\mu$ l of PCR mixture (12.5  $\mu$ l of SYBR Green Supermix, 1  $\mu$ l of each primer at 7  $\mu$ M, and 5.5  $\mu$ l of RNase-free water). Amplifications were performed with specific primers (Table 1) designed with Primer Express software (Applied Biosystems); primers specific for elongation factor thermo-unstable (*tufA*) and recombinase A (*recA*) genes were used as references. The cycling settings were those default-established by Applied Biosystems. For each condition, RT-qPCR analysis was performed on RNA purified from three independently grown cultures.

### Green fluorescence measurements

For whole-cell fluorescence measurements, equal amounts of cells were harvested, washed and subsequently resuspended in 50 mM KPi, pH 7.2, as previously described [40]. GFP emission was measured in a volume of 200  $\mu$ l of cells, using a Cary Eclipse fluorescence spectrophotometer (Varian Inc., Palo Alto, CA), at an excitation wavelength of 485 nm and an emission wavelength 530 nm. For direct comparison, all the GFP fluorescence data were normalized to the same  $A_{600}$ . Background fluorescence levels were assessed by measuring non-fluorescent control cells, and these values subtracted.

### Statistical analysis

Student *t* tests were used to evaluate the consistency of the data.

### Abbreviations

AGD: Agmatine deiminase; GFP: Green fluorescence protein; VRE: Vancomycin resistant *Enterococcus*; RT-qPCR: Reverse transcription quantitative PCR; CFU: Colony forming units.

### Competing interests

The authors declare that they have no competing interests.

### Authors' contributions

DML designed, carried out some experiments and drafted the manuscript; MP and BR performed some experiments; VL, BdR, MC and MF participated in study design and helped to write the manuscript; MAA provided the general concept and supervised the work and the manuscript. All authors contributed to the discussion of the research and approved the final manuscript.

### Acknowledgements

This work was funded by the Spanish Ministry of Economy and Competitiveness (AGL2013-45431-R). We acknowledge support of the publication fee by the CSIC Open Access Publication Support Initiative through its Unit of Information Resources for Research (URICI). We are also grateful to Bert Poolman for providing the GFP-derivative vectors. The authors thank Adrian Burton for linguistic assistance. Strain *L. lactis* NZ9000 and plasmid pNZ8048 were kindly provided by the NIZO Food Research. D.M.L. and B.d.R. were the beneficiaries of JAE DOC contracts (CSIC). M.P. is the recipient of an FPU fellowship from the Spanish Ministry of Education, Culture and Sport.

Received: 1 September 2014 Accepted: 18 November 2014

Published online: 04 December 2014

### References

- Centeno JA, Menendez S, Hermida M, Rodríguez-Otero JL: **Effects of the addition of *Enterococcus faecalis* in Cebreiro cheese manufacture.** *Int J Food Microbiol* 1999, **48**:97–111.
- Foulquié Moreno MR, Sarantinopoulos P, Tsakalidou E, De Vuyst L: **The role and application of enterococci in food and health.** *Int J Food Microbiol* 2006, **106**(1):1–24.
- Morandi S, Silveti T, Brasca M: **Biotechnological and safety characterization of *Enterococcus lactis*, a recently described species of dairy origin.** *Antonie Van Leeuwenhoek* 2013, **103**(1):239–249.
- Aymerich T, Artigas MG, Garriga M, Monfort JM, Hugas M: **Effect of sausage ingredients and additives on the production of enterocins A and B by *Enterococcus faecium* CTC492. Optimization of *in vitro* production and anti-listerial effect in dry fermented sausages.** *J Appl Microbiol* 2000, **88**:686–694.
- Leroy F, De VL: **Bacteriocin production by *Enterococcus faecium* RZS C5 is cell density limited and occurs in the very early growth phase.** *Int J Food Microbiol* 2002, **72**(1–2):155–164.
- De Vuyst L: **Bacteriocins of *Enterococcus*.** In *Bacteriocins of Lactic Acid Bacteria*. Edited by De Vuyst L, Vandamme EJ. London: Microbiology, Genetics and Applications, Blackie Academic & Professional; 1994:511–513.
- Giraffa G: **Functionality of enterococci in dairy products.** *Int J Food Microbiol* 2003, **88**(2–3):215–222.
- Franz CM, van Belkum MJ, Holzapfel WH, Abriouel H, Gálvez A: **Diversity of enterococcal bacteriocins and their grouping in a new classification scheme.** *FEMS Microbiol Rev* 2007, **31**:293–310.
- Franz CM, Holzapfel W, Stiles ME: **Enterococci at the crossroads of food safety?** *Int J Food Microbiol* 1999, **47**:1–24.
- Giraffa G: **Enterococcal bacteriocins: their potential as anti-*Listeria* factors in dairy technology.** *Food Microbiol* 1995, **12**:291–299.
- Stiles ME, Holzapfel WH: **Lactic acid bacteria of foods and their current taxonomy.** *Int J Food Microbiol* 1997, **36**:1–29.
- ACNFP: MAFF Advisory Committee on Novel Foods and Processes: *Report on Enterococcus faecium, strain K77D*. 17 Smith Square, London SW1 3JR, United Kingdom: Ergon House c/o Nobel House; 1996.
- Fritzenwanker M, Kuenne C, Billion A, Hain T, Zimmermann K, Goesmann A, Chakraborty T, Domann E: **Complete genome sequence of the probiotic *Enterococcus faecalis* Symbioflor 1 clone DSM 16431.** *Genome Announc* 2013, **1**(1):e00165–12.
- Commission E: **List of the authorized additives in feedingstuffs published in application of Article 9 t (b) of Council Directive 70/524/EEC concerning additives in feeding stuffs.** *Off J Eur Union* 2004, **C50**.
- Lojanica M, Manojlović M, Jeremić D, Petronijević S: **The effects of probiotic *Enterococcus faecium* DSM 7134 in the weaned pigs nutrition.** *Biotechnol Anim Husbandry* 2010, **26**(1–2):57–64.
- Franz CM, Huch M, Abriouel H, Holzapfel W, Gálvez A: **Enterococci as probiotics and their implications in food safety.** *Int J Food Microbiol* 2011, **151**(2):125–140.
- Kreuzer S, Machnowska P, Aßmus J, Sieber M, Pieper R, Schmidt MF, Brockmann GA, Scharek-Tedin L, Johne R: **Feeding of the probiotic bacterium *Enterococcus faecium* NCIMB 10415 differentially affects shedding of enteric viruses in pigs.** *Vet Res* 2012, **43**(1):58.
- Bellomo G, Mangiagale A, Nicastro L, Frigerio G: **A controlled double-blind study of SF68 strain as a new biological preparation for the treatment of diarrhoea in pediatrics.** *Curr Ther Res* 1980, **28**:927–936.
- EFSA Panel on Additives and products or substances used in animal feed (FEEDAP): **Scientific opinion on Cylactin® (*Enterococcus faecium*) for calves, lambs and kids for rearing and for fattening.** *EFSA J* 2013, **11**(2):3097–3110.
- Fernández M, Linares DM, Álvarez MA: **Sequencing of the tyrosine decarboxylase cluster of *Lactococcus lactis* IPLA 655 and the development of a PCR method for detecting tyrosine decarboxylating lactic acid bacteria.** *J Food Prot* 2004, **67**(11):2521–2529.
- Linares DM, Fernández M, Martín MC, Álvarez MA: **Tyramine biosynthesis in *Enterococcus durans* is transcriptionally regulated by the extracellular pH and tyrosine concentration.** *Microb Biotechnol* 2009, **2**(6):625–633.
- Llácer JL, Polo LM, Tavárez S, Alarcón B, Hilario R, Rubio V: **The gene cluster for agmatine catabolism of *Enterococcus faecalis*: study of recombinant putrescine transcarbamylase and agmatine deiminase and a snapshot of agmatine deiminase catalyzing its reaction.** *J Bacteriol* 2007, **189**(4):1254–1265.
- Ladero V, Fernández M, Calles-Enríquez M, Sánchez-Llana E, Cañedo E, Martín MC, Álvarez MA: **Is the production of the biogenic amines tyramine and putrescine a species-level trait in enterococci?** *Food Microbiol* 2012, **30**(1):132–138.
- Sundsfjord A, Willems R: ***Enterococcus* research: recent developments and clinical challenges.** *Clin Microbiol Infect* 2010, **16**(6):525–526.
- Arias CA, Murray BE: **The rise of the *Enterococcus*: beyond vancomycin resistance.** *Nat Rev Microbiol* 2012, **10**:266–278.
- Abebe W, Endris M, Tiruneh M, Moges F: **Prevalence of vancomycin resistant Enterococci and associated risk factors among clients with and without HIV in Northwest Ethiopia: a cross-sectional study.** *BMC Public Health* 2014, **14**:185–192.
- Cetinkaya Y, Falk P, Mayhall CG: **Vancomycin-resistant enterococci.** *Clin Microbiol Rev* 2000, **13**(4):686–707.
- Lenz CA, Hew Ferstl CM, Vogel RF: **Sub-lethal stress effects on virulence gene expression in *Enterococcus faecalis*.** *Food Microbiol* 2010, **27**(3):317–326.
- Linares DM, Fernández M, Del-Río B, Ladero V, Martín MC, Álvarez MA: **The tyrosyl-tRNA synthetase like gene located in the tyramine biosynthesis**



- cluster of *Enterococcus durans* is transcriptionally regulated by tyrosine concentration and extracellular pH. *BMC Microbiol* 2012, **12**:23.
30. Ryan MP, McAuliffe O, Ross RP, Hill C: Heterologous expression of lacticin 3147 in *Enterococcus faecalis*: comparison of biological activity with cytolsin. *Lett Appl Microbiol* 2001, **32**(2):71–77.
  31. Hasman H, Aarestrup FM, Dalsgaard A, Guardabassi L: Heterologous expression of glycopeptide resistance *vanHAX* gene clusters from soil bacteria in *Enterococcus faecalis*. *J Antimicrob Chemother* 2006, **57**(4):648–653.
  32. Hickey RM, Twomey DP, Ross RP, Hill C: Potential of the enterocin regulatory system to control expression of heterologous genes in *Enterococcus*. *J Appl Microbiol* 2003, **95**(2):390–397.
  33. Eichenbaum Z, Federle MJ, Marra D, de Vos WM, Kuipers OP, Kleerebezem M, Scott JR: Use of the lactococcal *nisA* promoter to regulate gene expression in gram-positive bacteria: comparison of induction level and promoter strength. *Appl Environ Microbiol* 1998, **64**(8):2763–2769.
  34. Bryan EM, Bae T, Kleerebezem M, Dunny GM: Improved vectors for nisin-controlled expression in gram-positive bacteria. *Plasmid* 2000, **44**(2):183–190.
  35. Kristich CJ, Wells CL, Dunny GM: A eukaryotic-type Ser/Thr kinase in *Enterococcus faecalis* mediates antimicrobial resistance and intestinal persistence. *Proc Natl Acad Sci U S A* 2007, **104**(9):3508–3513.
  36. D'Urzo N, Martinelli M, Nenci C, Brettoni C, Telford JL, Maione D: High-level intracellular expression of heterologous proteins in *Brevibacillus choshinensis* SP3 under the control of a xylose inducible promoter. *Microb Cell Fact* 2013, **12**:12.
  37. Suárez C, Espariz M, Blancato VS, Magni C: Expression of the agmatine deiminase pathway in *Enterococcus faecalis* is activated by the AguR regulator and repressed by CcpA and PTS(Man) systems. *PLoS One* 2013, **8**(10):e76170.
  38. Kuipers OP, de Ruyter PG, Kleerebezem M, de Vos WM: Controlled overproduction of proteins by lactic acid bacteria. *Trends Biotechnol* 1997, **15**:135–140.
  39. Kuipers OP, de Ruyter PG, Kleerebezem M, de Vos WM: Quorum sensing-controlled gene expression in lactic acid bacteria. *J Biotechnol* 1998, **64**:15–21.
  40. Linares DM, Geertsma ER, Poolman B: Evolved *Lactococcus lactis* strains for enhanced expression of recombinant membrane proteins. *J Mol Biol* 2010, **401**(1):45–55.
  41. Mayeur C, Veuillet G, Michaud M, Raul F, Blottière HM, Blachier F: Effects of agmatine accumulation in human colon carcinoma cells on polyamine metabolism, DNA synthesis and the cell cycle. *Biochim Biophys Acta* 2005, **1745**(1):111–123.
  42. Hancock L, Perego M: Two-component signal transduction in *Enterococcus faecalis*. *J Bacteriol* 2002, **184**(21):5819–5825.
  43. Jiménez E, Ladero V, Chico I, Maldonado-Barragán A, López M, Martín V, Fernández L, Fernández M, Alvarez MA, Torres C, Rodríguez JM: Antibiotic resistance, virulence determinants and production of biogenic amines among enterococci from ovine, feline, canine, porcine and human milk. *BMC Microbiol* 2013, **13**:288.
  44. Gardini F, Bover-Cid S, Tofalo R, Belletti N, Gatto V, Suzzi G, Torriani S: Modeling the aminogenic potential of *Enterococcus faecalis* EF37 in dry fermented sausages through chemical and molecular approaches. *Appl Environ Microbiol* 2008, **74**:2740–2750.
  45. Serio A, Paparella A, Chaves-Lopez C, Corsetti A, Suzzi G: *Enterococcus* populations in Pecorino Abruzzese cheese: biodiversity and safety aspects. *J Food Prot* 2007, **70**:1561–1568.
  46. Nallapareddy SR, Wenxiang H, Weinstock GM, Murray BE: Molecular characterization of a widespread, pathogenic, and antibiotic resistance-receptive *Enterococcus faecalis* lineage and dissemination of its putative pathogenicity island. *J Bacteriol* 2005, **187**(16):5709–5718.
  47. Abranches J, Martínez AR, Kajfasz JK, Chávez V, Garsin DA, Lemos JA: The molecular alarmone (p)ppGpp mediates stress responses, vancomycin tolerance, and virulence in *Enterococcus faecalis*. *J Bacteriol* 2009, **191**(7):2248–2256.
  48. Debroy S, van der Hoeven R, Singh KV, Gao P, Harvey BR, Murray BE, Garsin DA: Development of a genomic site for gene integration and expression in *Enterococcus faecalis*. *J Microbiol Methods* 2012, **90**(1):1–8.
  49. Li H, O'Sullivan DJ: Heterologous expression of the *Lactococcus lactis* bacteriocin, nisin, in a dairy *Enterococcus* strain. *Appl Environ Microbiol* 2002, **68**(7):3392–3400.
  50. Fujimoto S, Ike Y: pAM401-Based shuttle vectors that enable overexpression of promoterless genes and one-step purification of tag fusion proteins directly from *Enterococcus faecalis*. *Appl Environ Microbiol* 2001, **67**:1262–1267.
  51. Kuipers OP, Beerthuyzen MM, de Ruyter PG, Luesink EJ, de Vos WM: Autoregulation of nisin biosynthesis in *Lactococcus lactis* by signal transduction. *J Biological Chem* 1995, **270**:27299–27304.
  52. Kuipers OP, Rollema HS, Beerthuyzen MM, Siezen RJ, de Vos WM: Protein engineering and biosynthesis of nisin and regulation of transcription of the structural *nisA* gene. *Int Dairy J* 1995, **5**:785–795.
  53. Kleerebezem M, Beerthuyzen MM, Vaughan EE, de Vos WM, Kuipers OP: Controlled gene expression systems for lactic acid bacteria: transferable nisin-inducible expression cassettes for *Lactococcus*, *Leuconostoc*, and *Lactobacillus* spp. *Appl Environ Microbiol* 1997, **63**:4581–4584.
  54. Borrero J, Chen Y, Dunny GM, Kaznessis YN: Modified lactic acid bacteria detect and inhibit multiresistant Enterococci. *ACS Synth Biol* 2014, Epub ahead of print.
  55. Geertsma ER, Groeneveld M, Slotboom DJ, Poolman B: Quality control of overexpressed membrane proteins. *Proc Natl Acad Sci U S A* 2008, **105**(15):5722–5727.
  56. Piletz JE, Aricioglu F, Cheng JT, Fairbanks CA, Gilad VH, Haenisch B, Halaris A, Hong S, Lee JE, Li J, Liu P, Molderings GJ, Rodrigues AL, Satriano J, Seong GJ, Wilcox G, Wu N, Gilad GM: Agmatine: clinical applications after 100 years in translation. *Drug Discov Today* 2013, **18**(17–18):880–893.
  57. Redruello B, Ladero V, Cuesta I, Álvarez-Buylla JR, Martín MC, Fernández M, Alvarez MA: A fast, reliable, ultra high performance liquid chromatography method for the simultaneous determination of amino acids, biogenic amines and ammonium ions in cheese, using diethyl ethoxymethylenemalonate as a derivatising agent. *Food Chem* 2013, **139**(1–4):1029–1035.
  58. Sambrook JD, Russell D: *Molecular Cloning a Laboratory Manual*. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory; 2001.
  59. de Vos WM, Vos P, Dehaard H, Boeritger I: Cloning and expression of the *Lactococcus lactis* ssp *cremoris* SK11 gene encoding an extracellular serine proteinase. *Gene* 1989, **85**:169–176.

doi:10.1186/s12934-014-0169-1

Cite this article as: Linares *et al.*: An agmatine-inducible system for the expression of recombinant proteins in *Enterococcus faecalis*. *Microbial Cell Factories* 2014 **13**:169.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

