Images in Clinical Medicine Acta Medica Academica 2022;51(1):64-65 DOI: 10.5644/ama2006-124.372


Large Bowel Obstruction Secondary to Urinary Retention

Seyedeh Kimia Yavari, Leili Pourafkari

Catholic Health System, Sisters of Charity Hospital, Buffalo, New York, USA

Correspondence: *leili.p@gmail.com*; Tel.: + 1 716 34549760 Received: 12 January 2022; Accepted: 17 February 2022

Key Words: Bowel Obstruction ■ Urinary Retention ■ CT Scan.



A 54-year-old man with past medical history of schizophrenia maintained on olanzapine and trihexyphenidyl presented with abdominal pain and distension, constipation and difficulty urinating. In physical exam abdominal distension with hyperactive bowel sounds were noted. Abdominal X-ray followed by CT scan with contrast showed

gaseous distension throughout the colon with a transition point in proximal sigmoid colon in favor of large bowel obstruction and a remarkably distended bladder (Panel A and B). Bladder catheterization resulted in 3700 mL of urine. Nasogastric tube inserted and patient kept nilper-os. Patient's abdominal discomfort improved

after catheterization. He started having bowel movements and tolerated diet over the next couple days. He recovered without need for surgical intervention and was discharged with stool softener and tamsulosin. Severe bladder distension is a rare cause of bowel obstruction (1). Bladder is a less mobile organ with lower intrapelvic fixation and its distention tends to pinch the rectosigmoid colon at the prominence of sacral promontory, where diameter of pelvis is narrow (2). In this case anticholinergic medication and prostate enlargement were contributing to urinary retention.

Conflict of Interest: The authors declare that they have no conflict of interest.

References

- 1. Papes D, Altarac S, Arslani N, Rajkovic Z. Urinary retention presenting as complete bowel obstruction. Can Urol Assoc J. 2013;7(9-10):E637-9.
- 2. Fujisaki T, Fujita Y, Mizuta H, Niina N, Miyazaki N, Tashiro A, et al. Large bowel obstruction caused by urinary retention from benign prostate hyperplasia. Radiol Case Rep. 2019;14(2):213-6.