



# Hepatocyte transplantation and advancements in alternative cell sources for liver-based regenerative medicine

Charlotte A Lee<sup>1</sup> · Siddharth Sinha<sup>1</sup> · Emer Fitzpatrick<sup>2</sup> · Anil Dhawan<sup>2</sup>

Received: 20 November 2017 / Revised: 7 March 2018 / Accepted: 11 April 2018 / Published online: 24 April 2018  
© The Author(s) 2018

## Abstract

Human hepatocyte transplantation has been actively perused as an alternative to liver replacement for acute liver failure and liver-based metabolic defects. Current challenges in this field include a limited cell source, reduced cell viability following cryopreservation and poor engraftment of cells into the recipient liver with consequent limited life span. As a result, alternative stem cell sources such as pluripotent stem cells, fibroblasts, hepatic progenitor cells, amniotic epithelial cells and mesenchymal stem/stromal cells (MSCs) can be used to generate induced hepatocyte like cells (HLC) with each technique exhibiting advantages and disadvantages. HLCs may have comparable function to primary human hepatocytes and could offer patient-specific treatment. However, long-term functionality of transplanted HLCs and the potential oncogenic risks of using stem cells have yet to be established. The immunomodulatory effects of MSCs are promising, and multiple clinical trials are investigating their effect in cirrhosis and acute liver failure. Here, we review the current status of hepatocyte transplantation, alternative cell sources to primary human hepatocytes and their potential in liver regeneration. We also describe recent clinical trials using hepatocytes derived from stem cells and their role in improving the phenotype of several liver diseases.

**Keywords** Hepatocyte transplantation · Induced pluripotent stem cells · Fibroblast · Mesenchymal stem/stromal cell · Hepatic progenitor cells · Liver regeneration

## Abbreviations

ALF	Acute liver failure	fVII	Factor VII
BMM	Bone marrow-derived macrophages	FGF	Fibroblast growth factor
BMP4	Bone morphogenetic protein 4	FHF	Fulminant hepatic failure
CN-1	Crigler-Najjar syndrome type 1	hAECs	Hepatic differentiation of amniotic epithelial cells
CYP	Cytochrome p450	HPCs	Hepatic progenitor cells
DEX	Dexamethasone	HNF1A	Hepatocyte nuclear factor 1 homeobox alpha
EGF	Epidermal growth factor	HT	Hepatocyte transplantation
		HGF	Hepatocyte-growth factor
		HLCs	Hepatocyte-like cells
		iPSCs	Induced pluripotent stem cells
		PE	Plasma exchange
		EpCAM <sup>+</sup>	Primary human bile duct cells
		CLiPs	Proliferative bipotent cells
		MSCs	Mesenchymal stem cells
		MSC-CM	MSC-conditioned medium
		OSM	Oncostatin M
		OTC	Ornithine transcarbamylase deficiency
		OLT	Orthotopic liver transplantation
		UC-MSC	Umbilical cord–mesenchymal stem cell

---

Siddharth Sinha and Charlotte A Lee are joint first authors

---

✉ Anil Dhawan  
Anil.dhawan@kcl.ac.uk

<sup>1</sup> Dhawan Lab, Institute of Liver Studies, King's College London at King's College Hospital NHS Foundation trust, London, UK

<sup>2</sup> Paediatric Liver GI and Nutrition Centre, King's College London at King's College Hospital NHS Foundation Trust, London, UK

## Introduction

The liver is responsible for a diverse range of functions within the body ranging from xenobiotic metabolism to plasma protein synthesis [1]. However, the liver can become susceptible to acute failure (ALF) due to viral infection or drug effect amongst other aetiologies [2]. Liver-based metabolic disorders that are caused by single-gene defects can result in a lack of specific liver-based enzyme function which may result in damage to other organs most often irreversible neurodisability [3]. Orthotopic liver transplantation (OLT) is currently the only viable treatment for severe ALF and certain liver-based metabolic disorders [2, 4, 5]. Liver replacement is not futuristic for liver-based metabolic defects where gene therapy would be ideal. Notably, a shortage in healthy donor livers has led to research into alternative treatment options [6]. Hepatocyte transplantation (HT), where liver cells may provide a potential alternative to OLT or act as a bridge until a suitable organ becomes available, has been demonstrated, with over 100 cases published worldwide showing the safety and preliminary efficacy of the technique [7, 8].

This review will focus on the current status of HT, alternate cell sources to primary human hepatocytes and alternate methods of liver regeneration which can be used either independently or in combination with HT.

## The current status of hepatocyte transplantation

HT is a less invasive alternative to OLT. Hepatocytes are isolated from donor livers that have been rejected for organ transplantation in view of prolonged warm or cold ischaemia times, mild steatosis or aberrant anatomy. Hepatocytes isolated from one donor liver can yield a high quantity of cells and be used to treat multiple patients [9, 10]. The ability to cryopreserve these cells is one of the major advantages of HT, potentially providing an off-the-shelf treatment and enabling cells of all blood groups to be immediately available for emergency cases, which is particularly important for patients with ALF. Several routes of infusion for cell transplantation have been described including intraportal, intrasplenic and intraperitoneal. Depending on the age of the patient, infusion of hepatocytes into the portal vein can be undertaken via the umbilical vein under fluoroscopy to avoid a patent ductus venosus. In older children, radiological or surgical catheters are required using radiological guidance [11–13].

Hepatocyte transplantation can be used to treat patients with liver-based metabolic diseases and ALF [7, 14, 15]. Metabolic liver diseases treated using HT include Crigler-Najjar syndrome type 1 (CN-1), urea cycle defects and factor VII (fVII) deficiency [16]. Crigler-Najjar syndrome has an incidence of 1 in 1,000,000 births in the UK, and although

the primary long-term treatment is liver transplantation, the lack of an appropriately sized, blood matched donor organ can lead to irreversible neurological disability. There are reports of 8 patients that have received between 1.4 and 7.5 billion hepatocytes, which showed up to a 50% reduction in bilirubin and UGT activity and a decrease in the need for phototherapy. The majority of these patients went onto receive an OLT within a year as the effects of HT were not long lasting, with one patient avoiding OLT for 4 years [16–21].

Several urea cycle defects have also been shown to benefit from hepatocyte infusions. This includes ornithine transcarbamylase deficiency (OTC), which is an X-linked genetic condition affecting males that causes hyperammonemia and can also lead to neurological implications including developmental delay and learning disabilities. HT led to decreased ammonia levels and increased urea production in six of these patients, with four receiving an OLT within 7 months and two reported mortalities [14, 22–25]. Other metabolic liver diseases shown to improve following HT include familial hypercholesterolemia where there was up to a 20% decrease in LDL in three patients, fVII deficiency where there was a 70% decrease in the need for recombinant fVII for 6 months and ASL deficiency where there was a decrease in ammonia production and increased psychomotor abilities [26–28]. Nevertheless, the long-term benefits of HT for the treatment of patients with liver-based metabolic disease have yet to be shown.

HT can also be used in patients with ALF. Although liver transplantation is the primary treatment, paediatric patients may die whilst waiting for an appropriately size and blood-matched liver, with no other treatment being shown to improve survival [29]. Furthermore, existing listing criteria are not robust in terms of accurately predicting death and survival; hence, some patients receive a liver transplantation, even if their liver may spontaneously recover, leading to unnecessary surgeries with life-long immunosuppression. Thirty-seven patients with acute liver failure have received human hepatocytes for both drug induced, viral and idiopathic ALF. Ten of these patients underwent intraportal hepatocyte infusions, with two making a full recovery without the need for OLT and three were successfully bridged to transplantation [15, 30–33]. However, there is a high risk associated with placing intrahepatic invasive catheters in coagulopathic patients. As an alternative, it is possible to encapsulate hepatocytes in alginate microbeads made from purified alginate using an encapsulator [34]. The semi-permeable membrane within the microbeads allows exchange of metabolites, maintaining synthetic detoxification function, whilst also protecting against immunocompetent cells by preventing the entry of antibodies [35, 36]. Hepatocyte-containing microbeads are transplanted into the intraperitoneal cavity. The transplanted hepatocytes can assist the necessary liver functions, allowing the liver to recover. After full recovery, usually 1 month, the microbeads can be removed using a laparoscopic peritoneal lavage.

Despite extensive clinical data, long-term clinical outcomes of HT have yet to be established with any type of liver-based metabolic disease or ALF. There are still many limitations to the technique, and current research aims to overcome this. Firstly, the cryopreservation process requires further optimisation with the current protocol resulting in low viability and function of hepatocytes post-thawing. Furthermore, transplanted donor hepatocytes undergo immune rejection with up to 70% of engrafted cells cleared within the first 24 h post-transplantation [37]. Upon transplantation, donor hepatocytes are recognised and activate the instant blood-mediated inflammatory reaction, during which both the complement and coagulation pathways are activated. Innate immune cells such as Kupffer cells, natural killer cells and monocytes rapidly clear transplanted donor hepatocytes [38, 39]. A major limitation of HT is the lack of good quality donor organs from which to isolate cells from. Neonatal livers have been investigated as a potential cell source as they are infrequently used for OLT due to the high incidence of hepatic artery thrombosis [40]. However, they may be an ideal cell source for HT, with one study demonstrating their mature function of cytochrome P450 and ureagenesis enzymes, with increased attachment efficiency and viability compared to hepatocytes isolated from adult donor livers [41, 42]. Currently, the number of neonatal donations is not sufficient to maintain an active clinical HT programme [41, 43]. As a result, there is a requirement to investigate alternative cell sources to resolve the shortage of healthy donor organs and avoid recipient immune rejection (Fig. 1 and Table 1).

## Alternative cell sources for hepatocyte transplantation

### Induced pluripotent stem cells generated from somatic cells


Induced pluripotent stem cells (iPSCs) that can be differentiated into hepatocyte-like cells (HLCs) are being widely explored as an alternative to primary human hepatocytes. These cells offer an excellent alternative source of hepatocytes and are advantageous over primary human hepatocytes because of their unlimited cell source and their ability to avoid the immune system [44]. Methods to produce iPSC-derived HLCs are well established, with most protocols using a three-dimensional matrix such as Matrigel® to establish the generation of hepatocytes. However 2D matrices including collagen have also shown successful iPSC to HLC differentiation [45, 46]. An effective protocol for generating HLCs has been established via the use of specific growth factors such as activin A and bone morphogenetic protein 4 (BMP4), which are crucial for initiating the first step of hepatic maturation. Hepatocyte-growth factor (HGF) and oncostatin stimulate

hepatoblast formation and hepatocyte-like differentiation, respectively. Si-Tayeb et al. showed that iPSC-derived HLCs displayed hepatic functionality, including glycogen accumulation, lipoprotein uptake and urea synthesis [47]. iPSCs are advantageous over primary human hepatocytes as generation from somatic cells of an individual will prevent activation of the recipient's immune response, avoiding the use of immunosuppressive drugs [38, 45, 48, 49]. iPSCs can proliferate indefinitely, forming a limitless pool of HLCs allowing patients to receive multiple infusions if necessary [48, 50]. iPSC-derived HLCs can also be used as a model for several metabolic liver diseases including familial hypercholesterolemia,  $\alpha_1$ -antitrypsin deficiency and glycogen storage disease type 1a. By generating cells with similar phenotypes to those caused by inherited diseases, it is possible to study the mechanisms of dysregulated cellular functions and identify ways to treat or reverse the condition [51]. Currently, the functional ability of HLCs is not comparable to human primary hepatocytes. Yu et al. showed that HLCs co-express alpha-fetoprotein and albumin, suggesting that they are not fully mature. Levels of albumin synthesis, urea production, cytochrome p450 activity and mitochondrial function are also significantly lower than human primary hepatocytes [50]. Furthermore, there are genetic variations in donor cells that effect the differentiation propensity of iPSCs which has been attributed to genetic variation of the donor, cell culture conditions and iPSC generation protocols [52, 53]. Following the development of new hepatic differentiation protocols, Kajiwara et al. compared 28 iPSC lines derived from different somatic cells and showed that it was the origin of the donor cells that determined the variation in hepatic differentiation and not the derivation method [53]. With albumin used as a marker to assess functionality, HLCs derived from human dermal fibroblast-iPSCs and peripheral blood cell-iPSCs showed minimal variation in hepatic differentiation from the same donor; however, inter-donor hepatic variation was more prominent. This creates a complication when using iPSCs for therapeutic use as non-identical cell lines cannot guarantee the same quality of HLC production [48, 53].

In addition, the tumorigenic potential of these cells due to the presence of oncogenes such as c-Myc may raise a safety concern regarding the clinical application of these cells. Chen et al. reported no formation of teratomas or tumours in Gunn rats transplanted with HLCs, suggesting a loss of pluripotent characteristics within these cells [54]. Although HLCs represent an ideal cell source for HT with no risk of rejection, work is still ongoing to advance their functional capacity and fully validate the safety and efficacy of using these stem cells.

### Fibroblasts

Human fibroblasts offer another potential source of HLCs for HT. Fibroblasts are connective tissue cells found in all areas of



**Fig. 1** Potential alternative cell sources (induced pluripotent stem cells, fibroblasts, mesenchymal stem/stromal cells and hepatic progenitor cells) which can be used to generate hepatocytes. Gene transfer is used to convert somatic cells to iPSCs and fibroblasts to HLCs. All other transformations occur under culture conditions. *HLC* induced hepatocyte, *iPSC* induced pluripotent stem cells, *MSC* mesenchymal stem cells,

*HPC* hepatic progenitor cells, *hAEC* human amniotic epithelial cells, *BMP* bone morphogenetic protein, *OSM* oncostatin M, *HGF* hepatic growth factor, *HNF1A* hepatocyte nuclear factor 1 homeobox alpha, *HNF4A* hepatocyte nuclear factor 4 alpha, *FGF* fibroblast growth factor, *EGF* epidermal growth factor, *Dex* dexamethasone, *FBS* foetal bovine serum

the human body [55]. Huang et al. established the first reprogramming of human fibroblasts to hepatocytes using both foetal and adult connective tissue cells. Using lentiviruses as a vector for expressing transcription factors, hepatocyte nuclear factor 1 homeobox alpha (HNF1A), HNF4A and FOXA3, successful fibroblast to HLC transformation has been achieved, with HNF1A being crucial for the human

fibroblast reprogramming [56]. Simeonov et al. also achieved fibroblast transformation using exogenous HNF1A messenger RNA (mRNA) [57]. Within the same year, Du et al. derived HLCs from human fibroblasts, with the generated hepatocytes possessing phase I/II/III drug metabolic activity comparable to primary human hepatocytes [58]. Fibroblast-derived HLC production requires only a single transformation step and they

**Table 1** Summary of selected clinical trials globally, researching the therapeutic benefits of alternative cell sources in liver disease

Alternative cell sources	Advantages	Disadvantages
Induced pluripotent stem cells (iPSC)	<ul style="list-style-type: none"> <li>• Patient-specific cell generation</li> <li>• Reduced immune response</li> <li>• Disease modelling</li> <li>• Limitless pool of cells</li> </ul>	<ul style="list-style-type: none"> <li>• Inadequate long-term functionality</li> <li>• Potential tumour formation</li> </ul>
Fibroblasts	<ul style="list-style-type: none"> <li>• Patient-specific cell generation</li> <li>• Reduced immune response</li> <li>• Disease modelling</li> </ul>	<ul style="list-style-type: none"> <li>• Proliferation arrest</li> <li>• Residual epigenetic memory</li> <li>• Resistant to transformation</li> </ul>
Mesenchymal stem/stromal cells (MSC)	<ul style="list-style-type: none"> <li>• Patient-specific cell generation</li> <li>• Immune modulation</li> <li>• Easily accessible from several tissues of the body</li> </ul>	<ul style="list-style-type: none"> <li>• Could potentially lose functionality</li> </ul>
Hepatic progenitor cells (HPC)	<ul style="list-style-type: none"> <li>• Naturally differentiate into new hepatocytes</li> <li>• Can be used to generate a number of new hepatocytes</li> <li>• Considered to be involved in hepatocyte regeneration</li> </ul>	<ul style="list-style-type: none"> <li>• Could play no role in liver regeneration</li> <li>• May be involved in the progression of liver fibrosis</li> <li>• Acquired from donor livers so may cause immune rejection</li> <li>• Immunosuppressive drugs would be required</li> </ul>
Amniotic epithelial cells (hAEC)	<ul style="list-style-type: none"> <li>• Reduced risk of tumour formation</li> <li>• Easily accessible</li> <li>• Reduced ethical implications</li> <li>• Reduced immune reaction</li> </ul>	<ul style="list-style-type: none"> <li>• Gene expression similar to foetal cells rather than adult hepatocytes</li> </ul>

can be patient-specific, reducing the chances of immune rejection and avoiding the use of immunosuppressive drugs [38, 49, 56]. Despite these advantageous, fibroblast-derived HLCs have therapeutic limitations. They have a limited reproductive capability and cannot be used for repeat infusions in a single patient. Furthermore, human fibroblasts are resistant to hepatic transdifferentiation, thereby creating an additional barrier when generating HLCs [56]. Hepatocytes generated from reprogrammed fibroblasts may still retain epigenetic memory from the fibroblast cell of origin. This creates limitations when choosing fibroblasts for hepatic transformation, as cells with significant epigenetic differences to hepatocytes may be further resistant to reprogramming and reduced functionality [59].

### Hepatic differentiation of amniotic epithelial cells

It is also possible to generate hepatocytes from amniotic epithelial cells. These cells have stem cell markers such as OCT-4, Nanog, SOX-2 and Rex-1, and as they do not have telomerase reverse transcriptase, they show a stable phenotype without the risk of tumorigenic potential [60]. Such cells have minimal ethical implications, and there is no shortage of placental tissue from which to isolate the cells. Following culture in Matrigel® or liver-derived ECM, these cells had albumin and CYP3A4, 3A7, 2B6 and 2D6 mRNA levels which increased over time with a peak at day 21 [61]. Following transplantation into the SCID mouse, genes were expressed for human cytochrome p450 genes, metabolic enzymes and hepatocyte-enriched transcription factors and plasma proteins 6 months post-transplantation. It has now been suggested that hepatic differentiation of amniotic epithelial cells (hAECs) represent a promising non-controversial, unlimited source of cells for liver-based metabolic diseases.

### The regenerative capacity of resident liver cell hepatic progenitor cells

Hepatic progenitor cells (HPCs), also known as oval cells, are believed to differentiate into mature hepatocytes or cholangiocytes, upon liver damage and help in tissue restoration. HGF and EGF are critical in inducing the transformation of HPC into hepatocytes. HGF activates the MET receptor, which further upregulates the expressions of AKT and STAT3 driving hepatic transformation. A lack of MET receptors completely attenuates HPC to hepatocyte differentiation even in the presence of EGFR. However, EGFR-null HPCs were still able to sufficiently transform into hepatocytes with MET alone [62].

Zhang et al. established *in vitro* generation of HLCs from human foetal HPCs, under the influence of oncostatin M (OSM), DEX and HGF. These newly differentiated hepatocytes have functional glycogen storage, albumin secretion and cytochrome p450 activity with Khuu et al., suggesting that *in vitro*

differentiated adult hepatic stem cells are also capable of urea production and ammonium chloride metabolism [63, 64]. Unlike some other HLCs, hepatocytes formed from liver-derived progenitor cells have reached clinical application. Sokal et al. transplanted HPCs in a 3-year-old female patient suffering with OTC deficiency. Previous transplantation of cryopreserved hepatocytes failed to improve the patient's symptoms. Fourteen weeks post-infusion, biopsies showed 3% presence of donor cells and the patient showed some functional improvement with a reduction in disease-related anorexia. Unfortunately, 6 months post-infusion, the child underwent OLT and later died. These results suggest that HPCs could play a role in treatment of metabolic liver disease; however, longer-scale clinical trials are required to assess their full potential [65].

### Human bile duct cells

In addition to HPCs, mature hepatocytes can be derived from a number of other resident cell types within the liver. Huch et al. established a protocol differentiating primary human bile duct cells (EpCAM<sup>+</sup>) into genetically stable functional HLCs in both *in vitro* and *in vivo* transplantations. Organoids were formed using ductal cells, and using medium consisting of BMP7, EGF and HGF, successful hepatic differentiation was achieved. Newly formed HLCs demonstrated albumin production, CYP3A3/4/5 activity and bile acid secretion. Furthermore, organoids successfully engrafted into Balb/c nude mice with induced liver damage, sustaining albumin and  $\alpha$ -1-antitrypsin levels for up to 120 days in two out of five recipient mice. Debate remains over the genetic stability of fibroblasts and iPSCs as cell sources for HLCs. The expandable nature and genetic stability of HLCs derived from human bile ductal cells makes them a desirable alternate cell source [66].

### Chemically induced liver progenitors

Recently, it has been shown that mature hepatocytes convert to HLCs during chronic liver injury [67]. Katsuda et al. showed that a cocktail of small molecules Y-27632, A-83-01 and CHIR99021 could contribute to the induction and maintenance of bipotent chemically induced liver progenitor cells (CLiPs). These cells could either be differentiated into mature hepatocytes or biliary epithelial cells. Rat CLiPs were capable of repopulating immunodeficient mice with chronic liver injury. Albumin levels were used to assess liver functionality, which showed a consistent increase up until 8 weeks post-transplantation. Immunohistochemistry showed that up to 75–90% of the mouse liver had been replaced by rat hepatocytes, demonstrating a selective proliferative advantage for the healthy donor cells. Mouse ductal structures also showed CLiPs-derived cell replacement, displaying the biopotency of the lineage. If similar protocols could be established using human hepatocytes, this offers an additional source for HLCs for use in HT.



Furthermore, the bipotent properties of CLiPs suggest that they could be used to tackle diseases related to the biliary tree as well as the liver [68].

## The role of mesenchymal stem/stromal cells in liver-based regenerative medicine

Mesenchymal stem/stromal cells (MSCs) have been investigated as another cell source for hepatocyte differentiation but with limited and controversial results. More promising is their immunogenic effect, and now, MSCs are being investigated as an immunomodulatory therapy to treat liver a number of different liver diseases.

MSCs can be isolated from various tissues, such as bone marrow, adipose tissue and the umbilical cord. BMP and fibroblast growth factor (FGF) induction lead to the differentiation of MSCs to the hepatic lineage, with dexamethasone (DEX) and IL-16 inducing hepatic maturation. Single-step procedures are also commonly used with HGF and epidermal growth factor (EGF). Transformed cells generated from this procedure usually exhibit functional hepatic properties 2–3 weeks post-culturing but do lose functional capabilities when cultured for prolonged periods [69]. Furthermore, it is still debated whether MSC-derived hepatocytes are able to efficiently re-populate a host liver to provide adequate function and clinical application is still in its infancy [70]. Perhaps of greater potential is the immunomodulatory effect of MSCs. MSCs play a key role in immune modulation due to their lack of MHC-I/MHC-II receptors and are unlikely to trigger a T cell response. Furthermore, MSCs are considered to reduce T cell proliferation and cytotoxicity, as well as improving liver injury and enhancing liver regeneration [71, 72]. This immune evasion capability has resulted in MSCs adopting the title of “immunoprivileged” or “immunotolerant” cells [73]. Thus, MSCs could be used in conjunction with hepatocytes during HT to increase engraftment and reduce the immune response. Hwang et al. showed that intrasplenic transplantation of MSC-derived HLCs restored liver functionality in rat models with thioacetamide-induced liver cirrhosis. Naïve implanted MSCs firstly transdifferentiated into hepatic oval cells and later into HLCs. The presence of newly formed HLCs reduced inflammation, reversed fibrosis and repaired damaged hepatocytes. The exact mechanism by which MSCs induce hepatic recovery is unclear, but the authors suggest that activation of humoral factors could contribute to liver regeneration [74].

### Mechanisms of MSCs in liver regeneration

Recently, it has been suggested that MSCs modulate liver failure by several mechanisms including differentiation of MSCs to replace damaged cells, secretion of soluble factors by MSCs

to promote liver repair, MSC-mediated transfer of mitochondria by tunnelling nanotubules and by MSC-mediated transfer of proteins, RNA, hormones and chemicals by extracellular vesicles such as exosomes or microvesicles [75].

MSC conditioned medium (MSC-CM) can play an important role in attenuating liver disease with a wide range of soluble factors thought to be present within MSC-CM [76]. Interleukin-6 secreted by MSCs reduces apoptosis in liver injury [77]. Furthermore, MSC secreted TGF- $\beta$  and nerve growth factor resulted in apoptosis of hepatic stellate cells, a hallmark of liver fibrosis [78, 79]. Huang et al. showed that mice with fulminant hepatic failure (FHF) and chronic liver failure treated with MSCs or MSC-CM displayed reduced liver pathology. Only MSC treatment of FHF mice showed great reduction in pro-inflammatory T helper-1/17 cells and upregulation of T regulatory cells. This indicates that direct presence of MSCs is required to induce complete immunomodulatory effects [80].

In addition to soluble factors present in MSC-CM, recently, exosomes have been identified as an important component that may promote hepatic regeneration. Tan et al. (2014) showed that CCL4-induced liver injury reduced AST and ALT levels and decreased the number of necrotic cells in mice that were treated with MSC-derived exosomes. Furthermore, proliferation of hepatocytes was greater, which was associated with increased expression of proliferating cell nuclear antigen [81]. The authors suggest that MSC-derived exosomes may have a therapeutic potential in toxic liver injury. It has also been suggested that tunnelling nanotubules can form between cells that act as a transport network, allowing the transport of mitochondria and lysosomal vesicles [82]. Currently, the transfer of mitochondria through tunnelling nanotubules from umbilical cord-derived MSCs to hepatocytes is being investigated as a mechanism for their increased survival and function and may play a role in liver regeneration [83].

MSCs represent an ideal cell source for liver regeneration-based medicine due to their easily accessible source, their immunomodulatory properties and their potential of transdifferentiating into hepatocytes.

## Current clinical trials using cell therapy for liver-based diseases

There are now multiple phase I/II and III clinical trials using different types of stem cells to improve a number of liver diseases including cirrhosis, liver failure and liver-based metabolic disorders. In liver cirrhosis and end-stage liver disease, reports have proposed that MSCs can replace hepatocytes in the injured liver, stimulating liver regeneration (Table 2).

Shi et al. (2012) showed that transfusion of umbilical cord-MSC (UC-MSC) into 24 patients with acute-on-chronic liver failure showed marked increase in liver functionality when compared to the control of 19 patients transfused with saline.

**Table 2** A summary of the advantages and disadvantages of various cell sources which can be used to generate induced hepatocytes for hepatocyte transplantation

Study name	Cell source	Condition	Intervention	Primary outcome	Study phase	Location	Start and end date	References
Umbilical cord mesenchymal stem cell transplantation combined with plasma exchange for patients with liver failure	Umbilical mesenchymal stem cell (UC-MSC)	Liver failure	Conventional treatment only (antiviral drugs, lowering aminotransferase and jaundice medicine) plus UC-MSC transplantation (via peripheral vein slowly for 30 min $1 \times 10^5$ /kg, once a week, four times) or plasma exchange (2000 mL every 3 days, three times) or both	Survival rate and time (time frame 48 weeks)	Phase I and II	Department of Infectious Diseases, The Third Affiliated Hospital of Sun Yat-Sen University Guangzhou, Guangdong China	November 2012– March 2015	[84]
Safety and efficacy of human umbilical cord-derived mesenchymal stem cells for treatment of HBV-related liver cirrhosis	Umbilical mesenchymal stem cell (UC-MSC)	Liver cirrhosis End-stage liver disease	Conventional treatment or UC-MSC transplantation ( $1 \times 10^6$ cells/kg via hepatic artery)	One-year survival rate (time frame 1-year treatment)	Phase I and II	Xijing Hospital of Digestive Disease Xi'an, Shaanxi, China	September 2012– September 2015	[85]
Phase II safety study of two dose regimens of HepaStem in patients with ACLF (HEP101)	Human liver-derived mesenchymal stem cell (HepaStem)	Acute-on-chronic liver failure	Low-dose cohort—two dose regimens of HepaStem will be given, differing in cell quantity per infusion. The low dose regimen will be given to the first cohort (first six patients included in the study). High-dose cohort—given to the second cohort after evaluation of the safety of the first cohort (stepwise approach)	Occurrence of adverse events (AEs) up to day 28 of the active study period (time frame up to 28 days post-first infusion day)	Phase II	Hospital Erasme, Brussels, Belgium UZ Antwerpen, Edegem, Belgium KU Leuven, Leuven, Belgium CHU de Liège, Liège, Belgium Cliniques St. Luc, Woluwe-Saint Lambert, Belgium Hospital Beaujon, Clichy, France Hospital de la Croix Rousse, Lyon, France Hospital Paul Brousse, Villejuif, France Digestive Disease Research Center, Shariati Hospital, North Kargar Ave., Tehran, Iran, Islamic Republic	December 2016– September 2018	[86]
Bone marrow stem cells as a source of allogenic hepatocyte transplantation in homozygous familial hypercholesterolemia	Bone marrow stem cells	Familial hypercholesterolemia	Bone marrow stem cell transplantation. $6 \times 10^8$ to $1 \times 10^9$ cells infused through the portal vein over 30 min, done once	Serum cholesterol and LDL level (time frame 6 months)	Phase I	Digestive Disease Research Center, Shariati Hospital, North Kargar Ave., Tehran, Iran, Islamic Republic	June 2007– June 2008	[87]

Table 2 (continued)

Study name	Cell source	Condition	Intervention	Primary outcome	Study phase	Location	Start and end date	References
Study to evaluate the efficacy of HepaStem in urea cycle disorders of paediatric patients (HEP002)	Human liver-derived mesenchymal stem cell (HepaStem)	Urea cycle disorders	HepaStem administered in maximum four infusion days, spread over 8 weeks, with 2/3-week interval between infusions. Target total dose $5 \times 10^7$ /kg body weight	Efficacy as determined by de novo ureagenesis (C13 tracer method) (time frame 6 months post-first infusion day)	Phase II	Cliniques Universitaires Saint-Luc, Brussels, Belgium, Hpital Jeanne de Flandre, CHRU Lille, Lille, France Instytut-Pomnik Centrum Zdrowia Dziecka, Warszawa, Poland Hospital Materno Infantil de Badajoz, Badajoz, Spain Hospital Universitari Vall d'Hebron de Barcelona, Spain Hospital Materno Infantil de Málaga, Málaga, Spain	October 2014– March 2017	[88]
Safety and tolerance of immunomodulating therapy with donor-specific MSC in paediatric liver-donor liver transplantation (MYSTEP1)	Bone marrow-derived MSCs	Paediatric liver transplantation	Two doses of $1 \times 10^6$ MSCs/kg body weight	Number of participants with MYSTEP-score grade 3 and grade 2 (toxicity of MSC infusion), number of participants with occurrence of any severe adverse events, graft function after liver transplantation, number of participants with abnormal liver tests)	Phase I	University Children's Hospital, Tübingen, Germany	July 2013– January 2019	[89]
Therapeutic strategy and the role of mesenchymal stromal cells for ABO incompatible liver transplantation	Mesenchymal stem cells	Liver transplantation	Six doses of $1 \times 10^6$ /kg body weight MSCs are given, intravenously	Efficacy 1-year graft survival rate	Phase I	The Third Affiliated Hospital, Sun Yat-Sen University, Guangzhou, Guangdong, China	February 2014– March 2017	[90]
Human mesenchymal stem cell transfusion is safe and improves liver function in acute-on-chronic liver failure patients	Umbilical cord mesenchymal stem cell (UC-MSC)	Acute-on-chronic liver failure	Conventional treatment and $0.5 \times 10^6$ /kg body weight UC-MSCs are given, intravenously at baseline, 4 weeks, and 8 weeks	Liver functionality tested over 48 weeks 72-week survival rate	Phase I and II	Beijing 302 Hospital Beijing, Beijing, China	March 2009– March 2014	[91]



**Table 2** (continued)

Study name	Cell source	Condition	Intervention	Primary outcome	Study phase	Location	Start and end date	References
Safety study of HepaStem for the treatment of urea cycle disorders (UCD) and Crigler-Najjar syndrome (CN) (HEP001)	Human liver-derived mesenchymal stem cell (HepaStem)	Urea cycle disorders Crigler Najjar syndrome	Conventional treatment and saline was used for the control group HepaStem low dose $12.5 \times 10^6$ /kg body weight HepaStem intermediate dose $50 \times 10^6$ /kg body weight HepaStem high dose $200 \times 10^6$ /kg body weight	Safety of HepaStem in paediatric patients suffering from urea cycle disorder and Crigler-Najjar syndrome	Phase I and II	Saint Luc University Hospital, Brussels, Belgium Universitair Ziekenhuis (US) Antwerpen, Edegem Belgium CHU Bicetre, Le Kremlin Bicetre Cedex, France Hôpital Jeanne de Flandre, CHRU Lille, Lille Cedex, France Hôpital des Enfants, CHU de Toulouse, Toulouse, Cedex 9, France Rambam Medical Center, Meyer Children's Hospital, Haifa, Israel Hadassah Ein-Kerem Medical Center of Israel, Jerusalem, Israel Schneider Children's Medical Center of Israel, Petach Tikva, Israel Ospedale Pediatrico Bambino Gesù di Roma, Roma, Italy Birmingham Children's Hospital London, London, United Kingdom	March 2012–April 2015	[92]
Macrophage therapy for liver cirrhosis	Autologous macrophages	Liver cirrhosis	Autologous activated macrophages infusion via peripheral vein for 30 min. Standard medical care used as the control	Liver function (MELD score) at 3 months	Phase I and II	Edinburgh Royal Infirmary Little France Crescent Edinburgh EH16 4SA United Kingdom	August 2016–August 2021	[93]

Information from [Clinicaltrials.gov](http://Clinicaltrials.gov) [80]

Patients were monitored over 48 weeks, with the treatment group showing an increase in albumin secretion, platelet count and a reduced end-stage liver disease (MELD) score. Furthermore, survival rate after 72 weeks was also higher in the treatment group compared to the control, with 20.8 and 47.4% mortality rate, respectively. The author suggests that although the mechanism of improved liver function may be unclear, *in vivo* differentiation of UC-MSC into hepatocytes is unlikely due to the short period of hepatic recovery and with only one treatment patient showing increased alpha-fetoprotein levels. It is more likely that soluble factors produced by MSCs may enhance liver revascularization and proliferation [95].

One study has suggested that plasma exchange (PE) helps promote liver regeneration and recovery, leading to UC-MSC differentiation into HLCs. A phase I/II clinical trial is now in progress, transplanting UC-MSCs into patients with liver failure. Patients received either conventional treatment (anti-viral drugs) with UC-MSCs and/or PE treatment, and survival rates were assessed at 48 weeks [84]. For patients with acute-on-chronic liver failure, Promethera Biosciences have developed a product known as HepaStem, which are MSCs that have the potential to differentiate into HLCs. A phase IIa clinical trial is now in progress, transplanting these cells via IV injection to establish the safety and biological efficacy of these cells. Bilirubin, creatinine, INR and albumin values are being assessed at day 28, 2 months and 1 year post-infusion. In addition to using stem cells for liver failure, HLCs are now being used for clinical HT to replace primary hepatocytes in patients with liver-based metabolic disorders. Bone marrow-derived MSCs transdifferentiated into hepatocytes have been transplanted via the portal vein into patients with familial hypercholesterolemia. Serum cholesterol/LDL levels were assessed after 6 months to determine the efficacy of the technique. Furthermore, HepaStem cells are also being used to treat patients suffering from urea cycle disorders. Ureagenesis, ammonia values and amino acid levels are being monitored as well as behaviour, cognitive skills and health-related quality of life indicators for up to 12 months post-infusion [96].

MSCs are also being used clinically for immunomodulating therapy in many liver-based applications. One trial is currently investigating the use of MSCs to promote allograft tolerance and reduce the toxicity that results from exposure to calcineurin inhibitors. Paediatric patients receiving a liver transplantation undergo IV injection of bone marrow-derived MSCs. MSC toxicity is being monitored as well as graft function measured by aminotransferase and gamma glutamyl transferase activity, bilirubin, albumin and INR and the individual need for immunosuppressive medication. In addition, MSCs are being used as immunomodulators in ABO-incompatible liver transplantation. The study aims to determine if MSCs are safe and effective at reducing the primary non-function, acute rejection, ischaemic-type biliary lesions and morbidity in ABO-incompatible liver transplantation.

Another promising area could be the use of macrophage therapy to treat liver disease. Macrophages reduce scar tissue and stimulate the HPCs to expand and differentiate into mature hepatocytes. Thomas et al. showed that bone marrow-derived macrophages (BMM) administered to mice with advanced liver fibrosis resulted in a degradation of fibrillar collagen and reduced fibrogenesis. There was also upregulation of the liver progenitor cell mitogen tumour necrosis factor-like weak inducer of apoptosis that was associated with an expansion of the progenitor cell compartment [97]. There are ongoing clinical trials to assess the role macrophage therapy could play in liver cirrhosis [93]. Significant advances have been made to translate the use of stem cells to promote liver regeneration and mature hepatic differentiation into clinical use. Currently, most trials are in early phase I/II and results demonstrating the efficacy of these techniques are yet to be published. In the near future, the full potential of stem cells for liver regeneration in patients with liver disease may be better established.

## Conclusion

HT offers an alternative therapy to OLT with the aim of treating liver-based metabolic diseases or ALF. Advances in liver cell therapy are being researched to overcome the obstacles associated with HT, particularly the shortage of healthy donor hepatocytes. Although HLCs are promising, no alternative cell source can yet replace the functionality and efficacy of primary human hepatocytes. MSCs likely hold the greatest attribute as immunomodulators, and co-culturing with mature donor hepatocytes. More clinical trials assessing the safety and efficacy of HLCs is pivotal before they can be considered as a reliable cell source. In the future, this may allow for liver-based diseases to be effectively treated without the need for OLT.

## Compliance with ethical standards

**Conflict of interest** The authors declare that they have no conflict of interest.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

1. Mulaikal TA, Emond JC (2012) Physiology and anatomy of the liver. In: Liver anaesthesiology and critical care medicine. Springer New York, New York, NY, pp 3–20
2. Bernal W, Wendon J (2013) Acute liver failure. *N Engl J Med* 369: 2525–2534

3. Schilsky ML, Mistry P (2001) Metabolic liver disease. *Curr Opin Gastroenterol* 17:221–231
4. Simpson KJ, Bates CM, Henderson NC, Wigmore SJ, Garden OJ, Lee A, Pollok A, Masterton G, Hayes PC (2009) The utilization of liver transplantation in the management of acute liver failure: comparison between acetaminophen and non-acetaminophen etiologies. *Liver Transpl* 15:600–609
5. Sze YK, Dhawan A, Taylor RM, Bansal S, Mieli-Vergani G, Rela M, Heaton N (2009) Pediatric liver transplantation for metabolic liver disease: experience at King's college hospital. *Transplantation* 87:87–93
6. Jadowiec CC, Taner T (2016) Liver transplantation: current status and challenges. *World J Gastroenterol* 22:4438–4445
7. Dhawan A, Puppi J, Hughes RD, Mitry RR (2010) Human hepatocyte transplantation: current experience and future challenges. *Nat Rev Gastroenterol Hepatol* 7:288–298
8. Hansel MC, Gramignoli R, Skvorak KJ et al (2014) The history and use of human hepatocytes for the treatment of liver diseases: the first 100 patients. *Curr Protoc Toxicol* 62:14.12.1–14.1223
9. Enosawa S, Yuan W, Douzen M, Nakazawa A, Omasa T, Fukuda A, Sakamoto S, Shigeta T, Kasahara M (2012) Consideration of a safe protocol for hepatocyte transplantation using infantile pigs. *Cell Med* 3:13–18
10. Hossein Aghdaie M, Geramizadeh B, Azarpira N et al (2013) Hepatocyte isolation from unused/rejected livers for transplantation: initial step toward hepatocyte transplantation, the first experience from Iran. *Hepat Mon* 13:e10397. <https://doi.org/10.5812/hepatmon.10397>
11. Wang F, Zhou L, Ma X, Ma W, Wang C, Lu Y, Chen Y, An L, An W, Yang Y (2014) Monitoring of Intrasplenic hepatocyte transplantation for acute-on-chronic liver failure: a prospective five-year follow-up study. *Transplant Proc* 46:192–198
12. Fitzpatrick E, Mitry RR, Dhawan A (2009) Human hepatocyte transplantation: state of the art. *J Intern Med* 266:339–357
13. Meyburg J, Hoerster F, Weitz J, Hoffmann GF, Schmidt J (2008) Use of the middle colic vein for liver cell transplantation in infants and small children. *Transplant Proc* 40:936–937
14. Strom SC, Fisher RA, Rubinstein WS, Barranger JA, Towbin RB, Charron M, Miele L, Pizarov LA, Dorko K, Thompson MT et al (1997) Transplantation of human hepatocytes. *Transplant Proc* 29:2103–2106
15. Strom S, Chowdhury J, Fox I (1999) Hepatocyte transplantation for the treatment of human disease. *Semin Liver Dis* 19:39–48
16. Dhawan A, Mitry RR, Hughes RD (2006) Hepatocyte transplantation for liver-based metabolic disorders. *J Inherit Metab Dis* 29:431–435
17. Fox IJ, Chowdhury JR, Kaufman SS, Goertzen TC, Chowdhury NR, Warkentin PI, Dorko K, Sauter BV, Strom SC (1998) Treatment of the Crigler–Najjar syndrome type I with hepatocyte transplantation. *N Engl J Med* 338:1422–1427
18. Ambrosino G, Varotto S, Strom SC, Guariso G, Franchin E, Miotto D, Caenazzo L, Basso S, Carraro P, Valente ML et al (2005) Isolated hepatocyte transplantation for Crigler–Najjar syndrome type I. *Cell Transplant* 14:151–157
19. Lysy P-A, Najimi M, Stephenne X et al (2008) Liver cell transplantation for Crigler–Najjar syndrome type I: update and perspectives. *World J Gastroenterol* 14:3464–3470
20. Allen KJ, Mifsud NA, Williamson R, Bertolino P, Hardikar W (2008) Cell-mediated rejection results in allograft loss after liver cell transplantation. *Liver Transpl* 14:688–694
21. Darwish AA, Sokal E, Stephenne X, Najimi M, de Goyet JV, Reding R (2004) Permanent access to the portal system for cellular transplantation using an implantable port device. *Liver Transpl* 10:1213–1215
22. Horslen SP, McCowan TC, Goertzen TC et al (2003) Isolated hepatocyte transplantation in an infant with a severe urea cycle disorder. *Pediatrics* 111:1262–1267
23. Puppi J, Tan N, Mitry RR, Hughes RD, Lehec S, Mieli-Vergani G, Karani J, Champion MP, Heaton N, Mohamed R et al (2008) Hepatocyte transplantation followed by auxiliary liver transplantation—a novel treatment for ornithine transcarbamylase deficiency. *Am J Transplant* 8:452–457
24. Stephenne X, Najimi M, Smets F, Reding R, de Goyet JV, Sokal EM (2005) Cryopreserved liver cell transplantation controls ornithine transcarbamylase deficient patient while awaiting liver transplantation. *Am J Transplant* 5:2058–2061
25. Meyburg J, Das AM, Hoerster F, Lindner M, Kriegbaum H, Engelmann G, Schmidt J, Ott M, Pettenazzo A, Luecke T et al (2009) One liver for four children: first clinical series of liver cell transplantation for severe neonatal urea cycle defects. *Transplantation* 87:636–641
26. Grossman M, Rader DJ, Muller DW et al (1995) A pilot study of ex vivo gene therapy for homozygous familial hypercholesterolaemia. *Nat Med* 1:1148–1154
27. Dhawan A, Mitry RR, Hughes RD, Lehec S, Terry C, Bansal S, Arya R, Wade JJ, Verma A, Heaton ND et al (2004) Hepatocyte transplantation for inherited factor VII deficiency. *Transplantation* 78:1812–1814
28. Stéphenne X, Najimi M, Sibille C, Nassogne M–C, Smets F, Sokal EM (2006) Sustained engraftment and tissue enzyme activity after liver cell transplantation for argininosuccinate lyase deficiency. *Gastroenterology* 130:1317–1323
29. Bilir BM, Guinette D, Karrer F, Kumpe DA, Krysl J, Stephens J, McGavran L, Ostrowska A, Durham J (2000) Hepatocyte transplantation in acute liver failure. *Liver Transpl* 6:32–40
30. Fisher RA, Strom SC (2006) Human hepatocyte transplantation: worldwide results. *Transplantation* 82:441–449
31. Squires RH, Shneider BL, Bucuvalas J et al (2006) Acute liver failure in children: the first 348 patients in the pediatric acute liver failure study group. *J Pediatr* 148:652–658
32. Soriano HE, Wood RP, Kang D-C, Ozaki CF, Finegold MJ, Darlington GJ, Ferry G (1996) Hepatocellular transplantation (HCT) via portal vein catheter in a patient with fulminant liver failure. *† 746. Pediatr Res* 39:127–127
33. Habibullah CM, Syed IH, Qamar A, Taher-Uz Z (1994) Human fetal hepatocyte transplantation in patients with fulminant hepatic failure. *Transplantation* 58:951–952
34. Jitraruch S, Dhawan A, Hughes RD, Filippi C, Soong D, Philippeos C, Lehec SC, Heaton ND, Longhi MS, Mitry RR (2014) Alginate microencapsulated hepatocytes optimised for transplantation in acute liver failure. *PLoS One* 9:e113609
35. Sgroi A, Mai G, Morel P, Baertschiger RM, Gonelle-Gispert C, Serre-Beinier V, Buhler LH (2011) Transplantation of encapsulated hepatocytes during acute liver failure improves survival without stimulating native liver regeneration. *Cell Transplant* 20:1791–1803
36. Teramura Y, Oommen OP, Olerud J, Hilborn J, Nilsson B (2013) Microencapsulation of cells, including islets, within stable ultra-thin membranes of maleimide-conjugated PEG-lipid with multifunctional crosslinkers. *Biomaterials* 34:2683–2693
37. Gupta S, Rajvanshi P, Sokhi R, Slehra S, Yam A, Kerr A, Novikoff PM (1999) Entry and integration of transplanted hepatocytes in rat liver plates occur by disruption of hepatic sinusoidal endothelium. *Hepatology* 29:509–519
38. Lee CA, Dhawan A, Smith RA, Mitry RR, Fitzpatrick E (2016) Instant-blood mediated inflammatory reaction in hepatocyte transplantation: current status and future perspectives. *Cell Transplant* 25:1227–1236
39. Joseph B, Malhi H, Bhargava KK, Palestro CJ, McCuskey RS, Gupta S (2002) Kupffer cells participate in early clearance of

- syngeneic hepatocytes transplanted in the rat liver. *Gastroenterology* 123:1677–1685
40. Yokoyama I, Tzakis AG, Inventarza O, Todo S, Casavilla A, Leggio A, Starzl TE (1992) Pediatric liver transplantation from neonatal donors. *Transpl Int* 5:205–208
  41. Tolosa L, Pareja-Ibars E, Donato MT, Cortés M, López S, Jiménez N, Mir J, Castell JV, Gómez-Lechón MJ (2014) Neonatal livers: a source for the isolation of good-performing hepatocytes for cell transplantation. *Cell Transplant* 23:1229–1242
  42. Lee C, Dhawan A, Iansante V, et al (2018) Cryopreserved neonatal hepatocytes may be a source for human application; evaluation of functionality towards clinical use. *Liver Transpl* (accepted awaiting Publ)
  43. Tolosa L, López S, Pareja E, Donato MT, Myara A, Nguyen TH, Castell JV, Gómez-Lechón MJ (2015) Human neonatal hepatocyte transplantation induces long-term rescue of unconjugated hyperbilirubinemia in the Gunn rat. *Liver Transpl* 21:801–811
  44. Sauer V, Roy-Chowdhury N, Guha C, Roy-Chowdhury J (2014) Induced pluripotent stem cells as a source of hepatocytes. *Curr Pathobiol Rep* 2:11–20
  45. Roy-Chowdhury N, Wang X, Guha C, Roy-Chowdhury J (2017) Hepatocyte-like cells derived from induced pluripotent stem cells. *Hepatol Int* 11:54–69
  46. Nagamoto Y, Tashiro K, Takayama K, Ohashi K, Kawabata K, Sakurai F, Tachibana M, Hayakawa T, Furue MK, Mizuguchi H (2012) The promotion of hepatic maturation of human pluripotent stem cells in 3D co-culture using type I collagen and Swiss 3T3 cell sheets. *Biomaterials* 33:4526–4534
  47. Si-Tayeb K, Noto FK, Nagaoka M, Li J, Battle MA, Duris C, North PE, Dalton S, Duncan SA (2010) Highly efficient generation of human hepatocyte-like cells from induced pluripotent stem cells. *Hepatology* 51:297–305
  48. Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, Tomoda K, Yamanaka S (2007) Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell* 131:861–872
  49. Hsu D, Katelaris C (2009) Long-term management of patients taking immunosuppressive drugs. *Aust Prescr* 32:68–71
  50. Yu Y, Liu H, Ikeda Y, Amiot BP, Rinaldo P, Duncan SA, Nyberg SL (2012) Hepatocyte-like cells differentiated from human induced pluripotent stem cells: relevance to cellular therapies. *Stem Cell Res* 9:196–207
  51. Rashid ST, Corbineau S, Hannan N, Marciniak SJ, Miranda E, Alexander G, Huang-Doran I, Griffin J, Ahrlund-Richter L, Skepper J et al (2010) Modeling inherited metabolic disorders of the liver using human induced pluripotent stem cells. *J Clin Invest* 120:3127–3136
  52. Miura K, Okada Y, Aoi T, Okada A, Takahashi K, Okita K, Nakagawa M, Koyanagi M, Tanabe K, Ohnuki M et al (2009) Variation in the safety of induced pluripotent stem cell lines. *Nat Biotechnol* 27:743–745
  53. Kajiwara M, Aoi T, Okita K, Takahashi R, Inoue H, Takayama N, Endo H, Eto K, Toguchida J, Uemoto S et al (2012) Donor-independent variations in hepatic differentiation from human-induced pluripotent stem cells. *Proc Natl Acad Sci U S A* 109:12538–12543
  54. Chen Y, Li Y, Wang X, Zhang W, Sauer V, Chang CJ, Han B, Tchaikovskaya T, Avsar Y, Tafaleng E et al (2015) Amelioration of hyperbilirubinemia in Gunn rats after transplantation of human induced pluripotent stem cell-derived hepatocytes. *Stem Cell Rep* 5:22–30
  55. Baum J, Duffy HS (2011) Fibroblasts and myofibroblasts: what are we talking about? *J Cardiovasc Pharmacol* 57:376–379
  56. Huang P, Zhang L, Gao Y, He Z, Yao D, Wu Z, Cen J, Chen X, Liu C, Hu Y et al (2014) Direct reprogramming of human fibroblasts to functional and expandable hepatocytes. *Cell Stem Cell* 14:370–384
  57. Simeonov KP, Uppal H, Gao Y et al (2014) Direct reprogramming of human fibroblasts to hepatocyte-like cells by synthetic modified mRNAs. *PLoS One* 9:e100134
  58. Du Y, Wang J, Jia J et al (2014) Human hepatocytes with drug metabolic function induced from fibroblasts by lineage reprogramming. *Cell Stem Cell* 14:394–403
  59. Ivanov NA, Tao R, Chenoweth JG, Brandtjen A, Mighdoll MI, Genova JD, McKay RD, Jia Y, Weinberger DR, Kleinman JE et al (2016) Strong components of epigenetic memory in cultured human fibroblasts related to site of origin and donor age. *PLoS Genet* 12:e1005819
  60. Tahan AC, Tahan V (2014) Placental amniotic epithelial cells and their therapeutic potential in liver diseases. *Front Med* 1:48
  61. Marongiu F, Gramignoli R, Dorko K, Miki T, Ranade AR, Paola Serra M, Doratiotto S, Sini M, Sharma S, Mitamura K et al (2011) Hepatic differentiation of amniotic epithelial cells. *Hepatology* 53:1719–1729
  62. Kitade M, Factor VM, Andersen JB, Tomokuni A, Kaji K, Akita H, Holczbauer A, Seo D, Marquardt JU, Conner EA et al (2013) Specific fate decisions in adult hepatic progenitor cells driven by MET and EGFR signaling. *Genes Dev* 27:1706–1717
  63. Zhang W, Li W, Liu B, Wang P, Li W, Zhang H (2012) Efficient generation of functional hepatocyte-like cells from human fetal hepatic progenitor cells in vitro. *J Cell Physiol* 227:2051–2058
  64. Khuu DN, Scheers I, Ehnert S, Jazouli N, Nyabi O, Buc-Calderon P, Meulemans A, Nussler A, Sokal E, Najimi M (2011) In vitro differentiated adult human liver progenitor cells display mature hepatic metabolic functions: a potential tool for in vitro pharmacotoxicological testing. *Cell Transplant* 20:287–302
  65. Sokal EM, Stéphenne X, Ottolenghi C et al (2013) Liver engraftment and repopulation by in vitro expanded adult derived human liver stem cells in a child with ornithine carbamoyltransferase deficiency. In: *JIMD reports*. pp 65–72
  66. Huch M, Gehart H, van Boxtel R, Hamer K, Blokzijl F, Versteegen MMA, Ellis E, van Wenum M, Fuchs SA, de Ligt J, van de Wetering M et al (2015) Long-term culture of genome-stable bipotent stem cells from adult human liver. *Cell* 160:299–312
  67. Tarlow BD, Pelz C, Naugler WE, Wakefield L, Wilson EM, Finegold MJ, Grompe M (2014) Bipotential adult liver progenitors are derived from chronically injured mature hepatocytes. *Cell Stem Cell* 15:605–618
  68. Katsuda T, Kawamata M, Hagiwara K, Takahashi RU, Yamamoto Y, Camargo FD, Ochiya T (2017) Conversion of terminally committed hepatocytes to culturable bipotent progenitor cells with regenerative capacity. *Cell Stem Cell* 20:41–55
  69. Stock P, Brückner S, Ebensing S, Hempel M, Dollinger MM, Christ B (2010) The generation of hepatocytes from mesenchymal stem cells and engraftment into murine liver. *Nat Protoc* 5:617–627
  70. Wu X-B, Tao R (2012) Hepatocyte differentiation of mesenchymal stem cells. *Hepatobiliary Pancreat Dis Int* 11:360–371
  71. Meier RPH, Müller YD, Morel P, Gonnelle-Gispert C, Bühler LH (2013) Transplantation of mesenchymal stem cells for the treatment of liver diseases, is there enough evidence? *Stem Cell Res* 11:1348–1364
  72. Selmani Z, Naji A, Zidi I, Favier B, Gaiffe E, Obert L, Borg C, Saas P, Tiberghien P, Rouas-Freiss N et al (2008) Human leukocyte antigen-G5 secretion by human mesenchymal stem cells is required to suppress T lymphocyte and natural killer function and to induce CD4<sup>+</sup> CD25<sup>high</sup> FOXP3<sup>+</sup> regulatory T cells. *Stem Cells* 26:212–222
  73. Faiella W, Atoui R (2016) Immunotolerant properties of mesenchymal stem cells: updated review. *Stem Cells Int* 2016:1859567–1859567
  74. Hwang S, Hong H, Kim H et al (2012) Hepatogenic differentiation of mesenchymal stem cells in a rat model of thioacetamide-induced liver cirrhosis. *Cell Biol Int* 36:279–288



75. Spees JL, Lee RH, Gregory CA Mechanisms of mesenchymal stem/stromal cell function. <https://doi.org/10.1186/s13287-016-0363-7>
76. Gu J, Shi X, Zhang Y, Ding Y (2009) Heterotypic interactions in the preservation of morphology and functionality of porcine hepatocytes by bone marrow mesenchymal stem cells in vitro. *J Cell Physiol* 219:100–108
77. van Poll D, Parekkadan B, Cho CH, Berthiaume F, Nahmias Y, Tilles AW, Yarmush ML (2008) Mesenchymal stem cell-derived molecules directly modulate hepatocellular death and regeneration in vitro and in vivo. *Hepatology* 47:1634–1643
78. Wang J, Bian C, Liao L, Zhu Y, Li J, Zeng L, Zhao RC (2009) Inhibition of hepatic stellate cells proliferation by mesenchymal stem cells and the possible mechanisms. *Hepatol Res* 39:1219–1228
79. Lin N, Hu K, Chen S, Xie S, Tang Z, Lin J, Xu R (2009) Nerve growth factor-mediated paracrine regulation of hepatic stellate cells by multipotent mesenchymal stromal cells. *Life Sci* 85:291–295
80. Huang B, Cheng X, Wang H, Huang W, la Ga hu Z, Wang D, Zhang K, Zhang H, Xue Z, Da Y et al (2016) Mesenchymal stem cells and their secreted molecules predominantly ameliorate fulminant hepatic failure and chronic liver fibrosis in mice respectively. *J Transl Med* 14:45
81. Tan C, Lai R, Wong W, Dan Y, Lim SK, Ho H (2014) Mesenchymal stem cell-derived exosomes promote hepatic regeneration in drug-induced liver injury models. *Stem Cell Res Ther* 5:76
82. Rustom A, Saffrich R, Markovic I et al (2004) Nanotubular highways for intercellular organelle transport. *Science* 303:1007–1010
83. Dacosta RF, Lee C, Walker S et al (2015) Investigation of the mitochondrial transfer from human mesenchymal stem/stromal cells to human primary hepatocytes through tunnelling nanotubes as a potential mechanism for cell survival and function enhancement during hepatocyte transplantation. *Xenotransplantation* 22: S47
84. Gao Z-L (2012) Umbilical cord mesenchymal stem cells transplantation combined with plasma exchange for patients with liver failure. In: [Clinicaltrials.gov](http://Clinicaltrials.gov)
85. Ying H (2012) Safety and efficacy of human umbilical cord derived mesenchymal stem cells for treatment of HBV-related liver cirrhosis. In: [Clinicaltrials.gov](http://Clinicaltrials.gov)
86. Promethera Biosciences (2016) Phase II safety study of 2 dose regimens of HepaStem in patients with ACLF. In: [Clinicaltrials.gov](http://Clinicaltrials.gov)
87. Malekzadeh R (2007) Bone marrow stem cells as a source of allogenic hepatocyte transplantation in homozygous familial hypercholesterolemia. In: [Clinicaltrials.gov](http://Clinicaltrials.gov)
88. Promethera Biosciences (2015) Study to evaluate the efficacy of HepaStem in urea cycle disorders Paediatric patients (HEP002). In: [Clinicaltrials.gov](http://Clinicaltrials.gov)
89. Sturm E (2016) Safety and tolerance of Immunomodulating therapy with donor-specific MSC in pediatric living-donor liver transplantation (MYSTEP1). In: [Clinicaltrials.gov](http://Clinicaltrials.gov)
90. Yang Y (2016) Therapeutic strategy and the role of mesenchymal stromal cells for ABO incompatible liver transplantation. In: [Clinicaltrials.gov](http://Clinicaltrials.gov)
91. Wang F-S (2010) Safety and efficacy of human mesenchymal stem cells for treatment of liver failure. In: [Clinicaltrials.gov](http://Clinicaltrials.gov)
92. Promethera Biosciences (2013) Safety study of HepaStem for the treatment of urea cycle disorders (UCD) and Crigler-Najjar syndrome (CN) (HEP001). In: [Clinicaltrials.gov](http://Clinicaltrials.gov)
93. Moroni F (2016) Macrophages therapy for liver cirrhosis. In: [ISRCTN.Regist](http://ISRCTN.Regist)
94. Promethera Biosciences (2016) Phase II safety study of 2 dose regimens of HepaStem in patients with ACLF. In: [Clinicaltrials.gov](http://Clinicaltrials.gov)
95. Shi M, Zhang Z, Xu R, Lin H, Fu J, Zou Z, Zhang A, Shi J, Chen L, Lv S et al (2012) Human mesenchymal stem cell transfusion is safe and improves liver function in acute-on-chronic liver failure patients. *Stem Cells Transl Med* 1:725–731
96. [ClinicalTrials.gov](http://ClinicalTrials.gov). <https://clinicaltrials.gov/ct2/results?term=alpha-1+antitrypsin&cond=diabetes>. Accessed 22 Aug 2017
97. Thomas JA, Pope C, Wojtacha D, Robson AJ, Gordon-Walker TT, Hartland S, Ramachandran P, van Deemter M, Hume DA, Iredale JP et al (2011) Macrophage therapy for murine liver fibrosis recruits host effector cells improving fibrosis, regeneration, and function. *Hepatology* 53:2003–2015
98. Promethera Biosciences (2013) Safety Study of HepaStem for the Treatment of Urea Cycle Disorders (UCD) and Crigler-Najjar Syndrome (CN) (HEP001). In: [Clinicaltrials.gov](http://Clinicaltrials.gov).