

Editorial:

KEY MESSAGES OF RECENT PUBLICATIONS IN THE FIELD OF TOXICOLOGY

C. Cadenas, R. Marchan, P. Godoy✉, R. Reif, I. von Recklinghausen, N. Schöbel

Leibniz Institut für Arbeitsforschung an der TU Dortmund, Leibniz Research Centre for Working Environment and Human Factors (IfADo), Ardeystrasse 67, 44139 Dortmund, Germany

godoy@ifado.de

In order to address the field of tension between toxicology as well as basic and clinical sciences, we closely cooperate with our partner journal Archives of Toxicology. To give our readers a rapid overview of recent developments we have summarized the key messages of the most cited publications. For better orientation we grouped the articles into the categories inorganic compounds (Table 1), toxicogenomics (Table 2), molecular toxicology (Table 3), organ toxicity and mechanisms (Table 4), reproductive toxicology (Table 5), genotoxicity and carcinogenicity (Table 6), as well as reviews and short communications (Table 7). We hope that our service facilitates the overview over recent achievements in toxicology and stimulates further reading.

Table 1: Inorganic compounds

Key message	Reference
Flavonoids have been reported to provide neuroprotection. However, this article provides evidence of a more complex situation: Both quercetin and quercitrin protected the mitochondria of rat brain slices from MeHg-induced lipid peroxidation. In contrast, rutin was not protective. Ca^{2+} plays a central role in MeHg-induced toxicity.	Wagner et al., 2010
Silicia nanoparticles (SiO_2 -NPs) were found in the endosomes and the cytosol of HeLa cells. No accumulation in mitochondria or nuclei was seen. In contrast, the larger 'submicron particles' (SiO_2 -SMPs) accumulated in lysosomes.	Al-Rawi et al., 2011
The toxicokinetics of thiomersal is completely different from that of methylmercury. Therefore, toxicity data from methylmercury studies are not appropriate when assessing the risk from mercury released from the ethylmercury-releasing preservative, thiomersal.	Rodrigues et al., 2010
The aspect ratio (defined as the ratio length: diameter) of carbon nanotubes has no influence on genotoxicity.	Kim et al., 2011a
Inorganic arsenic induces apoptosis in the cerebrum of mice.	Yen et al., 2011
Transition metal ions induce lipid peroxidation in artificial phospholipid liposomes.	Repetto et al., 2010
Zinc oxide nanoparticles induce the release of pro-inflammatory cytokines in mouse and human cell systems.	Heng et al., 2011
DNA strand breaks induced by platinum nanoparticles are mediated by platinum ions released from the nanoparticles.	Gehrke et al., 2011

Table 1 (cont.): Inorganic compounds

Key message	Reference
This review deals with the description and comparison of cyclotron-based irradiation techniques for the generation of radiolabelled nanoparticles applicable in nanotoxicity tracing approaches.	Gibson et al., 2011
Possible estrogenic effects of cadmium were analyzed in the rat intestine. Cadmium exposure was shown to modulate molecular and functional parameters of estrogenicity such as proliferation and expression of the estrogen-regulated gene ER beta.	Höfer et al., 2010
Only minor pulmonary irritation and inflammation potencies were found for TiO ₂ nanoparticles in standardised mouse bioassays.	Leppänen et al., 2011
The importance of nanoparticle surface functionalization could be shown in an in vivo model of embryonic zebrafish, whereby two lead sulfide nanoparticles with identical core size but different surface functionality led to drastic differences in embryo mortality.	Truong et al., 2011
Studies on genetic susceptibility to carcinogenicity are fundamental for improving risk assessment strategies. This report shows that the polymorphic status at the locus of A35931C in XPD/ERCC2 might modulate the risk for arsenic-related skin lesions.	Lin et al., 2010
By assessing time-dependent pathological alterations in rats administered subtoxic or toxic MeHg doses for 3 weeks, the authors determine that MeHg at toxic doses causes selective functional alteration of choroid plexus before leading to pathological alterations in the brain.	Nakamura et al., 2011
This study shows that normal working conditions in a boric acid production plant present low risk of boron toxicity, as indicated by daily boron exposure levels lower than those associated with developmental and reproductive toxic effects in animals and absence of reproductive toxicity indicators in those workers.	Duydu et al., 2011
Welding fume is a complex aerosol of different metals. This study indicates that stainless steel welding fume induces more lung injury and inflammation compared with mild steel fume after a short-term inhalation exposure in rats.	Antonini et al., 2011
Sperm hyperactivity, needed for successful fertilization, is decreased when exposed to sodium fluoride in a dose dependent manner. Exposure to elevated fluoride levels decreased intracellular Ca ²⁺ and downstream Ca ²⁺ -dependent signalling pathways.	Sun et al., 2010
Silver nanoparticles (Ag NPs) have significant antimicrobial properties, and for the first time, the mechanism underlying their toxicity was investigated. Ag NPs caused death by inhibiting proliferation, upregulating pro apoptotic Bcl-2 members, and inhibiting PKC family members. PKC zeta was shown to be particularly important for Ag NP-mediated toxicity.	Lee et al., 2011a
Exposure to manganese chloride (MnCl) can lead to neurotoxicity with Parkinson-like symptoms. The increased oxidative damage induced by MnCl is accompanied by accumulation of NF-E2-related factor2 (Nrf2) in the cytoplasm and nucleus, which subsequently binds to the antioxidant response element (ARE) of antioxidant genes. The addition of an additional antioxidant, tert-butylhydroquinone (t-BHQ), prevents the cytotoxicity induced by MnCl.	Li et al., 2011a
The search for cancer-predictive biomarkers continues to be challenging as three potential biomarkers - SMRP, CA125 and CYFRA21-1 - proved to be highly specific, but not very sensitive, as predictors of lung cancer and malignant mesothelioma in a cohort of workers exposed to asbestos.	Gube et al., 2011

Table 1 (cont.): Inorganic compounds

Key message	Reference
Parallel application of glutathione during inorganic arsenic treatment can promote arsenic metabolism by methylation and might facilitate excretion of arsenicals.	Wang et al., 2011a
Investigation of inorganic arsenite on four different rat derived cell lines gives hints that Nrf2 might be the reason for cardiac toxicity of arsenite.	Sumi et al., 2011
Upon exposure to high fluoride and low iodine, rat brain proteins related to cell signaling, energy metabolism, and protein metabolism were found to be deregulated.	Ge et al., 2011
Maternal injection of cisplatin to pregnant guinea pigs lead to hearing loss and cochlear hair cell apoptosis in pups.	Hao et al., 2011
The dispersion stability of carbon nanotubes (CNTs) used in toxicity tests was evaluated. During 16 weeks dispersion stability decreased in the following order: BSA, Tween 80, DPPC, and DMSO for the MWCNTs and BSA, DPPC, Tween 80, and DMSO for the SWCNTs.	Kim et al., 2011b
For the first time it was reported that two codominant alleles at a genetic locus are responsible for the susceptibility to Cd-induced testicular toxicity in inbred rat strains.	Shimada et al., 2011
Methylation of iAs has been implicated in arsenic toxicity, however, the mechanism of methylation is not fully understood. The study shows that AS3MT expression generates dimethylated arsenicals which enhanced the cytotoxic effect of arsenite.	Watanabe et al., 2011
Mercuric chloride induced urinary excretion of Oat5 in rats, suggesting that it might be an early indicator of mercury-induced nephropathy, which predicts the perturbation before the manifestation of histopathological damages.	Di Giusto et al., 2010

Table 2: Toxicogenomics

Key message	Reference
Gene expression alterations in the brains of neonate mice exposed to methylmercury and polychlorinated biphenyls, alone or in combination, reveal not only toxicity effects but also a protective, detoxication response upon co-exposure.	Shimada et al., 2010
The mechanism underlying perfluorooctane (PFOS)-mediated decrease in circulating thyroid hormone levels was studied. It was shown that PFOS increased hepatic expression of OATP2 and MRP2 leading to enhanced hepatic uptake and metabolism of T4.	Yu et al., 2011
Using recombinant UGT2B15 enzymes (wild type and mutants) the authors establish that variants with D85Y substitution have a markedly reduced V_{max} and CL_{int} values, with no changes in K_m .	Hanioka et al., 2011
Ethanol can induce severe toxic effects in developmental brain. By gene array analysis in a model of neural stem cell differentiation in vitro, the authors identify two main gene clusters influenced by ethanol: a Wnt-Fzd and a cluster of heat shock proteins genes and the heat shock transcription factor HSF5.	Choi et al., 2011a

Table 2 (cont.): Toxicogenomics

Key message	Reference
Second-generation antihistamine drugs overdose or reduced metabolism may increase ventricular arrhythmia. The present study identified gene signatures indicative of toxicity, namely cardiotoxicity, oxidative stress, heat shock proteins, hypoxia and inflammation, suggesting that these sets of these genes could serve as biomarkers for cardiotoxicity.	Lee et al., 2010
This paper describes an analysis of the mutation pattern by benzene in the TP53 gene using the 'Functional Analysis of Separated Allels in Yeast' technique.	Billet et al., 2010
The authors found a strong correlation between proteasomal induction and lipid elevations in a rat model for HIV protease inhibitor-induced hyperlipidemia. The identification of proteasome gene regulation allowed them to develop a rapid screen for identifying novel PIs that do not induce the proteasome.	Waring et al., 2010
This publication is the first comprehensive gene expression profiling study conducted for welding fume exposure in monkeys. The expressed genes are expected to be useful in helping to understand transcriptional changes in monkey lungs after welding fume exposure.	Heo et al., 2010
The study shows that a genetic polymorphism of the insulin-like growth factor (IGFBP-3) binding protein-3 gene may be involved in the etiology of bladder transitional cell carcinoma (TCC). Larger studies are needed for further confirmation.	Safarinejad et al., 2011
The combination of hepatic transcriptomics and proteomics was used to investigate the molecular dynamics of glutathione depletion induced by diethyl maleate (DEM) in rat. The genes Trib3, Srxn1, Myc, Asns, Igfbp1, Txnrd1, or Hmox1, were suggested as robust mRNA biomarkers for evaluating hepatic glutathione depletion. L-FABP, MAWDBP, aldo-keto reductase family 1 member A1, catalase and ATP synthase subunit beta were suggested as potential protein biomarkers.	Yamauchi et al., 2011

Table 3: Molecular toxicology

Key message	Reference
Standardisation of the cell response to sodium nitroprusside (SNP) revealed that long-term culturing-associated resistance to SNP-induced cell toxicity was accompanied by higher levels of the stress protein Hsp70. Suggested mechanism of action of Hsp70 includes increase of CAT and GSH-Px activities as well as decrease in caspase-3 activation.	Romero et al., 2010
Based on its selective pro-apoptotic effect over the tumor cell line Hep G2 and not on the normal WRL68 liver cells, the novel <i>P. vulgaris</i> lectin has a potential therapeutic use on liver cancer.	Fang et al., 2011
This article describes detailed molecular mechanisms of ceramide induced apoptosis in stem cells, initiated by ROS and propagated by caspases.	Park et al., 2011a
Due to its antioxidant potential, diphenyl diselenide has been studied as a potential protective molecule. In this article, the authors present clear evidence on a cytotoxic potential of (PhSe) ₂ in a neuronal cell line, mediated by the ERK1/2 pathway.	Posser et al., 2011

Table 3 (cont.): Molecular toxicology

Key message	Reference
Methylation changes in mouse liver, in response to a methyl deficient diet with or without arsenic, were sex-dependent. This may account for the differences in susceptibility of males and females to various diseases that are dependent on epigenetic modifications.	Nohara et al., 2011
Inducing the drug transporter, p-glycoprotein (P-gp) with doxorubicin decreased paraquat-induced cytotoxicity; a protection reversed by inhibiting P-gp with a specific inhibitor. Caco-2 cells proved to be an efficient model to study the mechanisms of paraquat toxicity.	Silva et al., 2011
Lipopolysaccharide (LPS) can cause septic shock, a response regulated by alveolar type II epithelial cells. In this study, the authors showed that LPS induces a time and concentration dependent decrease in A549 cell viability, with a concurrent increase in reactive oxygen species (ROS) and apoptosis via the intrinsic mitochondria-dependent pathway. Pretreating the cells with the antioxidant, N-acetylcysteine attenuated LPS-induced ROS formation and cell death.	Chuang et al., 2011
Phosphate and tyrode buffer are suitable for inhibition and reactivation kinetics of human erythrocyte AChE, while TRIS and MOPS buffer markedly changed kinetics of AChE.	Wille et al., 2011
The shellfish toxins okadaic acid and dinophysin toxin 2 show different cell cycle arrests in HepG2 compared to clone9 cells; in addition different expressions of cyclins and cyclin dependent kinases had been observed.	Rubiolo et al., 2011
Entry pathway of tested nanoparticles shifted from direct penetration to endocytosis depending on size. 30 nm particles led to reduction in Zn ²⁺ and Mg ²⁺ content in nuclei and to increased p53 in the whole cell. While particles of 50 -100 nm localised in acidic vesicles and showed increased p53 in nuclei.	Zhang et al., 2011
Depleting regulatory T cells in mice aggravates 1,3-beta-glucan induced lung inflammation.	Liu et al., 2011a
Sub-chronic exposure of mice to perfluorooctanesulfonate altered the host's immune state towards a more T(H)2-like state with increased IL-4, IL-10, IgG, IgG1, and IgE, but decreased IL-2, IFN-gamma, and IgM levels.	Dong et al., 2011
A neurotoxin-like protein derived from Egyptian cobra venom was fractionated chromatographically. The venom fractions exerted various cytotoxic effects against different cancer cells.	El Hakim et al., 2011
(S)-dimethyl 2-(3-(phenyltellanyl) propanamido) succinate, a telluro-amino acid derivative showed glutathione peroxidase-like activity in rat brain homogenates, but genotoxic and mutagenic effects in adult male mice.	Meinerz et al., 2011
The authors developed a novel proteomic analysis platform which they named antigen-subtracted 2-DE/MS strategy. The article describes the new technique and includes a comparative proteomic analysis for 16HBE and 16HBE-C GO/G1 cell proteins. New proteins involved in carcinogenesis, oxidative stress and protein synthesis were identified and showed that this new technique improves resolution of protein samples.	Zhao et al., 2011a
The proteasome inhibitor MG132 reduces growth of As4.1 juxtaglomerular cells via caspase-independent apoptosis. The changes in ROS and GSH levels by MG132 and caspase inhibitors partially influenced the growth inhibition and death of As4.1 cells.	Han et al., 2010a

Table 3 (cont.): Molecular toxicology

Key message	Reference
In their in vivo assessment of antiemetic drugs in Beagle dogs, the authors found evidence that predominantly Neurokinin-1 (NK(1)) and to a lesser extent 5-hydroxytryptamine 3 (5-HT(3)) receptors are involved in lycorine-induced emesis facilitating a target-orientated therapy.	Kretzing et al., 2011
This paper describes a biochemical and pharmacological characterization of BpirPLA(2)-I, the first acidic Asp49-PLA(2) isolated from Bothrops pirajai snake venom. The authors suggest a possible relationship between the basic or acid character of PLA(2) enzymes and the functionality of the Ca ²⁺ ion binding loop.	Teixeira et al., 2011
'Designer steroids' are misused for doping purposes in the bodybuilding scene. Results suggest that the MHC expression shift (MHC II/d/x upregulated, MHC II/b downregulated) induced by NOR, madol and TP could serve as a molecular marker to determine anabolic activity of anabolic steroids at least in skeletal muscle of orchi rats.	Frese et al., 2011
The study shows that amlodipine exerts a significant inhibition on adriamycin-induced toxicity in rat mesangial cells by affecting the expression of TGF-β/Smad signaling intermediates p-Smad2 and Smad4.	Song et al., 2011
The authors demonstrate that PAHs found in the Chattanooga Superfund site induce apoptosis in vitro via activation of group IVC PLA2, and this may be the mechanism for vascular damage in mice inhabiting the Superfund site.	Tithof et al., 2011
Acrolein significantly downregulated cellular NF-κB activity as well as LPS- induced NO production in human skin cells without altering cellular levels of the phosphorylated and nonphosphorylated forms of Iκ-Bα, implying that the effect of acrolein on cellular NF-κB activity is independent of Iκ-Bα.	Moon, 2011
The authors report potent proteolytic and neurotoxic but no hemorrhagic activities in crude toxins from salivary excretion (SE) of H. angulatus. In addition, the authors purify a cysteine-rich secretory protein called helicopsin from the SE that exhibited robust neurotoxic activity.	Estrella et al., 2011
The authors report two main effects of Brevetoxin toxicity in mast cells: an initial degranulation and increase in IL-6 expression independent of IgE receptor, followed by cellular toxicity within 24 h by LDH release and Annexin-V staining.	Hilderbrand et al., 2011

Table 4: Organ toxicity and mechanisms

Key message	Reference
The pyrethroid insecticide, cypermethrin disrupts testosterone synthesis in testes of mice.	Wang et al., 2010
Sodium fluoride suppresses proliferation and induces apoptosis in cultivated osteoblasts. This effect was caused by decreased insulin-like growth factor-1 expression.	Wang et al., 2011b
Administration of silver nanoparticles to rats caused a dose-dependent accumulation of particles in the lamina propria of the small and large intestine, increased numbers of goblet cells, and altered mucus composition.	Jeong et al., 2010a

Table 4 (cont.): Organ toxicity and mechanisms

Key message	Reference
A single intratracheal instillation of carbon nanotubes may induce early lung fibrosis.	Park et al., 2011b
The genotoxic potential of dental composite components, such as BisGMA, TEGDMA, HEMA and MMA, was studied in gingival fibroblasts. It was found that DNA strand breaks comparable to those induced by irradiation are only achieved with concentrations that are unrealistic.	Durner et al., 2011
Elevated expression of Th2 cytokines and signal molecules during the inflammation response in silica-induced pulmonary fibrosis in mice is mediated by IL-6R alpha.	Tripathi et al., 2010
The mechanism by which fenvalerate negatively affects male reproduction and spermatogenesis was investigated. The results show that fenvalerate induces germ cell apoptosis in testes by upregulating expression of Fas and FasL.	Zhao et al., 2011b
A comparison of the effects of curcumin and resveratrol on aflatoxin B-1-induced liver injury in rats revealed that only curcumin has a hepatoprotective effect against damage by aflatoxin B-1.	El-Agamy, 2010
Pentachlorophenol (PCB) and its metabolites are very potent toxicants that are still in use worldwide. This article reveals detailed mechanisms of toxicity induced by these chemicals in human lymphocytes with implications on pathology and their use as a sensitive cell type for toxicity testing.	Michałowicz, 2010
Bisabololoxide A, a chamomile tea constituent, can induce apoptosis of thymocytes, rising important issues on herbal tea toxicity.	Ogata et al., 2010
Ziram is a widely used fungicide in agriculture and as an accelerating agent in latex production. The authors report a novel immunotoxicity effect of this compound namely induction of apoptosis/necrosis in U937 cells.	Li et al., 2011b
Neuropathy Target Esterase (NTE) is the primary target esterase of organophosphorus compound-induced delayed neuropathy. Some studies suggest it may perform a function in embryonic development. This study reports NTE expression in ESC and its variation during first stages of in vitro differentiation to embryonic bodies. The role of this activity and the meaning of the variations detected during differentiation requires further studies.	Pamies et al., 2010
Pentabromodiphenyl ether (PentaBDE) more dramatically altered the redox status in rat liver compared to the other commonly used flame retardant, decabromodiphenyl ether (DecaBDE), as seen by increased activity of reducing enzymes, fatty acid degeneration, liver mass, CYP activities and microsomal enzymes. These endpoints were significantly less or not detectable in rats treated with DecaBDE.	Bruchajzer et al., 2010
Kidney damage induced by adriamycin (ADR) is marked by increased levels of hydrogen sulfide (H ₂ S), which leads to increased inflammation and oxidative stress. DL-propargylglycine (PAG) was shown to inhibit ADR-induced nephrotoxicity in rats treated with different concentrations of ADR up to 15 days.	Francescato et al., 2011
Known nanoparticles, fullerenes (also known as fullerols or fullerenols) [C-60(OH)(n)] cause cytotoxicity by disrupting mitochondrial function as determined by mitochondrial permeability transition. Disruption to the mitochondria results in mitochondrial depolarization, decreased ATP synthesis, oxidized GSH and proteins, and lipid peroxidation. An interesting finding is that the number of hydroxyl groups may influence toxicity.	Nakagawa et al., 2011

Table 4 (cont.): Organ toxicity and mechanisms

Key message	Reference
Generation of hydroxyl radical ((OH)-O-aEuro cent) in the striatum due to carbon monoxide (CO) poisoning is independent of hypoxia, which is also generated after exposure to CO. There also appears to be a threshold concentration at which OH-O-aEuro cent is formed.	Hara et al., 2011
Instead of supporting the more expected hypothesis that metabolites of benzo-a-pyrene (BaP) lead to more oxidative stress, reduced vascular nitric acid bioactivity and increased blood pressure, the authors observed inflammation in the lung after 7-day intranasal exposure to BaP, and altered circadian rhythm of blood pressure.	Gentner and Weber, 2011
The timing and concentration of the carbamate, physostigime used to treat organophosphate (OP) poisoning should be re-visited due to the possibility of increased toxicity and even mortality observed in a guinea pig model after treatment with the OP, soman. Obidoxime, at low concentrations, was the most effective therapeutic agent, reversing many of the effects of soman-induced toxicity.	Joosen et al., 2011
This study is presenting a local lymph node assay to assign disperse dyes sensitising potency. With this assay the dyes can be arranged into four potency classes.	Ahuja et al., 2010a
There is a significant correlation between melamine intake and urolithiasis in infants.	Ke et al., 2010
3,3 '-diselenodipropionic acid can ameliorate the polarisation of Th1/Th2 immune balance towards Th2 during body irradiation. This might be the reason for the prevention of oxidative stress in small intestine.	Kunwar et al., 2011
Monocrotaline/lipopolysaccharide co-treatment caused renal toxicity in mice. Co-treatment with tissue factor (TF) antisense oligonucleotide partially prevented renal damage and improved animal survival.	Abdel-Bakky et al., 2011
Diindolylmethane, a natural compound derived from cruciferous vegetables, induced the phospholipase C-dependent release of intracellular Ca ²⁺ in HA59T human hepatoma cells and caused dose-dependent cell death.	Cheng et al., 2011
Acrylamide treatment similarly affected the nervous system of young and adult male rats but lead to more pronounced spermatid degeneration in younger animals.	Takahashi et al., 2011
In a model of human primary lung cells, DNA damage induced by H ₂ O ₂ treatment correlated well with PARP-1 activity.	Ahmad et al., 2011
The anthracycline antineoplastic drug daunorubicin does not cause cardiotoxicity by affecting the glutathione antioxidant system as seen in a rabbit model of intravenous injection, or treatment of H9c2 rat cardiac cells.	Vávrová et al., 2011
In mice, the intake of fluoride with drinking water induced histopathological lesions of liver and kidney that were correlated with synthesis of stress proteins.	Chattopadhyay et al., 2011
Cyclophosphamide injection in adult female rats reduced the expression of uroplakin, a mucosal barrier against toxic materials, in urinary bladder mucosa. The reduced expression of uroplakin was preserved by concomitant treatment with mesna.	Kyung et al., 2011
This study deals with the effects of Pyrogallol (PG) on As4.1 juxtaglomerular cells. MAPK inhibitors attenuated As4.1 cell growth inhibition and death mediated by PG treatment. The changes in O ₂ ^{•-} and GSH levels by PG and/or MAPK inhibitors appeared to affect the growth and death of As4.1 cells.	Han et al., 2010b

Table 4 (cont.): Organ toxicity and mechanisms

Key message	Reference
<p>The aim of our study was to assess irritative effects of SO₂ up to 2 ppm on the airways using non-invasive methods like exhaled breath condensate (EBC), nasal lavage fluid (NALF) and exhaled nitric oxide (FeNO). Acute low dose SO₂ exposure in not adapted subjects did not induce airway irritation or/and inflammation measured under the tested conditions.</p>	<p>Raulf-Heimsoth et al., 2010</p>
<p>This study describes the involvement of sensory nerves and TRPV1 receptors in the rat airway inflammatory response to two environment pollutants: diesel exhaust particles (DEP) and 1,2-naphthoquinone (1,2-NQ). The authors found that DEP-induced airways oedema is highly influenced by increased ambient levels of 1,2-NQ and takes place by neurogenic mechanisms involving upregulation of TRPV1 and tachykinin receptors.</p>	<p>Costa et al., 2010</p>
<p>In vivo cholinergic toxicity and lethality was studied in freely moving guinea pigs upon exposure to the organophosphorus nerve agents sarin (GB) and VX. Seizures generated from GB resulted in more rapid and severe cholinergic effects, elevation of ACh level and increased glutamate levels. Glutamate did not increase in the VX seizure group. The authors conclude that seizure activity per se contributes to the elevated levels of brain ACh observed after nerve agent exposure.</p>	<p>O'Donnell et al., 2011</p>
<p>Monolayer cultures of primary hepatocytes, isolated from freshly removed livers are as commonly used as in vitro tool. The present study elucidated two cell death peaks, shortly after cell seeding and in the final stages of the cultivation period. The authors explain the underlying molecular processes and give strategies for increasing cell survival in this in vitro system.</p>	<p>Vinken et al., 2011</p>
<p>Biodiesels contain biocides in order to inhibit and remove microbial growth. The toxicity of four candidate biocides were tested using the NR8383 cell line and freshly isolated rat alveolar macrophages, the latter being among the first cell type exposed to inhaled biodiesel aerosols. The study suggested the biocide CMIT to cause respiratory effects and the need for further investigations and further animal studies.</p>	<p>Poon et al., 2011</p>
<p>Isoline-induced oxidative stress injury to the livers of newly weaned and young mice is age-dependent and due to the different responses of the glutathione antioxidant system.</p>	<p>Liang et al., 2011</p>
<p>Cadmium nephrotoxicity was tested in combined enzymatic and ¹³C-NMR measurements with a mathematical model of lactate metabolism. The authors showed that cadmium chloride highly diminishes the entire flux through the gluconeogenic pathway without altering the fluxes through enzymes of the tricarboxylic acid cycle.</p>	<p>Faiz et al., 2011</p>
<p>In order to characterize idiosyncratic liver injury the authors injected monocrotaline (MCT) in combination with modest inflammatory conditions induced by LPS in mice. Their experiments clearly showed that the expression of ox-LDL, CXCL16, and TF represents an early event in the onset of hepatotoxicity induced by MCT/LPS.</p>	<p>Hammad et al., 2011</p>
<p>The toxicokinetics and the biodistribution profile of the Shiga toxin type 2 (Stx2) was studied in rat. High accumulation was found in the lungs, kidneys, nasal turbinates and sometimes the eyes.</p>	<p>Liu et al., 2011b</p>
<p>The repeated administration of the flame retardant pentabromodiphenyl (PentaBDE) effected heme biosynthesis and leads to strongly enhanced porphyrin concentrations in the liver. Also the urinary excretion of total porphyrins was augmented.</p>	<p>Bruchajzer, 2011</p>

Table 4 (cont.): Organ toxicity and mechanisms

Key message	Reference
This study identifies a novel mechanism of phthalate toxicity in reproductive organs, namely induction of PLD via pERK1/2 activation, leading to testicular Leydig cell damage.	Lee et al., 2011b
NaF and TCDD have potentiative, harmful effects on the formation of dental hard tissue formation, as evidenced by morphological changes in whole tooth photographs, histological tissue sections and impaired mineralization.	Salmela et al., 2011
The mechanism underlying DBAA toxicity is not fully understood. The study shows that DBAA in vivo induces downregulation of cytochrome p450c17a at mRNA and protein level, a key enzyme in the biosynthesis of androgens, revealing a potential mechanism for DBAA toxicity in spermatogenesis.	Carr et al., 2011
This study shows that species differences in FYX-051-induced nephropathy are produced by combined effects of purine metabolism, urinary xanthine solubility, and plasma concentrations of FYX-051.	Shimo et al., 2011
The idiosyncratic toxicity of Ticlopidine (TIC) is suspected to occur by reactive metabolites covalently binding to proteins. The authors demonstrate that GSH depletion in animals enhances TIC covalent binding to liver proteins which may be involved in hepatotoxicity induced by TIC.	Shimizu et al., 2011
Increasing number of adverse effect case reports suggests that KP transdermal formulation can cause photoallergic reactions. The authors demonstrated that KP transdermal patch with TiO ₂ - included backing provide improved photostability and photosafety over conventional fabric KP patch.	Choi et al., 2011b
Mebendazole has been reported to cause liver injury with inflammatory responses. The authors report that MBZ and structurally similar drugs induce release of TNF α and IL-8 from human monocytes via activation of ERK1/2 MAPK pathway.	Mizuno et al., 2011
The phytochemicals zerumbone and auraptene effectively reduced TCDD and DTT toxicity, by protecting U937 macrophages from TCDD-induced apoptosis and blocking TCDD and DTT-induction of COX-2 and PDGF mRNA.	Sciullo et al., 2010
The study shows the in vivo effects of GLA on dopaminergic neurotransmission, namely GLA induced release of dopamine, probably mediated by an exocytotic, action potential-dependent mechanism, independent of dopamine transporter.	Nunes et al., 2010
Adverse effects of ortho-phthalaldehyde (OPA) have been reported in patients. Subcutaneous administration of OPA in mice induced a strong acute immune response, and additionally induced specific IgE and IgG in the sera, suggesting that OPA acts as an hapten.	Morinaga et al., 2010

Table 5: Reproductive toxicology

Key message	Reference
The study successfully used two validated in vitro tests, the rat whole embryo culture and the mouse embryonic stem cell test, to confirm the embryotoxicity of the immunosuppressive drug, mycophenolate mofetil. These results support what was observed in previous in vivo studies, and support the use of in vitro tests to measure teratogenic compounds.	Eckardt and Stahlmann, 2010

Table 5 (cont.): Reproductive toxicology

Key message	Reference
Newborn rats with prenatal acrylamide intoxication show alterations in lipid metabolism and serum protein changes.	Allam et al., 2010
Fluoride exposure of male mice altered the expression of genes involved in several sperm biological processes. Furthermore, fluoride exposure seemed to trigger sperm apoptosis by oxidative stress thereby reducing male fertility.	Sun et al., 2011
Maternal exposure to acrylamide induced compensatory mechanisms in the fetal nervous system, probably as a correction of impaired neurogenesis and migration.	Ogawa et al., 2011
Injection of alkaline but not neutral or acidic solution into mouse embryos at embryonic day 15 caused a significant increase in the number of apoptotic cells in the trigeminal ganglia at embryonic day 16.	Mukai et al., 2011
Perinatal exposure to methylmercury (MeHg) and/or PCB153 affects cerebral dopamine D1- and D2-Rs in a gender-, time-, and brain area-dependent manner. Combined treatment does not exacerbate the neurochemical effects of the individual compounds.	Coccini et al., 2011
Moxifloxacin exerted clear embryotoxic effects in vitro using the murine limb bud culture, however, only at concentrations that are higher than plasma concentrations observed during therapy.	Bode et al., 2010

Table 6: Genotoxicity and carcinogenicity

Key message	Reference
A single nucleotide polymorphism, rs710521[A], located near TP63, and recently discovered in genome wide association studies, was associated with human bladder cancer risk in a case-control series of 1,425 cases and 1,740 controls.	Lehmann et al., 2010
This study applied a genotoxicity assay based on the detection of histone H2AX phosphorylation to compare bisphenol A and bisphenol F. Bisphenol A was not found to be genotoxic, whereas bisphenol F showed positive effects.	Audebert et al., 2011
The flavonoid quercetin protects against methylmercury induced DNA damage and oxidative stress in rats.	Barcelos et al., 2011
The antioxidants isoquercitrin and melatonin reduce oxidative stress-mediated liver tumour promotion by the benzimidazole anthelmintic, oxfendasole in rats.	Nishimura et al., 2010
Indole-3-carbinol and flutamide increased expression of CYP1a1 and induced liver cell foci in rats.	Shimamoto et al., 2011
The carotenoid lutein protects against cisplatin-induced DNA damage and chromosome instability in peripheral blood cells by improving antioxidant defense.	Serpeloni et al., 2010
The genotoxicity potential of beauvericin (BEA) and ochratoxin A (OTA) was evaluated in PK15 cells and human leukocytes using the alkaline comet assay. It was found that BEA is more toxic than OTA and that the combined genotoxic action is additive or both synergistic and additive depending on the cell line.	Klarić et al., 2010
Using a two stage hepatocarcinogenesis model, this article describes early mechanisms in liver tumor progression, mainly via metal-related molecules, suggesting a role for lipid peroxidation and GST-P/Tfrc induction but not for Mt-1/2.	Mizukami et al., 2010

Table 6 (cont.): Genotoxicity and carcinogenicity

Key message	Reference
The carcinogenic potential of Diuron, a widely used herbicide, remains controversial. This study indicates that Diuron is a promoting agent to the urinary bladder but not to the mammary gland.	de Moura et al., 2010
The precise mechanism of oxfendazole (OX)-induced hepatocarcinogenesis is still unclear. The authors report that OX not only altered xenobiotic and antioxidant functions but also promoted cell proliferation in neoplastic lesions, suggesting that dysregulation of cell proliferation and apoptosis play important roles in OX-induced hepatocarcinogenesis.	Dewa et al., 2011
Although 3-MCPD is known to be mutagenic, its carcinogenic potential remains controversial. The authors conclude that drinking water administration in rats of 3-MCPD for 104 weeks revealed no evidence of carcinogenic potential.	Jeong et al., 2010b
Gatifloxacin (GFX), a third generation fluoroquinolone (FQ) and component of a new antituberculosis drug, caused more genotoxic damage than earlier drugs when tested in human lymphocytes. Prolonged sampling time was needed to see the effects, and aberrations were observed with non-cytotoxic concentrations supporting the need to review the conventional setups of these studies.	Anupama et al., 2010
In human lymphocytes and a human acute lymphoblastic cell line, the monomer bisphenol A-glycidyl methacrylate (BisGMA), a component of dental fillings, caused broad spectrum DNA damage with severe DNA strand breaks. Such damage was shown to cause a delay in the S phase of the cell cycle, thus facilitating repair.	Drozd et al., 2011
Widely used fungicide and anthelmintic, thiabendazole, caused increased aneugenicity, as observed by an increase in micronucleus formation, in isolated human lymphocytes.	Santovito et al., 2011a
Wy-14,643 causes DNA damage in livers of rats which is repaired by DNA repair genes.	Suzuki et al., 2010
The short-chain chlorinated paraffin C500C causes a peroxisome proliferation and a downregulation of $\alpha 2u$ in male rat liver.	Warnasuriya et al., 2010
Enzymatically modified isoquercitrine suppresses the liver tumor-promoting activity of phenobarbital by inhibiting nuclear translocation of CAR, and not by suppression of oxidative stress.	Morita et al., 2011
The hepatocarcinogen aflatoxin B-1 produced dose-related DNA damage in fetal livers of turkey and chicken embryos after in ovo injection. Turkey embryos appeared to be slightly more susceptible to the treatment.	Williams et al., 2011a
Using the comet assay, differences between the genotoxicity of orally administered bisphenol A and octylphenol were evident only in rats that received higher doses of the substances.	Ulutaş et al., 2011
In ovo diethylnitrosamine injection induced hepatocellular transformation reflecting disruption of differentiation and proliferation as well as gallbladder agenesis in fetal turkeys.	Williams et al., 2011b
The concentration-dependent cytotoxic and antimutagenic effects of the natural antitumor quinone biflorin were tested on Salmonella, yeast and V79 mammalian cells. At lower concentrations, biflorin has significant antioxidant and protective effects against the cytotoxicity, genotoxicity, mutagenicity, and intracellular lipid peroxidation induced by H_2O_2 in yeast and mammalian cells, which can be attributed to its hydroxyl radical-scavenging property. However, at higher concentrations, biflorin is cytotoxic and genotoxic.	Vasconcellos et al., 2010

Table 6 (cont.): Genotoxicity and carcinogenicity

Key message	Reference
The authors studied EP4 upregulation of Ras signaling and feedback regulation of Ras in human colon tissues and cancer. They suggest that Ras overexpression leads to cell proliferation through activating Ras/PI3K/GSK3 beta/EP4 PGE ₂ receptor signals and caused a feedback regulation of Ras by EP4 in colorectal tumor progression.	Wu et al., 2010
The fibroblast growth factor 9 (FGF-9) has strong potential roles in benzo(a)pyrene-induced CL5 cell invasion and human lung adenocarcinoma metastasis. Benzo-(a)pyrene and motorcycle exhaust particulate (MEP) increased FGF-9 mRNA expression and induced invasive ability of CL5 cells. Immunohistochemistry detected FGF-9 protein in the adenocarcinoma cells but not in normal epithelium.	Ueng et al., 2010
The widely used antileishmanial drug Glucantime [®] had no genotoxic effect in vitro on human lymphocytes but induced DNA damage in vivo in mice. The authors conclude that Glucantime [®] is a pro-mutagenic compound that causes damage to DNA after reduction of pentavalent antimony (SbV) into the more toxic trivalent antimony (SbIII) in the antimonial drug meglumine antimoniate.	Lima et al., 2010
In this toxicogenomic analysis 68 deregulated hepatocarcinogen-specific genes were identified as promising candidate biomarkers of cancer in chemically induced hepatocarcinogenesis in rasH2 mice on days 7 and 91 after treatment. Biomarker filter analysis suggested that 28 signature genes are involved in early-stage hepatocarcinogenesis.	Park et al., 2011c
The cyanobacterial alkaloid cylindrospermopsin (CYN) is being increasingly identified in drinking water supplies worldwide. This thorough study describes the different aspects of genotoxicity of CYN in detail and recommends to consider it in human health risk assessment.	Straser et al., 2011
The DNA methylation status of the lysophosphatidic acid 3 (LPA3) gene was disturbed in established HCC's and rats fed on choline-deficient L-amino acid-deficient (CDAA) diet. LPA1 and LPA2 receptor genes were unmethylated in rats fed normal diet, but methylation increased time-dependently in rats fed with CDAA diet. Thus LPA receptor genes might be involved during hepatocarcinogenesis induced by the CDDA diet.	Okabe et al., 2011
The authors developed a model to determine the genotoxic potential of formaldehyde (FA) in occupationally exposed individuals. In FA exposed and control subjects the frequency of chromosomal aberrations (CAs) in peripheral blood lymphocytes and two metabolic gene polymorphisms were correlated. FA induces CAs even to low levels of daily exposure, indicating an increased risk of genetic damage for workers to this air pollutant.	Santovito et al., 2011b
The inhibition of Cyt-P450-linked mixed function oxidases and benzene-induced genotoxicity in bone marrow and lung cells by polyphenolic acetates are mediated by the action of calreticulin transacetylase.	Kumar et al., 2011
Although fenofibrate caused transient DNA damage in rat liver, it also induced expression of DNA repair genes (e.g. Apex1, Ogg1 and Mlh1). This compensatory DNA repair response might account for the lack of permanent genotoxic damage.	Tawfeeq et al., 2011
RasH2 transgenic mice receiving 1 % DIDP developed a higher number of hepatocellular adenomas compared to wild type and rasH2 control mice, an effect not observed in previous rodent studies, indicating that this model is suitable for short term carcinogenicity testing.	Cho et al., 2011

Table 6 (cont.): Genotoxicity and carcinogenicity

Key message	Reference
<p>Concentration and time dependent experiments with carcinogenic genotoxins showed a simultaneous increase in both superoxide anions and Ty1 retrotransposition rates, which were blocked by addition of the scavenger N-acetylcysteine. Thus, the carcinogen-induced high level of reactive oxygen species plays a key role in the response of Ty1 test to carcinogenic genotoxins.</p>	<p>Dimitrov et al., 2011</p>

Table 7: Reviews and short communications

Key message	Reference
<p>Many degenerative diseases and toxicological insults converge on iron dysregulation. This review summarises several concepts of auto-catalytic production of hydroxyl radicals - a process intensified by positive feedback loops. Systems biology approaches predict that interventions with for example, iron chelators and antioxidants, may prove most effective in diseases, such as Parkinson's, Huntington's, Alzheimer's, prions as well as various forms of intoxications. The comprehensive review is the most cited article of the current evaluation period (2010, 2011).</p>	<p>Kell, 2010</p>
<p>The trichothecene mycotoxin deoxynivalanol (DON) is produced by the fungus, <i>Fusarium</i> in wheat and corn. This review summarises the molecular mechanisms of DON; which include ribotoxic stress, disturbed protein synthesis, compromised cell signalling, differentiation and proliferation. Proinflammatory gene induction, disruption of the growth hormone axis, and altered gut integrity finally lead to gastroenteritis ("vomitoxin"), anorexia, immunotoxicity and impaired reproduction in experimental animals. This review is the second most cited article of this evaluation period.</p>	<p>Pestka, 2010</p>
<p>This review summarises the state of the art of in vitro toxicity tests in five critical fields of toxicity: toxicokinetics, repeated-dose toxicity, carcinogenicity, skin sensitisation, and reproductive toxicity. The background of this review stems from the prohibition of animal-tested cosmetics on the market in Europe after 2013. The status and perspectives of each field are carefully analysed. For example, for skin sensitisation, in vitro techniques may already be able to identify sensitisers ahead of 2017. However, in other fields, particularly carcinogenesis, repeated-dose toxicity and reproductive toxicity, a time frame for in vitro alternatives cannot yet be estimated. This comprehensive review ranked third in the current evaluation period.</p>	<p>Adler et al., 2011</p>
<p>Metabolism of inorganic arsenic (iAs) is critical for its toxicity. This study analysed the relevance of arsenic transporters on human hepatocytes for the generation of methylated metabolites from iAs. A major finding is that MRP2 expression inversely correlates with cellular retention of iAs, as well as methylated metabolites in hepatocytes. This suggests that MRP2 plays an important role in the efflux of iAs and its metabolites. This study is the second most cited original article from the evaluation period.</p>	<p>Drobná et al., 2010</p>

Table 7 (cont.): Reviews and short communications

Key message	Reference
The comprehensive review article gives an overview of mutagenicity and carcinogenicity studies on selenium, and discusses the molecular mechanisms involved. At low concentrations, selenium shows anti-carcinogenic effects. However, at concentrations higher than needed for nutrition, selenium proves to be genotoxic and carcinogenic. This study may help regulate the use of selenium in nutrition.	Valdiglesias et al., 2010
Metabolomics have successfully identified novel biomarkers of disease prognosis and drug efficacy as well as toxicity. This review summarises how novel biomarkers discovered by metabolomics should be verified and introduced into clinical practice.	Mamas et al., 2011
This review focusses on the following aspects of selenium toxicity: (i) The majority of epidemiological studies suggest a cancer-preventing activity. (ii) In cancer treatment, selenium acts as a pro-oxidant by inducing apoptosis. (iii) The use of <i>Saccharomyces cerevisiae</i> is reviewed as a powerful tool to study the mode of action of selenium.	Brozmanová et al., 2010
Formaldehyde causes nasal cancer and lymphohematopoietic malignancies (LHM) in laboratory animals. Nasal cancer seems to be associated with cytotoxicity-induced proliferation. LHM occurs at even higher doses than nasal cancer. This study discusses the guideline value of 0.08 ppm formaldehyde to prevent carcinogenic effects.	Nielsen and Wolkoff, 2010
This review discusses the future perspectives of organoselenium as pharmacological agents. It also focuses on epidemiological evidence that selenium overexposure leads to chronic degenerative diseases.	Nogueira and Rocha, 2011
This is a comprehensive review on the protective network controlled by the Keap 1-Nrf2 axis, focussing on proliferation, angiogenesis and apoptosis.	Baird and Dinkova-Kostova, 2011
This review on Nrf2 focusses on the relevance of Nrf2-disruption in colon, bladder, lung, stomach, breast, skin and liver cancers.	Slocum and Kensler, 2011
This review gives an overview on how arsenate and arsenite interfere with intracellular signal transduction networks.	Druwe and Vaillancourt, 2010
This review gives a comprehensive update of the micronucleus assay, including its toxicological relevance, protocols, application as a high-throughput assay, and mechanisms of micronucleus formation.	Kirsch-Volders et al., 2011
The comprehensive review gives an overview of the use of human pluripotent stem cells, embryonic stem cells and induced pluripotent stem cells in developmental, cardio- and hepatotoxicity testing.	Wobus and Löser, 2011
Oxidative stress alone is not sufficient to explain specific mechanisms induced by nanoparticles. This article addresses nanoparticle-induced activation of MAP kinase cascades, p38, JNK, NF kappa B and Nrf-2 signalling pathways.	Marano et al., 2011
The capping material of nanocrystal quantum dots, and not the material of the core, determines toxicity.	Hoshino et al., 2011
This review discussed the current possibilities and perspectives of in vitro test systems for nanotoxicology.	Clift et al., 2011
This review gives an update on the mechanisms of action and cellular targets of toxic metals, as well as the use of chelating agents for pharmaceutical treatment.	Sinicropi et al., 2010
This review summarises epidemiologic studies on maternal exposure to particulate matters and adverse pregnancy outcomes. Overall, there is no convincing evidence of an association.	Bosetti et al., 2010

Table 7 (cont.): Reviews and short communications

Key message	Reference
The author critically discusses Hermann J. Muller's well-known Nobel lecture where a linear dose-response for radiation-induced germ cell mutations was presented. In contrast to this concept, Calabrese presents arguments speaking against the linear no-threshold model.	Calabrese, 2011
In urinary bladder cancer all known validated individual SNPs are associated with only moderate risk that is too low to justify preventive measures. The authors review this issue and propose that these so-called wimp-SNPs may interact and therefore collectively result in much higher risk with preventive relevance.	Golka et al., 2011
An overview of currently available metabolic databases is given with the MetaCyc family being described in particular detail.	Karp and Caspi, 2011
Mimicking a high fat diet by treating wild type mice with the bile acid, deoxycholic acid (DOC) at high physiologic levels, induced colonic tumours in 17 of 18 treated mice. The dietary antioxidant, chlorogenic acid, reduced tumour formation.	Bernstein et al., 2011
This review article summarises the protective effect of the antioxidant and mucolytic agent, N-acetylcysteine on contrast-induced acute kidney injury.	Briguori et al., 2011
This review expresses the need of alternative testing systems and herein the expectations on stem cells.	Seiler et al., 2011
This paper provides an up-to-date review of the progress made in the field of development of in vivo animal models for testing the allergic potential of novel proteins which resemble existing allergens. Finding and validating a strain of an animal species that is prone for allergic disorders equivalent to human individuals with a predisposed background is very desirable.	Ahuja et al., 2010b
This review provides a compendium of the results of genotoxicity and carcinogenicity assays performed on 70 marketed antihistamines.	Brambilla et al., 2011
Toxic effects on the Caco-2 cell line induced by Microcystin-LR, a toxin produced by toxic cyanobacteria, were higher at 48 h compared to those observed at 24 h. Differentiated Caco-2 cells were slightly more sensitive than undifferentiated cells. The most sensitive endpoint for the cell line was the reduction of total protein content.	Puerto et al., 2010

REFERENCES

- Abdel-Bakky MS, Hammad MA, Walker LA, Ashfaq MK. Silencing of tissue factor by antisense deoxyoligonucleotide prevents monocrotaline/LPS renal injury in mice. *Arch Toxicol* 2011;85(10):1245-56.
- Adler S, Basketter D, Creton S, Pelkonen O, van Benthem J, Zuang V et al. Alternative (non-animal) methods for cosmetics testing: current status and future prospects-2010. *Arch Toxicol* 2011;85(5):367-485.
- Ahmad M, Torky A, Glahn F, Scheubel RJ, Foth H. PARP-1 expression and activity in primary human lung cells. *Arch Toxicol* 2011;85(6):669-79.
- Ahuja V, Platzek T, Fink H, Sonnenburg A, Stahlmann R. Study of the sensitising potential of various textile dyes using a biphasic murine local lymph node assay. *Arch Toxicol* 2010a;84(9):709-18.

Ahuja V, Quatchadze M, Ahuja V, Stelter D, Albrecht A, Stahlmann R. Evaluation of biotechnology-derived novel proteins for the risk of food-allergic potential: advances in the development of animal models and future challenges. *Arch Toxicol* 2010b;84(12): 909-17.

Allam AA, El-Ghareeb AW, Abdul-Hamid M, Bakery AE, Gad M, Sabri M. Effect of prenatal and perinatal acrylamide on the biochemical and morphological changes in liver of developing albino rat. *Arch Toxicol* 2010;84(2):129-41.

Al-Rawi M, Diabaté S, Weiss C. Uptake and intracellular localization of submicron and nano-sized SiO₂ particles in HeLa cells. *Arch Toxicol* 2011;85(7):813-26.

Antonini JM, Roberts JR, Stone S, Chen BT, Schwegler-Berry D, Chapman R et al. Persistence of deposited metals in the lungs after stainless steel and mild steel welding fume inhalation in rats. *Arch Toxicol* 2011; 85(5):487-98.

Anupama M, Seiler JP, Murthy PB. A comparative analysis of chromosomal aberrations in cultured human lymphocytes due to fluoroquinolone drugs at different expression periods. *Arch Toxicol* 2010;84(5): 411-20.

Audebert M, Dolo L, Perdu E, Cravedi JP, Zalko D. Use of the γ H2AX assay for assessing the genotoxicity of bisphenol A and bisphenol F in human cell lines. *Arch Toxicol* 2011;85(11):1463-73.

Baird L, Dinkova-Kostova AT. The cytoprotective role of the Keap1-Nrf2 pathway. *Arch Toxicol* 2011;85(4):241-72.

Barcelos GR, Grotto D, Serpeloni JM, Angeli JP, Rocha BA, de Oliveira Souza VC et al. Protective properties of quercetin against DNA damage and oxidative stress induced by methylmercury in rats. *Arch Toxicol* 2011;85(9):1151-7.

Bernstein C, Holubec H, Bhattacharyya AK, Nguyen H, Payne CM, Zaitlin B et al. Carcinogenicity of deoxycholate, a secondary bile acid. *Arch Toxicol* 2011;85(8):863-71.

Billet S, Paget V, Garçon G, Heutte N, André V, Shirali P et al. Benzene-induced mutational pattern in the tumour suppressor gene TP53 analysed by use of a functional assay, the functional analysis of separated alleles in yeast, in human lung cells. *Arch Toxicol* 2010;84(2):99-107.

Bode K, Kunz A, Baumann-Wilschke I, Shakibaei M, Stahlmann R. Effects of moxifloxacin and clinafloxacin on murine limb buds cultured in regular and in magnesium-deficient medium. *Arch Toxicol* 2010;84(3):221-6.

Bosetti C, Nieuwenhuijsen MJ, Gallus S, Cipriani S, La Vecchia C, Parazzini F. Ambient particulate matter and preterm birth or birth weight: a review of the literature. *Arch Toxicol* 2010;84(6):447-60.

Brambilla G, Mattioli F, Robbiano L, Martelli A. Genotoxicity and carcinogenicity studies of antihistamines. *Arch Toxicol* 2011;85(10):1173-87.

Briguori C, Quintavalle C, De Micco F, Condorelli G. Nephrotoxicity of contrast media and protective effects of acetylcysteine. *Arch Toxicol* 2011;85(3):165-73.

Brozmanová J, Mániková D, Vlčková V, Chovanec M. Selenium: a double-edged sword for defense and offence in cancer. *Arch Toxicol* 2010;84(12):919-38.

Bruchajzer E, Frydrych B, Sporny S, Szymańska JA. Toxicity of penta- and decabromodiphenyl ethers after repeated administration to rats: a comparative study. *Arch Toxicol* 2010;84(4):287-99.

Bruchajzer E. Porphyrogenic effect of pentabromodiphenyl ether after repeated administration to rats. *Arch Toxicol* 2011;85(8):965-74.

Calabrese EJ. Muller's Nobel lecture on dose-response for ionizing radiation: ideology or science? *Arch Toxicol* 2011;85(12):1495-8.

Carr TL, Ciurlionis R, Milicic I, Whitney K, Liguori MJ, Warder SE et al. Role of cytochrome P450c17 α in dibromoacetic acid-induced testicular toxicity in rats. *Arch Toxicol* 2011;85(5):513-23.

Chattopadhyay A, Podder S, Agarwal S, Bhattacharya S. Fluoride-induced histopathology and synthesis of stress protein in liver and kidney of mice. *Arch Toxicol* 2011;85(4):327-35.

Cheng JS, Shu SS, Kuo CC, Chou CT, Tsai WL, Fang YC et al. Effect of diindolylmethane on Ca(2+) movement and viability in HA59T human hepatoma cells. *Arch Toxicol* 2011;85(10):1257-66.

Cho WS, Jeong J, Choi M, Park SN, Han BS, Son WC. 26-Week carcinogenicity study of di-isodecyl phthalate by dietary administration to CB6F1-rasH2 transgenic mice. *Arch Toxicol* 2011;85(1):59-66.

Choi MR, Jung KH, Park JH, Das ND, Chung MK, Choi IG et al. Ethanol-induced small heat shock protein genes in the differentiation of mouse embryonic neural stem cells. *Arch Toxicol* 2011a;85(4):293-304.

Choi YG, Lee JH, Bae IH, Ah YC, Ki HM, Bae JH et al. Titanium dioxide inclusion in backing reduce the photoallergenicity of ketoprofen transdermal patch. *Arch Toxicol* 2011b;85(3):219-26.

Chuang CY, Chen TL, Cherng YG, Tai YT, Chen TG, Chen RM. Lipopolysaccharide induces apoptotic insults to human alveolar epithelial A549 cells through reactive oxygen species-mediated activation of an intrinsic mitochondrion-dependent pathway. *Arch Toxicol* 2011;85(3):209-18.

Clift MJ, Gehr P, Rothen-Rutishauser B. Nanotoxicology: a perspective and discussion of whether or not in vitro testing is a valid alternative. *Arch Toxicol* 2011;85(7):723-31.

Coccini T, Roda E, Castoldi AF, Poli D, Goldoni M, Vettori MV et al. Developmental exposure to methylmercury and 2,2',4,4',5,5'-hexachlorobiphenyl (PCB153) affects cerebral dopamine D1-like and D2-like receptors of weanling and pubertal rats. *Arch Toxicol* 2011;85(10):1281-94.

Costa SK, Kumagai Y, Brain SD, Teixeira SA, Varriano AA, Barreto MA et al. Involvement of sensory nerves and TRPV1 receptors in the rat airway inflammatory response to two environment pollutants: diesel exhaust particles (DEP) and 1,2-naphthoquinone (1,2-NQ). *Arch Toxicol* 2010;84(2):109-117.

de Moura NA, Grassi TF, Rodrigues MA, Barbisan LF. Potential effects of the herbicide Diuron on mammary and urinary bladder two-stage carcinogenesis in a female Swiss mouse model. *Arch Toxicol* 2010;84(2):165-73.

Dewa Y, Nishimura J, Jin M, Kawai M, Saegusa Y, Kenmochi S et al. Immunohistochemical analyses at the late stage of tumor promotion by oxfendazole in a rat hepatocarcinogenesis model. *Arch Toxicol* 2011;85(2):155-62.

Di Giusto G, Torres AM. Organic anion transporter 5 renal expression and urinary excretion in rats exposed to mercuric chloride: a potential biomarker of mercury-induced nephropathy. *Arch Toxicol* 2010; 84(10):741-9.

Dimitrov M, Venkov P, Pesheva M. The positive response of Ty1 retrotransposition test to carcinogens is due to increased levels of reactive oxygen species generated by the genotoxins. *Arch Toxicol* 2011;85(1):67-74.

Dong GH, Liu MM, Wang D, Zheng L, Liang ZF, Jin YH. Sub-chronic effect of perfluorooctanesulfonate (PFOS) on the balance of type 1 and type 2 cytokine in adult C57BL6 mice. *Arch Toxicol* 2011;85(10): 1235-44.

Drobná Z, Walton FS, Paul DS, Xing W, Thomas DJ, Stýblo M. Metabolism of arsenic in human liver: the role of membrane transporters. *Arch Toxicol* 2010;84(1):3-16.

Drozd K, Wysokinski D, Krupa R, Wozniak K. Bisphenol A-glycidyl methacrylate induces a broad spectrum of DNA damage in human lymphocytes. *Arch Toxicol* 2011; 85(11):1453-61.

Druwe IL, Vaillancourt RR. Influence of arsenate and arsenite on signal transduction pathways: an update. *Arch Toxicol* 2010; 84(8):585-96.

Durner J, Dębiak M, Bürkle A, Hickel R, Reichl FX. Induction of DNA strand breaks by dental composite components compared to X-ray exposure in human gingival fibroblasts. *Arch Toxicol* 2011;85(2):143-8.

Duydu Y, Başaran N, Üstündağ A, Aydin S, Ündeğer Ü, Ataman OY et al. Reproductive toxicity parameters and biological monitoring in occupationally and environmentally boron-exposed persons in Bandırma, Turkey. *Arch Toxicol* 2011;85 (6):589-600.

Eckardt K, Stahlmann R. Use of two validated in vitro tests to assess the embryotoxic potential of mycophenolic acid. *Arch Toxicol* 2010;84(1):37-43.

El-Agamy DS. Comparative effects of curcumin and resveratrol on aflatoxin B(1)-induced liver injury in rats. *Arch Toxicol* 2010;84(5):389-96.

El Hakim AE, Gamal-Eldeen AM, Shahein YE, Mansour NM, Wahby AF, Abouelella AM. Purification and characterization of a cytotoxic neurotoxin-like protein from *Naja haja haja* venom that induces mitochondrial apoptosis pathway. *Arch Toxicol* 2011;85 (8):941-52.

Estrella A, Sánchez EE, Galán JA, Tao WA, Guerrero B, Navarrete LF et al. Characterization of toxins from the broad-banded water snake *Helicops angulatus* (Linnaeus, 1758): isolation of a cysteine-rich secretory protein, Helicopsin. *Arch Toxicol* 2011;85(4):305-13.

Faiz H, Conjard-Duplany A, Boghossian M, Martin G, Baverel G, Ferrier B. Cadmium chloride inhibits lactate gluconeogenesis in isolated human renal proximal tubules: a cellular metabolomic approach with ¹³C-NMR. *Arch Toxicol* 2011;85(9): 1067-77.

Fang EF, Pan WL, Wong JH, Chan YS, Ye XJ, Ng TB. A new *Phaseolus vulgaris* lectin induces selective toxicity on human liver carcinoma Hep G2 cells. *Arch Toxicol* 2011;85(12):1551-63.

Francescato HD, Marin EC, Cunha Fde Q, Costa RS, Silva CG, Coimbra TM. Role of endogenous hydrogen sulfide on renal damage induced by adriamycin injection. *Arch Toxicol* 2011;85(12):1597-606.

- Frese S, Velders M, Schleipen B, Schänzer W, Bloch W, Diel P. Myosin heavy chain expression pattern as a marker for anabolic potency: desoxymethyltestosterone (madol), norandrostenedione and testosterone repress MHC-IIb expression and stimulate MHC-IIId/x expression in orchietomized rat gastrocnemius muscle. *Arch Toxicol* 2011;85(6):635-43.
- Ge Y, Niu R, Zhang J, Wang J. Proteomic analysis of brain proteins of rats exposed to high fluoride and low iodine. *Arch Toxicol* 2011;85(1):27-33.
- Gentner NJ, Weber LP. Intranasal benzo[a]pyrene alters circadian blood pressure patterns and causes lung inflammation in rats. *Arch Toxicol* 2011;85(4):337-46.
- Gehrke H, Pelka J, Hartinger CG, Blank H, Bleimund F, Schneider R et al. Platinum nanoparticles and their cellular uptake and DNA platination at non-cytotoxic concentrations. *Arch Toxicol* 2011;85(7):799-812.
- Gibson N, Holzwarth U, Abbas K, Simionelli F, Kozempel J, Cydzik I et al. Radiolabelling of engineered nanoparticles for in vitro and in vivo tracing applications using cyclotron accelerators. *Arch Toxicol* 2011;85(7):751-73.
- Golka K, Selinski S, Lehmann ML, Blaszkewicz M, Marchan R, Ickstadt K et al. Genetic variants in urinary bladder cancer: collective power of the "wimp SNPs". *Arch Toxicol* 2011;85(6):539-54.
- Gube M, Taeger D, Weber DG, Pesch B, Brand P, Johnen G et al. Performance of biomarkers SMRP, CA125, and CYFRA 21-1 as potential tumor markers for malignant mesothelioma and lung cancer in a cohort of workers formerly exposed to asbestos. *Arch Toxicol* 2011;85(3):185-92.
- Hammad MA, Abdel-Bakky MS, Walker LA, Ashfaq MK. Oxidized low-density lipoprotein and tissue factor are involved in monocrotaline/lipopolysaccharide-induced hepatotoxicity. *Arch Toxicol* 2011;85(9):1079-89.
- Han YH, Park WH. Proteasome inhibitor MG132 reduces growth of As4.1 juxtaglomerular cells via caspase-independent apoptosis. *Arch Toxicol* 2010a;84(9):689-98.
- Han YH, Park WH. Pyrogallol-induced As4.1 juxtaglomerular cell death is attenuated by MAPK inhibitors via preventing GSH depletion. *Arch Toxicol* 2010b;84(8):631-40.
- Hanioka N, Oka H, Nagaoka K, Ikushiro S, Narimatsu S. Effect of UDP-glucuronosyltransferase 2B15 polymorphism on bisphenol A glucuronidation. *Arch Toxicol* 2011;85(11):1373-81.
- Hao S, Jiang X, Yan A, Yang B. Perinatal cisplatin exposure induces cochlear apoptosis in newborn guinea pigs. *Arch Toxicol* 2011;85(1):19-25.
- Hara S, Mizukami H, Kurosaki K, Kuriwa F, Mukai T. Existence of a threshold for hydroxyl radical generation independent of hypoxia in rat striatum during carbon monoxide poisoning. *Arch Toxicol* 2011;85(9):1091-9.
- Heng BC, Zhao X, Tan EC, Khamis N, Asodani A, Xiong S et al. Evaluation of the cytotoxic and inflammatory potential of differentially shaped zinc oxide nanoparticles. *Arch Toxicol* 2011;85(12):1517-28.
- Heo JD, Oh JH, Lee K, Kim CY, Song CW, Yoon S et al. Gene expression profiling in the lung tissue of cynomolgus monkeys in response to repeated exposure to welding fumes. *Arch Toxicol* 2010;84(3):191-203.

Hilderbrand SC, Murrell RN, Gibson JE, Brown JM. Marine brevetoxin induces IgE-independent mast cell activation. *Arch Toxicol* 2011;85(2):135-41.

Höfer N, Diel P, Wittsiepe J, Wilhelm M, Kluxen FM, Degen GH. Investigations on the estrogenic activity of the metallo-hormone cadmium in the rat intestine. *Arch Toxicol* 2010;84(7):579-81.

Hoshino A, Hanada S, Yamamoto K. Toxicity of nanocrystal quantum dots: the relevance of surface modifications. *Arch Toxicol* 2011;85(7):707-20.

Jeong GN, Jo UB, Ryu HY, Kim YS, Song KS, Yu IJ. Histochemical study of intestinal mucins after administration of silver nanoparticles in Sprague-Dawley rats. *Arch Toxicol* 2010a;84(1):63-9.

Jeong J, Han BS, Cho WS, Choi M, Ha CS, Lee BS et al. Carcinogenicity study of 3-monochloropropane-1, 2-diol (3-MCPD) administered by drinking water to B6C3F1 mice showed no carcinogenic potential. *Arch Toxicol* 2010b;84(9):719-29.

Joosen MJ, Smit AB, van Helden HP. Treatment efficacy in a soman-poisoned guinea pig model: added value of physostigmine? *Arch Toxicol* 2011;85(3):227-37.

Karp PD, Caspi R. A survey of metabolic databases emphasizing the MetaCyc family. *Arch Toxicol* 2011;85(9):1015-33.

Ke Y, Duan X, Wen F, Xu X, Tao G, Zhou L et al. Association of melamine exposure with urinary stone and oxidative DNA damage in infants. *Arch Toxicol* 2010;84(4):301-7.

Kell DB. Towards a unifying, systems biology understanding of large-scale cellular death and destruction caused by poorly liganded iron: Parkinson's, Huntington's, Alzheimer's, prions, bactericides, chemical toxicology and others as examples. *Arch Toxicol* 2010;84(11):825-89.

Kim JS, Lee K, Lee YH, Cho HS, Kim KH, Choi KH et al. Aspect ratio has no effect on genotoxicity of multi-wall carbon nanotubes. *Arch Toxicol* 2011a;85(7):775-86.

Kim JS, Song KS, Lee JH, Yu IJ. Evaluation of biocompatible dispersants for carbon nanotube toxicity tests. *Arch Toxicol* 2011b; 85(12):1499-508.

Kirsch-Volders M, Plas G, Elhajouji A, Lukamowicz M, Gonzalez L, Vande Loock K et al. The in vitro MN assay in 2011: origin and fate, biological significance, protocols, high throughput methodologies and toxicological relevance. *Arch Toxicol* 2011; 85(8):873-99.

Klarić MS, Darabos D, Rozgaj R, Kasuba V, Pepeljnjak S. Beauvericin and ochratoxin A genotoxicity evaluated using the alkaline comet assay: single and combined genotoxic action. *Arch Toxicol* 2010;84(8): 641-50.

Kretzing S, Abraham G, Seiwert B, Unge-mach FR, Krügel U, Teichert J et al. In vivo assessment of antiemetic drugs and mechanism of lycorine-induced nausea and emesis. *Arch Toxicol* 2011;85(12):1565-73.

Kumar A, Sushama A, Rohil V, Manral S, Gangopadhyay S, Prasad AK et al. Prevention of benzene-induced genotoxicity in bone marrow and lung cells: superiority of polyphenolic acetates to polyphenols. *Arch Toxicol* 2011;85(9):1141-50.

- Kunwar A, Bag PP, Chattopadhyay S, Jain VK, Priyadarsini KI. Anti-apoptotic, anti-inflammatory, and immunomodulatory activities of 3,3'-diselenodipropionic acid in mice exposed to whole body γ -radiation. *Arch Toxicol* 2011;85(11):1395-405.
- Kyung YS, Park HY, Lee G. Preservation of uroplakins by 2-mercaptoethanesulfonate in cyclophosphamide-induced rat cystitis. *Arch Toxicol* 2011;85(1):51-7.
- Lee EH, Oh JH, Park HJ, Kim DG, Lee JH, Kim CY et al. Simultaneous gene expression signature of heart and peripheral blood mononuclear cells in astemizole-treated rats. *Arch Toxicol* 2010;84(8):609-18.
- Lee YS, Kim DW, Lee YH, Oh JH, Yoon S, Choi MS et al. Silver nanoparticles induce apoptosis and G2/M arrest via PKC ζ -dependent signaling in A549 lung cells. *Arch Toxicol* 2011a;85(12):1529-40.
- Lee YJ, Ahn MY, Kim HS, Kwack SJ, Park KL, Yoon S et al. Role of phospholipase D in regulation of testicular Leydig cell hyperplasia in Sprague-Dawley rats treated with di(2-ethylhexyl) phthalate. *Arch Toxicol* 2011b;85(8):975-85.
- Lehmann ML, Selinski S, Blaszkewicz M, Orlich M, Ovsiannikov D, Moormann O et al. Rs710521[A] on chromosome 3q28 close to TP63 is associated with increased urinary bladder cancer risk. *Arch Toxicol* 2010;84(12):967-78.
- Leppänen M, Korpi A, Miettinen M, Leskinen J, Torvela T, Rossi EM et al. Nanosized TiO₂ caused minor airflow limitation in the murine airways. *Arch Toxicol* 2011;85(7):827-39.
- Li H, Wu S, Shi N, Lin W, You J, Zhou W. NF-E2-related factor 2 activation in PC12 cells: its protective role in manganese-induced damage. *Arch Toxicol* 2011a;85(8):901-10.
- Li Q, Kobayashi M, Kawada T. Ziram induces apoptosis and necrosis in human immune cells. *Arch Toxicol* 2011b;85(4):355-61.
- Liang QN, Sheng YC, Jiang P, Ji LL, Xia YY, Min Y et al. The difference of glutathione antioxidant system in newly weaned and young mice liver and its involvement in isoleucine-induced hepatotoxicity. *Arch Toxicol* 2011;85(10):1267-79.
- Lima MI, Arruda VO, Alves EV, de Azevedo AP, Monteiro SG, Pereira SR. Genotoxic effects of the antileishmanial drug Glucantime. *Arch Toxicol* 2010;84(3):227-32.
- Lin G, Du H, Chen J, Lu H, Guo W, Golka K et al. Association of XPD/ERCC2 G (23591) A and A (35931) C polymorphisms with skin lesion prevalence in a multiethnic, arseniasis-hyperendemic village exposed to indoor combustion of high arsenic coal. *Arch Toxicol* 2010;84(1):17-24.
- Liu F, Weng D, Chen Y, Song L, Li C, Dong L et al. Depletion of CD4+CD25+ Foxp3+ regulatory T cells with anti-CD25 antibody may exacerbate the 1,3- β -glucan-induced lung inflammatory response in mice. *Arch Toxicol* 2011a;85(11):1383-94.
- Liu YN, Wang SH, Li T, Wang Q, Tu W, Cai K et al. Shiga toxin type 2 (Stx2), a potential agent of bioterrorism, has a short distribution and a long elimination half-life, and induces kidney and thymus lesions in rats. *Arch Toxicol* 2011b;85(9):1133-40.
- Mamas M, Dunn WB, Neyses L, Goodacre R. The role of metabolites and metabolomics in clinically applicable biomarkers of disease. *Arch Toxicol* 2011;85(1):5-17.
- Marano F, Hussain S, Rodrigues-Lima F, Baeza-Squiban A, Boland S. Nanoparticles: molecular targets and cell signalling. *Arch Toxicol* 2011;85(7):733-41.

Meinerz DF, Sudati JH, dos Santos DB, Frediani A, Alberto EE, Allebrandt J et al. Evaluation of the biological effects of (S)-dimethyl 2-(3-(phenyltellanyl) propanamido) succinate, a new telluroamino acid derivative of aspartic acid. *Arch Toxicol* 2011;85(1):43-9.

Michałowicz J. Pentachlorophenol and its derivatives induce oxidative damage and morphological changes in human lymphocytes (in vitro). *Arch Toxicol* 2010;84(5):379-87.

Mizukami S, Ichimura R, Kemmochi S, Taniai E, Shimamoto K, Ohishi T et al. Induction of GST-P-positive proliferative lesions facilitating lipid peroxidation with possible involvement of transferrin receptor up-regulation and ceruloplasmin down-regulation from the early stage of liver tumor promotion in rats. *Arch Toxicol* 2010;84(4):319-31.

Mizuno K, Toyoda Y, Fukami T, Nakajima M, Yokoi T. Stimulation of pro-inflammatory responses by mebendazole in human monocytic THP-1 cells through an ERK signaling pathway. *Arch Toxicol* 2011;85(3):199-207.

Moon KY. Acrolein, an I- κ B α -independent downregulator of NF- κ B activity, causes the decrease in nitric oxide production in human malignant keratinocytes. *Arch Toxicol* 2011;85(5):499-504.

Morinaga T, Hasegawa G, Koyama S, Ishihara Y, Nishikawa T. Acute inflammation and immunoresponses induced by orthophthalaldehyde in mice. *Arch Toxicol* 2010;84(5):397-404.

Morita R, Shimamoto K, Ishii Y, Kuwata K, Ogawa B, Imaoka M et al. Suppressive effect of enzymatically modified isoquercitrin on phenobarbital-induced liver tumor promotion in rats. *Arch Toxicol* 2011;85(11):1475-84.

Mukai S, Nakagawa H, Ichikawa H, Miyazaki S, Nishimura K, Matsuo S. Effects of extracellular acidic-alkaline stresses on trigeminal ganglion neurons in the mouse embryo in vivo. *Arch Toxicol* 2011;85(2):149-54.

Nakagawa Y, Suzuki T, Ishii H, Nakae D, Ogata A. Cytotoxic effects of hydroxylated fullerenes on isolated rat hepatocytes via mitochondrial dysfunction. *Arch Toxicol* 2011;85(11):1429-40.

Nakamura M, Yasutake A, Fujimura M, Hachiya N, Marumoto M. Effect of methylmercury administration on choroid plexus function in rats. *Arch Toxicol* 2011;85(8):911-8.

Nielsen GD, Wolkoff P. Cancer effects of formaldehyde: a proposal for an indoor air guideline value. *Arch Toxicol* 2010;84(6):423-46.

Nishimura J, Saegusa Y, Dewa Y, Jin M, Kawai M, Kemmochi S et al. Antioxidant enzymatically modified isoquercitrin or melatonin supplementation reduces oxidative stress-mediated hepatocellular tumor promotion of oxfendazole in rats. *Arch Toxicol* 2010;84(2):143-53.

Nogueira CW, Rocha JB. Toxicology and pharmacology of selenium: emphasis on synthetic organoselenium compounds. *Arch Toxicol* 2011;85(11):1313-59.

Nohara K, Baba T, Murai H, Kobayashi Y, Suzuki T, Tateishi Y et al. Global DNA methylation in the mouse liver is affected by methyl deficiency and arsenic in a sex-dependent manner. *Arch Toxicol* 2011;85(6):653-61.

- Nunes BV, Durán R, Alfonso M, de Oliveira IM, Ferreira Faro LR. Evaluation of the effects and mechanisms of action of glufosinate, an organophosphate insecticide, on striatal dopamine release by using in vivo microdialysis in freely moving rats. *Arch Toxicol* 2010;84(10):777-85.
- O'Donnell JC, McDonough JH, Shih TM. In vivo microdialysis and electroencephalographic activity in freely moving guinea pigs exposed to organophosphorus nerve agents sarin and VX: analysis of acetylcholine and glutamate. *Arch Toxicol* 2011;85(12):1607-16.
- Ogata I, Kawanai T, Hashimoto E, Nishimura Y, Oyama Y, Seo H. Bisabololoxide A, one of the main constituents in German chamomile extract, induces apoptosis in rat thymocytes. *Arch Toxicol* 2010;84(1):45-52.
- Ogawa B, Ohishi T, Wang L, Takahashi M, Taniai E, Hayashi H et al. Disruptive neuronal development by acrylamide in the hippocampal dentate hilus after developmental exposure in rats. *Arch Toxicol* 2011;85(8):987-94.
- Okabe K, Hayashi M, Yoshida I, Nishimura K, Fukushima N, Tsujiuchi T. Distinct DNA methylation patterns of lysophosphatidic acid receptor genes during rat hepatocarcinogenesis induced by a choline-deficient L-amino acid-defined diet. *Arch Toxicol* 2011;85(10):1303-10.
- Pamies D, Reig JA, Vilanova E, Sogorb MA. Expression of neuropathy target esterase in mouse embryonic stem cells during differentiation. *Arch Toxicol* 2010;84(6):481-91.
- Park JY, Kim MJ, Kim YK, Woo JS. Ceramide induces apoptosis via caspase-dependent and caspase-independent pathways in mesenchymal stem cells derived from human adipose tissue. *Arch Toxicol* 2011a;85(9):1057-65.
- Park EJ, Roh J, Kim SN, Kang MS, Han YA, Kim Y et al. A single intratracheal instillation of single-walled carbon nanotubes induced early lung fibrosis and subchronic tissue damage in mice. *Arch Toxicol* 2011b;85(9):1121-31.
- Park HJ, Oh JH, Park SM, Cho JW, Yum YN, Park SN et al. Identification of biomarkers of chemically induced hepatocarcinogenesis in rasH2 mice by toxicogenomic analysis. *Arch Toxicol* 2011c;85(12):1627-40.
- Pestka JJ. Deoxynivalenol: mechanisms of action, human exposure, and toxicological relevance. *Arch Toxicol* 2010;84(9):663-79.
- Poon R, Rigden M, Edmonds N, Charman N, Lamy S. Effects of 5-chloro-2-methyl-4-isothiazolin-3-one and other candidate bio-diesel biocides on rat alveolar macrophages and NR8383 cells. *Arch Toxicol* 2011;85(11):1419-27.
- Posser T, de Paula MT, Franco JL, Leal RB, da Rocha JB. Diphenyl diselenide induces apoptotic cell death and modulates ERK1/2 phosphorylation in human neuroblastoma SH-SY5Y cells. *Arch Toxicol* 2011;85(6):645-51.
- Puerto M, Pichardo S, Jos A, Cameán AM. Microcystin-LR induces toxic effects in differentiated and undifferentiated Caco-2 cells. *Arch Toxicol* 2010;84(5):405-10.
- Raulf-Heimsoth M, Hoffmeyer F, van Thriel C, Blaszkewicz M, Bünger J, Brüning T. Assessment of low dose effects of acute sulphur dioxide exposure on the airways using non-invasive methods. *Arch Toxicol* 2010;84(2):121-7.
- Repetto MG, Ferrarotti NF, Boveris A. The involvement of transition metal ions on iron-dependent lipid peroxidation. *Arch Toxicol* 2010;84(4):255-62.

Rodrigues JL, Serpeloni JM, Batista BL, Souza SS, Barbosa F Jr. Identification and distribution of mercury species in rat tissues following administration of thimerosal or methylmercury. *Arch Toxicol* 2010;84(11):891-6.

Romero C, Benedí J, Villar A, Martín-Aragón S. Involvement of Hsp70, a stress protein, in the resistance of long-term culture of PC12 cells against sodium nitroprusside (SNP)-induced cell death. *Arch Toxicol* 2010;84(9):699-708.

Rubiolo JA, López-Alonso H, Vega FV, Vieytes MR, Botana LM. Okadaic acid and dinophysin toxin 2 have differential toxicological effects in hepatic cell lines inducing cell cycle arrest, at G0/G1 or G2/M with aberrant mitosis depending on the cell line. *Arch Toxicol* 2011;85(12):1541-50.

Safarinejad MR, Shafiei N, Safarinejad SH. The association between bladder cancer and a single nucleotide polymorphism (rs2854744) in the insulin-like growth factor (IGF)-binding protein-3 (IGFBP-3) gene. *Arch Toxicol* 2011;85(10):1209-18.

Salmela E, Lukinmaa PL, Partanen AM, Sahlberg C, Alaluusua S. Combined effect of fluoride and 2,3,7,8-tetrachlorodibenzo-p-dioxin on mouse dental hard tissue formation in vitro. *Arch Toxicol* 2011;85(8):953-63.

Santovito A, Cervella P, Delpero M. In vitro aneugenic effects of the fungicide thiabendazole evaluated in human lymphocytes by the micronucleus assay. *Arch Toxicol* 2011a;85(6):689-93.

Santovito A, Schilirò T, Castellano S, Cervella P, Bigatti MP, Gilli G et al. Combined analysis of chromosomal aberrations and glutathione S-transferase M1 and T1 polymorphisms in pathologists occupationally exposed to formaldehyde. *Arch Toxicol* 2011b;85(10):1295-302.

Sciullo EM, Vogel CF, Wu D, Murakami A, Ohigashi H, Matsumura F. Effects of selected food phytochemicals in reducing the toxic actions of TCDD and p,p'-DDT in U937 macrophages. *Arch Toxicol* 2010;84(12):957-66.

Seiler A, Oelgeschläger M, Liebsch M, Pirow R, Riebeling C, Tralau T et al. Developmental toxicity testing in the 21st century: the sword of Damocles shattered by embryonic stem cell assays? *Arch Toxicol* 2011;85(11):1361-72.

Serpeloni JM, Grotto D, Mercadante AZ, de Lourdes Pires Bianchi M, Antunes LM. Lutein improves antioxidant defense in vivo and protects against DNA damage and chromosome instability induced by cisplatin. *Arch Toxicol* 2010;84(10):811-22.

Shimada M, Kameo S, Sugawara N, Yaginuma Sakurai K, Kurokawa N, Mizukami-Murata S et al. Gene expression profiles in the brain of the neonate mouse perinatally exposed to methylmercury and/or polychlorinated biphenyls. *Arch Toxicol* 2010;84(4):271-86.

Shimada H, Hata I, Hashiguchi T, Imamura Y. Genetic background of resistance to cadmium-induced testicular toxicity in inbred Wistar-Imamichi rats. *Arch Toxicol* 2011;85(10):1195-9.

Shimamoto K, Dewa Y, Kemmochi S, Taniai E, Hayashi H, Imaoka M et al. Relationship between CYP1A induction by indole-3-carbinol or flutamide and liver tumor-promoting potential in rats. *Arch Toxicol* 2011;85(9):1159-66.

Shimizu S, Atsumi R, Nakazawa T, Izumi T, Sudo K, Okazaki O et al. Ticlopidine-induced hepatotoxicity in a GSH-depleted rat model. *Arch Toxicol* 2011;85(4):347-53.

Shimo T, Ashizawa N, Moto M, Iwanaga T, Nagata O. Study on species differences in nephropathy induced by FYX-051, a xanthine oxidoreductase inhibitor. *Arch Toxicol* 2011;85(5):505-12.

Silva R, Carmo H, Dinis-Oliveira R, Cordeiro-da-Silva A, Lima SC, Carvalho F et al. In vitro study of P-glycoprotein induction as an antidotal pathway to prevent cytotoxicity in Caco-2 cells. *Arch Toxicol* 2011;85(4):315-26.

Sinicropi MS, Amantea D, Caruso A, Saturnino C. Chemical and biological properties of toxic metals and use of chelating agents for the pharmacological treatment of metal poisoning. *Arch Toxicol* 2010;84(7):501-20.

Slocum SL, Kensler TW. Nrf2: control of sensitivity to carcinogens. *Arch Toxicol* 2011;85(4):273-84.

Song YJ, Li J, Xie XF, Wang H, Li QX. Effects of amlodipine on TGF- β -induced Smad2, 4 expressions in adriamycin toxicity of rat mesangial cells. *Arch Toxicol* 2011;85(6):663-8.

Straser A, Filipič M, Zegura B. Genotoxic effects of the cyanobacterial hepatotoxin cylindrospermopsin in the HepG2 cell line. *Arch Toxicol* 2011;85(12):1617-26.

Sumi D, Sasaki T, Miyataka H, Himeno S. Rat H9c2 cardiac myocytes are sensitive to arsenite due to a modest activation of transcription factor Nrf2. *Arch Toxicol* 2011;85(12):1509-16.

Sun Z, Niu R, Su K, Wang B, Wang J, Zhang J et al. Effects of sodium fluoride on hyperactivation and Ca²⁺ signaling pathway in sperm from mice: an in vivo study. *Arch Toxicol* 2010;84(5):353-61.

Sun Z, Niu R, Wang B, Jiao Z, Wang J, Zhang J et al. Fluoride-induced apoptosis and gene expression profiling in mice sperm in vivo. *Arch Toxicol* 2011;85(11):1441-52.

Suzuki T, Jin M, Dewa Y, Ichimura R, Shimada Y, Mizukami S et al. Evaluation of in vivo liver genotoxic potential of Wy-14,643 and piperonyl butoxide in rats subjected to two-week repeated oral administration. *Arch Toxicol* 2010;84(6):493-500.

Takahashi M, Inoue K, Koyama N, Yoshida M, Irie K, Morikawa T et al. Life stage-related differences in susceptibility to acrylamide-induced neural and testicular toxicity. *Arch Toxicol* 2011;85(9):1109-20.

Tawfeeq MM, Suzuki T, Shimamoto K, Hayashi H, Shibutani M, Mitsumori K. Evaluation of in vivo genotoxic potential of fenofibrate in rats subjected to two-week repeated oral administration. *Arch Toxicol* 2011;85(8):1003-11.

Teixeira SS, Silveira LB, da Silva FM, Marchi-Salvador DP, Silva FP Jr, Izidoro LF et al. Molecular characterization of an acidic phospholipase A(2) from Bothrops pirajai snake venom: synthetic C-terminal peptide identifies its antiplatelet region. *Arch Toxicol* 2011;85(10):1219-33.

Tithof PK, Richards SM, Elgayyar MA, Menn FM, Vulava VM, McKay L et al. Activation of group IVC phospholipase A(2) by polycyclic aromatic hydrocarbons induces apoptosis of human coronary artery endothelial cells. *Arch Toxicol* 2011;85(6):623-34.

Tripathi SS, Mishra V, Shukla M, Verma M, Chaudhury BP, Kumar P et al. IL-6 receptor-mediated lung Th2 cytokine networking in silica-induced pulmonary fibrosis. *Arch Toxicol* 2010;84(12):947-55.

- Truong L, Moody IS, Stankus DP, Nason JA, Lonergan MC, Tanguay RL. Differential stability of lead sulfide nanoparticles influences biological responses in embryonic zebrafish. *Arch Toxicol* 2011;85(7):787-98.
- Ueng TH, Chang YL, Tsai YY, Su JL, Chan PK, Shih JY et al. Potential roles of fibroblast growth factor-9 in the benzo(a) pyrene-induced invasion in vitro and the metastasis of human lung adenocarcinoma. *Arch Toxicol* 2010;84(8):651-60.
- Ulutaş OK, Yildiz N, Durmaz E, Ahab MA, Barlas N, Çok İ. An in vivo assessment of the genotoxic potential of bisphenol A and 4-tert-octylphenol in rats. *Arch Toxicol* 2011;85(8):995-1001.
- Valdiglesias V, Pásaro E, Méndez J, Laffon B. In vitro evaluation of selenium genotoxic, cytotoxic, and protective effects: a review. *Arch Toxicol* 2010;84(5):337-51.
- Vasconcellos MC, Moura DJ, Rosa RM, Machado MS, Guecheva TN, Villela I et al. Evaluation of the cytotoxic and antimutagenic effects of biflorin, an antitumor 1,4 naphthoquinone isolated from *Capraria biflora* L. *Arch Toxicol* 2010;84(10):799-810.
- Vávrová A, Popelová O, Stěrba M, Jirkovský E, Hašková P, Mertlíková-Kaiserová H et al. In vivo and in vitro assessment of the role of glutathione antioxidant system in anthracycline-induced cardiotoxicity. *Arch Toxicol* 2011;85(5):525-35.
- Vinken M, Decrock E, Doktorova T, Ramboer E, De Vuyst E, Vanhaecke T et al. Characterization of spontaneous cell death in monolayer cultures of primary hepatocytes. *Arch Toxicol* 2011;85(12):1589-96.
- Wagner C, Vargas AP, Roos DH, Morel AF, Farina M, Nogueira CW et al. Comparative study of quercetin and its two glycoside derivatives quercitrin and rutin against methylmercury (MeHg)-induced ROS production in rat brain slices. *Arch Toxicol* 2010;84(2):89-97.
- Wang H, Wang Q, Zhao XF, Liu P, Meng XH, Yu T et al. Cypermethrin exposure during puberty disrupts testosterone synthesis via downregulating StAR in mouse testes. *Arch Toxicol* 2010;84(1):53-61.
- Wang Y, Zhao F, Jin Y, Zhong Y, Yu X, Li G et al. Effects of exogenous glutathione on arsenic burden and NO metabolism in brain of mice exposed to arsenite through drinking water. *Arch Toxicol* 2011a;85(3):177-84.
- Wang Z, Yang X, Yang S, Ren G, Ferreri M, Su Y et al. Sodium fluoride suppress proliferation and induce apoptosis through decreased insulin-like growth factor-I expression and oxidative stress in primary cultured mouse osteoblasts. *Arch Toxicol* 2011b;85(11):1407-17.
- Waring JF, Ciurlionis R, Marsh K, Klein LL, DeGoey DA, Randolph JT et al. Identification of proteasome gene regulation in a rat model for HIV protease inhibitor-induced hyperlipidemia. *Arch Toxicol* 2010;84(4):263-70.
- Warnasuriya GD, Elcombe BM, Foster JR, Elcombe CR. A mechanism for the induction of renal tumours in male Fischer 344 rats by short-chain chlorinated paraffins. *Arch Toxicol* 2010;84(3):233-43.
- Watanabe T, Ohta Y, Mizumura A, Kobayashi Y, Hirano S. Analysis of arsenic metabolites in HepG2 and AS3MT-transfected cells. *Arch Toxicol* 2011;85(6):577-88.

Wille T, Thiermann H, Worek F. Effect of different buffers on kinetic properties of human acetylcholinesterase and the interaction with organophosphates and oximes. *Arch Toxicol* 2011;85(3):193-8.

Williams JG, Deschl U, Williams GM. DNA damage in fetal liver cells of turkey and chicken eggs dosed with aflatoxin B1. *Arch Toxicol* 2011a;85(9):1167-72.

Williams GM, Brunnemann KD, Iatropoulos MJ, Smart DJ, Enzmann HG. Production of liver preneoplasia and gallbladder agenesis in turkey fetuses administered diethylnitrosamine. *Arch Toxicol* 2011b; 85(6): 681-7.

Wobus AM, Löser P. Present state and future perspectives of using pluripotent stem cells in toxicology research. *Arch Toxicol* 2011;85(2):79-117.

Wu CH, Shih YW, Chang CH, Ou TT, Huang CC, Hsu JD et al. EP4 upregulation of Ras signaling and feedback regulation of Ras in human colon tissues and cancer cells. *Arch Toxicol* 2010;84(9):731-40.

Yamauchi S, Kiyosawa N, Ando Y, Watanabe K, Niino N, Ito K et al. Hepatic transcriptome and proteome responses against diethyl maleate-induced glutathione depletion in the rat. *Arch Toxicol* 2011;85 (9):1045-56.

Yen CC, Ho TJ, Wu CC, Chang CF, Su CC, Chen YW et al. Inorganic arsenic causes cell apoptosis in mouse cerebrum through an oxidative stress-regulated signaling pathway. *Arch Toxicol* 2011;85(6): 565-75.

Yu WG, Liu W, Liu L, Jin YH. Perfluorooctane sulfonate increased hepatic expression of OAPT2 and MRP2 in rats. *Arch Toxicol* 2011;85(6):613-21.

Zhang M, Li J, Xing G, He R, Li W, Song Y et al. Variation in the internalization of differently sized nanoparticles induces different DNA-damaging effects on a macrophage cell line. *Arch Toxicol* 2011;85(12): 1575-88.

Zhao P, Zhang L, Zhong W, Ying X, Huang H, Yao B et al. Antigen-subtracted 2-DE/MS strategy, a novel proteomic analysis platform. *Arch Toxicol* 2011a;85(1): 35-41.

Zhao XF, Wang Q, Ji YL, Wang H, Liu P, Zhang C et al. Fenvalerate induces germ cell apoptosis in mouse testes through the Fas/FasL signaling pathway. *Arch Toxicol* 2011b;85(9):1101-8.