



Metabolomics, a New Promising Technology for Toxicological Research

Kyu-Bong Kim¹ and Byung Mu Lee²

¹National Institute of Toxicological Research, Korea Food and Drug Administration, Seoul 122-704

²Division of Toxicology, School of Pharmacy, Sungkyunkwan University, Suwon 440-746, Korea

(Received May 4, 2009; Revised May 20, 2009; Accepted May 21, 2009)

Metabolomics which deals with the biological metabolite profile produced in the body and its relation to disease state is a relatively recent research area for drug discovery and biological sciences including toxicology and pharmacology. Metabolomics, based on analytical method and multivariate analysis, has been considered a promising technology because of its advantage over other toxicogenomic and toxicoproteomic approaches. The application of metabolomics includes the development of biomarkers associated with the pathogenesis of various diseases, alternative toxicity tests, high-throughput screening (HTS), and risk assessment, allowing the simultaneous acquisition of multiple biochemical parameters in biological samples. The metabolic profile of urine, in particular, often shows changes in response to exposure to xenobiotics or disease-induced stress, because of the biological system's attempt to maintain homeostasis. In this review, we focus on the most recent advances and applications of metabolomics in toxicological research.

Key words: Metabolomics, Biomarker, Toxicology, High-throughput screening (HTS), Risk assessment

INTRODUCTION

Metabolomics (or metabonomics), a newly categorized '-omics', is derived from the Greek roots 'meta' meaning change and 'nomos' meaning rules or laws, to provide insight into the generation of pattern recognition-based models that are able to classify or predict changes in biological metabolism (Nicholson *et al.*, 1999; Watkins and German, 2002; Lindon *et al.*, 2004). Metabolomics is generally defined as the systemic investigation of the unique metabolite network or fingerprint which explains specific biological or etiological status with change of metabolome, the collection of all metabolites produced in biological systems (Lindon *et al.*, 2003). Metabolomes, relatively small molecular weight proteins of 100-1000 and the end products of biological organism's gene expression, are generally analyzed by mass spectrometry. Nicholson *et al.* (1999) defined 'metabonomics' as the quantitative measurement of the dynamic multi-parametric metabolic response of living systems to patho-physiological stimuli or genetic

modification. There is no significant difference between the terms metabolomics and metabonomics, but the term metabolomics rather than metabonomics has been used because 'metabolomics' is more commonly used. Metabolomic strategies aim to detect changes in the distribution and concentration of a broad range of endogenous metabolites and can be applied to multiple levels of biological system: from single cell to whole organism (Celia *et al.*, 2002). The '-omics' suffix has come to signify the determination of a targeted level of biological molecules and information (Yang *et al.*, 2009). Therefore, genomics measures the entire genetic makeup of an organism, while proteomics analyzes all the proteins expressed under given conditions (Hrmova and Fincher, 2009; Rampitsch *et al.*, 2009). Metabolomics is not exceptional. Its relationship to the other '-omics' (toxicogenomics and toxicoproteomics) is represented in Fig. 1. Metabolism is the biochemical modification of chemical compounds by interaction with catalytic enzymes in living organisms or cells. This includes the biosynthesis of complex organic molecules (anabolism) and their breakdown (catabolism) into small ones. Metabolism usually consists of sequences of enzymatic steps, also called metabolic pathways. The total metabolisms are all biochemical processes of an organism,

Correspondence to: Byung Mu Lee, Division of Toxicology, College of Pharmacy, Sungkyunkwan University, Chunchung-dong 300, Changan-ku, Suwon, Gyeonggi-do 440-746, Korea
E-mail: bmlee@skku.ac.kr




Fig. 1. The “OMICS” technologies involved in toxicological research. The integration of “omics” sciences might lead to a better comprehensive understanding of toxicological sciences.

whereas the cellular metabolism includes all chemical processes in a cell. Important metabolic pathways are composed of general pathways (carbohydrate, fatty acid, and citric acid metabolisms), catabolism, anabolism, and drug metabolism (Nealson and Conrad, 1999; Raman *et al.*, 2005; Testa and Kramer, 2006). A metabolome is generally termed as the whole set of metabolic small entities in a cell, tissue, organ, organisms, and species

(Allen *et al.*, 2003). It includes small circuits of pathway networks. The very large portion of metabolome study has been metabolic engineering to produce industrially meaningful compounds (Wu *et al.*, 2005). Metabolomics is the method of studying, profiling and fingerprinting metabolites in various physiologic states (Fiehn *et al.*, 2002) (Table 1). Metabolite profiling is a main tool for the analysis of a class of metabolites. Metabolomics aims to include all classes of endogenous metabolites and utilizes metabolic fingerprinting of them to maintain a rapid classification of biological samples according to their origin and biochemical relevance (Nicholson *et al.*, 1999; Lindon *et al.*, 2004). In order to optimize and utilize metabolomics, a stable and reproducible metabolite fingerprint must be established (Bino *et al.*, 2004). Metabolomic techniques are looking for a way to analyze changes to those endogenous biomolecules caused by xenobiotic toxicity or drug efficacy. In toxicological research, metabolomics is also viewed as holding great promise, including use in specific biomarker discovery for clinical diagnostics and drug discovery. In this review, the brief analytical technologies for metabolomics are summarized, the current toxicological applications of metabolomics are described, and then the prospective future of metabolomics for toxicology is discussed.

Table 1. General terminology for metabolite analysis

General areas	Description
Target compound analysis	- The quantification of specific metabolites
Metabolic profiling	- Quantitative or qualitative determination of a group of related compounds or of members of specific metabolic pathways
Metabolomics	- Qualitative and quantitative analysis of all metabolites
Metabolic fingerprinting	- Sample classification by rapid and global analysis without extensive compound identification

Table 2. Analytical methods used for characterization of metabolites

Analytical methods	Feature	Ref.
NMR spectroscopy	- Cheap after initial purchase - Robust and reliable - Minimal sample preparation - High throughput - Significant metabolite overlap - Large initial outlay	Reo <i>et al.</i> , 2002 Nicholson <i>et al.</i> , 1989 Raamsdonk <i>et al.</i> , 2001
GC-MS	- Excellent sensitivity - No need to derivatise - More global than NMR or GC-MS - Either specific or global - LC reproducibility is less than GC - Ion suppression can impede some metabolite detection	Fiehn <i>et al.</i> , 2002 Fiehn <i>et al.</i> , 2000
LC-MS	- Good sensitivity - Cheap to purchase - Good identification software - Good chromatograms compared to LC-MS	Wilson <i>et al.</i> , 2003

ANALYTICAL TECHNOLOGIES

Metabolomics is a multi-disciplinary technology, requiring cooperation between toxicologists, pharmacologists, chemists, biologists and informaticians. Current metabolomics practice has mainly relied on mass spectrometry (MS) and nuclear magnetic resonance (NMR) spectroscopy. The MS requires a pre-separational procedure of metabolites using gas chromatography (GC) or liquid chromatography (LC) (Table 2). It's almost impossible to detect the whole population of metabolites in a system with single analytical method. Isolation of metabolites from biological samples requires the preparation of an extract. The choice of solvent used for this initial extraction process directly affects the chemical classes of compounds present in that extract. Furthermore, no spectroscopic method is available for the detection of all classes of metabolites.

Therefore, a variety of comprehensive and targeted




Fig. 2. Schematic diagram of the procedure for metabolomic approach.

methods need to be applied and the data integrated in order to provide a complete profile of metabolic status. A schematic representation of the sample preparation process for metabolite analysis, from tissue treatment to data analysis, is given in Fig. 2. In any case, comprehensive quantitative and qualitative determination of metabolites in biological samples may require parallel implementation of several processing and detection methods (Dunn and Ellis, 2005). All metabolomics studies produce complex multivariate data sets that need to be interpreted using chemometric and bioinformatic methods (Nicholson *et al.*, 1999).

NMR. Proton (^1H) NMR can detect any metabolites containing hydrogen (Table 3). Signals can be assigned by comparison with libraries of reference compounds, or by two-dimensional NMR. The ^1H NMR spectra of crude biological tissue extracts are inevitably crowded with many overlapped signals, not only because there is a large number of contributing compounds, but also because of the low overall chemical shift dispersion. ^1H NMR spectra are also complicated by spin-spin couplings which add to signal multiplicity, although they are an important source of structural information (Griffin *et al.*, 2003). In ^{13}C NMR, the chemical shift dispersion is twenty times greater and spin-spin interactions are removed by decoupling. Despite these advantages, the low sensitivity of ^{13}C NMR prevents its routine use with complex extracts (Kenney *et al.*, 2003; Bundy *et al.*, 2003).

Gas chromatography. Gas chromatography (GC) provides compound separations with high-resolution and can be used in conjunction with a flame ionization detector (GC/FID) or a mass spectrometer (GC/MS). Both detection methods are highly sensitive and able to detect almost any organic compound, regardless of its class or structure. However, some of the metabolites found in biological samples are too involatile to be anal-

Table 3. NMR-based metabolomics and potential biomarkers

Type	Sample	Biomarkers	Ref.
Neurochemicals TTX(tetrodotoxin): rat frontal cortex.	Cerebrospinal fluid (CSF)	- Glutamate, isoleucine, valine, alanine, α - and β -hydroxybutyrate \downarrow	Khandelwal <i>et al.</i> , 2004
Myocardial ischemia.	Tissues	- Plasma: lactate, acetate, acetone - Brain: glycerol, succinate, propionate, lactate	Price <i>et al.</i> , 2005
Dominant-submissive relationships: rats	Urine	- Milk sugar consumption, urinary galactose normalized to creatinine	Leo <i>et al.</i> , 2005
Dietary influence (healthy British, Swedish subjects)	Urine	- Trimethylamine-N-oxide (TMAO) \uparrow taurine \uparrow	Lenz <i>et al.</i> , 2004

ysed directly by GC methods. Therefore, the compounds have to be converted to less polar, more volatile derivatives before they are applied to the GC column.

High performance liquid chromatography (HPLC).

HPLC, with UV detection, is a common method used for targeted analysis of biological samples and for metabolic profiling of individual classes (Fan *et al.*, 2005). Derivatisation is not essential. Selection of compounds arises initially from the type of solvent used for extraction and then from the type of column and detector. For example, HPLC/UV will only detect compounds with a suitable chromophore; a column selected for its ability to separate one class of compounds will not generally be useful for other types. HPLC profiling methods all rely, to a great extent, on comparisons with reference compounds. The full UV spectrum (measured for each peak when UV-diode array detectors are used) gives some useful information on the nature of compounds in complex profiles, but often indicates the class of the compound rather than its exact identity (Plumb *et al.*, 2002).

LC/MS, LC/MS/MS and LC/NMR. LC/MS, LC/MS/MS and LC/NMR are powerful solutions to the problems of detector generality and structure determination. LC/MS can be used to detect compounds that are not well characterized by other methods (those that are not easily derivatised), lie above the available GC/MS mass range, or do not contain good chromophores for conventional HPLC (Wilson *et al.*, 2005). The electrospray ionization (ESI) technique has made polar molecules accessible to direct analysis by MS, as well as being compatible with HPLC separations (Buchholz *et al.*, 2001). Quantification of multiple compounds in crude extracts can, in principle, be achieved the same way as GC/MS, with high sensitivity. However, automation of the procedure presents greater practical difficulties (Jonsson *et al.*, 2004). LC/MS/MS provides additional structural information that can be a very useful aid in the identification of new or unusual metabolites, or in the characterization of known metabolites in cases where ambiguity exists. LC/NMR combines the superior structure-determining power of NMR with HPLC in a flow system (Exarchou, 2005).

Direct injection into high-resolution MS. It is possible to obtain metabolite 'mass profiles' without any chromatographic separation. Such profiles are obtained by injecting crude extracts into the source of a high-resolution mass spectrometer (Dunn and Ellis, 2005). Electrospray ionization (ESI) or atmospheric pressure chemical

ionization (APCI) generates mainly protonated, deprotonated or adduct molecules, such as $[M+H]^+$, $[M+cation]^+$ or $[M-H]^-$ for each species present in the mixture, with little or no fragmentation (Nordstrom *et al.*, 2008). Thus, a fingerprint spectrum is obtained with a single or a few peaks for each metabolite, which are separated from other metabolites according to (accurate) molecular mass. The fingerprint can be used as a classification tool. Some mass analyzers (e.g., fourier transform ion cyclotron resonance instruments, FT-ICR-MS) are capable of ultra-high resolution and permit the mass to be determined to four or five decimal places (Brown *et al.*, 2005). This allows empirical formulae to be assigned to peaks. Additionally, the coupling of high sensitivity with high resolution provides a rapid method of estimating the number of metabolites present and a valuable first indication, from the formulae, of their possible identities. Its main weakness is the inability to separate isomers of the same molecular mass (Pitt *et al.*, 2002).

FT-ICR mass spectrometry. Fourier transform (FT)-ion cyclotron resonance (ICR) mass spectrometry (MS) has traditionally not been widely used. FT-ICR spectrum can measure many mass spectra per second, making it increasingly attractive in the pharmaceutical industry because of its ability to deliver more information per measurement (Brown *et al.*, 2005). FT-ICR provides ultra-high mass resolution and mass accuracy, non-destructive detection, high sensitivity and multi-stage MS_n (Brown *et al.*, 2005). It has undergone rapid development and is now applied in many fields. High mass accuracy ensures rapid protein identification with high confidence based on single peptide mass measurements (Brown *et al.*, 2005). Many metabolites are in a mass range where FT-ICR can give immediate elemental composition, enabling direct identification without MS/MS, by comparison with public or locally generated databases (Zhang *et al.*, 2005).

Multivariate statistical analysis (pattern recognition methods).

Pattern recognition (PR) and related multivariate statistical approaches can be used to discern significant patterns in complex multivariate data sets and are particularly appropriate in situations where there are more variables than samples in the data set (Lindon *et al.*, 2004). The general aim of PR is to classify objects or to predict the origin of objects based on identification of inherent patterns in a set of measurements (Dunn *et al.*, 2005). PR methods can reduce the dimensionality of complex data sets via 2 dimensions (2D) or 3 dimensions (3D) mapping procedures, thereby facilitating the visualization of inherent patterns in the

data. Methods such as principal components analysis (PCA) are termed 'unsupervised' techniques, in that no *a priori* knowledge of the class of the samples is required, and they are based on the calculation of latent variables (Weckwerth and Morgenthal, 2005). Principal components are linear combinations of the original descriptors, such that they are uncorrelated, and describe decreasing amounts of data variance (that is, PC1 > PC2 > PC3 and so on). Use of PCA enables the 'best' representation, in terms of biochemical variation in the data set, to be shown in 2D or 3D. In addition, multi-parametric data can be modeled, so that the class of a sample from an independent data set can be predicted on the basis of a series of mathematical models that are derived from the original data or 'training' set. These methods are known as 'supervised' methods, and use class information to maximize the separation between classes (Weckwerth and Morgenthal, 2005). Supervised methods, such as soft independent modeling of classification analogy (SIMCA), partial least squares (PLS) analysis and PLS discriminant analysis (PLS-DA), can be used to predict objects that are unknown to the system on the basis of their NMR spectral properties or MS profiles, and are therefore valuable for generating models for predicting chemical toxicity, drug efficacy, disease status, and so on (Kim *et al.*, 2008; Quinones and Kaddurah-Daouk, 2009; Um *et al.*, 2009).

Artificial neural networks (ANN). Artificial neural (or neuronal) networks (ANN) are simplified mathematical models of a biological neuronal system (Mao *et al.*, 1995). They 'learn' from existing data sets and are 'optimized' by specific algorithms (Mao *et al.*, 1995). Neurons can 'activate' their neighbors according to certain mathematical rules such as those developed by Hopfield (1982). They may then recognize familiar patterns, correct errors and remember sequences of events. An ANN takes the input data (e.g., gene expression data)

and builds a network to predict either the categorical or continuous responses (Ripley *et al.*, 1996). They are 'robust' to a moderate amount of 'noise' in the data, but if the number of input variables (genes) > 1000, then the technique requires intensive computer-aid.

Linear discriminant analysis (LDA). LDA is a statistical technique that can be used for the classification of individuals into mutually exclusive and exhaustive groups based on a set of independent variables. The LDA involves finding a linear combination of the independent variables that minimizes the probability of misclassifying the individuals into their respective groups (Goodacre, 2005).

TOXICOLOGICAL APPLICATIONS

Using metabolomic technology, researchers are able to systematically determine metabolite concentration in a sample. This new technology has the potential for application in the areas of drug discovery/development and preventive screening/diagnostics (Table 4). Research continues to refine this technology in an effort to put these applications to use as quickly as possible (Lindon *et al.*, 2004). Metabolic profiling (of biological samples such as urine or blood plasma) can be generally used to determine the physiological changes induced by toxic effect of a chemical (or mixture of chemicals) (Kim *et al.*, 2008, 2009). The observed metabolic changes can be closely related to specific toxicity, e.g. specific hepatic or renal lesions (Nicholson *et al.*, 1985; Kim *et al.*, 2008; Park *et al.*, 2009). This is of particular interest to pharmaceutical companies that want to evaluate the toxicity of new drug candidates. If a new compound can be screened for adverse toxicity before it reaches clinical trials, then companies gain the advantage of being able to save the enormous cost of these trials (Lindon *et al.*, 2004). Metabolomics is emerging as an

Table 4. Application of metabolomics

Application		Ref.
Drug Discovery	- The major aim is to increase compound attrition in drug discovery.	Zhang <i>et al.</i> , 2005 Wang <i>et al.</i> , 2009
	- Reduce compound attrition in development and clinical analysis.	
	- Help explain reasons for toxicity.	
	- Produce expert system to identify toxicity.	
Biomarker identification	- Markers of drug toxicity.	Kim <i>et al.</i> , 2008 Nicholson <i>et al.</i> , 1985 Park <i>et al.</i> , 2009
	- Markers of drug efficacy.	
Human disease diagnosis and personalized medicine		Yang <i>et al.</i> , 2004a, b Clayton <i>et al.</i> , 2006
Agriculture	- Measurement and identification of unintended effects of genetic modification.	Roessner <i>et al.</i> , 2001
Nutrition	- Naturally functional foods.	Gidley <i>et al.</i> , 2004

excellent tool for elucidating the phenotype induced by a genetic modification, such as gene insertion or deletion. Metabolomics has the unique ability to relate direct phenotypic change with metabolic profile rather than genomics or proteomics (Holmes and Antti, 2002). It is more exciting to predict the function of unknown genes by comparing the metabolic perturbations induced by insertion/deletion of known genes. *Saccharomyces cerevisiae* and *Arabidopsis thaliana* are good model organisms for such advances in metabolomics and functional genomics (Allen *et al.*, 2003; Saito and Matsuda, 2008).

Biomarker discovery for candidate drugs. One of the most interesting goals of pharmaceutical industry is to select robust new drug candidates for development or screen them for adverse effects in order to save money and time. Early preclinical screening of candidate drug toxicity is cost-effective and can suggest expected harmful side effects in clinical trials in drug development (Lindon *et al.*, 2004). Therefore, an important issue has been emerging to assess and screen toxicity of new molecular entities in early stages of new drug development. Biomarker discovery has been one of representative applications using metabolomics (Table 5). Consortium for Metabonomic Toxicology (COMET) has recently explored the evaluation of xenobiotic toxicity by biomarkers using metabolomics (Lindon *et al.*, 2003). COMET was formed between Imperial College London, UK and 6 major pharmaceutical companies to apply metabolomics data produced using ^1H NMR spectroscopy of urine and blood serum samples to preclinical toxicological screening of candidate drugs (Ebbels *et al.*, 2007). Kim *et al.* (2008) reported that endoge-

nous metabolites of allantoin, citrate, taurine, 2-oxoglutarate, acetate, lactate, phenylacetyl glycine, succinate, phenylacetate, 1-methylnicotinamide, hippurate, and benzoate were closely related to hepatotoxicity caused by CCl_4 , acetaminophen, and D-galactosamine using 600 M ^1H NMR spectroscopy. These results suggested that these endogenous metabolites could be used as putative biomarkers for preclinical hepatotoxicity. Recently, high throughput toxicity and safety screening methods were developed through a combination of metabolite profiles (Ebbels *et al.*, 2007). Schoonen *et al.* (2007) reported that the total NMR dataset of rat urine showed more sensitivity to liver toxicity caused by bromobenzene and paracetamol than histopathology and clinical chemistry.

In addition, phospholipid metabolic profiles were studied and potential biomarkers were identified in rat plasma using HPLC-MS after γ -irradiation exposure (Wang *et al.*, 2009). Phosphatidylethanolamine and phosphatidylserine were assigned to be biomarkers for exposure to gamma-rays. Metabolomic investigation of toxicity caused by doxorubicin (adriamycin) was performed in rats using NMR spectroscopy (Park *et al.*, 2009). Presence of trimethylamine N-oxide (TMAO), glucose, lactate, alanine, and valine were related to renal toxicity induced by doxorubicin and creatine, phenylacetyl glycine, N-methylnicotinic acid, and hippurate levels were suggestive of liver injury (Park *et al.*, 2009). Metabolic profiling of realgar in rats was recently studied for toxicological effects using NMR spectroscopy (Wei *et al.*, 2009). Various endogenous metabolites were suggested as biomarkers correlated to liver and kidney toxicity caused by realgar. Metabolomic studies for

Table 5. Metabolomic biomarkers associated with target organ toxicity

Target organ	Toxicants	Biomarkers	Ref.
Liver toxicity	Allyl alcohol	↑ creatinine, lactate, phenylacetyl glycine, ↓ N-methyl nicotinamide, taurine	Beckwith-Hall <i>et al.</i> , 1998
	Bromobenzene	↑ 5-oxoproline, glucose, acetate, lactate ↓ citrate, α -ketoglutarate, succinate	Waters <i>et al.</i> , 2006
	α -Naphthylisocyanate	↑ taurine, creatine, glucose ↓ citrate, α -ketoglutarate, succinate	Waters <i>et al.</i> , 2001
	Methapyrilene	↑ succinate, triglyceride, dimethylglycine, trimethylamine-N-oxide ↓ glucose, glycogen	Craig <i>et al.</i> , 2006
	Hydrazine	↑ β -alanine, 3-D-hydroxybutyrate, citrulline, N-acetyl-citrulline ↓ trimethylamine-N-oxide	Bollard <i>et al.</i> , 2005
Renal toxicity	Gentamicin	↑ glucose ↓ trimethylamine-N-oxide	Lenz <i>et al.</i> , 2005
	Cisplatin	↑ alanine, leucine, glucose, 2-oxoglutarate, pyruvate, valine ↓ trimethylamine	Garrod <i>et al.</i> , 2001
	Mercuric chloride	↑ acetate, amino acids, glucose, organic acids ↓ citate, creatinine, hippurate, α -ketoglutarate, succinate.	Nicholson <i>et al.</i> , 1985

investigation of biomarker discovery have been carried out and several endogenous metabolites were useful for prediction or screening of specific toxic effects caused by chemicals.

Connections to genomics and proteomics. Metabolomics can be very useful for functional genomics and proteomics. Many companies that invested in genomic and proteomic approaches have augmented their work with metabolomic technologies (Nicholson *et al.*, 1999). For example, if a company is interested in a specific gene, but it is not sure what the gene product does, it can create a gene knock-out and compare its metabolic profile against that of the wild-type. This will give direct information as to the function of a protein and its coding gene. There is a linear progression of biological events, from genomics to metabolomics, which opens a new post-genomic era and a new approach to elucidate mechanisms of toxicities in toxicological research: e.g. the analysis of genetic variation (toxicogenomics), gene expression (transcriptomics), gene products (toxicoproteomics) and their metabolic effects (metabolomics) (Kell, 2004). The genetic approach to unravel the complexity inherent within human biology is very limited. But, it provides only one layer of complexity. To understand common human disease traits, the integration of multiple 'omics' datasets is required (Zhu *et al.*, 2007). As seen in recent literature, there have been attempts to combine two datasets related to clinical phenotype, such as pharmacogenomics, toxicogenomics, epigenetics, and metagenomics. Recently, combining genetics and metabolomics has been applied. Cholesterol-associated susceptible genes for Alzheimer's disease were identified by combining genetics with sterol metabolic profiling (Papassotiropoulos *et al.*, 2005). The ratio of metabolites was used to identify the function of putative genes (Gieger *et al.*, 2008). Stylianou and colleagues (2008) linked proteomics to quantitative trait loci (QTL) to identify changes in function rather than the protein quantification. To elucidate complex biological phenomenon, combining multiple types of technologies, including genetics, transcriptomics, proteomics and metabolomics, should be performed.

Disease diagnosis. Recently, metabolomics studies have been used to screen the patients for general metabolic disorders using NMR analysis (van Doorn *et al.*, 2007). There has been a report that analysis of serum lipid profiles using NMR could discriminate tumors (83%) and non-tumors (8%) from 52 patients diagnosed as coronary heart disease (Brindle *et al.*, 2002). Colon cancer and coronary heart disease could be pre-

cisely diagnosed from healthy volunteers through neural network analysis and classification of NMR spectra of lipoprotein fractions (Bathen *et al.*, 2000). Moreover, testosterone and its metabolites, epitestosterone and dihydrotestosterone were measured from human hair using GC-MS (Choi *et al.*, 2001). Hair samples from bald men presented separate metabolic profiles from healthy volunteers and testosterone/epitestosterone ratio was suggested as biochemical biomarker useful for diagnosis of early baldness (Choi *et al.*, 2001). Research on the correlation between cervical cancer and urinary polyamines and endogenous steroids using MS, showed that the ratios of 16 α -hydroxyestrone/2-hydroxyestrone, 5 β -tetrahydrocortisol/5 α -tetrahydrocortisol, and putresine/N-acetylspermidine were very crucial for diagnosis of cervical cancer (Lee *et al.*, 2003). After liver fibrosis and cirrhosis were progressively induced by thioacetamide in rats, liver extracts and serum were analyzed using ¹H NMR to characterize the stage of fibrosis (Constantinou *et al.*, 2007). PCA profiles from lipid liver extract and serum showed the progress of fibrosis and cirrhosis. Serum leucine, isoleucine, valine, lactate, alanine, acetate, acetoacetate, glutamine, trimethylamine, creatine, and glucose were sensitively influenced during liver fibrosis and cirrhosis caused by thioacetamide and this study could be extended in clinical diagnosis for management of cirrhotic patients (Constantinou *et al.*, 2007). Cerebrospinal fluid (CSF), a clear bodily fluid that occupies the subarachnoid space and the ventricular system around and inside the brain, can serve as a metabolomic sample for investigating Alzheimer's disease, meningitis, and so on. CSF samples were used to distinguish normal subjects from those with meningitis using NMR spectroscopy (Coen *et al.*, 2005). Early detection of oral cancer using NMR spectroscopy was studied in patients' serum (Tiziani *et al.*, 2009). Serum metabolite profile completely discriminated cancer patients from control group and also between different stages of oral cancer (Tiziani *et al.*, 2009). Normal colon tissues and colorectal cancer tissues were analyzed with GC-TOF (time-of-flight)-MS to investigate metabolomic profiling of human colon carcinoma (Denkert *et al.*, 2008). In cancer, intermediates of the TCA cycle and lipids were down-regulated, whereas urea cycle metabolites, purines, pyrimidines and amino acids were up-regulated compared to normal samples (Denkert *et al.*, 2008).

Risk assessment. Metabolomics can be of great interest to risk assessment of toxic substances. There are two advantages of metabolomics for risk assessment: understanding molecular mechanisms of toxicity

and sensitive exposure assessment. Kim *et al.* (2009) studied the exposure assessment of methoxychlor, an organochlorine pesticide, to rats using urinary NMR spectral data and compared it with traditional exposure of methoxychlor, using uterotrophic assay, in ovariectomized female rats for 3 consecutive days. Metabolomics dataset was much more sensitive to methoxychlor rather than traditional uterotrophic assay and the endogenous metabolites of acetate, alanine, benzoate, lactate and glycine were suggested as putative exposure biomarkers for an endocrine disruptor of methoxychlor (Kim *et al.*, 2009). Nonylphenol is an environmental contaminant and endocrine disrupting chemical. Lee *et al.* (2007) investigated the metabolomic profiles for endocrine toxic effects caused by nonylphenol, using GC-MS. Urinary tetrahydrocorticosterone and 5 α -tetrahydrocorticosterone were suggested as possible biomarkers of nonylphenol-induced toxicity or exposure.

Future prospects and concluding remarks. Metabolomics has been considered an emerging and promising technology in toxicological research, although its validation is required. Using metabolomics, the data has shown that combinational biomarkers for toxicity or disease were identified, which were able to monitor the toxicity or efficacy of chemicals or drugs in preclinical or clinical trials. Also, new biochemical assays for disease diagnosis were derived using NMR or MS analysis. As one part of multiple 'omics' technologies, metabolomics plays an important role in understanding biological phenomenon because it's very close to the phenotype of biological effects. Risk assessment is a very promising area to explore toxic mechanism or exposure assessment using metabolomics. One thing that needs to be pointed out for metabolomics is totally based on analytical method and multivariate analysis. Therefore, the analytical procedures should be stable, robust, and highly reproducible. Multivariate analysis can turn complex dataset into readable and interpretable. In contrast to other 'omics', metabolomics has the advantage of having a good level of biological reproducibility, low cost of per sample, minimal invasion of sampling, and direct identification of phenotypes with real biological endpoints. However, it requires a further research for validation, specificity, and sensitivity for the prediction of toxic manifestations. In addition, a comparative relationship between conventional biomarkers and new biomarkers derived from metabolomics should be needed.

REFERENCES

Allen, J., Davey, H.M., Broadhurst, D., Heald, J.K., Rowland,

- J.J., Oliver, S.G. and Kell, D.B. (2003). High-throughput classification of yeast mutants for functional genomics using metabolic footprinting. *Nat. Biotechnol.*, **21**, 692-696.
- Bathen, T.F., Engan, T., Krane, J. and Axelson, D. (2000). Analysis and classification of proton NMR spectra of lipoprotein fractions from healthy volunteers and patients with cancer or CHD. *Anticancer Res.*, **20**, 2393-2408.
- Beckwith-Hall, B.M., Nicholson, J.K., Nicholls, A.W., Foxall, P.J., Lindon, J.C., Connor, S.C., Abdi, M., Connelly, J. and Holmes, E. (1998). Nuclear magnetic resonance spectroscopic and principal components analysis investigations into biochemical effects of three model hepatotoxins. *Chem. Res. Toxicol.*, **11**, 260-272.
- Bino, R.J., Hall, R.D., Fiehn, O., Kopka, J., Saito, K., Draper, J., Nikolau, B.J., Mendes, P., Roessner-Tunali, U., Beale, M.H., Trethewey, R.N., Lange, B.M., Wurtele, E.S. and Sumner, L.W. (2004). Potential of metabolomics as a functional genomics tool. *Trends Plant Sci.*, **9**, 418-425.
- Bollard, M.E., Keun, H.C., Beckonert, O., Ebbels, T.M., Antti, H., Nicholls, A.W., Shockcor, J.P., Cantor, G.H., Stevens, G., Lindon, J.C., Holmes, E. and Nicholson, J.K. (2005). Comparative metabolomics of differential hydrazine toxicity in the rat and mouse. *Toxicol. Appl. Pharmacol.*, **204**, 135-151.
- Brindle, J.T., Antti, H., Holmes, E., Tranter, G., Nicholson, J.K., Bethell, H.W., Clarke, S., Schofield, P.M., McKilligan, E., Mosedale, D.E. and Grainger, D.J. (2002). Rapid and noninvasive diagnosis of the presence and severity of coronary heart disease using 1H-NMR-based metabolomics. *Nat. Med.*, **8**, 1439-1444.
- Brown, S.C., Kruppa, G. and Dasseux, J.L. (2005). Metabolomics applications of FT-ICR mass spectrometry. *Mass Spectrom. Rev.*, **24**, 223-231.
- Buchholz, A., Takors, R. and Wandrey, C. (2001). Quantification of intracellular metabolites in *Escherichia coli* K12 using liquid chromatographic-electrospray ionization tandem mass spectrometric techniques. *Anal. Biochem.*, **295**, 129-137.
- Bundy, J.G., Willey, T.L., Castell, R.S., Ellar, D.J. and Brindle, K.M. (2005). Discrimination of pathogenic clinical isolates and laboratory strains of *Bacillus cereus* by NMR-based metabolomic profiling. *FEMS Microbiol. Lett.*, **242**, 127-136.
- Celia, M.H. (2002). The "metabonome" provides real-world information about drug toxicity, gene function. *Chem. Eng. News*, **80**, 66-70.
- Choi, M.H., Yoo, Y.S. and Chung, B.C. (2001). Biochemical roles of testosterone and epitestosterone to 5 alpha-reductase as indicators of male-pattern baldness. *J. Invest. Dermatol.*, **116**, 57-61.
- Clayton, T.A., Lindon, J.C., Cloarec, O., Antti, H., Charuel, C., Hanton, G., Provost, J.P., Le Net, J.L., Baker, D., Walley, R.J., Everett, J.R. and Nicholson, J.K. (2006). Pharmacometabolomic phenotyping and personalized drug treatment. *Nature*, **440**, 1073-1077.
- Coen, M., O'Sullivan, M., Bubb, W.A., Kuchel, P.W. and Sorrell, T. (2005). Proton nuclear magnetic resonance-based metabolomics for rapid diagnosis of meningitis and ven-

- triculitis. *Clin. Infect. Dis.*, **41**, 1582-1590.
- Constantinou, M.A., Theocharis, S.E. and Mikros, E. (2007). Application of metabolomics on an experimental model of fibrosis and cirrhosis induced by thioacetamide in rats. *Toxicol. Appl. Pharmacol.*, **218**, 11-19.
- Craig, A., Sidaway, J., Holmes, E., Orton, T., Jackson, D., Rowlinson, R., Nickson, J., Tonge, R., Wilson, I. and Nicholson, J. (2006). Systems toxicology: integrated genomic, proteomic and metabolomic analysis of methapyrilene induced hepatotoxicity in the rat. *J. Proteome Res.*, **5**, 1586-1601.
- Denkert, C., Budczies, J., Weichert, W., Wohlgemuth, G., Scholz, M., Kind, T., Niesporek, S., Noske, A., Buckendahl, A., Dietel, M. and Fiehn, O. (2008). Metabolite profiling of human colon carcinoma—deregulation of TCA cycle and amino acid turnover. *Mol. Cancer*, **7**, 72.
- Dunn, W.B., Bailey, N.J.C. and Johnson, H.E. (2005). Measuring the metabolome: current analytical technologies. *Analyst*, **130**, 606-625.
- Dunn, W.B. and Ellis, D.I. (2005). Metabolomics: current analytical platforms and methodologies. *Trac-trends In Analytical Chemistry*, **24**, 285-294.
- Ebbels, T.M., Keun, H.C., Beckonert, O.P., Bollard, M.E., Lindon, J.C., Holmes, E. and Nicholson, J.K. (2007). Prediction and classification of drug toxicity using probabilistic modeling of temporal metabolic data: the consortium on metabolomic toxicology screening approach. *J. Proteome Res.*, **6**, 4407-4422.
- Exarchou, V., Krucker, M., van Beek, T.A., Vervoort, J., Gerotheranassis, I.P. and Albert, K. (2005). LC-NMR coupling technology: recent advancements and applications in natural products analysis. *Magn. Reson. Chem.*, **43**, 681-687.
- Fan, X., Bai, J. and Shen, P. (2005). Diagnosis of breast cancer using HPLC metabolomics fingerprints coupled with computational methods. *Conf. Proc. IEEE Eng. Med. Biol. Soc.*, **6**, 6081-6084.
- Fiehn, O. (2002). Metabolomics—the link between genotypes and phenotypes. *Plant Mol. Biol.*, **48**, 155-171.
- Fiehn, O., Kopka, J., Dormann, P., Altmann, T., Trethewey, R. N. and Willmitzer, L. (2000). Metabolite profiling for plant functional genomics. *Nat. Biotechnol.*, **18**, 1157-1161.
- Gidley, M., Wahlqvist, M., Okada, A., Samman, S. and Sullivan, D. (2004). Naturally functional foods—challenges and opportunities. *Proc. Nutri. Soc. Australia*, **13**, 531.
- Gieger, C., Geistlinger, L., Altmayer, E., Hrabec de Angelis, M., Kronenberg, F., Meitinger, T., Mewes, H.W., Wichmann, H.E., Weinberger, K.M., Adamski, J., Illig, T. and Suhre, K. (2008). Genetics meets metabolomics: a genome-wide association study of metabolite profiles in human serum. *PLoS Genet.*, **4**, e1000282.
- Goodacre, R. (2005). Making sense of the metabolome using evolutionary computation: seeing the wood with the trees. *J. Exp. Bot.*, **56**, 245-254.
- Griffin, J.L. (2003). Metabolomics: NMR spectroscopy and pattern recognition analysis of body fluids and tissues for characterisation of xenobiotic toxicity and disease diagnosis. *Curr. Opin. Chem. Biol.*, **7**, 648-654.
- Holmes, E. and Antti, H. (2002). Chemometric contributions to the evolution of metabolomics: mathematical solutions to characterising and interpreting complex biological NMR spectra. *Analyst*, **127**, 1549-1557.
- Holmes, E., Nicholls, A.W., Lindon, J.C., Ramos, S., Spraul, M., Neidig, P., Connor, S.C., Connelly, J., Damment, S.J., Haselden, J. and Nicholson, J.K. (1998). Development of a model for classification of toxin-induced lesions using ¹H NMR spectroscopy of urine combined with pattern recognition. *NMR Biomed.*, **11**, 235-244.
- Hopfield, J.J. (1982). Neural networks and physical systems with emergent collective computational abilities. *Proc. Natl. Acad. Sci. U.S.A.*, **79**, 2554-2558.
- Hrmova, M. and Fincher, G.B. (2009). Functional genomics and structural biology in the definition of gene function. *Methods Mol. Biol.*, **513**, 199-227.
- Jonsson, P., Gullberg, J., Nordstrom, A., Kusano, M., Kowalczyk, M., Sjostrom, M. and Moritz, T. (2004). A strategy for identifying differences in large series of metabolomic samples analyzed by GC/MS. *Anal. Chem.*, **76**, 1738-1745.
- Kell, D.B. (2004). Metabolomics and systems biology: making sense of the soup. *Curr. Opin. Microbiol.*, **7**, 296-307.
- Kenney, B. and Shockcor, J.P. (2003). Complementary NMR and LC-MS technologies for metabolomic studies. *PharmaGenomics*, 56-63.
- Khandelwal, P., Beyer, C.E., Lin, Q., Schechter, L.E. and Bach, A.C., 2nd. (2004). Studying rat brain neurochemistry using nanoprobe NMR spectroscopy: a metabolomics approach. *Anal. Chem.*, **76**, 4123-4127.
- Kim, K.B., Chung, M.W., Um, S.Y., Oh, J.S., Kim, S.H., Na, M.A., Oh, H.Y., Cho, W.S. and Choi, K.H. (2008). Metabolomics and biomarker discovery: NMR spectral data of urine and hepatotoxicity by carbon tetrachloride, acetaminophen, and D-galactosamine in rats. *Metabolomics*, **4**, 377-392.
- Kim, K.B., Kim, S.H., Um, S.Y., Chung, M.W., Oh, J.S., Jung, S.C., Kim, T.S., Moon, H.J., Han, S.Y., Oh, H.Y., Lee, B.M. and Choi, K.H. (2009). Metabolomics approach to risk assessment: methoxychlor exposure to rats. *J. Toxicol. Environ. Health A* (in print).
- Lee, S.H., Woo, H.M., Jung, B.H., Lee, J., Kwon, O.S., Pyo, H.S., Choi, M.H. and Chung, B.C. (2007). Metabolomic approach to evaluate the toxicological effects of nonylphenol with rat urine. *Anal. Chem.*, **79**, 6102-6110.
- Lee, S.H., Yang, Y.J., Kim, K.M. and Chung, B.C. (2003). Altered urinary profiles of polyamines and endogenous steroids in patients with benign cervical disease and cervical cancer. *Cancer Lett.*, **201**, 121-131.
- Lenz, E.M., Bright, J., Knight, R., Westwood, F.R., Davies, D., Major, H. and Wilson, I.D. 2005. Metabolomics with ¹H-NMR spectroscopy and liquid chromatography-mass spectrometry applied to the investigation of metabolic changes caused by gentamicin-induced nephrotoxicity in the rat. *Biomarkers*, **10**, 173-187.
- Lenz, E.M., Bright, J., Knight, R., Westwood, F.R., Davies, D., Major, H. and Wilson, I.D. 2005. Metabolomics with ¹H-NMR spectroscopy and liquid chromatography-mass spectrometry applied to the investigation of metabolic changes caused by gentamicin-induced nephrotoxicity in the rat. *Biomarkers*, **10**, 173-187.

- Lenz, E.M., Bright, J., Wilson, I.D., Hughes, A., Morrisson, J., Lindberg, H. and Lockton, A. (2004). Metabonomics, dietary influences and cultural differences: a ^1H NMR-based study of urine samples obtained from healthy British and Swedish subjects. *J. Pharm. Biomed. Anal.*, **36**, 841-849.
- Leo, G.C., Caldwell, G.W., Croke, J., Malatynska, E., Cotto, C., Hastings, B., Scowcroft, J., Hall, J., Browne, K. and Hageman, W. (2005). The application of nuclear magnetic resonance-based metabonomics to the dominant-submissive rat behavioral model. *Anal. Biochem.*, **339**, 174-178.
- Lindon, J.C., Holmes, E., Bollard, M.E., Stanley, E.G. and Nicholson, J.K. (2004). Metabonomics technologies and their applications in physiological monitoring, drug safety assessment and disease diagnosis. *Biomarkers*, **9**, 1-31.
- Lindon, J.C., Nicholson, J.K., Holmes, E., Antti, H., Bollard, M.E., Keun, H., Beckonert, O., Ebbels, T.M., Reilly, M.D., Robertson, D., Stevens, G.J., Luke, P., Breaux, A.P., Cantor, G.H., Bible, R.H., Niederhauser, U., Senn, H., Schlotterbeck, G., Sidelmann, U.G., Laursen, S.M., Tymiak, A., Car, B.D., Lehman-McKeeman, L., Colet, J.M., Loukaci, A. and Thomas, C. (2003). Contemporary issues in toxicology the role of metabonomics in toxicology and its evaluation by the COMET project. *Toxicol. Appl. Pharmacol.*, **187**, 137-146.
- Mao, J., Jain, A.K., Center, I.B.M.A.R. and San Jose, C. (1995). Artificial neural networks for feature extraction and multivariate data projection. *IEEE Trans. Neural Networks*, **6**, 296-317.
- Nealson, K.H. and Conrad, P.G. (1999). Life: past, present and future. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.*, **354**, 1923-1939.
- Nicholson, J.K. and Wilson, I.D. (1989). High resolution proton magnetic resonance spectroscopy of biological fluids. *Prog. NMR Spectrosc.*, **21**, 449-501.
- Nicholson, J.K., Lindon, J.C. and Holmes, E. (1999). 'Metabonomics': understanding the metabolic responses of living systems to pathophysiological stimuli via multivariate statistical analysis of biological NMR spectroscopic data. *Xenobiotica*, **29**, 1181-1189.
- Nicholson, J.K., Timbrell, J.A. and Sadler, P.J. (1985). Proton NMR spectra of urine as indicators of renal damage. Mercury-induced nephrotoxicity in rats. *Mol. Pharmacol.*, **27**, 644-651.
- Nordstrom, A., Want, E., Northen, T., Lehtio, J. and Siuzdak, G. (2008). Multiple ionization mass spectrometry strategy used to reveal the complexity of metabolomics. *Anal. Chem.*, **80**, 421-429.
- Papassotiropoulos, A., Wollmer, M.A., Tzolaki, M., Brunner, F., Molyva, D., Lutjohann, D., Nitsch, R.M. and Hock, C. (2005). A cluster of cholesterol-related genes confers susceptibility for Alzheimer's disease. *J. Clin. Psychiatry*, **66**, 940-947.
- Park, J.C., Hong, Y.S., Kim, Y.J., Yang, J.Y., Kim, E.Y., Kwack, S.J., Ryu do, H., Hwang, G.S. and Lee, B.M. (2009). A metabonomic study on the biochemical effects of doxorubicin in rats using (^1H) -NMR spectroscopy. *J. Toxicol. Environ. Health A*, **72**, 374-384.
- Pitt, J.J., Eggington, M. and Kahler, S.G. (2002). Comprehensive screening of urine samples for inborn errors of metabolism by electrospray tandem mass spectrometry. *Clin. Chem.*, **48**, 1970-1980.
- Plumb, R.S., Stumpf, C.L., Gorenstein, M.V., Castro-Perez, J.M., Dear, G.J., Anthony, M., Sweatman, B.C., Connor, S.C. and Haselden, J.N. (2002). Metabonomics: the use of electrospray mass spectrometry coupled to reversed-phase liquid chromatography shows potential for the screening of rat urine in drug development. *Rapid Commun. Mass Spectrom.*, **16**, 1991-1996.
- Price, K.E., Vandaveer, S.S., Lunte, C.E. and Larive, C.K. (2005). Tissue targeted metabonomics: metabolic profiling by microdialysis sampling and microcoil NMR. *J. Pharm. Biomed. Anal.*, **38**, 904-909.
- Quinones, M.P. and Kaddurah-Daouk, R. (2009). Metabolomics tools for identifying biomarkers for neuropsychiatric diseases. *Neurobiol. Dis.* (Epub ahead of print) doi:10.1016/j.nbd.2009.02.019.
- Raamsdonk, L.M., Teusink, B., Broadhurst, D., Zhang, N., Hayes, A., Walsh, M.C., Berden, J.A., Brindle, K.M., Kell, D.B., Rowland, J.J., Westerhoff, H.V., van Dam, K. and Oliver, S.G. (2001). A functional genomics strategy that uses metabolome data to reveal the phenotype of silent mutations. *Nat. Biotechnol.*, **19**, 45-50.
- Raman, R., Raguram, S., Venkataraman, G., Paulson, J.C. and Sasisekharan, R. (2005). Glycomics: an integrated systems approach to structure-function relationships of glycans. *Nat. Methods*, **2**, 817-824.
- Rampitsch, C. and Bykova, N.V. (2009). Methods for functional proteomic analyses. *Methods Mol. Biol.*, **513**, 93-110.
- Reo, N.V. (2002). NMR-based metabolomics. *Drug Chem. Toxicol.*, **25**, 375-382.
- Ripley, B.D. (1996). Pattern recognition and neural networks. 1st Ed. Cambridge university press, UK.
- Roessner, U., Willmitzer, L. and Fernie, A.R. (2001). High-resolution metabolic phenotyping of genetically and environmentally diverse potato tuber systems. Identification of phenocopies. *Plant Physiol.*, **127**, 749-764.
- Saito, K. and Matsuda, F. (2008). Metabolomics for Functional Genomics, Systems Biology, and Biotechnology. *Annu. Rev. Plant Biol.* (Epub ahead of print).
- Schoonen, W.G., Kloks, C.P., Ploemen, J.P., Horbach, G.J., Smit, M.J., Zandberg, P., Mellema, J.R., Zuylen, C.T., Tas, A.C., van Nesselrooij, J.H. and Vogels, J.T. (2007). Sensitivity of (^1H) NMR analysis of rat urine in relation to toxicometabonomics. Part I: dose-dependent toxic effects of bromobenzene and paracetamol. *Toxicol. Sci.*, **98**, 271-285.
- Stylianou, I.M., Affourtit, J.P., Shockley, K.R., Wilpan, R.Y., Abdi, F.A., Bhardwaj, S., Rollins, J., Churchill, G.A. and Paigen, B. (2008). Applying gene expression, proteomics and single-nucleotide polymorphism analysis for complex trait gene identification. *Genetics*, **178**, 1795-1805.
- Testa, B. and Kramer, S.D. 2006. The biochemistry of drug metabolism—an introduction: part 1. Principles and overview. *Chem. Biodivers*, **3**, 1053-1101.
- Tiziani, S., Lopes, V. and Gunther, U.L. (2009). Early stage

- diagnosis of oral cancer using ¹H NMR-based metabolomics. *Neoplasia*, **11**, 269-276.
- Um, S.Y., Chung, M.W., Kim, K.B., Kim, S.H., Oh, J.S., Oh, H.Y., Lee, H.J. and Choi, K.H. (2009). Pattern recognition analysis for the prediction of adverse effects by NSAIDs using ¹H-NMR based metabolomics in rats. *Anal. Chem.* (in print).
- van Doorn, M., Vogels, J., Tas, A., van Hoogdalem, E.J., Burggraaf, J., Cohen, A. and van der Greef, J. (2007). Evaluation of metabolite profiles as biomarkers for the pharmacological effects of thiazolidinediones in Type 2 diabetes mellitus patients and healthy volunteers. *Br. J. Clin. Pharmacol.*, **63**, 562-574.
- Wang, C., Yang, J. and Nie, J. (2009). Plasma phospholipid metabolic profiling and biomarkers of rats following radiation exposure based on liquid chromatography-mass spectrometry technique. *Biomed. Chromatogr.* (Epub ahead of print) 10.1002/bmc.1226.
- Waters, N.J., Holmes, E., Williams, A., Waterfield, C.J., Farrant, R.D. and Nicholson, J.K. (2001). NMR and pattern recognition studies on the time-related metabolic effects of alpha-naphthylisothiocyanate on liver, urine, and plasma in the rat: an integrative metabonomic approach. *Chem. Res. Toxicol.*, **14**, 1401-1412.
- Waters, N.J., Waterfield, C.J., Farrant, R.D., Holmes, E. and Nicholson, J.K. (2006). Integrated metabonomic analysis of bromobenzene-induced hepatotoxicity: novel induction of 5-oxoprolinosis. *J. Proteome Res.*, **5**, 1448-1459.
- Watkins, S.M. and German, J.B. (2002). Metabolomics and biochemical profiling in drug discovery and development. *Curr. Opin. Mol. Ther.*, **4**, 224-228.
- Weckwerth, W. and Morgenthal, K. (2005). Metabolomics: from pattern recognition to biological interpretation. *Drug Discov. Today*, **10**, 1551-1558.
- Wei, L., Liao, P., Wu, H., Li, X., Pei, F., Li, W. and Wu, Y. (2009). Metabolic profiling studies on the toxicological effects of realgar in rats by (¹H) NMR spectroscopy. *Toxicol. Appl. Pharmacol.*, **234**, 314-325.
- Wilson, I.D., Plumb, R., Granger, J., Major, H., Williams, R. and Lenz, E.M. (2005). HPLC-MS-based methods for the study of metabonomics. *J. Chromatogr. B. Analyt. Technol. Biomed. Life Sci.*, **817**, 67-76.
- Wu, L., van Winden, W.A., van Gulik, W.M. and Heijnen, J.J. (2005). Application of metabolome data in functional genomics: a conceptual strategy. *Metab. Eng.*, **7**, 302-310.
- Yang, J., Xu, G., Hong, Q., Liebich, H.M., Lutz, K., Schmulding, R.M. and Wahl, H.G. (2004). Discrimination of Type 2 diabetic patients from healthy controls by using metabonomics method based on their serum fatty acid profiles. *J. Chromatogr. B. Analyt. Technol. Biomed. Life. Sci.*, **813**, 53-58.
- Yang, J., Xu, G., Zheng, Y., Kong, H., Pang, T., Lv, S. and Yang, Q. (2004). Diagnosis of liver cancer using HPLC-based metabonomics avoiding false-positive result from hepatitis and hepatocirrhosis diseases. *J. Chromatogr. B. Analyt. Technol. Biomed. Life. Sci.*, **813**, 59-65.
- Yang, Y., Adelstein, S.J. and Kassis, A.I. (2009). Target discovery from data mining approaches. *Drug Discov. Today*, **14**, 147-154.
- Zhang, J., McCombie, G., Guenat, C. and Knochenmuss, R. (2005). FT-ICR mass spectrometry in the drug discovery process. *Drug Discov. Today*, **10**, 635-642.
- Zhu, J., Wiener, M.C., Zhang, C., Fridman, A., Minch, E., Lum, P.Y., Sachs, J.R. and Schadt, E.E. (2007). Increasing the power to detect causal associations by combining genotypic and expression data in segregating populations. *PLoS Comput. Biol.*, **3**, e69.