





Article

Evaluation of the Cytotoxicity of Ayahuasca Beverages

Ana Y. Simão^{1,2}, Joana Gonçalves^{1,2}, Ana Gradillas³ , Antonia García³, José Restolho¹, Nicolás Fernández⁴, Jesus M. Rodilla⁵ , Mário Barroso⁶, Ana Paula Duarte^{1,2} , Ana C. Cristóvão^{1,7,*} and Eugenia Gallardo^{1,2,*} 

- ¹ Centro de Investigação em Ciências da Saúde (CICS-UBI), Universidade da Beira Interior, Avenida Infante D. Henrique, 6200-506 Covilhã, Portugal; ana.simao@ubi.pt (A.Y.S.); janitagoncalves@hotmail.com (J.G.); jose.restolho@ubi.pt (J.R.); apcd@ubi.pt (A.P.D.)
- ² Laboratório de Fármaco-Toxicologia, UBIMedical, Universidade da Beira Interior, Estrada Municipal 506, 6200-284 Covilhã, Portugal
- ³ CEMBIO, Center for Metabolomics and Bioanalysis, Facultad de Farmacia, Universidad San Pablo CEU, CEU Universities, Campus Monteprincipe, Boadilla del Monte, 28668 Madrid, Spain; gradini@ceu.es (A.G.); antogar@ceu.es (A.G.)
- ⁴ Cátedra de Toxicología y Química Legal, Laboratorio de Asesoramiento Toxicológico Analítico (CENATOXA), Facultad de Farmacia y Bioquímica, Universidad de Buenos Aires, Junín 956, Ciudad Autónoma de Buenos Aires (CABA), Buenos Aires C1113AAD, Argentina; nfernandez@ffyb.uba.ar
- ⁵ Materiais Fibrosos e Tecnologias Ambientais—FibEnTech, Departamento de Química, Universidade da Beira Interior, Rua Marquês D'Ávila e Bolama, 6201-001 Covilhã, Portugal; rodilla@ubi.pt
- ⁶ Instituto Nacional de Medicina Legal e Ciências Forenses, Serviço de Química e Toxicologia Forenses, Delegação do Sul, Rua Manuel Bento de Sousa n.º3, 1169-201 Lisboa, Portugal; mario.j.barroso@inmlcf.mj.pt
- ⁷ NEUROSOV, UBIMedical, Universidade da Beira Interior, Estrada Municipal 506, 6200-284 Covilhã, Portugal
- * Correspondence: aclaracristovao@gmail.com (A.C.C.); egallardo@fcsaude.ubi.pt (E.G.); Tel.: +351-275-329-002/3 (A.C.C. & E.G.)

Academic Editor: Francesco Cacciola

Received: 31 October 2020; Accepted: 26 November 2020; Published: 28 November 2020



Abstract: Ayahuasca is a beverage consumed at shamanic ceremonies and currently has gained popularity on recreational scenarios. It contains beta-carboline alkaloids and *N,N*-dimethyltryptamine, which possesses hallucinogenic effects. Only a few studies have elicited the psychoactive effects and the dose of such compounds on neurological dopaminergic cells or animals. In this work, we aimed to study the cytotoxic effects of these compounds present in ayahuasca beverages and on five different teas (*Banisteriopsis caapi*, *Psychotria viridis*, *Peganum harmala*, *Mimosa tenuiflora* and Dc Ab (commercial name)) preparations on dopaminergic immortalized cell lines. Moreover, a characterization of the derivative alkaloids was also performed. All the extracts were characterized by chromatographic systems and the effect of those compounds in cell viability and total protein levels were analyzed in N27 dopaminergic neurons cell line. This is the first article where cytotoxicity of ayahuasca tea is studied on neurological dopaminergic cells. Overall, results showed that both cell viability and protein contents decreased when cells were exposed to the individual compounds, as well as to the teas and to the two mixtures based on the traditional ayahuasca beverages.

Keywords: ayahuasca; cytotoxicity; mesencephalic dopaminergic neurons; *N,N*-dimethyltryptamine; beta-carboline alkaloids

1. Introduction

Ayahuasca is a psychoactive beverage prepared traditionally from a brew mixture of the leaves and stems of *Psychotria viridis* and *Banisteriopsis caapi*, respectively, being originally consumed by indigenous Amazonian tribes for ritual and medicinal purposes [1–4]. Over the years, its use has

spread to other groups not only as a source of personal growth and spiritual connection, but also in recreational scenarios, hence becoming a serious concern for public health due to its toxicity. In addition, the hypothesis of its use as a potential therapeutic drug is being discussed [4–7]. The main compounds of this tea-like preparation are *N,N*-dimethyltryptamine (DMT) and beta-carboline alkaloids or harmala alkaloids, such as harmine (HMN), tetrahydroharmine (THH) and harmaline (HML) [8–11]. Due to the fact that these compounds are present in several different plants, it is possible to replace those used in the traditional preparation for other plants, such as *P. harmala* and *Mimosa tenuiflora* also known as *M. hostilis*, amongst others [3,12–15]. Beta-carboline alkaloids are monoamine-oxidase (MAO) inhibitors, thus allowing DMT, which is the main psychoactive compound in the ayahuasca beverages, to exert its psychoactive and hallucinogenic effects on the central nervous system (CNS) [11,16]. Since DMT presents a similar structure to serotonin [17], its main biochemical effects are on the serotonergic receptors, such as 5HT_{2A}, which are known to be the site of action of hallucinogenic drugs [18]. In addition, the consumption of ayahuasca beverages with high contents of DMT, increases the levels of serotonergic receptors, but also the intake of MAO inhibitors leading to accumulation of dopamine, which is associated to the sensations of pleasure and reward, but also addiction [18]. Nevertheless, the accumulation of dopamine may contribute to its auto-oxidation, which in turn it is well known to account for the formation of neuromelanin and consequent neurodegeneration of dopaminergic neurons [19,20]. Although consumers defend the use, its effects on the CNS are not fully understood yet. Currently, there are few studies on the toxicological effects of ayahuasca [11] and to date no study has evaluated the cytotoxic effects of ayahuasca teas on dopaminergic neuronal cells; also, the majority of investigations performed so far are based on clinical trials or even on surveys. In the present study, the aim was to evaluate the toxic effects on dopaminergic rat cells of HMN, HML, THH, DMT and five different plants, which are used to prepare ayahuasca beverages, and, the traditional mixtures of *P. viridis*/*B. caapi* and *P. viridis*/*P. harmala*. To the best of our knowledge, this is the first study to investigate cytotoxicity of ayahuasca compounds and commercial teas on dopaminergic neuronal cells.

2. Results and Discussion

2.1. Characterization of Alkaloid Derivatives and Other Compounds

All five plants (*P. viridis*, *B. caapi*, *P. harmala*, *M. tenuiflora* and Dc Ab (commercial name)) were characterized in terms of alkaloid derivatives present in them.

Through GC–MS analysis in scan (Figure 1) and SIM modes it was possible to identify and determine the concentrations of DMT, HMN, THH and HML. The concentrations found were 2.06 µg/mL of DMT and 0.20 µg/mL of HML for *P. viridis*. In the case of *M. tenuiflora*, the concentration of DMT was inferior (1.70 µg/mL), and also 0.74 µg/mL of HMN was found. As for *B. caapi*, *P. harmala* and Dc Ab, all plant extracts contained THH, HML and HMN on different concentration levels. The first, contained 1.22, 0.99 and 12.81 µg/mL, respectively. As for *P. harmala*, the levels were as follows, 7.46, 26.95 and 24.27 µg/mL, it is important to note that this plant extract was the one that presented the highest levels of beta-carboline alkaloids. Finally, Dc Ab contained 1.11, 0.30 and 0.20 µg/mL of THH, HML and HMN, respectively [21].

However, due to the characteristics and limitations of GC, a more complete characterization was carried out through UHPLC/QTOF-MS through a non-targeted metabolomic approach mainly focused on secondary metabolites such as alkaloids and phenolic compounds.

The major phenolic compounds were identified by comparison with authentic reference standards (Table 1). Putative identification of the alkaloids detected (Table 1 and Figures 2 and 3) was carried out by comparing their retention times, accurate masses (± 5 ppm error) against online databases, such as FooDB (<http://foodb.ca>), CEU massmediator (<http://ceumass.eps.uspceu.es/mediator>) and scientific bibliography. The in-source fragmentation gives a pattern of fragments based on the dissociation of the protonated pseudomolecular ions $[M + H]^+$ when a 175 V fragmentor voltage is applied. The characteristic fragment ions were key to study and understand the fragmentation mechanisms

and to support metabolite annotation with a higher confidence level (Figures 2 and 3). Besides they are in agreement with previous studies [22–24].




Figure 1. Chromatogram of *Psychotria viridis*.




Figure 2. (a) Structures of the tryptamine alkaloids detected from the methanolic extracts. (b) Extracted ion chromatograms (EICs) at m/z 205.1341. A complete chromatographic separation was observed for bufotenine (5-HO-DMT) and its structural isomer, psilocin (4-HO-DMT). (c) Detail of the main fragment ions and the fragmentation pathway proposed.

Table 1. Groups of the main tryptamine and harmala alkaloids and phenolic compounds identified in the methanol extracts by UHPLC-Q/TOF-MS.

Compounds	tR	Molecular Formula	Monoisotopic Mass	[M – H] [–]	[M + H] ⁺	In-Source Fragments
Tryptamine Alkaloids						
bufotenine (5-HO-DMT)	6.4	C ₁₂ H ₁₆ N ₂ O	204.1263	–	205.1341	160.0762 134.0604
psilocin (4-HO-DMT)	7.3	C ₁₂ H ₁₆ N ₂ O	204.1263	–	205.1341	160.0762 134.0604
<i>N,N</i> -dimethyltryptamine (DMT)	8.1	C ₁₂ H ₁₆ N ₂	188.1313	–	189.1392	144.0829 118.0648
5-methoxybufotenine (5-MeO-DMT)	8.5	C ₁₃ H ₁₈ N ₂ O	218.1419	–	219.1491	174.0912 130.1588
Harmala Alkaloids						
tetrahydroharmol	7.3	C ₁₂ H ₁₄ N ₂ O	202.1106	–	203.1190	174.0919
harmalol	8.8	C ₁₂ H ₁₂ N ₂ O	200.0949	–	201.1028	160.0757
harmol	9.2	C ₁₂ H ₁₀ N ₂ O	198.0793	–	199.0960	–
tetrahydroharmine (THH)	9.4	C ₁₃ H ₁₆ N ₂ O	216.1263	–	217.1331	200.0951, 188.1070
harmane	9.7	C ₁₂ H ₁₀ N ₂	182.0844	–	183.0941	–
harmaline (HML)	9.8	C ₁₃ H ₁₄ N ₂ O	214.1106	213.1069	215.1167	200.0943, 174.0908
harmine (HMN)	10.4	C ₁₃ H ₁₂ N ₂ O	212.0950	211.0910	213.1037	198.0788, 170.0845
Phenolic Compounds						
Hydroxybenzoic Acids						
gallic acid	3.0	C ₇ H ₆ O ₅	170.0215	169.0149	–	–
protocatechuic acid	6.5	C ₇ H ₆ O ₄	154.0266	153.0201	–	–
4-hydroxybenzoic acid	8.1	C ₇ H ₆ O ₃	138.0317	137.0271	–	–
gentisic acid	8.5	C ₇ H ₆ O ₄	154.0266	153.0197	–	–
salicylic acid	11.5	C ₇ H ₆ O ₃	138.0317	137.0271	–	–
Hydroxycinnamic Acids						
3-chlorogenic acid	9.0	C ₁₆ H ₁₈ O ₉	354.0951	353.0803	–	–
5-chlorogenic acid	9.7	C ₁₆ H ₁₈ O ₉	354.0951	353.0803	–	–
Flavonoids-Flavanols						
(+)-catechin	8.6	C ₁₅ H ₁₄ O ₆	290.079	289.0723	–	–
(–)-epicatechin	9.5	C ₁₅ H ₁₄ O ₆	290.079	289.0723	–	–
quercetin-3- <i>O</i> -galactoside	10.9	C ₂₁ H ₂₀ O ₁₂	464.0955	463.0888	–	–
quercetin-3- <i>O</i> -glucoside	11.1	C ₂₁ H ₂₀ O ₁₂	464.0955	463.0888	–	–
quercetin-3- <i>O</i> -rutinoside	11.1	C ₂₇ H ₃₀ O ₁₆	610.1534	609.1461	–	–
kaempferol-3- <i>O</i> -glucoside	11.6	C ₂₁ H ₂₀ O ₁₁	448.1006	447.0933	–	–
kaempferol-3- <i>O</i> -rutinoside	11.5	C ₂₇ H ₃₀ O ₁₅	594.1615	593.1542	–	–
Flavonoids-Dihydrochalcone						
phlorizin	11.3	C ₂₁ H ₂₄ O ₁₀	436.1369	435.1302	–	–

tR: retention time (min).

Results show that *P. viridis*, *M. hostilis* and *Dc Ab* are the ones that contain the highest levels of tryptamine alkaloids, mainly DMT and 5-OH-DMT [25]. On the other hand, *B. caapi* and *P. harmala* contain the highest levels of beta-carboline harmala alkaloids, mainly HML and HMN (Figure 3), as it was expected [25].

The main fragmentation pathway proposed for tetrahydroharmol and THH involves a retro-Diels-Alder (RDA) mechanism to the tetrahydro ring, Figure 3 [23,24]. A characteristic methyl radical loss is also observed for THH, HML and HMN, Figure 3.

Table 1 includes also relevant information about phenolic compounds. The profiles of these compounds in each plant variety of ayahuasca beverages was recently published by our group, where, in addition to determining the concentration of phenolic compounds, the antioxidant activity, anti-inflammatory activity and antimicrobial properties were also evaluated [26].




Figure 3. (a) Structures of the beta-carboline alkaloids detected from the methanolic extracts. (b–d). Detail of the main fragment ions and the fragmentation pathways proposed.

2.2. Cytotoxicity Evaluation

Firstly, a cytotoxic evaluation of the individual compounds (DMT, HMN, HML and THH) was performed. Since they are water insoluble, DMSO was used as a vehicle. We used 0.2% of DMSO in order to guarantee the solubility of the standard compounds. Therefore, a preliminary study was carried out in order to assess if the solvent could induce cytotoxicity on its own (Figure 4). It is known that high amounts of this solvent ($\geq 1\text{--}5\%$) can induce cell death [27]. It was tested the toxic effect on N27 cells of the highest percentage of DMSO, i.e., 0.2%, that was used as a vehicle. As observed on Figure 4, despite existing a decrease in cell viability, when compared to untreated cells (control), no significant difference is observed. Therefore, this suggest that 0.2% of DMSO did not induce a reduction of cell viability, ruling out its contribution to the putative toxic effect of the compounds under study. For this reason, it could be assumed that, if a compound or even the teas induce an increase of cell death, this may be caused by the compound itself and not due to the presence of the vehicle.

After discarding the possibility of vehicle-induced toxicity, the cytotoxic evaluation was performed for all the individual compounds, that is, HMN, HML, THH and DMT at different concentrations. Results are shown in Figure 5.




Figure 4. The effect of 0.2% of DMSO on N27 cell viability (24 h). The highest concentration used as a vehicle in the evaluation of the respective compounds was 0.2% of DMSO (Control—untreated cells (100 ± 0.00), DMT ($1 \mu\text{M}$; 84.86 ± 6.29), HMN ($10 \mu\text{M}$; 78.04 ± 20.13), HML ($80 \mu\text{M}$; 91.40 ± 5.89) and THH ($10 \mu\text{M}$; 90.49 ± 4.36); $n = 3$, values are shown as mean \pm SEM; no significant difference was observed when performing ANOVA followed by Dunnett's multiple comparison test).




Figure 5. The effects of (A) HMN, (B) HML, (C) THH and (D) DMT on N27 cell line viability (24 h), ($n = 3$, values are shown as mean \pm SEM ** indicates values that are significantly different from control $p < 0.01$, one-way analysis of variance followed by Bonferroni's multiple comparison test). CTR: Control.

Regarding the response induced by HMN on N27 cells, a range of concentrations from 0.00064 to 10 μM was assessed. It was observed that from 0.00064 to 2 μM , cell viability was not affected during

the time of cell incubation (24 h) and no statistical differences were noticed. Nonetheless, for cells incubated with 10 μM , we were able to see a decrease in cell viability to $62.85\% \pm 4.78\%$ when compared to the control (0 μM) culture (Figure 5A). The same concentrations were studied for THH and the results were similar, that is, when cells were incubated with 10 μM of this compound, a decrease in cell viability was observed ($48.25\% \pm 16.21\%$) in comparison to the control. For all other concentrations (0.00064–2 μM), cell viability was not affected (Figure 5C). As for HML, a range of concentration levels from 0.00512 to 80 μM was evaluated. Once again, only at the highest concentration level, cell viability was affected, namely when cells were exposed to 80 μM of HML (Figure 5B). As a whole, the results suggest that a higher concentration of these compounds may induce dopaminergic neuronal toxicity. Finally, for DMT the concentration levels studied covered were from 0.0008 to 1 μM . In this case, for all concentrations no statistical differences were observed when cells were exposed to this compound (Figure 5D). These results are similar according to the literature, once the psychotropic effects of ayahuasca are only achieved when there is an interaction between DMT and beta-carbolines [11], on the other hand, recent studies showed that beta-carbolines have neuroprotective effects [28].

Once obtaining the results for the individual chemical compounds present in tea, the dopaminergic neuronal cytotoxic effect of tea extracts was analyzed. The concentrations were chosen based on the results of the cytotoxic evaluation of each compound (the highest concentration causing less cell viability and the previous). A total of five tea plants (*P. viridis*, *B. caapi*, *P. harmala*, *M. tenuiflora* (also known as *M. hostilis*) and Dc Ab (commercial name)) and two different mixtures of plants were incubated with N27 cells to evaluate their effects on cell viability (Figure 4). Each plant extract quantity was chosen according to the concentrations of the standards that caused less cell viability (Figure 5) and the concentration of compounds present in each of them after GC–MS analysis. The cytotoxic evaluation of the tea extracts is present on Figure 6.

The evaluation of *P. harmala*'s effect on N27 cells, at two different concentrations (16 and 80 μM of HML), showed a significant decrease in cell viability, suggesting a neurotoxic effect. In fact, at the highest concentration studied, the observed cell viability was of $17.72\% \pm 2.29\%$ when compared to the control culture. This cytotoxic effect, when compared with the effects observed with the individual compounds, may result from the interaction between the beta-carbolines present in this tea (THH, HMN and HML), exerting synergetic and negative effects for these cells.




Figure 6. The effects of *P. harmala* (16 and 80 μM), *B. caapi* (2 and 10 μM), *M. tenuiflora* (1 μM), *P. viridis* (1 μM) and Dc Ab (2 and 10 μM) on N27 cell viability (24 h); CTR—control, ($n = 3$, values are shown as mean \pm SEM ** indicates values that are significantly different from control $p < 0.01$, one-way analysis of variance followed by Dunnett's multiple comparison test).

The same happened for *B. caapi*, i.e., when cells were exposed to 10 μM of HMN (highest concentration) a significant decrease in cell viability was noted. From the quantification results

obtained by Simão et al., [21], this tea presents high content in HMN (12.80 µg/mL), still it also contains low traces of THH (1.12 µg/mL) and HML (0.99 µg/mL). Once again, these results suggest that the possible interaction between these compounds may cause the cytotoxic observed effects on N27 cells.

In the case of *M. tenuiflora*, it is known that DMT is part of its [3,14], still on our previous study we were able to quantify 0.74 µg/mL of HMN in this tea [21]. This may suggest the obtained result of incubation cells with this tea at 1 µM of DMT. The same was observed for *P. viridis*, only this time apart from DMT, the tea was also composed of HML [21].

Finally, for Dc Ab, only when cells were exposed to 10 µM of THH, a significant decrease of N27 cell viability was observed.

In addition, when exposing N27 cells were exposed to tea mixtures, a clear decrease in cell viability occurred (Figure 7).




Figure 7. The effects of tea plant extracts mixtures (*P. viridis* plus *B. caapi* and *P. viridis* plus *P. harmala*) on N27 cell viability ($n = 3$). The presented values are presented as mean \pm SEM. ** indicates values that are significantly different from control ($p < 0.05$, one-way analysis of variance (ANOVA)). CTR—control.

2.3. Total Protein Quantification

In order to understand the effect of the studied compounds in the cellular protein metabolism, the total protein levels quantification was performed for the highest and the lowest concentration of each compound individually (Figure 8). This assay allows measuring the total amount of intracellular protein levels, to estimate the total biomass contents of the cells that remained adherent (therefore, viable).

These results were in agreement with those obtained from the cytotoxicity assays. In other words, when cell viability decreased, so did the total cellular proteic contents, namely at the highest concentration levels. Although the number of cells decreased, we did not observe the same relation in the reduction of the total protein contents of the surviving cells. We were able to observe a higher decrease in total protein levels, when compared to the decrease in cell viability. This may be a reflection of a reduction of cell function in the presence of HML. The reduction of functionality is often associated with the decrease in transcription and transduction, which is reflected in a lower amount of total protein production [19]. Moreover, it is also known that oxidated dopamine may interfere with the normal protein degradation system, leading to an increase of degradation [29,30].




Figure 8. Protein levels when N27 cells are exposed during 24 h to HMN, THH, HML and DMT; CTR (control with untreated cells; 0.59 ± 0.08), HMN $0.00064 \mu\text{M}$ (0.65 ± 0.04), HMN $10 \mu\text{M}$ (0.19 ± 0.02), THH $0.00064 \mu\text{M}$ (0.56 ± 0.09), THH $10 \mu\text{M}$ (0.32 ± 0.05), HML $0.00512 \mu\text{M}$ (0.59 ± 0.04) and HML $10 \mu\text{M}$ (0.14 ± 0.0006); DMT $0.0008 \mu\text{M}$ (0.73 ± 0.02); DMT $1 \mu\text{M}$ (0.59 ± 0.03). $n = 3$, values are shown as mean \pm SEM * indicates values that are significantly different from control $p < 0.05$, one-way analysis of variance followed by Dunnett's multiple comparison test. CTR—control.

Cellular protein levels is determined both by the amounts that are synthesized and the rates of degradation, which in the latter case there are different pathways that explain how this may occur: the ubiquitin–proteasome system, autophagy or by degradation mediated by proteolysis in the lysosome [31]. Although this remains to be evaluated in future studies, the results obtained on the reduction of the total protein content in N27 cells exposed to each compound may be explained by the putative inhibitory effect on the protein synthesis machinery or activator effect on the protein degradation systems, exerted by each compound. For instance, on a study by Valente and coworkers [32], it was shown a relationship between the neurotoxic effects caused by two β -keto amphetamines on SH-SY5Y human dopaminergic cells and autophagy dysregulation. The results obtained by these authors suggest that the neurotoxicity of such compounds lead to the production of reactive oxygen species and consequently to an activation of autophagy, both in a time and concentration-dependent manner. Despite being different compounds from those studied in this work, one can speculate that psychoactive compounds may induce autophagy and downregulate protein levels and ultimately induce cellular apoptosis [32]. Nevertheless, future studies need to be performed to understand if the compounds studied herein affect the autophagy flux.

3. Materials and Methods

3.1. Sample Preparation

Samples of *P. viridis*, *B. caapi*, *M. hostilis* and *P. harmala*, and the commercial mixture Dc Ab were purchased online from the Shayana Shop (<https://www.shayanashop.com>, Amsterdam, The Netherlands). Five decoctions of each individual plant were prepared, for which 0.210 g of each sample was macerated in a mortar with a few drops of water. The mixture was transferred to a flask and 250 mL of ultrapure water was added. The flasks were boiled at $100 \text{ }^\circ\text{C}$ for 4 h. Similarly, two decoctions were prepared, in which two vegetal samples were mixed (*P. viridis* and *B. caapi*; *P. viridis* and *P. harmala*; *M. hostilis* and *B. caapi*; *M. hostilis* and *P. harmala*).

On the one hand, the sample preparation method previously described by Simão et al. [21] was carried out to perform the determination of the major metabolites on the five different plants and in the tea mixtures. The method was performed as follows: after solid phase extraction, the obtained extracts were evaporated to dryness under a stream of nitrogen. After this step, the residues were dissolved in 50 μ L of methanol, vortex mixed and 2 μ L was injected into the GC–MS system in the SCAN mode.

On the other hand, after the decoctions were prepared and let to cool at room temperature and filtered, 50 mL of each sample was collected into falcon tubes and stored at -20 °C for 12 h and the stored at -80 °C for 24 h. Afterwards, samples were freeze-dried for 5 days.

The analysis of UHPLC-QTOF/MS was performed as follows, addition of 300 μ L of methanol to the 30 mg of the tea plant lyophilized powder, followed by a 2 min agitation in a vortex. The mixture was then sonicated for 15 min and centrifuged at $10,000\times g$ for 5 min at 4 °C. Supernatants were collected and transferred to Chromacol vials (Thermo Fisher Scientific, Madrid, Spain) for LC/MS analysis. The whole procedure was performed in duplicate.

3.2. GC–MS and UHPLC-QTOF/MS Analysis

This analysis was based on the study by Simão et al., [21] and was accomplished using a gas chromatographer (GC) model HP7890B coupled to a mass spectrometer (MS) model 5977A from Agilent Technologies. The separation of the analytes was achieved using a 5% phenylmethylsiloxane (HP-5MS) capillary column (30 m \times 0.25 mm; 0.25 μ m I.D.) from Agilent Technologies. The carrier gas, helium, was set at a constant flow rate of 0.8 mL/min. The injection volume was 2 μ L in the splitless mode. The temperatures of the injector and detector were set at 250 and 280 °C, respectively. Ion source temperature was set at 230 °C and quadrupole at 150 °C. The oven temperature started at 90 °C for 3 min increasing at 15 °C/min up to 300 °C, held for 8 min. The total analysis time was 25 min. The mass spectrometer was operated at 70 eV with a filament current of 300 μ A in the positive electron ionization mode. Data was acquired in the scan mode (m/z 40–550) and the selected ion monitoring (SIM) mode using the ChemStation from Agilent Technologies.

Samples were analyzed using a 1290 Infinity series UHPLC system coupled to a 6545 iFunnel QTOF/MS system (Agilent Technologies, Waldbronn, Germany). For compound separation, 2 μ L of extract was injected in a reversed-phase column (Zorbax Eclipse XDB-C18 4.6 mm \times 50 mm, 1.8 μ m, Agilent Technologies) at 40 °C. The flow rate was 0.5 mL/min, and the mobile phase consisted of 0.1% formic acid in ultrapure water (solvent A) and methanol (solvent B). Gradient elution consisted of 2% B (0–6 min), 2–50% B (6–10 min), 50–95% B (11–18 min), 95% B for 2 min (18–20 min), and returned to the initial conditions from 20–21 min. The total analysis time was 25 min. The detector was operated in full scan mode (m/z 50–1500), at a rate of 1 scan/s. Accurate mass measurement was assured through an automated calibrator delivery system that continuously introduced a reference solution containing masses of m/z 121.0509 (purine) and m/z 922.0098 (HP-921) in the positive ESI mode; and of m/z 112.9856 (TFA) and m/z 922.009798 (HP-921) in the negative ESI mode. The capillary voltage was ± 4000 V for positive and negative ionization modes. The source temperature was 225 °C. The nebulizer and gas flow rates were 35 psig and 11 L/min respectively; the fragmentor voltage was set to 75 V and a radiofrequency voltage in the octopole (OCT RF Vpp) of 750 V was used. Chromatographic analysis was performed using optimized conditions developed by the Center of Metabolomics and Bioanalysis (CEMBIO) [26,33,34].

LC–MS grade solvents were used, and ultrapure water was obtained from a Milli-Q Plus™ System from Millipore (Milford, MA, USA). Formic acid was purchased from Aldrich (St. Louis, MO, USA).

The MassHunter Workstation Software LC/MS Data Acquisition version B.07.00 (Agilent Technologies) was used to control and for data acquisition. The secondary metabolic profile of methanol extracts was analyzed in both ESI negative and positive modes in the range of m/z 50–1500 Da.

3.3. Biological Activities Evaluation

3.3.1. Cell Treatments

The immortalized rat mesencephalic dopaminergic neuron (N27) cell line was maintained in Roswell Park Memorial Institute (RPMI)-1640 culture medium supplemented with 10% fetal bovine serum (FBS), and a mixture of antibiotics (100 U/mL penicillin and 100 µg/mL streptomycin). Subsequently, the cells were incubated at 37 °C with a humidified atmosphere with 5% CO₂.

When cells were approximately 90–95% confluent, they were trypsinized and seeded in 96-well plates (1×10^4 cells/well) and were left to adhere for 12–16 h. Then, after reaching 80–90% confluence in the well plates, the medium was replaced and cells were incubated with certain amounts of the most abundant compound in each plant material, assuming that toxicity is due mainly to its action. The used concentrations were 16 and 80 µM of HML (*P. harmala*), 2 and 10 µM of HMN (*B. caapi* and Dc Ab) and 1 µM of DMT (*P. viridis* and *M. tenuiflora*) for 24 h, in the appropriate medium. The tea mixtures (*P. viridis/B. caapi* and *P. viridis/P. harmala*) were prepared by weighing plant material, assuring that the concentrations of DMT and HMN were 1 and 10 µM respectively for *P. viridis/B. caapi* and 1 and 80 µM of DMT and HML for *P. viridis/P. harmala*. The DMSO vehicle present in each extract was kept below 0.2% to prevent solvent-induced cytotoxicity. Each working solution was prepared in NaCl 0.9%. Each experiment was repeated at least three times, independently.

3.3.2. Cellular Viability Assay

The cytotoxicity of the chosen samples was determined by measuring the levels of cell viability using the cell counting kit-8 (CCK-8) assay (Tebu-bio, Lisboa, Portugal) after 24 h of incubation. This assay was based on the reduction of a slightly yellow water-soluble tetrazolium salt, 2-(2-methoxy-4-nitrophenyl)-3-(4-nitrophenyl)-5-(2,4-disulfophenyl)-2H-tetrazolium and monosodium salt (WST-8), into an orange colored product (WST-formazan), which is soluble in the culture medium. Moreover, the quantity of formazan dye formed was directly proportional to the number of living cells, thus allowing determining the number of viable cells through absorbance measurements.

The cells were seeded in 96-well plates (1×10^4 cells/well), which after reaching confluence were exposed to the tea samples dissolved in RPMI-1640 culture medium. At the end of the incubation time, the cell culture medium was removed from the wells and replaced by a mixture of 5 µL of CCK-8 solution and 95 µL of culture medium and incubated again at 37 °C, light protected, for 3 h. Afterwards, the absorbances were measured using a microplate spectrophotometer (xMark™ Microplate Absorbance Spectrophotometer, BIO-RAD) at 450 nm.

3.3.3. Total Protein Extraction and Quantification

N27 cells were maintained following the same criteria as explained on Section 3.3.1. The cells were then seeded in 6-well plates (2×10^5 cells/well). After 24 h incubation with the highest and lowest concentrations of each tea preparation, the cell medium was removed and cells extracts were collected. In order to obtain cell extracts, cells were scraped and collected in PBS from the culture dish. Afterwards, the resulting suspension was centrifuged at $18,000 \times g$ for 1 min. The resulting pellet was suspended in 20 µL of radioimmunoprecipitation (RIPA) lysis buffer and cells were let to rest on ice for 15 min. Afterwards, cells were centrifuged at $14,000 \times g$ at 4 °C for 20 min; the new resulting supernatant was collected for total protein quantification using the Pierce™ bicinchoninic acid (BCA) protein assay kit.

A calibration curve of bovine serum albumin (BSA) was used to determine the total protein amount in each sample. Therefore, from 100 µL of BSA (2 µg/µL), serial dilutions with water were made (the final concentrations were of 1 µg/µL, 0.5 µg/µL, 0.25 µg/µL, 0.125 µg/µL, 0.0625 µg/µL and 0 µg/µL). A solution of RIPA was diluted 11 times in water on a different Eppendorf tube. All protein samples were also diluted 11 times in water, that is, for 5 µL of protein, 50 µL of water was added into the respective tubes. The estimated volume of BCA reagent need was obtained according to the

number of wells, so knowing that each well uses a total volume of 200 μL , the ratio used was 49 parts of reagent A (mixture of sodium carbonate, sodium bicarbonate, BCA and sodium tartrate in 0.1 M of sodium hydroxide) and 1 part of the B reagent (4% of cupric sulphate).

In a 96-plate, for the standard curve 25 μL of RIPA + 25 μL of BSA standards was added to each well. As for our samples, 25 μL of water + 25 μL of each diluted sample was added to each well. All standards and samples were analyzed in duplicate. Following this, 200 μL of the mix (reagent A + B) was added to each well. Then, the plate was incubated for 30 min at 37 °C and kept in the dark. Subsequently, the absorbance was read at 570 nm.

3.4. Statistical Analysis

Experiments were performed in three sets of independent experiments (n), and four replicates. The results are presented in percentage of controls, according to the following expression:

$$\% \text{ CTR} = (\text{mean values of absorbance after compounds' incubation} / \text{mean values of controls' absorbance}) \times 100$$

Statistical analysis of the biological activities was performed using GraphPad Prism version 5.00 for Windows (GraphPad® Software, San Diego, CA, USA). Statistical significance was determined using a one-way analysis of variance (ANOVA) followed by a Dunnett's or Bonferroni's multiple comparison test and was considered statistically significant when differences between experimental groups had p values < 0.05. Data are shown as mean \pm standard error of mean (SEM). Sigmoid dose-response curves for cell lines were plotted to calculate the concentration, in each incubation period, which kills 50% of cells (LC50): data were fit to a one-site model with a non-linear regression analysis using GraphPad Prism software (version 5; GraphPad Software Inc., La Jolla, CA, USA).

4. Conclusions

The short-term effects of the five studied plants (*P. viridis*, *B. caapi*, *P. harmala*, *M. tenuiflora* and Dc Ab) on the central nervous system remain, to the best of our knowledge, largely unknown. In fact, this paper clarified, for the first time, new findings regarding the phytochemical composition of the decoctions of the previously mentioned plants and of the mixture of *P. viridis* and *B. caapi* (traditional ayahuasca preparation) and *P. viridis* and *P. harmala*, along with their in vitro cytotoxicity in dopaminergic neuronal cells, being dose dependent. These results suggest that the synergetic effect of compounds present in each plant exert neurotoxicity. This is significant since one of the target organs affected by the intake of such substances is the brain. Nonetheless, more studies would be important to assess chronic effects of the compounds and the cellular mechanisms responsible for their cytotoxicity, even the determination of these compounds in biological samples.

Author Contributions: Conceptualization, E.G., A.C.C.; methodology, A.Y.S., J.G.; plant materials, J.R.; sample preparation, N.F.; Chromatographic analysis, A.Y.S., J.G., M.B., E.G., A.P.D; J.M.R.; A.G. (Ana Gradillas) and A.G. (Antonia García); cytotoxicity assays, A.Y.S; A.C.C.; formal analysis, A.C.C., E.G.; resources E.G., A.C.C.; writing—original draft preparation, A.Y.S.; writing—review and editing, A.Y.S, J.G., M.B., J.R., N.F., A.C.C., E.G., A.P.D., A.G. (Ana Gradillas) and A.G. (Antonia García); All authors have read and agreed to the published version of the manuscript.

Funding: This work was partially supported by CICS-UBI, which is financed by National Funds from Fundação para a Ciência e a Tecnologia (FCT) and by Fundo Europeu de Desenvolvimento Regional (FEDER) under the scope of PORTUGAL 2020 and Programa Operacional do Centro (CENTRO 2020), with the project reference UIDB/00709/2020. Ana Y. Simão acknowledges the PhD fellowship from Program Santander-Totta Universidades (Reference: Bolsa BID/UBI-Santander Universidades/2019) Joana Gonçalves acknowledges the PhD fellowship from FCT (Reference: SFRH/BD/149360/2019). Ana Gradillas and Antonia García express their gratitude to the Spanish Ministry of Science, Innovation and Universities MICINN, co-funded by European Regional Development Fund (FEDER) with the project reference RTI2018-095166-B-I00.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Dos Santos, R.G.; Bouso, J.C.; Hallak, J.E.C. Ayahuasca, dimethyltryptamine, and psychosis: A systematic review of human studies. *Ther. Adv. Psychopharmacol.* **2017**, *7*, 141–157. [[CrossRef](#)] [[PubMed](#)]
2. Andrade, T.S.; Oliveira, R.; Da Silva, M.L.; Von Zuben, M.V.; Grisolia, C.; Domingues, I.; Caldas, E.D.; Pic-Taylor, A.; Lopes, M. Exposure to ayahuasca induces developmental and behavioral alterations on early life stages of zebrafish. *Chem. Interact.* **2018**, *293*, 133–140. [[CrossRef](#)] [[PubMed](#)]
3. Gaujac, A.; Dempster, N.; Navickiene, S.; Brandt, S.D.; De Andrade, J.B. Determination of *N,N*-dimethyltryptamine in beverages consumed in religious practices by headspace solid-phase microextraction followed by gas chromatography ion trap mass spectrometry. *Talanta* **2013**, *106*, 394–398. [[CrossRef](#)] [[PubMed](#)]
4. Labate, B.C.; Feeney, K. Ayahuasca and the process of regulation in Brazil and internationally: Implications and challenges. *Int. J. Drug Policy* **2012**, *23*, 154–161. [[CrossRef](#)] [[PubMed](#)]
5. Anderson, B.T.; Labate, B.C.; Meyer, M.; Tupper, K.W.; Barbosa, P.C.; Grob, C.S.; Dawson, A.; McKenna, D. Statement on ayahuasca. *Int. J. Drug Policy* **2012**, *23*, 173–175. [[CrossRef](#)] [[PubMed](#)]
6. Malcolm, B.J.; Lee, K.C. Ayahuasca: An ancient sacrament for treatment of contemporary psychiatric illness? *Ment. Health Clin.* **2017**, *7*, 39–45. [[CrossRef](#)]
7. Pic-Taylor, A.; Da Motta, L.G.; De Moraes, J.A.; Junior, W.M.; Santos, A.D.F.A.; Campos, L.A.; Mortari, M.R.; Von Zuben, M.V.; Caldas, E.D. Behavioural and neurotoxic effects of ayahuasca infusion (*Banisteriopsis caapi* and *Psychotria viridis*) in female Wistar rat. *Behav. Process.* **2015**, *118*, 102–110. [[CrossRef](#)]
8. Callaway, J.C.; Brito, G.S.; Neves, E.S. Phytochemical analyses of *Banisteriopsis caapi* and *Psychotria viridis*. *J. Psychoact. Drugs* **2005**, *37*, 145–150. [[CrossRef](#)]
9. Gable, R.S. Risk assessment of ritual use of oral dimethyltryptamine (DMT) and harmala alkaloids. *Addiction* **2006**, *102*, 24–34. [[CrossRef](#)]
10. Morales-Garcia, J.A.; Revenga, M.D.L.F.; Alonso-Gil, S.; Rodríguez-Franco, M.I.; Feilding, A.; Perez-Castillo, A.; Riba, J. The alkaloids of *Banisteriopsis caapi*, the plant source of the Amazonian hallucinogen Ayahuasca, stimulate adult neurogenesis in vitro. *Sci. Rep.* **2017**, *7*, 1–13. [[CrossRef](#)]
11. Simão, A.Y.; Gonçalves, J.; Duarte, A.P.; Barroso, M.; Cristovão, A.C.; Gallardo, E. Toxicological Aspects and Determination of the Main Components of Ayahuasca: A Critical Review. *Medicines* **2019**, *6*, 106. [[CrossRef](#)] [[PubMed](#)]
12. Cameron, L.P.; Olson, D.E. Dark Classics in Chemical Neuroscience: *N,N*-Dimethyltryptamine (DMT). *ACS Chem. Neurosci.* **2018**, *9*, 2344–2357. [[CrossRef](#)] [[PubMed](#)]
13. Halpern, J.H. Hallucinogens and Dissociative Agents Naturally Growing in the United States. *Pharmacol. Ther.* **2004**, *102*, 131–138. [[CrossRef](#)]
14. Frison, G.; Favretto, D.; Zancanaro, F.; Fazzin, G.; Ferrara, S.D. A case of β -carboline alkaloid intoxication following ingestion of *Peganum harmala* seed extract. *Forensic Sci. Int.* **2008**, *179*, e37–e43. [[CrossRef](#)] [[PubMed](#)]
15. Sklerov, J.; Levine, B.; Moore, K.A.; King, T.; Fowler, D. A Fatal Intoxication Following the Ingestion of 5-Methoxy-*N,N*-Dimethyltryptamine in an Ayahuasca Preparation. *J. Anal. Toxicol.* **2005**, *29*, 838–841. [[CrossRef](#)] [[PubMed](#)]
16. McKenna, D.J.; Towers, G.H.N. Biochemistry and pharmacology of tryptamines and beta-carbolines a minireview. *J. Psychoact. Drugs* **1984**, *16*, 347–358. [[CrossRef](#)] [[PubMed](#)]
17. Strassman, R. *DMT: The Spirit Molecule: A Doctor's Revolutionary Research into the Biology of Near-Death and Mystical Experiences*, 1st ed.; Park Street Press: Rochester, VT, USA, 2001; pp. 33–35.
18. Stahl, S.M. Impulsivity, compulsivity and additions. In *Stahl's Essential Psychopharmacology: Neuroscientific Basis and Practical Applications*, 4th ed.; Stahl, S.M., Ed.; Cambridge University Press: Cambridge, UK, 2013; pp. 889–948.
19. Muñoz, P.; Huenchuguala, S.; Paris, I.; Segura-Aguilar, J. Dopamine Oxidation and Autophagy. *Park. Dis.* **2012**, *2012*, 920953. [[CrossRef](#)] [[PubMed](#)]
20. Goldstein, D.S.; Jinsmaa, Y.; Sullivan, P.; Holmes, C.; Kopin, I.J.; Sharabi, Y. Comparison of Monoamine Oxidase Inhibitors in Decreasing Production of the Autotoxic Dopamine Metabolite 3,4-Dihydroxyphenylacetaldehyde in PC12 Cells. *J. Pharmacol. Exp. Ther.* **2016**, *356*, 484–493. [[CrossRef](#)]

21. Simão, A.Y.; Gonçalves, J.; Caramelo, D.; Rosado, T.; Barroso, M.; Restolho, J.; Fernández, N.; Rodilla, J.; Duarte, A.P.; Cristóvão, A.C.; et al. Determination of *N,N*-dimethyltryptamine and beta-carbolines in plants used to prepare ayahuasca beverages by means of solid-phase extraction and gas-chromatography–mass spectrometry. *SN Appl. Sci.* **2020**, *2*, 1–11. [[CrossRef](#)]
22. Queiroz, M.M.F.; Marti, G.; Queiroz, E.F.; Marcourt, L.; Castro-Gamboa, I.; Bolzani, V.S.; Wolfender, J.-L. LC-MS/MS quantitative determination of tetraptyrys mucronata alkaloids, a plant occasionally used in ayahuasca preparation. *Phytochem. Anal.* **2015**, *26*, 183–188. [[CrossRef](#)]
23. Liu, L.; Zhao, T.; Cheng, X.; Wang, C.; Wang, Z. Characterization and determination of trace alkaloids in seeds extracts from *Peganum harmala* Linn. using LC-ESI-MS and HPLC. *Acta Chromatogr.* **2013**, *25*, 221–240. [[CrossRef](#)]
24. Coutts, R.T.; Locock, R.A.; Slywka, G.W.A. Mass spectra of selected beta-carbolines [β -9H-pyrido(3,4-b)indoles]. *Org. Mass Spectrom.* **1970**, *3*, 879–889. [[CrossRef](#)]
25. Estrella-Parra, E.A.; Almanza-Pérez, J.C.; Alarcón-Aguilar, F.J. Ayahuasca: Uses, Phytochemical and Biological Activities. *Nat. Prod. Bioprospecting* **2019**, *9*, 251–265. [[CrossRef](#)] [[PubMed](#)]
26. Gonçalves, J.; Luís, Â.; Gradillas, A.; García, A.; Restolho, J.; Fernández, N.; Domingues, F.; Gallardo, E.; Duarte, A.P. Ayahuasca beverages: Phytochemical analysis and biological properties. *Antibiotics* **2020**, *9*, 731. [[CrossRef](#)] [[PubMed](#)]
27. Zhang, C.; Deng, Y.; Dai, H.; Zhou, W.; Tian, J.; Bing, G.; Zhao, L. Effects of dimethyl sulfoxide on the morphology and viability of primary cultured neurons and astrocytes. *Brain Res. Bull.* **2017**, *128*, 34–39. [[CrossRef](#)] [[PubMed](#)]
28. Fisher, R.; Lincoln, L.; Jackson, M.J.; Abbate, V.; Jenner, P.; Hider, R.; Lees, A.; Rose, S. The effect of *Banisteriopsis caapi* (*B. caapi*) on the motor deficits in the MPTP-treated common marmoset model of Parkinson's disease. *Phytother. Res.* **2018**, *32*, 678–687. [[CrossRef](#)]
29. Butler, M.; Spearman, M.; Braasch, K. Monitoring cell growth, viability, and apoptosis. *Methods Mol. Biol.* **2014**, *1104*, 169–192. [[CrossRef](#)]
30. Zhang, S.; Wang, R.; Wang, G. Impact of dopamine oxidation on dopaminergic neurodegeneration. *ACS Chem. Neurosci.* **2019**, *10*, 945–953. [[CrossRef](#)]
31. Protein Degradation—The Cell—NCBI Bookshelf. Available online: <https://www.ncbi.nlm.nih.gov/books/NBK9957/> (accessed on 13 October 2020).
32. Valente, M.J.; Amaral, C.; Correia-Da-Silva, G.; Duarte, J.A.; Bastos, M.D.L.; Carvalho, F.; De Pinho, P.G.; Carvalho, M. Methylone and MDPV activate autophagy in human dopaminergic SH-SY5Y cells: A new insight into the context of β -keto amphetamines-related neurotoxicity. *Arch. Toxicol.* **2017**, *91*, 3663–3676. [[CrossRef](#)]
33. Acero, N.; Gradillas, A.; Beltran, M.; Garcia, A.; Mingarro, D.M. Comparison of phenolic compounds profile and antioxidant properties of different sweet cherry (*Prunus avium* L.) varieties. *Food Chem.* **2019**, *279*, 260–271. [[CrossRef](#)]
34. Gutiérrez-Albanchez, E.; Gradillas, A.; García, A.; García-Villaraco, A.; Gutierrez-Mañero, F.J.; Solano, B.R. Elicitation with *Bacillus* QV15 reveals a pivotal role of F3H on flavonoid metabolism improving adaptation to biotic stress in blackberry. *PLoS ONE* **2020**, *15*, e0232626. [[CrossRef](#)]

Sample Availability: Not available.

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).