


Drug Discovery | *Hot Paper*

Total Synthesis of a Partial Structure from Arabinogalactan and Its Application for Allergy Prevention

 Matthias Krumb,^[b] Maximilian Jäger,^[a] Alice Voss,^[c] Loreen Immig,^[a] Karin Peters,^[a] Danuta Kowalczyk,^[b] Albrecht Bufe,^[a] Till Opatz,^{*,[b]} Otto Holst,^[d] Christian Vogel,^[c] and Marcus Peters^{*,[a, e]}

Abstract: Arabinogalactan, a microheterogeneous polysaccharide occurring in plants, is known for its allergy-protective activity, which could potentially be used for preventive allergy treatment. New treatment options are highly desirable, especially in a preventive manner, due to the constant rise of atopic diseases worldwide. The structural origin of the allergy-protective activity of arabinogalactan is, however, still unclear and isolation of the polysaccharide is not feasible for pharmaceutical applications due to a variation of the activity of the natural product and contaminations with endotoxins. Therefore, a pentasaccharide partial structure was selected for total synthesis and subsequently coupled to a carrier protein to form a neoglycoconjugate. The allergy-protective activity of arabinogalactan could be reproduced with the partial structure in subsequent in vivo experiments. This is the first example of a successful simplification of arabinogalactan with a single partial structure while retaining its allergy-preventive potential.

The negative impact on human health of allergic diseases, often overrepresented in first world countries, has been increasing unimpeded. However, studies have shown that growing up and living in a traditional farming environment can decrease the risk of developing these allergies significantly, which is known as the “farm effect” in literature.^[1–4] Nevertheless, the connection between living in a farming environment

and allergy protection is not yet fully understood.^[5] Arabinogalactan (AG), an immune-modulating natural polysaccharide, has recently received considerable attention as one potential origin for this phenomenon.^[5,6]

The polysaccharide occurs frequently in grasses that are present in a farming environment and can be isolated therefrom.^[6,7] Unfortunately, the composition of the structurally microheterogeneous AG is very complex and difficult to analyze.^[8,9] There is literature evidence that the allergy-protective activity of AG could originate from recognition of particular (terminal) arabinose containing epitopes by the immune system, since it is known that a cleavage of these significantly reduced the allergy-protective activity of the polysaccharide.^[6] Moreover, it was shown in the same work that AG from other plants, namely *Acacia* or *Larix*, does not show allergy-protective activity. This is probably based on a differing molecular structure, however, the exact chemical structure of the different AG is unknown. Fortunately, pioneering work by Brecker et al. shed light on the structural composition of AG isolated from Timothy grass (*Phleum pratensis*) by deciphering arabinose and galactose containing partial structures (Figure 1 A).^[7] The latter consist of a β -(1 \rightarrow 6)-linked D-galactopyranosyl backbone with L-arabinofuranosyl- α -(1 \rightarrow 5)- or α -(1 \rightarrow 3)- and β -(1 \rightarrow 3)-linked side chains. It was revealed in previous studies that isolated AG from meadow foxtail (*Alopecurus pratensis*), which contains these partial structures, is allergy-protective.^[6]

Preventive and prophylactic approaches with a promising perspective are still rare, despite a huge research effort in the search for new treatment options for allergic diseases.^[11] While

[a] Dr. M. Jäger, L. Immig, Dr. K. Peters, Prof. Dr. A. Bufe, Dr. M. Peters
 Department of Experimental Pneumology
 Ruhr-University Bochum
 Universitätsstr. 150, 44801 Bochum (Germany)
 E-mail: marcus.peters@rub.de



[b] M. Krumb, D. Kowalczyk, Prof. Dr. T. Opatz
 Department of Chemistry
 Johannes Gutenberg-University
 Duesbergweg 10–14, 55128 Mainz (Germany)
 E-mail: opatz@uni-mainz.de


[c] A. Voss, Prof. Dr. C. Vogel
 Department of Chemistry
 University of Rostock
 Albert-Einstein-Str. 3a, 18051 Rostock (Germany)

[d] Prof. Dr. O. Holst
 Division of Structural Biochemistry
 Research Center Borstel

Leibniz Lung Center
 Airway Research Center North (ARCN)
 Parkallee 1–40, 23845 Borstel (Germany)

[e] Dr. M. Peters
 Department of Molecular Immunology
 Ruhr-University Bochum
 Universitätsstr. 150, 44801 Bochum (Germany)

 Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under:
 <https://doi.org/10.1002/chem.202002287>.

 © 2020 The Authors. Published by Wiley-VCH GmbH. This is an open access article under the terms of Creative Commons Attribution NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.





Figure 1. Illustration of partial structures of AG identified by Brecker and co-workers using symbol nomenclature for glycans (A).^[7,10] Partial structure of arabinogalactan selected for total synthesis and its retrosynthetic analysis (B).

the use of natural polysaccharides for prophylaxis could be envisioned, the size and complexity of AG precludes the production of chemically well-defined species with consistent biological activity. In addition, the separation of natural polysaccharides from other immune-modulating contaminants, such as endotoxins, is, at least, highly challenging, especially under the good manufacturing practice conditions required for pharmaceutical applications. Gratifyingly, the prime example of fondaparinux, a synthetic pentasaccharide used as an anticoagulant, showcases that the total synthesis of carbohydrate pharmaceuticals can be cost-effective, despite their intrinsic high complexity and polyfunctionality.^[12–15] In terms of AG, there are a few examples for the synthesis of related oligosaccharides but structure–activity relationships are unfortunately missing.^[16–21]

Thus, we selected a partial structure of arabinogalactan, namely a pentasaccharide containing two L-arabinose and three D-galactose units (Araf₂Galp₃) as the target for total synthesis to match one of the partial structures in AG (Figure 1A (structure II) and Figure 1B)^[7] which were shown to be biologically active in earlier studies.^[6] Our overall goal for this project was to gain a better understanding of the role of glyco-epitopes in AG and to achieve a strategic simplification of the polysaccharide while the useful allergy-protective activity is retained.


The pentasaccharide selected requires construction of a trigalactose (Galp₃) backbone with two β-(1→6)-glycosylations and an additional β-(1→3)-glycosylation for the attachment of the arabinose side chain. The latter required the construction of an α-(1→5)-glycosidic linkage. Additionally, a reactive isothiocyanate spacer was needed for covalent coupling to a carrier protein. The synthesis of the pentasaccharide Araf₂Galp₃ started with the construction of the trigalactose (Galp₃) backbone; the (1→6)-connections desired were furnished using a



Scheme 1. Synthesis of Galp₃ intermediate **9** over nine linear steps starting from peracetylated D-galactose (**1**).

combination of regioselective enzymatic acetylation and chemoselective deacetylation in the presence of benzoyl groups (Scheme 1). An orthogonally stable Fmoc-protecting group was introduced to enable glycosylation of the Araf₂-disaccharide later in the synthesis. In detail, the synthetic route for Galp₃ (**9**) started from peracetylated D-galactose (**1**) to give the 6-O-acetyl-protected thioglycoside **2** through a reaction with thiophenol, Zemplén deprotection and regioselective acetylation using a lipase to selectively furnish the 6-O-protected galactose **2** in an 83% yield over three steps.^[22] Subsequently, galactose **2** was benzoylated and glycosylated with spacer **4**, which was synthesized in six steps from 5-phenylvaleric acid (**3**, details can be found in the Supporting Information). The 6-O-acetyl group was then removed using methanolic HCl to give intermediate **5**. The free hydroxyl group obtained in **5** was then glycosylated with an orthogonally protected galactose **6**. The latter was synthesized from galactose **2** through orthoester formation with triethyl orthobenzoate, benzoylation, regioselective cleavage of the orthoester intermediate **2a** and, subsequent Fmoc-protection in a 40% yield over three steps.^[23] Both building blocks, galactose **5** and galactose **6**, were glycosylated, 6-O-deprotected and again glycosylated with thioglycoside **8** to synthesize a Galp₃ building block **9** in four further steps.

The synthesis of the Araf₂ side chain **14** required a (1→5) linkage between the L-arabinose units, which was furnished using the established tritylation-detritylation approach (Scheme 2).^[24] The *cis*-glycosidic linkage in the first galactose-connected arabinose required the use of non-participating *O*-protection, while the latter arabinose has a *trans*-glycosidic linkage and, therefore, requires participating *O*-protection for synthesis.^[25] According to a report by Lowary and co-workers, the *cis*-glycosylation is possible when a conformationally restricted *O*-xylylene-protected donor is used.^[26,27] We adapted




Scheme 2. Synthesis of Araf₂ intermediate 13 over eight linear steps starting from L-arabinose (9).

this approach and started the synthesis from L-arabinose (10) in a Fischer glycosylation in methanol, the product of which was subsequently benzoylated. Thioarabinoside 11 was subsequently prepared from a 1-O-acetyl-intermediate in four steps and a 26% yield. Next, the 5-O-deprotected arabinose 12 was produced in a sequence of debenzoylation, 5-O-tritylation, introduction of the 2,3-O-xylylene ether, according to Lowary et al., and detritylation to give glycosyl acceptor 12 in a 31% yield over four steps. The thioglycoside in arabinose 12 should be used afterwards for glycosylation to trigalactose 9. Thus, an orthogonal glycosylation, which leaves the thioglycoside untouched, was required to introduce the second arabinose unit. Gratifyingly, using the *N*-phenyltrifluoroacetimidate donor 13,^[28] synthesized through cleavage of thioglycoside 11 to its hemi-acetal and reaction with 2,2,2-trifluoro-*N*-phenylacetimidoyl chloride, furnished the Araf₂ intermediate 14 desired

in a 62% yield in a modified procedure based on Andersen et al.^[29]

The final steps of the synthesis started with the glycosylation of Araf₂ donor 14 with Galp₃ acceptor 9 (Scheme 3). After a successful *cis*-glycosylation and Zemplén deprotection, protected Araf₂Galp₃ 15 was synthesized in a 27% yield. In the two final steps, the 2,3-O-xylylene protecting group was removed under Birch conditions using liquid ammonia and sodium. This also led to the reduction desired of the azide to an amino group at the same time. Conversion of the latter to the reactive isothiocyanate with thiophosgene under mildly basic conditions using BaCO₃ as a mild base in a water/ethanol mixture gave the target structure Araf₂Galp₃ 16 in a 71% yield over two steps.

The pentasaccharide was coupled to bovine serum albumin (BSA) as the carrier protein to create a neoglycoconjugate (Araf₂Galp₃-BSA, Figure S1) for biological evaluation. Coupling was accomplished using the reactive isothiocyanate group, which reacts readily with BSA's free amino groups. The neoglycoconjugate created was subsequently evaluated for allergy-protective activity in a mouse model of allergic asthma (Figure 2). The treatment of mice with Araf₂Galp₃-BSA led to reduced allergic inflammation in the airways upon sensitization and challenge with ovalbumin (OVA) as a model allergen. This can be seen by the reduction of eosinophilic granulocytes, lymphocytes and macrophages (Figure 2A). Moreover, immunoglobulin E (IgE) production (Figure 2B) and the T_H2 response determined by measuring the *in vitro* production of interleukin (IL)-5 and IL-13 by spleen cells is significantly reduced (Figure 2C), as well as the goblet cell metaplasia (Figure 2D and Figure S2). The efficacy of treatment was comparable to AG, shown to be allergy-protective in former studies.^[6] Another hallmark of allergic asthma is airway hyperreactivity (AHR) towards provocation



Scheme 3. Synthesis of the AG partial structure Araf₂Galp₃ 15 over four linear steps from the key intermediates Galp₃ 8 and Araf₂ 13.





Figure 2. Araf₂Galp₃-BSA reduces the allergic airway inflammation and sensitization. (A). immunoglobulin E (IgE) was measured in BAL fluid (B). Production of IL-5 and IL-13 was determined in spleen cell supernatants after the restimulation of cells with OVA (C). Goblet cells were counted from five different bronchi of each mouse and expressed as the number of cells per 100 μm basement membrane (D). One-way ANOVA was used to analyse the data for statistical differences. **p* < 0.05 and ***p* < 0.01.

with cholinergic stimuli, such as methacholine. Therefore, the AHR of mice was tested in the asthma model by measuring the airway mechanics. As seen for airway inflammation, the AHR is also significantly reduced in mice treated with Araf₂Galp₃-BSA (Figure S4).

Next, a neutral control structure was needed to clarify that the biological activity of pentasaccharide **16** observed results from the actual recognition of the structure by the immune system and not from unspecific glycosylation. Hence, we synthesized a glucose pentamer (Glc_{p5}) equal in size and presence of multivalent carbohydrate ligands to pentasaccharide **12** to form a robust control structure (Scheme 4). D-glucose (**17**) was converted to intermediate **18** over five steps in a 21% yield using standard operations. Glycosylation with thioglycoside **19** and deprotection gave disaccharide **20** over two steps. This process was repeated to obtain the pentaglycose **21** over five steps in 46% in an iterative manner (details can be found in the Supporting Information). After global deprotection, reduction and conversion of the amine to the isothiocyanate using the established procedure, pentaglycose **22** was synthesized in a 39% yield. It was subsequently coupled to BSA to give the neoglycoconjugate Glc_{p5}-BSA (Figure S1). No significant protection against allergic airway disease was found using Glc_{p5}-BSA as a neutral control structure (Figure S3). These results underline that the allergy-protective activity depends on the specific composition of the pentasaccharide and is not due to the unspecific glycosylation of BSA.

Following the initial observation of the allergy-protective effect of Araf₂Galp₃-BSA, the mechanism of protection was investigated. During the treatment of mice with Araf₂Galp₃-BSA, a significant increase of BSA-specific immunoglobulin G1 (IgG1) in serum was observed, which was absent when mice were treated with unmodified BSA. By contrast, treatment with Glc_{p5}-BSA did not lead to a significantly increased antibody production (Figure 3A). Interestingly, the titre of BSA-specific IgG1 shows a strong negative correlation with the number of eosinophilic granulocytes and OVA-specific IgE, suggesting a causative relationship between these parameters (Figure 3B and C). The galactosylation of the IgG produced was analysed since it is known that the modification of the Fc-part of IgG with galactose confers immune-regulatory properties.^[30–32] It was found that the treatment of mice with Araf₂Galp₃-BSA causes a significant increase of galactosylated IgG1 (Figure 3D). The IgG produced was, therefore, suspected of being involved in the regulation of airway inflammation. However, an interference with the induction of eosinophilic airway inflammation by adoptive transfer of IgG1 was not observed (details can be found in the Supporting Information, Figure S5), which hints at a more complex mechanism. In addition to antibodies, B-lymphocytes produce cytokines that may affect immune regulation. The so-called regulatory B cells are releasing IL-10 and provide several other mechanisms of down-regulation of inflammatory immune responses.^[33,34] Thus, IL-10 production of B cells from the lung tissue of mice treated with Araf₂Galp₃-BSA was confirmed (Figure S6). The activity of IL-10 was blocked by intranasal administration of a blocking antibody binding to the IL-10 receptor to analyse whether the IL-10 produced is involved in the regulation of the allergic immune response. As expected, mice that were treated with Araf₂Galp₃-BSA and an isotype control antibody showed a significantly reduced allergic inflammation (Figure 3E). By contrast, the allergic airway inflammation and sensitization in mice that received



Scheme 4. Synthesis of the glucose pentamer (**22**, Glcp₃).

Araf₂Galp₃-BSA and an IL-10R blocking antibody showed no significant difference from the phenotype observed in mice that were not treated with the glycoconjugate. This shows that IL-10 signaling is probably involved in the mechanism of immune modulation mediated by Araf₂Galp₃-BSA (discussion can be found in the Supporting Information).

In summary, a convergent total synthesis of a partial structure of AG was accomplished by synthesis of a trigalactose backbone and an arabinose disaccharide sidechain and their subsequent combination. Late-stage introduction of a reactive isothiocyanate and covalent coupling to BSA led to a neoglycoconjugate. It was shown for the first time that this construct can reduce the allergic airway inflammation in a mouse model of allergic asthma in comparable potency as natural AG. The allergy-protective activity was specific for the composition of the partial structure and does not result from unspecific glycosylation of BSA, as shown by the synthesis and investigation of the pentaglyucose control structure. Additionally, the mode of action of the synthesized glyco-epitope was investigated, where induction of B cells that produce galactosylated antibodies and IL-10 was identified, which was found to regulate the allergic response. Gratifyingly, we did not observe any negative side effects accompanying the treatment with Araf₂Galp₃-BSA and, therefore, believe strongly that this compound is an interesting candidate for a pharmaceutical approach to prevent allergic sensitization in future work.

Acknowledgements

This study was supported by Deutsche Forschungsgemeinschaft (INST 213/950-1 FUGG, PE 1813/2-1), collaborative research centre 1066 of Johannes Gutenberg-University, intramural research funding of Ruhr-University Bochum F659-2009 and F797-2013 and a grant provided by the Gesellschaft für Pädiatrische Allergologie. We thank the Verein zur Förderung der Glycowissenschaften e.V., Hamburg, Germany, for financial support. We thank Dr. Johannes C. Liermann (Mainz) for NMR Spectroscopy, Dr. Christopher Kampf (Mainz) for mass spectrometry, and Petra Bonowitz (Bochum) and Sandra Busse (Bochum) at Experimental Pneumology, Ruhr-University Bochum, for excellent technical assistance. We like to thank Mr. Philip Saunders (Language Support Services, Berlin) for proof-reading our manuscript. Open access funding enabled and organized by Projekt DEAL.

Conflict of interest

MP, AV, OH and CV are inventors in a patent related to this study (WO2017093276A1). The other authors declared that they have no conflict of interest.

Keywords: airway inflammation · allergy protection · arabinogalactan · carbohydrates · total synthesis




Figure 3. $\text{Araf}_2\text{Galp}_3\text{-BSA}$ -induced production of specific IgG1. The BSA-specific IgG1 antibodies were detected by ELISA (A). Antibody titre was correlated to the number of eosinophilic granulocytes (B) and the concentration of IgE (C). Galactose modification of IgG was measured in a lectin-binding test by using the galactose-specific *Erythrina crista-galli* lectin (D). The immunoregulatory activity of IL-10 is also involved in $\text{Araf}_2\text{Galp}_3\text{-BSA}$ -mediated allergy protection. This was shown by blocking the IL-10 receptor by using an antibody, leading to the inhibition of the anti-inflammatory action of $\text{Araf}_2\text{Galp}_3\text{-BSA}$ (E). One-way ANOVA was used to analyse the data for statistical differences. Correlation analysis was done by Spearman's rank correlation. * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$.

- [1] R. P. Lauener, T. Birchler, J. Adamski, C. Braun-Fahrlander, A. Bufe, U. Herz, E. von Mutius, D. Nowak, J. Riedler, M. Waser, F. H. Sennhauser, *Lancet* **2002**, *360*, 465.
- [2] M. J. Ege, I. Herzum, G. Büchele, S. Krauss-Etschmann, R. P. Lauener, M. Roponen, A. Hyvärinen, D. A. Vuitton, J. Riedler, B. Brunekreef, J.-C. Dalphin, C. Braun-Fahrlander, J. Pekkanen, H. Renz, E. von Mutius, *J. Allergy Clin. Immunol.* **2008**, *122*, 407.
- [3] O. S. Von Ehrenstein, E. Von Mutius, S. Illi, L. Baumann, O. Böhm, R. Von Kries, *Clin. Exp. Allergy* **2000**, *30*, 187.
- [4] J. Riedler, C. Braun-Fahrlander, W. Eder, M. Schreuer, M. Waser, S. Maisch, D. Carr, R. Schierl, D. Nowak, E. Von Mutius, *Lancet* **2001**, *358*, 1129.
- [5] L. Gorelik, M. Kauth, K. Gehlhar, A. Bufe, O. Holst, M. Peters, *Innate Immun.* **2008**, *14*, 345.
- [6] M. Peters, M. Kauth, O. Scherner, K. Gehlhar, I. Steffen, P. Wentker, E. von Mutius, O. Holst, A. Bufe, *J. Allergy Clin. Immunol.* **2010**, *126*, 648.
- [7] L. Breckler, D. Wicklein, H. Moll, E. C. Fuchs, W.-M. Becker, A. Petersen, *Carbohydr. Res.* **2005**, *340*, 657.
- [8] G. B. Fincher, B. A. Stone, A. E. Clarke, *Annu. Rev. Plant Physiol.* **1983**, *34*, 47.
- [9] W. Qi, C. Fong, D. T. A. Lamport, *Plant Physiol.* **1991**, *96*, 848.
- [10] K. Cheng, Y. Zhou, S. Neelamegham, *Glycobiology* **2017**, *27*, 200.
- [11] A. Nieto, U. Wahn, A. Bufe, P. Eigenmann, S. Halken, G. Hedlin, A. Høst, J. Hourihane, J. Just, G. Lack, *Pediatr. Allergy Immunol.* **2014**, *25*, 516.
- [12] C.-H. Chang, L. S. Lico, T.-Y. Huang, S.-Y. Lin, C.-L. Chang, S. D. Arco, S.-C. Hung, *Angew. Chem. Int. Ed.* **2014**, *53*, 9876; *Angew. Chem.* **2014**, *126*, 10034.
- [13] Y. Ding, C. V. N. S. Vara Prasad, H. Bai, B. Wang, *Bioorg. Med. Chem. Lett.* **2017**, *27*, 2424.
- [14] T. Li, H. Ye, X. Cao, J. Wang, Y. Liu, L. Zhou, Q. Liu, W. Wang, J. Shen, W. Zhao, P. Wang, *ChemMedChem* **2014**, *9*, 1071.
- [15] F. Lin, G. Lian, Y. Zhou, *Carbohydr. Res.* **2013**, *371*, 32.
- [16] M. P. Bartetzko, F. Schuhmacher, H. S. Hahm, P. H. Seeberger, F. Pfeingler, *Org. Lett.* **2015**, *17*, 4344.
- [17] F. W. D'Souza, T. L. Lowary, *Org. Lett.* **2000**, *2*, 1493.
- [18] A. Fekete, A. Borbás, S. Antus, A. Lipták, *Carbohydr. Res.* **2009**, *344*, 1434.
- [19] H. Huang, L. Han, Y.-M. Lan, L.-L. Zhang, *J. Asian Nat. Prod. Res.* **2014**, *16*, 640.
- [20] K. S. Kim, B.-Y. Lee, S. H. Yoon, H. J. Jeon, J. Y. Baek, K.-S. Jeong, *Org. Lett.* **2008**, *10*, 2373.
- [21] Y. Wu, D.-C. Xiong, S.-C. Chen, Y.-S. Wang, X.-S. Ye, *Nat. Commun.* **2017**, *8*, 14851.
- [22] X. Zhang, T. Kamiya, N. Otsubo, H. Ishida, M. Kiso, *J. Carbohydr. Chem.* **1999**, *18*, 225.
- [23] R. Lemieux, H. Driguez, *J. Am. Chem. Soc.* **1975**, *97*, 4069.
- [24] B. Helferich in *Advances in Carbohydrate Chemistry, Vol. 3* (Eds.: W. W. Pigman, M. L. Wolfrom, S. Peat), Academic Press, Cambridge, MA, **1948**, pp. 79–111.
- [25] D. R. Mootoo, P. Konradsson, U. Udodong, B. Fraser-Reid, *J. Am. Chem. Soc.* **1988**, *110*, 5583.
- [26] L. Zhang, K. Shen, H. A. Taha, T. L. Lowary, *J. Org. Chem.* **2018**, *83*, 7659.
- [27] A. Imamura, T. L. Lowary, *Org. Lett.* **2010**, *12*, 3686.
- [28] B. Yu, H. Tao, *Tetrahedron Lett.* **2001**, *42*, 2405.
- [29] M. C. Andersen, S. K. Kračun, M. G. Rydahl, W. G. Willats, M. H. Clausen, *Chem. Eur. J.* **2016**, *22*, 11543.
- [30] R. M. Anthony, F. Nimmerjahn, D. J. Ashline, V. N. Reinhold, J. C. Paulson, J. V. Ravetch, *Science* **2008**, *320*, 373.
- [31] N. Washburn, I. Schwab, D. Ortiz, N. Bhatnagar, J. C. Lansing, A. Me-deiros, S. Tyler, D. Mekala, E. Cochran, H. Sarvaiya, *Proc. Natl. Acad. Sci. USA* **2015**, *112*, E1297.
- [32] G. Dekkers, T. Rispens, G. Vidarsson, *Front. Immunol.* **2018**, *9*, 553.
- [33] B. Peng, Y. Ming, C. Yang, *Cell Death Dis.* **2018**, *9*, 109.
- [34] T. F. Tedder, *J. Immunol.* **2015**, *194*, 1395.

Manuscript received: May 8, 2020

Accepted manuscript online: June 24, 2020

Version of record online: October 27, 2020