



BMJ Open ED to EPI: protocol for a pragmatic randomised controlled trial of an SMS (text) messaging intervention to improve the transition from the emergency department to early psychosis intervention for young people with psychosis

Alexia Polillo,^{1,2} George Foussias,^{1,2} Albert H C Wong,^{2,3} Augustina Ampofo,⁴ Vicky Stergiopoulos,^{2,4} Kelly K Anderson,⁵ Sarah Bromley,¹ Jessica D'Arcey,^{1,6} Claire de Oliveira,^{7,8} Lillian Duda,⁴ Joanna Henderson ,^{2,9} Sean Kidd,^{2,4} Paul Kurdyak,^{2,7} Wei Wang,^{2,4} Juveria Zaheer,^{2,7} Aristotle N Voineskos,^{1,2} Nicole Kozloff ^{1,2}

To cite: Polillo A, Foussias G, Wong AHC, *et al.* ED to EPI: protocol for a pragmatic randomised controlled trial of an SMS (text) messaging intervention to improve the transition from the emergency department to early psychosis intervention for young people with psychosis. *BMJ Open* 2020;**10**:e042751. doi:10.1136/bmjopen-2020-042751

► Prepublication history and supplemental material for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-042751>).

Received 15 July 2020

Revised 19 November 2020

Accepted 25 November 2020



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Nicole Kozloff;
n.kozloff@mail.utoronto.ca

ABSTRACT

Introduction While nearly half of all new psychotic disorders are diagnosed in the emergency department (ED), most young people who present to the ED with psychosis do not receive timely follow-up with a psychiatrist, and even fewer with evidence-based early psychosis intervention (EPI) services. We aim to test an intervention delivered using short message service (SMS), a low-cost, low-complexity, youth-friendly approach, to improve transitions from the ED to EPI services.

Methods and analysis This is a protocol for a pragmatic randomised, single blind, controlled trial with accompanying economic and qualitative evaluations conducted at the Centre for Addiction and Mental Health (CAMH) in Toronto, Canada. A consecutive series of 186 participants aged 16–29 referred by the ED to CAMH's EPI programme will be recruited for a trial of a two-way intervention involving reminders, psychoeducation and check-ins delivered via SMS. The primary outcome will be attendance at the first consultation appointment within 30 days of study enrolment assessed through chart reviews in the electronic health record. We will also extract routine clinical measures, including the Brief Psychiatric Rating Scale, Clinical Global Impression and Service Engagement Scale, and link with provincial health administrative data to examine system-level outcomes, including ED visits and psychiatric hospitalisations, 6 months and up to 2 years after baseline. We will perform a cost-effectiveness analysis of the primary study outcome and costs incurred, calculating an incremental cost effectiveness ratio. Web-based surveys and qualitative interviews will explore intervention user experience. Patients and families with lived experience will be engaged in all aspects of the project.

Ethics and dissemination Research Ethics Board approval has been obtained. Findings will be reported in scientific journal articles and shared with key stakeholders including youth, family members, knowledge users and decision makers.

Strengths and limitations of this study

- Pragmatic randomised controlled trial leveraging mobile health technology, chart reviews, routinely collected administrative data, and economic and qualitative evaluations.
- Intervention well-positioned for local adoption as well as scale and spread to other early psychosis intervention programmes and youth mental health services more broadly.
- Collaboration with health system decision-makers, clinical stakeholders, knowledge users, team members with clinical and research expertise, people with lived experience of psychosis, and key relationships with organisations well-positioned to support widespread implementation.
- Conducting the trial at a single site will support streamlined recruitment but may limit generalisability of our findings.
- The pragmatic data collection methods being used, particularly chart review, may be subject to unreliable extraction and missing data.

Trial registration number NCT04298450.

INTRODUCTION

Psychosis, characterised by delusions and/or hallucinations, typically manifests during adolescence or early adulthood. It is the characteristic presentation of schizophrenia and schizoaffective disorder, and often occurs in bipolar disorder.¹ These disorders can cause significant dysfunction: in disability weighting surveys used to establish global disease burden, participants rated schizophrenia as



the disorder most disabling for individuals.² Young people experience greater mortality by up to 24-fold in the year following a first psychotic disorder diagnosis compared with peers in the general population.³ In the long term, psychotic disorders are associated with ongoing increased mortality particularly by suicide,⁴ substance use disorders, homelessness, victimisation, acts of violence⁵ and high economic costs due to healthcare use as well as lost productivity.^{6,7} Early psychosis intervention (EPI) is a model developed to provide treatment early in the course of illness to improve patients' long-term trajectories and reduce the burden on individuals and their families. The rationale for EPI has been strengthened by consistent findings that long duration of untreated psychosis is associated with greater symptom burden, lower likelihood of remission and poor social functioning and global outcomes.⁸ Members of our team have shown that EPI service use is associated with a fourfold reduction in all-cause mortality for young people with psychosis compared with those who do not access EPI services.⁹ EPI services have also been associated with improved access to psychiatric care, reduced risk of relapse, fewer hospital readmissions and increased employment rates.^{9–13} EPI can be a lifechanging and lifesaving intervention for young people with psychosis.

EPI programmes are well-established in Ontario and provide services to young people with early psychosis across the province.¹⁴ Despite this, and the clear mandate for EPI programmes to promote their services and minimise barriers to care, many youth with psychosis in Ontario either never access these services, or enter them far later than indicated.¹⁵ In Canada, nearly half of all new psychotic disorders are diagnosed in the emergency department (ED).¹⁶ We recently found that among young people across Ontario presenting with psychotic disorders to the ED for the first time, 40% received no outpatient mental health follow-up within 30 days and only 45% saw a psychiatrist.¹⁷ The reasons young people with psychosis discharged from the ED did not receive follow-up were unclear in this study: both issues of access (availability and awareness of services) and engagement (youth following through on referrals) are potential explanations. Our experience at the Centre for Addiction and Mental Health (CAMH) in Toronto, where our EPI programme sees referred patients for consultation within 2 weeks on average, suggests a problem of engagement: according to clinic data, 50% of the youth referred from the ED do not attend their first EPI consultation appointment (compared with approximately 30% from all other referral sources). It is clear that new approaches are required to engage this population in accessing evidence-based care that is life-saving and improves illness outcomes. In surveys of patients and families in EPI services, appointment reminders are cited as a top factor that would improve service engagement, with a preference for email and text communication.^{18–20}

Mobile health technologies are increasingly being tested to improve outcomes, including symptoms,

appointment attendance and medication adherence, among young people with mental illness, particularly psychosis.^{21–23} Short message service (SMS) or text message is a commonly used mode of communication by adolescents and young adults: in a survey of users of community mental health services, access to mobile phones approached 100%.²⁴ SMS is associated with low user and financial burden. SMS does not require people to own a smartphone, have data plans, or have access to wireless internet. In a study of people with psychosis, participants were found to be highly engaged with an SMS intervention.²⁵ SMS reminders have been associated with improved service engagement in psychosis across studies,²² including twice the attendance rates for initial appointments in an EPI programme.²⁶ An ongoing pilot study at CAMH investigating the effect of a weekly two-way SMS intervention on service engagement during the first year of EPI treatment found this approach to be feasible and valued by participants.²⁷ We are unaware of any studies examining interventions specifically to improve the transition in care from the ED to EPI for young people with psychosis, using mobile health technologies or otherwise.

Objectives and hypotheses

'ED to EPI', a pragmatic randomised, single blind, controlled trial²⁸ with accompanying economic and qualitative evaluations, aims to improve the transition from the ED to EPI services for youth with psychosis using an SMS text messaging intervention. It is pragmatic in its participant eligibility criteria (broad and inclusive), comparison intervention (usual care), follow-up intensity (low), primary trial outcome (objective, meaningful and assessed under usual conditions), measurement of participant compliance and practitioner adherence to study protocol (unobtrusive) and analysis of primary outcome (inclusive, ie, intention-to-treat).²⁹ We also leverage linked routinely collected data through ICES (previously known as the Institute for Clinical Evaluative Sciences), which holds data on all hospital and physician visits for the province. We will also evaluate the cost effectiveness of the intervention and explore young people's perspectives on its various components. Our study team, in addition to clinicians and researchers, includes a patient and family member with lived experience of using EPI services and key decision-makers to increase the relevance and uptake of the intervention. Specifically, our primary objectives are to:

1. Evaluate the effect of an SMS intervention on attendance at the first consultation appointment within 30 days of study enrolment.
Hypothesis 1: The SMS intervention will increase rate of attendance at the consultation appointment.
2. Assess indicators of longer-term service engagement 6 months following study enrolment.
Hypothesis 2: The SMS intervention will lead to improved indicators of longer-term service engagement

(Service Engagement Scale (SES) scores and dropout rates).

3. Determine system-level outcomes, including ED visits and psychiatric hospitalisations, as a function of receiving the SMS intervention, and its cost effectiveness, factoring costs of the intervention and cost offsets of health service utilisation.

Hypothesis 3: The SMS intervention will lead to decreased use of acute care services (ED visits and psychiatric hospitalisations) and will be cost-effective relative to the control condition, based on improved rate of transition from the ED to EPI services and anticipated reductions in use of costly acute care services.

4. Explore young people's experiences of the intervention and their perspectives on its various components.

This is an exploratory research question that seeks to understand how young people experience the SMS intervention and how they perceive its various components impact their service engagement.

METHODS AND ANALYSIS

Study setting

The study setting will be the Gerald Sheff and Shanitha Kachan Emergency Department at CAMH, Ontario's only 24 hours stand-alone psychiatric emergency service. The ED also houses a drop-in 'Bridging Clinic' which provides care to less acute patients who are diverted after ED triage, and rapid follow-up care for patients discharged from the ED and CAMH inpatient units. Together, they serve approximately 1200 patients each month. The EPI programme at CAMH receives over 600 referrals for suspected psychosis annually, approximately 25% of which are from the ED and Bridging Clinic. Reflecting the Ontario EPI Programme Standards, CAMH EPI services are delivered by multidisciplinary teams, employ strategies to promote early entry and ongoing engagement, and provide pharmacotherapy and psychosocial therapies for an average of 3 years.¹⁴ Patients are assigned to the next available and/or most appropriate psychiatrist and case manager (nurse, social worker or occupational therapist) for a joint consultation appointment, typically within 2 weeks, and are contacted by phone by the EPI programme administrator to book and confirm their appointment. After the initial appointment is confirmed, patients receive a phone call reminder the day before their scheduled appointment. As part of routine care, patients who do not attend their scheduled first appointment receive follow-up calls to reschedule an appointment for up to 30 days from the initial referral.

Eligibility criteria

Study inclusion criteria mirror the intake criteria for the CAMH EPI programme. Participants will be eligible for the study if they: (i) are between 16 and 29 years old and (ii) have been referred by the CAMH ED to CAMH EPI services for suspected psychosis. Our only exclusion criterion is inability to communicate in basic written English.

In our pilot SMS study at CAMH, fewer than 5% of potential participants were excluded for lacking a phone; we have budgeted to offer 5% of participants access to a prepaid cellphone for the duration of the study.²⁷

Intervention procedures

Study participants will be recruited at the time of EPI referral for a trial of an SMS intervention designed to engage them during the waiting period for their consultation appointment. They will be randomised to receive either sham or active SMS intervention.

Sham SMS will consist of one message sent just after enrolment indicating that they will be contacted for an appointment. The sham SMS group will not be denied any part of usual clinical care. Thus, the clinic administrator will call patients in both groups to book and remind them of the consultation appointment. Active SMS intervention will include the initial message sent to the control group, plus a series of subsequent messages. These will include the following content: (i) appointment reminders and instructions, (ii) psychoeducational material and (iii) two-way communication check-ins to rate distress (figure 1). Intervention components were developed based on feedback from a survey of youth in the same EPI programme,³⁰ as well as psychosocial interventions with evidence in early psychosis, including cognitive-behavioural therapy (psychoeducation, behavioural activation) and illness self-management (reminders, distress check-in).^{31 32} See online supplemental file 1 for a comprehensive description of the intervention. Messages sent back by participants will be monitored by the research team, and the case manager assigned to their consultation appointment will be notified and will respond accordingly if there are indicators of elevated distress. The intervention will continue until the patient attends the first consultation appointment, or for up to 30 days if the patient does not attend, which reflects the programme's practice of closing referrals for non-attending patients.

Messages will be sent through CAMH's in-house Research Electronic Data Capture (REDCap) platform via a third-party plug-in, Twilio, which supports routing of SMS messages to participant devices.³³⁻³⁵ All data are stored securely on CAMH's REDCap servers or in a locked office and password-protected database on CAMH's secure network. The purpose of the sham intervention is to separate out the content of the intervention (ie, reminders, psychoeducation, two-way check-ins) from the effect of simply receiving SMS messages, decreasing participant bias.

Assignment of intervention

Immediately after informed consent, participants will be randomised by REDCap to the active or sham intervention. Generation of the randomisation sequence will be managed by the study biostatistician who is not involved in enrolling participants or assigning intervention arms. Randomisation will be stratified by sex (male or female) and referral source (ED or Bridging Clinic), using a




Figure 1 Study intervention schedule. This figure represents a summary of the intervention schedule and is not exhaustive of text message content. Left arrow, outgoing text messages sent by participants; right arrow, incoming text messages received by participants; SMS, short message service.

computer algorithm to determine a randomised, blocked allocation of participants into intervention groups within strata. Once randomised, treatment assignment will be known only by the study personnel involved in managing the SMS intervention and the case manager monitoring text message responses; the principal investigator and study personnel involved in the chart review and the ICES analysis will remain blind to treatment assignment. Study personnel involved in the qualitative interviews and analysis will also be aware of treatment assignment since only individuals receiving the active intervention will participate in this component of the project.

Sample size

Our sample size calculations are based on the primary outcome of rate of attendance at the EPI programme consultation appointment. The current rate of attendance at the first consultation appointment for patients referred from the ED and Bridging Clinic is 50%; we have powered our study to detect the treatment effect with an anticipated rate of attendance of 70%, which is the average for all referral sources other than the ED and

Bridging Clinic. A total of 186 participants (93 per group) will provide >80% power to detect a change in attendance from 50% to 70% at $\alpha < 0.05$. Non-compliance and loss to follow-up are not a concern for our primary outcome since these are counted in the outcome as non-attendance at the consultation. Adjusting analyses for covariates expected to affect attendance (eg, age, sex) is expected to further increase power.

Study procedures

We will recruit 186 patients consecutively referred from the CAMH ED and Bridging Clinic to the CAMH EPI programme at the time of discharge. During business hours and into the evening, research staff will be on call to the ED to recruit patients as soon as they are identified for the study. Eligible participants who present after hours may be identified by ED staff and referred to the research team who will send them an e-consent form via SMS or email. See online supplemental file 2 for patient consent form. Additionally, potential participants who are missed in the ED/Bridging Clinic may be identified by the EPI clinic administrator who will approach them

over the phone and send them an e-consent form via SMS or email. We have used the approach of having clinical and administrative staff obtain verbal consent to send e-consent forms in other studies with this population.¹⁹

As part of informed consent, participants will be asked to consent to a review of their chart, a follow-up web-based survey, and linkage of their information to data held at ICES. They will also be given the option to provide consent to be re-contacted for participation in a qualitative interview. There will be no additional in-person assessments for the quantitative component of this study. CAMH uses an electronic health record, which facilitates data abstraction from multiple clinical programmes (ie, both the ED and Bridging Clinic and EPI programme). Research staff and students will be trained by the clinician principal investigators to abstract data into a structured database.

All study participants will receive a \$10 e-giftcard once the baseline e-visit is complete. Participants in the active intervention group who complete a web-based survey will receive another \$10 e-giftcard and those who complete a qualitative interview will receive a \$50 e-giftcard. It will be clarified through the consent process that honoraria are to compensate participants for their time and will not be tied to clinical appointment attendance. For the majority of participants, their initial recruitment and consent will be their only interaction with the research team, with a small subgroup completing qualitative interviews.

Outcome measures

Chart review

Outcome measures are shown in [table 1](#). CAMH uses many standardised assessment forms which increases the completeness of patient data. Demographic variables, clinical diagnoses, substance use, duration of untreated psychosis (measured as the period of time from first onset of psychotic symptoms to initiation of EPI services and initiation of treatment with an antipsychotic or mood-stabilising medication), characteristics of the ED visit from which they were referred (urgent presentation—brought by police, involuntary status; timing of visit), and family involvement in care are routinely recorded in the clinical chart by clinicians and will be abstracted from the chart at the time of consultation. Additional variables will likely be available but only for patients accepted into the EPI programme, and this will be reflected in the data analysis. These include several assessments that are performed routinely in the EPI programme. The SES³⁶ is a brief validated tool designed to measure engagement with community mental health services. In 14 items, it assesses patients' availability for treatment, collaboration, help-seeking behaviours and treatment adherence on a four-point Likert scale with higher scores indicating difficulties in service engagement. The Brief Psychiatric Rating Scale³⁷ is a clinician or interviewer-rated measure of psychiatric symptoms commonly used as an outcome measure for psychotic disorders and collected monthly in CAMH's EPI programme. It includes items related

to suicidality and hostility. The Clinical Global Impression³⁸ is a clinician-rated measure of the patient's global severity of illness prior to and after initiating a medication. It includes subscales for severity and improvement. Medication and appointment non-adherence will also be assessed over 6 months of treatment. Finally, after 6 months, current EPI enrolment status will be assessed and categorised as: not offered or enrolled in treatment (eg, because they did not ultimately have psychosis), enrolled but disengaged prematurely, accepted for treatment but transitioned to other services or continued in treatment. Data abstractors will undergo rigorous training and monitoring, use standardised extraction forms, and calculate inter-rater reliability.

Administrative data

Primary data collected for the study will be linked deterministically to data sources held at ICES via participants' unique health card numbers. The information available for each participant will be de-identified, stored and analysed onsite at ICES following procedures approved by Ontario's Information and Privacy Commissioner. The following ICES data sources will be used: the Ontario Mental Health Reporting System, capturing hospitalisations on adult inpatient mental health units,³⁹ the Canadian Institute of Health Information Discharge Abstract Database, capturing all hospital admissions including hospitalisations on child and adolescent inpatient mental health units,^{40 41} National Ambulatory Care Reporting System which captures all ED visits,⁴² Ontario Health Insurance Plan claims database, which captures outpatient physician visits,⁴⁰ Registered Persons Database, which contains health card numbers, demographic information, and deaths, and Ontario Drug Benefits claims database, which provides information on all covered prescriptions (based on financial need for those under age 65 and for young people up to age 25 who lack private insurance). These data will also be used for cost effectiveness analysis. Outcomes examined in the linked ICES data are listed in [table 1](#).

Statistical analysis

Primary analysis

Descriptive and graphical statistics will be used to summarise the data on all randomised participants and to confirm that there are no group differences in baseline demographics and clinical characteristics. Distributional assumptions will be inspected and appropriate transformations or non-parametric methods will be applied as necessary. In general, generalised linear models⁴³ will be used throughout. These models account for deviation from normal assumption of the outcome variables and control for covariates.

Our analysis of the primary outcome will be a logistic regression to examine the likelihood of attendance at the EPI consultation with treatment assignment using risk ratios.⁴⁴ We will carefully select demographic variables and factors known to influence treatment engagement

**Table 1** Summary of outcome measures and covariates

Type	Variables	Data source	Timing
Demographic characteristics	Age	Chart review for all demographic characteristics (CAMH Health Equity form and notes)	Baseline*
	Sex and gender		
	Sexual orientation		
	Race/ethnicity		
	Born in Canada		
	Religious/spiritual affiliation		
	Highest level of education		
	Source of income and family income		
	Number of people supported by income		
	Employment status		
	Legal history		
	Housing status		
	Living situation		
	Experience of homelessness		
Relationship status			
Clinical characteristics	Clinical diagnoses	Chart review for all clinical characteristics (consultation and progress notes)	Baseline
	Substance use		Baseline
	DUP		Baseline
	Family involvement in care		Baseline
	Urgent status at ED visit (brought by police, involuntary)		
	Timing of ED visit		
	BPRS ³⁷		Baseline and 6 months
	CGI ³⁸		Baseline and 6 months
Service engagement	Attendance at consultation appointment	Chart review for all service engagement measures	30 days
	SES ³⁶		6 months (completed around 3 months in treatment)
	Medication and appointment non-adherence		6 months
	EPI enrolment status		6 months
System-level outcomes†	Number of ED visits	NACRS ⁴²	6 months and up to 2 years
	Number of inpatient mental health hospitalisations	OMHRS, ³⁹ CIHI-DAD ^{40 41}	
	Number of days in inpatient mental health hospitalisations	OMHRS, ³⁹ CIHI-DAD ^{40 41}	
	Number of outpatient mental health visits	OHIP ⁴⁰	
	Prescription drugs—psychiatric medications, continuously vs non-continuously prescribed	ODB	
	Mortality including cause of death	Registered Persons Database	

*Items may be extracted from the ED note or EPI consultation note.

†Administrative data held at Institute for Clinical Evaluative Sciences.

BPRS, Brief Psychiatric Rating Scale; CAMH, Centre for Addiction and Mental Health; CGI, Clinical Global Impression; CIHI-DAD, Canadian Institute of Health Information Discharge Abstract Database; DUP, duration of untreated psychosis; ED, emergency department; EPI, early psychosis intervention; NACRS, National Ambulatory Care Reporting System; ODB, Ontario Drug Benefits; OHIP, Ontario Health Insurance Plan; OMHRS, Ontario Mental Health Reporting System; SES, Service Engagement Scale.

(eg, substance use, family involvement in care)⁴⁵ to be included in the model as covariates. A difference in attendance between groups will be declared at a significance level of 0.05. Similar models will be used to address the secondary hypotheses, with specific types of models appropriate to each outcome, including time-to-event analysis to examine premature disengagement from services. Administrative data outcomes will be examined using generalised linear models with proper distribution assumptions.

Additional analyses: moderation and generalisability

We plan to conduct two additional exploratory analyses. First, we will run moderation analyses on potential effect modifiers by adding an interaction term between the potential moderator and the treatment assignment indicator in the generalised linear models. We are specifically interested in the moderation effects of health equity factors including gender, race/ethnicity and housing status. A significant interaction will provide evidence that the treatment effects may be different in the subgroups. A second exploratory analysis will be conducted to evaluate the impact of selection bias of the study sample and estimate the population average treatment effects by employing weighted analysis using propensity scores.⁴⁶

Missing data

The risk of missing data is mitigated through the use of chart review and analysis of administrative data. While the primary outcome will not suffer from attrition, other outcomes will, as some follow-up data will only be available for participants who attended their consultation appointment and those who are enrolled in CAMH EPI services. For these additional outcomes, we plan to use multiple imputation methods developed by Schafer to correct potential bias that could be introduced by missing data.⁴⁷

Economic evaluation

Full details of the economic evaluation appear in online supplemental file 3. We will undertake a cost-effectiveness analysis, where the outcome of interest is consultation appointment attendance, adopting the perspective of the public third-party payer (ie, the Ontario Ministry of Health). We will collect data on the costs of delivering both arms of the intervention. In addition, using a costing algorithm available at ICES,⁴⁸ we will estimate all direct patient-level healthcare costs incurred by the public third-party payer for the intervention and control groups, which will include costs of hospitalisations, ED visits, physician services (ie, primary care, psychiatry and other) and diagnostic tests, outpatient prescription drugs for individuals covered under the provincial public drug insurance plan, and other hospital-based care. We will calculate the incremental cost-effectiveness ratio as the difference in discounted mean costs between the intervention and control groups, divided by the difference in attendance rates.

We will use a net benefit regression approach to model probabilities of cost effectiveness for each additional patient referred who attends their consultation appointment in the intervention compared with control group. In addition, we will undertake relevant sensitivity analyses to test the robustness of findings by varying relevant parameters, such as the discount rate. Finally, we will examine the real-world budget impact of implementing the intervention across Ontario, to estimate the cost to the Ministry of Health of implementing this model of care across the province and the potential cost-savings to the system associated with this.

Understanding patient experiences: survey and qualitative analysis

Participants in the active intervention group will receive a one-time survey sent as a web-link to their phone or email address. Survey topics include user experience, attitudes toward the SMS intervention, its perceived benefits and challenges, acceptability and suggestions for improvements. Those who consent to participate in the qualitative research component will be recontacted by phone or email to participate in semi-structured interviews to gain a more in-depth understanding of survey topics. A subsample of 10–15 participants in the active intervention group will be purposively selected to maximise diversity of age, gender and service attendance. We will use critical realist theory as an underlying framework to guide our interviews, surveys and analysis.⁴⁹ Interviews will be completed until thematic saturation is achieved, estimated at 12 participants. Interviews will be digitally audio-recorded and transcribed verbatim. Surveys will be completed and stored in REDCap. Transcriptions and survey responses will be analysed using thematic content analysis in NVivo-11. Research participants will be invited to assist with member checking to confirm that themes reflect their experiences. The analysis can inform future improvements to the intervention and considerations for broader implementation, privileging the experiences of the patients attending these programmes.

Patient and public involvement

A youth and family member who previously received EPI services have been engaged in helping shape the intervention and study design from project inception. They are active members of the project's Steering Committee that meets monthly to inform study design, implementation, evaluation, and dissemination of results. They will have key roles in the plan to spread the intervention, if successful, to other EPI programmes by working with patients and families to adapt the intervention to local contexts. Additional youth with lived experience of receiving EPI services have been consulted on an ad hoc basis through the CAMH Youth Engagement Initiative to provide detailed feedback on the SMS intervention. Patient and family representatives on the research team are compensated for their time.

ETHICS AND DISSEMINATION

The study was approved by the Research Ethics Board (REB) at the CAMH. The study protocol was prepared according to Standard Protocol Items: Recommendations for Interventional Trials guidelines⁵⁰ and registered with clinicaltrials.gov on 6 March 2020 (<https://clinicaltrials.gov/ct2/show/NCT04298450?term=ed+to+epi&draw=2&rank=1>). REB-approved protocol amendments will be posted on the site. The principal investigators and study team will meet regularly to review accrued data, data confidentiality, any adverse events, adherence to protocol design, recruitment and implementation. This intervention has been designed to have high likelihood of adoption and readiness for spread and scale-up because it responds to a critical need, has a strong evidentiary basis, has advantages over existing practice and is both low complexity and low cost.⁵¹ The study team is well-positioned to support widespread implementation of the intervention if successful. We have used an integrated knowledge translation approach that leverages input from stakeholders, including patients, clinicians (both from the ED and EPI services), policymakers and relevant organisations throughout the study to champion the spread of the intervention to other EPI programmes and youth mental health services more broadly. This trial focuses on a particularly vulnerable population—young people transitioning from adolescence to adulthood and from the ED to EPI services—but the basic intervention is widely applicable. The software platform used to coordinate this intervention is available at no charge, and the SMS functionality for sending and receiving messages carries a nominal fee, supporting broad uptake.

The results of the trial will be reported in scientific journal articles and shared with key stakeholders as they become available. Our study team includes the co-chair of the Early Psychosis Intervention Ontario Network, a network of over 50 EPI programmes across Ontario, and several members of the Canadian Consortium for Early Intervention in Psychosis, providing a durable and established community of practice for immediate spread. De-identified participant data will be available on reasonable request other than system-level data held at ICES. Requests can be made by contacting the principal investigator Dr Nicole Kozloff at nicole.kozloff@camh.ca and will be managed by the Steering Committee.

This pragmatic randomised-controlled trial of a low-cost, low-complexity SMS intervention aims to improve the transition from the ED to EPI services for young people with psychosis. It targets a brief but critical period: if young people cannot even get in the door to EPI services, there is no way for them to reap the many known benefits of EPI care. Improving the ED to EPI transition has the potential to result in more young people with psychosis getting appropriate treatment earlier. The proposed intervention is also likely to be easily adaptable to other referral pathways to EPI services and youth mental health services more broadly. At potentially lower cost to the health system, applying this SMS intervention to the ED

to EPI transition has the potential to lead to improved short-term symptoms and functioning, long-term disease trajectories, decreased burden on patients and families, and fewer deaths among young people with psychosis.

Author affiliations

- ¹Slaight Family Centre for Youth in Transition, Centre for Addiction and Mental Health, Toronto, Ontario, Canada
²Psychiatry, University of Toronto, Toronto, Ontario, Canada
³Campbell Family Mental Health Research Institute, Centre for Addiction and Mental Health, Toronto, Ontario, Canada
⁴Centre for Addiction and Mental Health, Toronto, Ontario, Canada
⁵Epidemiology & Biostatistics, Western University, London, Ontario, Canada
⁶Institute of Medical Science, University of Toronto, Toronto, Ontario, Canada
⁷Institute for Mental Health Policy Research, Centre for Addiction and Mental Health, Toronto, Ontario, Canada
⁸Institute of Health Policy, Management and Evaluation, University of Toronto, Toronto, Ontario, Canada
⁹Margaret and Wallace McCain Centre for Child, Youth and Family Mental Health, Centre for Addiction and Mental Health, Toronto, Ontario, Canada

Twitter Alexia Polillo @alexiaPolillo, Kelly K Anderson @anderskk3, Claire de Oliveira @clairede0, Juveria Zaheer @juveriazaheer and Nicole Kozloff @nicole_kozloff

Contributors NK is the principal investigator who conceived the original study design and obtained funding, with most of the current authors having contributed to the funding application, and all authors having participated in revisions to the study design for important intellectual content. AP, GF, AHCW, AA, VS, JD, LD, ANV and NK sit on the project's Steering Committee. AP, GF, JD and NK form the Data Management Committee. AP leads the survey and qualitative analysis. GF, AHCW, SB and ANV have administrative roles in the clinical programmes and will support the acquisition and interpretation of data. AA and LD act as patient and public consultants. VS acts as a Health System Decision-Maker on the project. KKA, CdO, PK and NK consult on the ICES analysis. CdO is a health economist who consults on the economic analysis. VS, JH and SK provide knowledge translation expertise. JZ provides consultation on the qualitative interviews and analysis. WW acts as the biostatistical consultant. AP and NK drafted the protocol. All authors read, revised and approved the final version of the manuscript.

Funding This work was supported by the Canadian Institutes of Health Research Transitions in Care Team Grant #165726 as well as by the CAMH Foundation (Award/Grant Number Not Applicable). The funders were not involved in study design; collection, management, analysis, and interpretation of data; writing of the report; and the decision to submit the report for publication, including whether they will have ultimate authority over any of these activities.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Joanna Henderson <http://orcid.org/0000-0002-9387-5193>
 Nicole Kozloff <http://orcid.org/0000-0003-1389-1351>

REFERENCES

- 1 American Psychiatric Association. *Diagnostic and statistical manual of mental disorders: DSM-5*. 5th edn. Arlington, VA: American Psychiatric Publishing, 2013.
- 2 Whiteford HA, Degenhardt L, Rehm J, *et al*. Global burden of disease attributable to mental and substance use disorders: findings from the global burden of disease study 2010. *Lancet* 2013;382:1575–86.
- 3 Schoenbaum M, Sutherland JM, Chappel A, *et al*. Twelve-month health care use and mortality in commercially insured young people with incident psychosis in the United States. *Schizophr Bull* 2017;43:1262–72.
- 4 Zaheer J, Olfson M, Mallia E, *et al*. Predictors of suicide at time of diagnosis in schizophrenia spectrum disorder: a 20-year total population study in Ontario, Canada. *Schizophr Res* 2020;222:382–8.
- 5 Lieberman JA, First MB. Psychotic disorders. *N Engl J Med* 2018;379:270–80.
- 6 de Oliveira C, Cheng J, Rehm J, *et al*. The economic burden of chronic psychotic disorders in Ontario. *J Ment Health Policy Econ* 2016;19:181–92.
- 7 Goeree R, Farahati F, Burke N, *et al*. The economic burden of schizophrenia in Canada in 2004. *Curr Med Res Opin* 2005;21:2017–28.
- 8 Penttilä M, Jääskeläinen E, Hirvonen N, *et al*. Duration of untreated psychosis as predictor of long-term outcome in schizophrenia: systematic review and meta-analysis. *Br J Psychiatry* 2014;205:88–94.
- 9 Anderson KK, Norman R, MacDougall A, *et al*. Effectiveness of early psychosis intervention: comparison of service users and nonusers in population-based health administrative data. *Am J Psychiatry* 2018;175:443–52.
- 10 Craig TKJ, Garety P, Power P, *et al*. The Lambeth early onset (LEO) team: randomised controlled trial of the effectiveness of specialised care for early psychosis. *BMJ* 2004;329:1067.
- 11 Petersen L, Jeppesen P, Thorup A, *et al*. A randomised multicentre trial of integrated versus standard treatment for patients with a first episode of psychotic illness. *BMJ* 2005;331:602.
- 12 Malla A, Schmitz N, Norman R, *et al*. A multisite Canadian study of outcome of first-episode psychosis treated in publicly funded early intervention services. *Can J Psychiatry* 2007;52:563–71.
- 13 Bond GR, Drake RE, Luciano A. Employment and educational outcomes in early intervention programmes for early psychosis: a systematic review. *Epidemiol Psychiatr Sci* 2015;24:446–57.
- 14 Ontario Ministry of Health and Long-Term Care. Early psychosis intervention program standards. Toronto, ON; 2011. http://www.health.gov.on.ca/english/providers/pub/mental/epi_program_standards.pdf
- 15 Anderson KK, Norman R, MacDougall AG, *et al*. Estimating the incidence of first-episode psychosis using population-based health administrative data to inform early psychosis intervention services. *Psychol Med* 2019;49:1–9.
- 16 Anderson KK, Fuhrer R, Wynant W, *et al*. Patterns of health services use prior to a first diagnosis of psychosis: the importance of primary care. *Soc Psychiatry Psychiatr Epidemiol* 2013;48:1389–98.
- 17 Kozloff N, Jacob B, Voineskos AN, *et al*. Care of youth in their first emergency presentation for psychotic disorder: a population-based retrospective cohort study. *J Clin Psychiatry* 2018;79 doi:10.4088/JCP.17m11947
- 18 Lal S, Dell'Elce J, Tucci N, *et al*. Preferences of young adults with first-episode psychosis for receiving specialized mental health services using technology: a survey study. *JMIR Ment Health* 2015;2:e18.
- 19 Voineskos AN, Foussias G, *et al*, Kozloff N. Patient and family perspectives on improving engagement in early psychosis intervention. *Congress of the Schizophrenia International Research Society*; April 10–14, 2019, Orlando, FL, 2019.
- 20 Polillo A, Voineskos A, *et al*, Sav A. Using REDCap to evaluate youth and family perspectives on engagement in early psychosis intervention. *REDCap Con December 2019*; December 12, 2019, Toronto, ON, Canada, 2019.
- 21 Firth J, Torous J. Smartphone apps for schizophrenia: a systematic review. *JMIR Mhealth Uhealth* 2015;3:e102.
- 22 D'Arcey J, Collaton J, Kozloff N, *et al*. The use of text messaging to improve clinical engagement for individuals with psychosis: systematic review. *JMIR Ment Health* 2020;7:e16993.
- 23 Alvarez-Jimenez M, Alcazar-Corcoles MA, González-Blanch C, *et al*. Online, social media and mobile technologies for psychosis treatment: a systematic review on novel user-led interventions. *Schizophr Res* 2014;156:96–106.
- 24 Ennis L, Rose D, Denis M, *et al*. Can't surf, won't surf: the digital divide in mental health. *J Ment Health* 2012;21:395–403.
- 25 Kauppi K, Kannisto KA, Hätönen H, *et al*. Mobile phone text message reminders: measuring preferences of people with antipsychotic medication. *Schizophr Res* 2015;168:514–22.
- 26 Thomas IF, Lawani AO, James BO. Effect of short message service reminders on clinic attendance among outpatients with psychosis at a psychiatric hospital in Nigeria. *Psychiatr Serv* 2017;68:75–80.
- 27 Using short message service as a means of engagement in early psychosis. IEPA 11th International Conference on Early Intervention in Mental Health - 'Prevention and Early Intervention: Broadening the Scope'; October 7–10 2018. Boston, Massachusetts, USA. Early Interv Psychiatry.
- 28 Meinecke A-K, Welsing P, Kafatos G, *et al*. Series: pragmatic trials and real world evidence: paper 8. data collection and management. *J Clin Epidemiol* 2017;91:13–22.
- 29 Thorpe KE, Zwarenstein M, Oxman AD, *et al*. A pragmatic-explanatory continuum indicator summary (PRECIS): a tool to help trial designers. *J Clin Epidemiol* 2009;62:464–75.
- 30 Voineskos AN, Foussias G, *et al*, Kozloff N. Improving youth engagement in early psychosis intervention programs: preliminary results from a youth survey. *66th Annual Meeting of the American Academy of Child and Adolescent Psychiatry*; October 14–19, 2019, Chicago, IL, USA, 2019:S334–5. doi:<https://doi.org/10.1016/j.jaac.2019.07.791>
- 31 Lecomte T, Abidi S, Garcia-Ortega I, *et al*. Canadian treatment guidelines on psychosocial treatment of schizophrenia in children and youth. *Can J Psychiatry* 2017;62:648–55.
- 32 Keepers GA, Fochtmann LJ, Anzia JM, *et al*. The American psychiatric association practice guideline for the treatment of patients with schizophrenia. *Am J Psychiatry* 2020;177:868–72.
- 33 Vanderbilt University. REDCap (research electronic data capture), 2018. Available: <https://www.project-redcap.org> [Accessed 2 Apr 2020].
- 34 Harris PA, Taylor R, Thielke R, *et al*. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009;42:377–81.
- 35 Harris PA, Taylor R, Minor BL, *et al*. The REDCap Consortium: building an international community of software platform partners. *J Biomed Inform* 2019;95:103208.
- 36 Tait L, Birchwood M, Trower P. A new scale (SES) to measure engagement with community mental health services. *J Ment Health* 2002;11:191–8.
- 37 Flemenbaum A, Zimmermann RL. Inter- and intra-rater reliability of the brief psychiatric rating scale. *Psychol Rep* 1973;32:783–92.
- 38 Guy W. *Clinical global impressions. ECDEU assessment manual for psychopharmacology*. Rockville, Maryland: National Institute for Mental Health, 1976: 218–22.
- 39 Canadian Institute for Health Information. Ontario mental health reporting system, data quality documentation, 2015–2016. Ottawa, ON, Canada CIHI; 2017. <https://www.cihi.ca/sites/default/files/document/omhrs-dq-external-report-2015-2016-en.pdf>
- 40 Williams JI, Young W. A summary of studies on the quality of health care administrative databases in Canada. In: Goel V, Williams JI, Anderson GM, *et al*, eds. *Patterns of health care in Ontario the ICES practice atlas*. 2nd edn. Ottawa, ON: Canadian Medical Association, 1996. <https://www.ices.on.ca/flip-publication/patterns-of-health-care-2d-edition/files/assets/common/downloads/ICES%20.pdf>
- 41 Juurlink DN, Preyra C, Croxford R. Canadian Institute for Health Information Discharge Abstract Database: A Validation Study. Toronto, ON, Canada Institute for Clinical Evaluative Sciences; 2006. <https://www.ices.on.ca/flip-publication/canadian-institute-for-health-information-discharge/files/assets/basic-html/index.html#1>
- 42 Canadian Institute for Health Information. CIHI data quality study of Ontario emergency department visits for 2004–2005. Ottawa, ON, Canada CIHI; 2008. https://secure.cihi.ca/free_products/vol1_nacrs_executive_summary_nov2_2007.pdf
- 43 McCullagh P, Nelder JA. *Generalized linear models*. Second Edition. London, UK: Chapman and Hall, 1989.
- 44 McNutt L-A, Wu C, Xue X, *et al*. Estimating the relative risk in cohort studies and clinical trials of common outcomes. *Am J Epidemiol* 2003;157:940–3.
- 45 Doyle R, Turner N, Fanning F, *et al*. First-episode psychosis and disengagement from treatment: a systematic review. *Psychiatr Serv* 2014;65:603–11.
- 46 Wang W, Ma Y, Huang Y, *et al*. Generalizability analysis for clinical trials: a simulation study. *Stat Med* 2017;36:1523–31.
- 47 Schafer JL. *Analysis of incomplete multivariate data*. London, UK: Chapman & Hall, 1997.
- 48 Wodchis WP, Austin PC, Henry DA. A 3-year study of high-cost users of health care. *CMAJ* 2016;188:182–8.



- 49 Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol* 2006;3:77–101.
- 50 Chan A-W, Tetzlaff JM, Altman DG, et al. SPIRIT 2013 statement: defining standard protocol items for clinical trials. *Ann Intern Med* 2013;158:200–7.
- 51 Damschroder LJ, Aron DC, Keith RE, et al. Fostering implementation of health services research findings into practice: a consolidated framework for advancing implementation science. *Implement Sci* 2009;4:50.