



## Correction to: Cost-Utility Analysis of Discontinuing Antidepressants in England Primary Care Patients Compared with Long-Term Maintenance: The ANTLER Study

Caroline S. Clarke<sup>1</sup> · Larisa Duffy<sup>2</sup> · Glyn Lewis<sup>2</sup> · Nick Freemantle<sup>3</sup> · Simon Gilbody<sup>4</sup> · Tony Kendrick<sup>5</sup> · David Kessler<sup>6</sup> · Michael King<sup>2</sup> · Paul Lanham<sup>7</sup> · Derelie Mangin<sup>8</sup> · Michael Moore<sup>5</sup> · Irwin Nazareth<sup>1</sup> · Nicola Wiles<sup>9</sup> · Louise Marston<sup>1</sup> · Rachael Maree Hunter<sup>1</sup>

Published online: 11 December 2021  
© The Author(s) 2021

**Correction to: Applied Health Economics and Health Policy**  
<https://doi.org/10.1007/s40258-021-00693-x>

In this article the text in Sect. 3.2 and the legend for Fig. 1 has been published with inadvertent errors.

The final sentence of Sect. 3.2, which previously read as

“The trend in utilities can also be seen in Fig. 1, which shows the mean adjusted imputed utility scores at each time point for the three different methods of calculating utilities.”

Should read as

“The trend in utilities can also be seen in Fig. 1, which shows the mean unadjusted complete-case utility scores at each time point for the three different methods of calculating utilities.”

The legend to Fig. 1, which previously read as

“Fig. 1 Adjusted imputed mean utility scores at each time-point, by arm, for each of the three methods: EQ-5D-5L

VSE, EQ-5D-5L crosswalk and SF-12/SF-6D. *SF-12* 12-Item Short-Form survey, *VSE* value set for England”

Should read as

“Fig. 1 Unadjusted complete-case mean utility scores at each timepoint, by arm, for each of the three methods: EQ-5D-5L VSE, EQ-5D-5L crosswalk and SF-12/SF-6D. *SF-12* 12-Item Short-Form survey, *VSE* value set for England”

The original article has been corrected.

**Open Access** This article is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License, which permits any non-commercial use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc/4.0/>.

The original article can be found online at <https://doi.org/10.1007/s40258-021-00693-x>.

✉ Caroline S. Clarke  
caroline.clarke@ucl.ac.uk

<sup>1</sup> Research Department of Primary Care and Population Health, University College London, London, UK

<sup>2</sup> Division of Psychiatry, University College London, London, UK

<sup>3</sup> Institute of Clinical Trials and Methodology, University College London, London, UK

<sup>4</sup> Department of Health Sciences and Hull York Medical School, University of York, York, UK

<sup>5</sup> Primary Care Population Sciences and Medical Education, Faculty of Medicine, University of Southampton, Southampton, UK

<sup>6</sup> Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

<sup>7</sup> Patient and Public Involvement Collaborator, London, UK

<sup>8</sup> Department of Family Medicine, McMaster University, Hamilton, Canada

<sup>9</sup> Centre for Academic Mental Health, Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK