

RESEARCH ARTICLE

Open Access

A novel locus (*CORD12*) for autosomal dominant cone-rod dystrophy on chromosome 2q24.2-2q33.1

Gaël Manes^{1,2,3*}, Maxime Hebrard^{1,2,3}, Béatrice Bocquet^{1,2,3}, Isabelle Meunier⁴, Delphine Coustes-Chazalette⁴, Audrey Sénéchal^{1,2,3}, Anne Bolland-Augé⁵, Diana Zelenika⁵ and Christian P Hamel^{1,2,3,4}

Abstract

Background: Rod-cone dystrophy, also known as retinitis pigmentosa (RP), and cone-rod dystrophy (CRD) are degenerative retinal dystrophies leading to blindness. To identify new genes responsible for these diseases, we have studied one large non consanguineous French family with autosomal dominant (ad) CRD.

Methods: Family members underwent detailed ophthalmological examination. Linkage analysis using microsatellite markers and a whole-genome SNP analysis with the use of Affymetrix 250 K SNP chips were performed. Five candidate genes within the candidate region were screened for mutations by direct sequencing.

Results: We first excluded the involvement of known adRP and adCRD genes in the family by genotyping and linkage analysis. Then, we undertook a whole-genome scan on 22 individuals in the family. The analysis revealed a 41.3-Mb locus on position 2q24.2-2q33.1. This locus was confirmed by linkage analysis with specific markers of this region. The maximum LOD score was 2.86 at $\theta = 0$ for this locus. Five candidate genes, *CERKL*, *BBS5*, *KLHL23*, *NEUROD1*, and *SF3B1* within this locus, were not mutated.

Conclusion: A novel locus for adCRD, named *CORD12*, has been mapped to chromosome 2q24.2-2q33.1 in a non consanguineous French family.

Background

Retinitis pigmentosa (RP, [MIM 268000]) is a genetically heterogeneous group of retinal photoreceptor degeneration characterized by night blindness and loss in the peripheral visual field, slowly progressing towards total blindness after several decades [1]. RP accounts for about 2/3 of the inherited retinal dystrophy cases [2]. In contrast to typical RP, also called rod-cone dystrophies (RCDs) because of primary involvement of rods, inverse RP or cone-rod dystrophies (CRDs) are pigmentary retinopathies characterized by first decrease in visual acuity and loss in the central visual field and lately by night blindness and loss in the peripheral visual field. CRDs are due to the primary degeneration of cone photoreceptors, followed by the secondary, or, sometimes,

concomitant loss of rod photoreceptors [3]. Forty nine genes and loci are responsible for non syndromic RP and 18 for non syndromic CRD (including 6 in common with RP and 4 with Leber congenital amaurosis) <http://www.sph.uth.tmc.edu/Retnet>. The three types of Mendelian inheritance are encountered in both RP and CRD.

Among the 18 CRD genes, ten (*GUCY2D*, *PITPNM3*, *GUCA1A*, *HRG4/UNC119*, *CRX*, *AIPL1*, *RIMS1*, *SEMA4A*, *PROM1* and *PRPH2/RDS*) are found in autosomal dominant (ad) CRD, six (*ABCA4*, *RPGRIP1*, *RAX2*, *CORD8*, *ADAM9* and *CERKL*) in autosomal recessive (ar) CRD and two (*RPGR* and *CACNA1F*) in X-linked CRD <http://www.sph.uth.tmc.edu/Retnet>. The prevalence of mutations for each gene in the CRD population is highly variable. *ABCA4*, which causes Stargardt macular dystrophy, is also a major gene for CRD, being responsible for 30-60% of arCRD cases [4-6]. In contrast, the overall prevalence of adCRD genes remains low, many of them being described in only one or a few

* Correspondence: gael.manes@inserm.fr

¹INSERM U1051, Institute for Neurosciences of Montpellier, (80 rue Augustin Fliche), Montpellier, (34091), France

Full list of author information is available at the end of the article

cases. Only *CRX*, *GUCY2D* and *PRPH2/RDS* have been consistently reported in adCRD [7-10]. Yet, *CRX* was estimated to account for only 5-10% of adCRD cases and the prevalence of *GUCY2D* and *PRPH2/RDS* is unknown [11,12]. Therefore, there are probably other genes remaining to be discovered in adCRD.

In search for new genes responsible for pigmentary retinopathies, we recruited one large non-consanguineous French family with adCRD. This family was unlinked to any known adRP or adCRD locus and SNP genotyping revealed that it was linked to a new locus on chromosome 2, designated *CORD12*.

Methods

Clinical examination

Members of this large French non-consanguineous family (RP470) were identified with CRD which segregated as a dominant trait (adCRD). There were 9

affected patients out of 22 in 4 generations (Figure 1). Examination included assessment of visual acuity, slit lamp biomicroscopy, direct funduscopy and full field electroretinography. There was no evidence of extra-ocular signs of disease indicating that CRD was non syndromic.

Genotyping of microsatellite markers and linkage analysis

Informed written consent and peripheral blood samples were obtained from 22 examined family members. The investigators followed the tenets of the Declaration of Helsinki. Genomic DNA was isolated from 10 ml peripheral blood leucocytes using standard salting out procedure [13]. The DNA samples were quantified by a spectrophotometer and diluted to 25 ng/μl for PCR amplification. PCR was carried out in a 25 μl final volume containing 50 ng genomic DNA, 5 picomoles of each primer, 0.2 mM dNTPs (MP Biochemicals), 2 mM




Figure 1 Pedigree of family RP470 with autosomal dominant inherited cone-rod dystrophy (adCRD). Arrow indicates the index patient. Filled symbols represent members with adCRD and empty symbols represent unaffected patients. Haplotypes of microsatellite markers spanning the locus 2q24.2-2q33.1 are shown. Question marks indicate unknown alleles. Solid bars denote the haplotype that segregated with the disease phenotype.

MgCl₂, PCR buffer and 1 unit of DNA polymerase (AmpliTaq Gold; Applied Biosystems, Foster city, CA). Initial denaturation at 95°C for 10 minutes was followed by 35 cycles of denaturation at 94°C for 30 seconds, specific annealing temperature for 30 seconds, and extension at 72°C for 1 minute. A final extension step was performed at 72°C for 10 minutes. The PCR products were diluted and mixed with Genescan 400HD ROX size standard and subsequently analysed on an Applied Biosystems 3130xL genetic analyser (Applied Biosystems, Foster city, CA).

Genotyping was performed using 2 to 3 polymorphic commercially available microsatellite markers from ABI PRISM Linkage Mapping Set version 2.5 (Applied Biosystems, Foster city, CA), within or contiguous to known adRP and adCRD genes, and within the locus *CORD12*. Results were analysed with GeneMapper software (version 4.0, Applied Biosystems, Foster city, CA). Segregation of the markers among the family members was examined.

Two-point LOD scores were calculated with Superlink-online <http://bioinfo.cs.technion.ac.il/superlink-online/>. The phenotype was analyzed as an autosomal dominant and fully penetrant trait with an affected allele frequency of 0.0001. Family and haplotype data were generated using Cyrillic software (version 2.1.3; Cherwell Scientific, Oxford, UK).

SNP genotyping and analysis

To map the disease locus, a genome-wide scan was performed by the Centre National de Génotypage (CNG, <http://www.cng.fr>) by genotyping 262,264 SNPs (GeneChip Mapping 250 K Nsp Array, Affymetrix, Santa Clara, CA). Results were analyzed using TASE (Transmitted Allele Search Engine) a home-made software which compared every SNP between each individuals in the family.

The first test, named Common Allele to All Affected individuals (C3A), highlighted the common allele to all affected patients within the family. The second test, Transmitted Allele to All Children (TAAC), estimated the specific allele carried by the affected parent in a nuclear family (parents + child) and transmitted to the affected child. Two consecutive mismatched SNPs limited the size of the locus. Only the regions longer than 1 Mb were considered.

Mutation screening

Coding exons and adjacent intronic sequences of candidate genes were sequenced with an Applied Biosystems 3130xL genetic analyser (Applied Biosystems, Foster city, CA) using BigDye Terminator cycle sequencing ready reaction kit V3.1 (Applied Biosystems, Foster city, CA) following manufacturer's instructions. Primer pairs

and PCR conditions are available on request. Sequence analysis and mutation identification were performed using Collection and Sequence Analysis software package (Applied Biosystems, Foster city, CA).

Ethics Committee

Statement about Conformity with Author Information: Informed and written consent was obtained for all patients participating to the study. The study was done in adherence to the tenets of the Declaration of Helsinki.

The authors confirm that they are in compliance with their Institutional Review Boards (IRBs) as the Department of Ophthalmology of the Hospital of Montpellier has the authorization # 11018S from the French Ministry of Health for biomedical research in the field of physiology, pathophysiology, epidemiology and genetics in ophthalmology.

Results

Clinical description

The pedigree of the four generations family is shown in Figure 1. The 9 affected patients revealed features of adCRD with intra-familial variable phenotype including progressive loss of the visual acuity, typical bone spicule-shaped pigmentary deposits in the macular area or macular atrophy, moderate night blindness and reduced electroretinogram (ERG) responses (Table 1). The proband (III:1) showed patches of atrophy in the macular area with a few pigment deposits, attenuation of retinal arterioles and temporal pallor of the optic disc (Figure 2).

Mapping to *CORD12*

Microsatellite markers for the 21 adRP genes <http://www.sph.uth.tmc.edu/Retnet>, the 3 most frequent adCRD genes (*CRX*, *GUCY2D* and *PRPH2/RDS*) [7-10] and a fourth adCRD gene, *GUCA1A*, were used to genotype family members, and to search for co-segregation of the markers with the disease phenotype. All these candidate genes were excluded. We then performed a genome wide scan using Affymetrix 250 K microarrays and genotypes were analysed with the TASE software. No linkage was found for most chromosomal regions except for a large region located on chromosome 2q24.2-2q33.1. The boundaries of the locus were determined by SNP exclusion between SNPs rs174240 and rs4619591 and encompassed a 41.3-Mb region (Figure 3).

Microsatellite markers were then used to confirm linkage with the locus. We genotyped all 22 members of the family with 8 microsatellite markers located on 2q24.2-2q33.1 (Figure 1). All affected patients had a common haplotype and the boundaries of the region were determined by recombination events that occurred in affected individuals III:1, III:4, III:9, IV:4, IV:9 and healthy individual III:7. The proximal boundary was defined by the

Table 1 Clinical features of patients with cone-rod dystrophy

Patient	Sex	Age at onset	Symptoms	Age at exam.	Visual acuity OD/OS	Fundus	Visual field	ERG OD/OS Scotopic dim blue Photopic single white flash Light adapted 30-Hz flickers
II:2	F		None	64	20/20 20/16	Mild attenuation of retinal vessels	NA	40 μ V/23 μ V 181 μ V/175 μ V 90 μ V/94 μ V
II:3	F	40	Nystagmus Night blindness Photophobia	70	20/40 20/32	Mild attenuation of retinal vessels. Macular atrophy	OD: relative 20° central scotoma OS: absolute 20-30° central scotoma Normal PVF on both eyes	124 μ V/173 μ V 56 μ V/60 μ V 46 μ V/56 μ V
III:1	F	32	Nystagmus No photophobia No night blindness	44	20/100 20/100	Severe macular atrophy Rare bone spicule-shaped pigment deposits	Absolute 30° central scotoma and normal PVF on both eyes	48 μ V/35 μ V 44 μ V/42 μ V 32 μ V/41 μ V
III:2	M		None	38	20/25 20/20	Normal	Normal	253 μ V/275 μ V 30 μ V/41 μ V 70 μ V/84 μ V
III:9	F	35	Nystagmus Night blindness Photophobia	45	20/25 20/25	Mild attenuation of retinal vessels	Normal	130 μ V/121 μ V 34 μ V/46 μ V 41 μ V/40 μ V
III:10	M	Early childhood	Night blindness Mild photophobia	38	20/32 20/32	Posterior pole atrophy Mild attenuation of retinal vessels	Absolute 10° central scotoma and normal PVF on both eyes	157 μ V/160 μ V 51 μ V/45 μ V 88 μ V/77 μ V
IV:1	M		Photophobia	1	NA	Mild attenuation of retinal vessels. Abnormal pigmentation of the macular area.	NA	NA
IV:4	M		No photophobia No night blindness	11	20/32 20/50	Posterior pole atrophy Attenuated retinal vessels	Normal	NA/139 μ V NA/24 μ V NA/66 μ V
IV:10	M	Early childhood	Nystagmus Photophobia No night blindness	19	20/200 20/200	Moderate pallor of the optic discs, and macular atrophy	Relative 20° central scotoma and normal PVF on both eyes	91 μ V/89 μ V 20 μ V/13 μ V 39 μ V/42 μ V

PVF = Peripheral Visual Field
 OD/OS = oculus dexter/oculus sinister
 NA = Not available
 Normal value ranges are:
 Scotopic dim blue: 160 μ V - 250 μ V
 Photopic single white: 70 μ V - 150 μ V
 Light adapted 30-Hz flickers: > 110 μ V





Figure 2 Fundus photographs of the patient III:1. at 47 years of age showing area of macular atrophy, rare pigment deposits and attenuation of retinal vessels.



recombination event between markers D2S142 and D2S306 in affected patient IV:9 and distal boundary by the recombination event between markers D2S117 and D2S325 in healthy individual III:7 (Figure 3). Using Superlink software, we found a maximum LOD score of 2.86 at $\theta = 0$ for the marker D2S118, defining a new

locus named *CORD12*. The markers D2S142, D2S325, D2S2361 and D2S126, outside the locus, gave negative LOD scores (Table 2).

The *CORD12* 41.3-Mb interval contains 280 genes. None of them were previously reported in adCRD or adRP. However the interval does contain two previously

Table 2 Two-point LOD score for microsatellite markers of family RP470 calculated at different recombination fractions θ

Marker	Position (Mb)	Recombination fraction θ						
		0.00	0.01	0.05	0.10	0.20	0.30	0.40
D2 S142	156,283,230	- ∞	- 0.9396	- 0.3168	- 0.1121	- 0.0002	0.0069	- 0.0000
D2 S306	160,562,440	2.6182	2.5720	2.3834	2.1389	1.6185	1.0548	0.4722
D2 S364	183,034,534	2.3172	2.2753	2.1047	1.8840	1.4168	0.9190	0.4192
D2 S118	191,606,469	2.8588	2.8094	2.6077	2.3459	1.7872	1.1789	0.5432
D2 S117	195,618,799	2.0280	1.9949	1.8600	1.6860	1.3181	0.9208	0.4865
D2 S325	208,270,870	- ∞	1.1956	1.6867	1.7184	1.4494	1.0188	0.5190
D2 S2361	216,478,443	- 3.8589	- 0.5393	0.1857	0.4554	0.5836	0.4992	0.2977
D2 S126	222,016,968	- ∞	0.2379	0.7858	0.8926	0.7898	0.5512	0.2719

described autosomal recessive RP genes, namely *CERKL* and *BBS5*, which cause autosomal recessive RP and Bardet-Biedl syndrome, respectively [14,15]. All exons and flanking intron regions were sequenced but no mutation was found. Within the *CORD12* locus, three other candidate genes were also sequenced. *KLHL23* has strong similarities with the recently described gene *KLHL7* responsible for adRP [16]. *NEUROD1* regulates development and maintenance in the visual system [17]. *SF3B1* is a splicing factor [18]. Other essential components of the spliceosome, *PRPF31*, *PRPF3*, *PRPF8*, *PAP1* and *SNRNP200*, have been associated with adRP [19-22]. No disease causing mutations were detected in *KLHL7*, *NEUROD1* and *SF3B1*.

Discussion

In this study, a novel locus, *CORD12*, for autosomal dominant cone-rod dystrophy (adCRD) was identified and localized to chromosome 2q24.2-2q33.1. With *CORD8* assigned to chromosome 1q23.1-q23.3, it is the second CRD locus for which the causative gene remains unknown [23]. To date, the total number of known adCRD genes and loci, including *CORD12*, is eleven.

A maximum two-point LOD score of 2.86 at $\theta = 0$ for the marker D2S118, close to theoretical significance, was obtained. The common haplotype for affected patients in the family was flanked by SNPs between rs174240 and rs4619591, which defined the 41.3-Mb *CORD12* locus. Two other retinal dystrophy loci are mapped on chromosome 2. *RP54*, a 19.98-Mb autosomal recessive RP interval flanked by D2S149 and D2S367 on chromosome 2p22.3-p24.1 [24] and *RP28*, a 14-Mb autosomal recessive RP interval flanked by D2S1337 and D2S286 on chromosome 2p11-p15 [25,26]. The causative genes have recently been reported for both regions in September 2010, respectively *ZNF513* [27] for *RP54* and *FAM161A* for *RP28* [28,29]. A third gene, *C2ORF71*, was identified earlier this year next to *ZNF513*, by homozygosity mapping in two independent

studies in an 8-Mb locus on chromosome 2p24.1-p23.1 and in a 6.8-Mb locus on chromosome 2p23.1-p24.1 [30,31]. None of these 3 regions overlap with *CORD12*.

The *CORD12* 41.3-Mb interval contains 280 annotated genes. We sequenced five possible candidate genes. *CERKL* and *BBS5* which cause autosomal recessive RP and Bardet-Biedl syndrome, respectively, [14,15] *KLHL23*, which has strong similarities with the recently described gene *KLHL7* responsible for adRP, [16] *NEUROD1* which regulates development and maintenance in the visual system [17] and the splicing factor *SF3B1* [18]. No mutation was found in the coding region and splice sites junctions, indicating that these genes do not cause *CORD12*. However, mutations in other parts of the gene cannot be excluded. Indeed, a single-base substitution in dominant retinitis pigmentosa disease-causing gene, *PRPF31*, located deep within intron 13 was recently identified [32]. No other obvious candidate genes have been identified in *CORD12* based on tissue expression pattern and function of gene products similar to known CRD genes. The comparison with additional families with cone-rod dystrophy showing linkage to this locus will be necessary to narrow the interval and to help the identification of a novel gene.

Conclusions

In summary, we report on the identification of a novel locus for adCRD in chromosome 2q24.2-2q33. Identification of the disease causing gene in the interval will increase our understanding of the causes of cone-rod dystrophy.

Acknowledgements

We thank the family members. The work was supported by private foundations (Information Recherche sur la Rétinite Pigmentaire, Retina France, SOS Rétinite and UNADEV), Centre National de Génotypage and INSERM. Special thanks to UNADEV which supports fellowship for GM and MH.

Author details

¹INSERM U1051, Institute for Neurosciences of Montpellier, (80 rue Augustin Fliche), Montpellier, (34091), France. ²Université Montpellier 1, (2 rue Ecole de

Médecine), Montpellier, (34060), France. ³Université Montpellier 2, (Place Eugène Bataillon), Montpellier, (34095), France. ⁴Genetics of Sensory Diseases, CHRU Gui de Chaulliac, (80 rue Augustin Fliche), Montpellier, (34295), France. ⁵Centre National de Génotypage, (2 rue Gaston Crémieux), Evry, (91057), France.

Authors' contributions

GM carried out the molecular genetic studies and the sequence alignment, participated in the design of the study and drafted the manuscript. MH performed the genotyping analysis. BB participated in the molecular genetic studies. IM participated in the medical examinations. DCC participated in the molecular genetic studies. AS participated in the sequence alignment. ABA performed the genotyping analysis. DZ performed the genotyping analysis. CPH carried out the medical examinations, conceived of the study, and participated in its design and coordination and helped to draft the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 1 July 2010 Accepted: 15 April 2011 Published: 15 April 2011

References

- Hartong DT, Berson EL, Dryja TP: **Retinitis pigmentosa**. *Lancet* 2006, **368**:1795-1809.
- Daiger SP, Bowne SJ, Sullivan LS: **Perspective on genes and mutations causing retinitis pigmentosa**. *Arch Ophthalmol* 2007, **125**:151-158.
- Hamel CP: **Cone rod dystrophies**. *Orphanet J Rare Dis* 2007, **2**:7.
- Maugeri A, Klevering BJ, Rohrschneider K, Blankenagel A, Brunner HG, Deutman AF, Hoyng CB, Cremers FP: **Mutations in the ABCA4 (ABCR) gene are the major cause of autosomal recessive cone-rod dystrophy**. *Am J Hum Genet* 2000, **67**:960-966.
- Ducroq D, Rozet J, Gerber S, Perrault I, Barbet D, Hanein S, Hakiki S, Dufier J, Munnich A, Hamel C, Kaplan J: **The ABCA4 gene in autosomal recessive cone-rod dystrophies**. *Am J Hum Genet* 2002, **71**:1480-1482.
- Fishman GA, Stone EM, Eliason DA, Taylor CM, Lindeman M, Derlacki DJ: **ABCA4 gene sequence variations in patients with autosomal recessive cone-rod dystrophy**. *Arch Ophthalmol* 2003, **121**:851-855.
- Nakazawa M, Naoi N, Wada Y, Nakazaki S, Maruiwa F, Sawada A, Tamai M: **Autosomal dominant cone-rod dystrophy associated with a Val200Glu mutation of the peripherin/RDS gene**. *Retina (Philadelphia, Pa)* 1996, **16**:405-410.
- Kelsell RE, Gregory-Evans K, Payne AM, Perrault I, Kaplan J, Yang RB, Garbers DL, Bird AC, Moore AT, Hunt DM: **Mutations in the retinal guanylate cyclase (RETGC-1) gene in dominant cone-rod dystrophy**. *Hum Mol Genet* 1998, **7**:1179-1184.
- Perrault I, Rozet JM, Gerber S, Kelsell RE, Souied E, Cabot A, Hunt DM, Munnich A, Kaplan J: **A retGC-1 mutation in autosomal dominant cone-rod dystrophy**. *Am J Hum Genet* 1998, **63**:651-654.
- Swaroop A, Wang QL, Wu W, Cook J, Coats C, Xu S, Chen S, Zack DJ, Sieving PA: **Leber congenital amaurosis caused by a homozygous mutation (R90W) in the homeodomain of the retinal transcription factor CRX: direct evidence for the involvement of CRX in the development of photoreceptor function**. *Hum Mol Genet* 1999, **8**:299-305.
- Freund CL, Gregory-Evans CY, Furukawa T, Papaioannou M, Looser J, Ploder L, Bellingham J, Ng D, Herbrick JA, Duncan A, Scherer SW, Tsui LC, Loutradis-Anagnostou A, Jacobson SG, Cepko CL, Bhattacharya SS, McInnes RR: **Cone-rod dystrophy due to mutations in a novel photoreceptor-specific homeobox gene (CRX) essential for maintenance of the photoreceptor**. *Cell* 1997, **91**:543-553.
- Swain PK, Chen S, Wang QL, Affatigato LM, Coats CL, Brady KD, Fishman GA, Jacobson SG, Swaroop A, Stone E, Sieving PA, Zack DJ: **Mutations in the cone-rod homeobox gene are associated with the cone-rod dystrophy photoreceptor degeneration**. *Neuron* 1997, **19**:1329-1336.
- Miller SA, Dykes DD, Polesky HF: **A simple salting out procedure for extracting DNA from human nucleated cells**. *Nucleic Acids Res* 1988, **16**:1215.
- Tuson M, Marfany G, González-Duarte R: **Mutation of CERKL, a novel human ceramide kinase gene, causes autosomal recessive retinitis pigmentosa (RP26)**. *Am J Hum Genet* 2004, **74**:128-138.
- Young TL, Penney L, Woods MO, Parfrey PS, Green JS, Hefferton D, Davidson WS: **A fifth locus for Bardet-Biedl syndrome maps to chromosome 2q31**. *Am J Hum Genet* 1999, **64**:900-904.
- Friedman JS, Ray JW, Waseem N, Johnson K, Brooks MJ, Hugosson T, Breuer D, Branham KE, Krauth DS, Bowne SJ, Sullivan V, Ponjavic V, Gränse L, Khanna R, Trager EH, Gieser LM, Hughbanks-Wheaton D, Cojocararu RI, Ghiasvand NM, Chakarova CF, Abrahamson M, Göring HHH, Webster AR, Birch DG, Abecasis GR, Fann Y, Bhattacharya SS, Daiger SP, Heckenlively JR, Andréasson S, Swaroop A: **Mutations in a BTB-Kelch protein, KLHL7, cause autosomal-dominant retinitis pigmentosa**. *Am J Hum Genet* 2009, **84**:792-800.
- Pennesi ME, Cho J, Yang Z, Wu SH, Zhang J, Wu SM, Tsai M: **BETA2/NeuroD1 null mice: a new model for transcription factor-dependent photoreceptor degeneration**. *J Neurosci* 2003, **23**:453-461.
- Isono K, Abe K, Tomaru Y, Okazaki Y, Hayashizaki Y, Koseki H: **Molecular cloning, genetic mapping, and expression of the mouse Sf3b1 (SAP155) gene for the U2 snRNP component of spliceosome**. *Mamm Genome* 2001, **12**:192-198.
- Vithana EN, Abu-Safieh L, Allen MJ, Carey A, Papaioannou M, Chakarova C, Al-Maghteh M, Ebenezer ND, Willis C, Moore AT, Bird AC, Hunt DM, Bhattacharya SS: **A human homolog of yeast pre-mRNA splicing gene, PRP31, underlies autosomal dominant retinitis pigmentosa on chromosome 19q13.4 (RP11)**. *Mol Cell* 2001, **8**:375-381.
- Chakarova CF, Hims MM, Bolz H, Abu-Safieh L, Patel RJ, Papaioannou MG, Inglehearn CF, Keen TJ, Willis C, Moore AT, Rosenberg T, Webster AR, Bird AC, Gal A, Hunt D, Vithana EN, Bhattacharya SS: **Mutations in HPRP3, a third member of pre-mRNA splicing factor genes, implicated in autosomal dominant retinitis pigmentosa**. *Hum Mol Genet* 2002, **11**:87-92.
- van Lith-Verhoeven JJC, van der Velde-Visser SD, Sohocki MM, Deutman AF, Brink HMA, Cremers FPM, Hoyng CB: **Clinical characterization, linkage analysis, and PRPC8 mutation analysis of a family with autosomal dominant retinitis pigmentosa type 13 (RP13)**. *Ophthalmic Genet* 2002, **23**:1-12.
- Zhao C, Bellur DL, Lu S, Zhao F, Grassi MA, Bowne SJ, Sullivan LS, Daiger SP, Chen LJ, Pang CP, Zhao K, Staley JP, Larsson C: **Autosomal-dominant retinitis pigmentosa caused by a mutation in SNRNP200, a gene required for unwinding of U4/U6 snRNAs**. *Am J Hum Genet* 2009, **85**:617-627.
- Khaliq S, Hameed A, Ismail M, Anwar K, Leroy BP, Mehdi SQ, Payne AM, Bhattacharya SS: **Novel locus for autosomal recessive cone-rod dystrophyCORD8 mapping to chromosome 1q12-Q24**. *Invest Ophthalmol Vis Sci* 2000, **41**:3709-3712.
- Naz S, Riazuddin SA, Li L, Shahid M, Kousar S, Sieving PA, Hejtmancik JF, Riazuddin S: **A novel locus for autosomal recessive retinitis pigmentosa in a consanguineous Pakistani family maps to chromosome 2p**. *Am J Ophthalmol* 2010, **149**:861-866.
- Gu S, Kumaramanickavel G, Srikumari CR, Denton MJ, Gal A: **Autosomal recessive retinitis pigmentosa locus RP28 maps between D2S1337 and D2S286 on chromosome 2p11-p15 in an Indian family**. *J Med Genet* 1999, **36**:705-707.
- Kumar A, Shetty J, Kumar B, Blanton SH: **Confirmation of linkage and refinement of the RP28 locus for autosomal recessive retinitis pigmentosa on chromosome 2p14-p15 in an Indian family**. *Mol Vis* 2004, **10**:399-402.
- Li L, Nakaya N, Chavali VRM, Ma Z, Jiao X, Sieving PA, Riazuddin S, Tomarev SI, Ayyagari R, Riazuddin SA, Hejtmancik JF: **A mutation in ZNF513, a putative regulator of photoreceptor development, causes autosomal-recessive retinitis pigmentosa**. *Am J Hum Genet* 2010, **87**:400-409.
- Langmann T, Di Gioia SA, Rau I, Stöhr H, Maksimovic NS, Corbo JC, Renner AB, Zrenner E, Kumaramanickavel G, Karlstetter M, Arsenijevic Y, Weber BHF, Gal A, Rivolta C: **Nonsense mutations in FAM161A cause RP28-associated recessive retinitis pigmentosa**. *Am J Hum Genet* 2010, **87**:376-381.
- Bandah-Rozenfeld D, Mizrahi-Meissonnier L, Farhy C, Obolensky A, Chowers I, Pe'er J, Merin S, Ben-Yosef T, Ashery-Padan R, Banin E, Sharon D: **Homozygosity mapping reveals null mutations in FAM161A as a cause of autosomal-recessive retinitis pigmentosa**. *Am J Hum Genet* 2010, **87**:382-391.
- Nishimura DY, Baye LM, Perveen R, Searby CC, Avila-Fernandez A, Pereiro I, Ayuso C, Valverde D, Bishop PN, Manson FDC, Urquhart J, Stone EM,

- Slusarski DC, Black GCM, Sheffield VC: **Discovery and functional analysis of a retinitis pigmentosa gene, C2ORF71.** *Am J Hum Genet* 2010, **86**:686-695.
31. Collin RWJ, Safieh C, Littink KW, Shalev SA, Garzozi HJ, Rizel L, Abbasi AH, Cremers FPM, den Hollander AI, Klevering BJ, Ben-Yosef T: **Mutations in C2ORF71 cause autosomal-recessive retinitis pigmentosa.** *Am J Hum Genet* 2010, **86**:783-788.
32. Rio Frio T, McGee TL, Wade NM, Iseli C, Beckmann JS, Berson EL, Rivolta C: **A single-base substitution within an intronic repetitive element causes dominant retinitis pigmentosa with reduced penetrance.** *Hum Mutat* 2009, **30**:1340-1347.

Pre-publication history

The pre-publication history for this paper can be accessed here:
<http://www.biomedcentral.com/1471-2350/12/54/prepub>

doi:10.1186/1471-2350-12-54

Cite this article as: Manes *et al.*: A novel locus (*CORD12*) for autosomal dominant cone-rod dystrophy on chromosome 2q24.2-2q33.1. *BMC Medical Genetics* 2011 **12**:54.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

