




CORRIGENDUM

Transanal total mesorectal excision for rectal cancer: a multicentric cohort study

Liang Kang¹, Yuan-Guang Chen², Hao Zhang³, Hong-Yu Zhang⁴, Guo-Le Lin⁵, Ying-Chi Yang⁶, Wen-Hao Chen⁷, Shuang-Ling Luo¹, Ning Chen⁸, Wei-Dong Tong⁹, Zhan-Long Shen¹⁰, De-Hai Xiong¹¹, Yi Xiao ^{5,*}, Zhong-Tao Zhang^{6,*} and Jian-Ping Wang^{1,*}

¹Department of Colorectal Surgery, The Sixth Affiliated Hospital of Sun Yat-sen University, Guangdong Provincial Key Laboratory of Colorectal and Pelvic Floor Diseases, Guangzhou, Guangdong, P. R. China;

²Department of Gastrointestinal Surgery, The First Affiliated Hospital of Guangzhou Medical University, Guangzhou, Guangdong, P. R. China; ³Department of General Surgery, Dongguan Kanghua Hospital, Dongguan, Guangdong, P. R. China;

⁴Department of Gastrointestinal Surgery, The First Affiliated Hospital of Chongqing Medical University, Chongqing, P. R. China; ⁵Department of General Surgery, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, P. R. China;

⁶Department of General Surgery, Beijing Friendship Hospital, Beijing, P. R. China; ⁷Department of Colorectal Surgery, Zhongnan Hospital of Wuhan University, Wuhan, Hubei, P. R. China; ⁸Department of General Surgery, Peking University Third Hospital, Beijing, P. R. China;

⁹Department of Gastrointestinal Surgery, Daping Hospital, Third Military Medical University, Chongqing, P. R. China; ¹⁰Department of Gastrointestinal Surgery, Peking University People's Hospital, Beijing, P. R. China;

¹¹Department of General Surgery, Chongqing Three Gorges Center Hospital, Chongqing, P. R. China

*Corresponding authors. Department of General Surgery, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College; No. 1 Wangfujing, Shuaifuyuan Hutong, Dongcheng, Beijing 100730, P. R. China. Tel: 86-10-69152212; Email: xiaoy@pumch.cn; Department of General Surgery, Beijing Friendship Hospital; No. 95 Yong'an Road, Xicheng, Beijing 100050, P. R. China. Tel: +86-10-63138712; Email: zhangzht@ccmu.edu.cn; Department of Colorectal Surgery, the Sixth Affiliated Hospital of Sun Yat-sen University, Guangdong Provincial Key Laboratory of Colorectal and Pelvic Floor Diseases, No. 26, Yuancun Erheng Road, Tianhe, Guangzhou, Guangdong 510655, P. R. China. Tel: +86-20-38254020; Email: wangjpngz@126.com

Gastroenterology Report, 8(1), 2020, 36–41.

<https://doi.org/10.1093/gastro/goz049>

In the original version of this manuscript, the following line in the abstract read: 'among the 211 patients, 175 (82.9%) had

complete TME and 33 (15.6%) had incomplete TME'. This has now been corrected to: 'among the 211 patients, 175 (82.9%) had complete TME and 33 (15.6%) had near complete TME'. The author apologises for the error.