

Poly[[diaqua(μ_4 -benzene-1,2,4,5-tetracarboxylato)tetrakis(1*H*-imidazole- κ N³)-dicopper(II)] *N,N*-dimethylformamide monosolvate]

Xiao-Fei Zhou, Hong-Ping Xiao, Ya-Juan Zhao and Xin-Hua Li*

College of Chemistry and Materials Engineering, Wenzhou University, Wenzhou, Zhejiang 325035, People's Republic of China
Correspondence e-mail: lixinhua01@126.com


Received 18 March 2013; accepted 14 April 2013

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.040; wR factor = 0.153; data-to-parameter ratio = 12.8.

The asymmetric unit of the polymeric title compound, $\{[\text{Cu}_2(\text{C}_{10}\text{H}_2\text{O}_8)(\text{C}_3\text{H}_4\text{N}_2)_4(\text{H}_2\text{O})_2] \cdot \text{C}_3\text{H}_7\text{NO}\}_n$, contains two independent Cu^{II} ions, each coordinated by one water molecule, two imidazole N atoms and two carboxylate O atoms from benzene-1,2,4,5-tetracarboxylate anions in a distorted square-pyramidal geometry. The benzene-1,2,4,5-tetracarboxylate anion bridges four Cu^{II} ions, forming a polymeric sheet parallel to (010). In the crystal, extensive $\text{N}-\text{H} \cdots \text{O}$ and $\text{O}-\text{H} \cdots \text{O}$ hydrogen bonds link the polymeric sheets and dimethylformamide solvent molecules into a three-dimensional supramolecular structure.

Related literature

For background to the benzene-1,2,4,5-tetracarboxylate ligand in coordination polymers, see: Andruh *et al.* (2011); Clarke *et al.* (2012); Jiang *et al.* (2008); Aghabozorg *et al.* (2007); Chu *et al.* (2001); Liu & Ding (2007); Wu *et al.* (2006). For related structures, see: Zhan & Li (2010); Luo *et al.* (2007); Yang *et al.* (2004). For the synthesis, see: Zhao *et al.* (2010).



Experimental

Crystal data

$[\text{Cu}_2(\text{C}_{10}\text{H}_2\text{O}_8)(\text{C}_3\text{H}_4\text{N}_2)_4 \cdot (\text{H}_2\text{O})_2] \cdot \text{C}_3\text{H}_7\text{NO}$
 $M_r = 758.65$
Monoclinic, $P2_1/c$
 $a = 8.999$ (4) Å
 $b = 19.296$ (8) Å
 $c = 18.926$ (7) Å

$\beta = 110.590$ (18)°
 $V = 3076$ (2) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 1.46$ mm⁻¹
 $T = 298$ K
 $0.23 \times 0.21 \times 0.15$ mm

Data collection

Bruker APEXII area-detector diffractometer
Absorption correction: multi-scan (SADABS; Bruker, 2001)
 $T_{\text{min}} = 0.731$, $T_{\text{max}} = 0.811$

17289 measured reflections
5445 independent reflections
4149 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.038$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.153$
 $S = 1.11$
5445 reflections

426 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.51$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.60$ e Å⁻³

Table 1

Selected bond lengths (Å).

Cu1—N1	2.003 (4)	Cu2—N5	1.985 (3)
Cu1—N3	2.025 (4)	Cu2—N7	1.980 (3)
Cu1—O1	1.984 (3)	Cu2—O3	2.243 (3)
Cu1—O5 ⁱ	1.985 (3)	Cu2—O7 ⁱⁱ	2.003 (3)
Cu1—O9	2.344 (3)	Cu2—O10	2.043 (3)

Symmetry codes: (i) $x + 1, -y + \frac{3}{2}, z + \frac{1}{2}$; (ii) $x, -y + \frac{3}{2}, z + \frac{1}{2}$.

Table 2

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N2—H2N \cdots O3 ⁱⁱⁱ	0.86	2.01	2.807 (5)	153
N4—H4N \cdots O7 ⁱⁱ	0.86	2.35	3.099 (5)	146
N6—H6N \cdots O6 ⁱ	0.86	2.03	2.733 (5)	138
N8—H8N \cdots O2 ^{iv}	0.86	2.35	2.952 (5)	127
N8—H8N \cdots O4 ^{iv}	0.86	2.26	3.033 (6)	149
O9—H9B \cdots O11 ^v	0.85	2.06	2.909 (5)	177
O9—H9C \cdots O11 ^{iv}	0.85	2.02	2.805 (6)	153
O10—H10A \cdots O8 ^{vi}	0.85	1.90	2.655 (4)	147
O10—H10B \cdots O4	0.85	2.24	2.703 (5)	114

Symmetry codes: (i) $x+1, -y+\frac{3}{2}, z+\frac{1}{2}$; (ii) $x, -y+\frac{3}{2}, z+\frac{1}{2}$; (iii) $x+1, y, z$; (iv) $x-1, y, z$; (v) $-x+2, -y+1, -z+1$; (vi) $-x, y+\frac{1}{2}, -z+\frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINTE* (Bruker, 2007); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

The work was supported by the National Natural Science Foundation of China (grant Nos. 21171133 and 21271143) and the Opening Foundation of Zhejiang Provincial Top Key Discipline (20121112).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5689).

References

- Aghabozorg, H., Bahrami, Z., Tabatabaie, M., Ghadermazi, M. & Attar Gharamaleki, J. (2007). *Acta Cryst.* **E63**, m2022–m2023.
- Andruh, M., Ruiz-Perez, C., Fokin, A. A., Gerbig, D. & Schreiner, P. R. (2011). *J. Am. Chem. Soc.* **133**, 20036–20039.
- Bruker (2001). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2007). *APEX2* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chu, D.-Q., Xu, J.-Q., Duan, L.-M., Wang, T.-G., Tang, A.-Q. & Ye, L. (2001). *Eur. J. Inorg. Chem.* pp. 1135–1137.
- Clarke, H. D., Hickey, M. B., Moulton, B., Perman, J. A., Peterson, M. L., Wojtas, L., Almarsson, O. & Zaworotko, M. J. (2012). *Cryst. Growth Des.* **12**, 4194–4201.
- Jiang, X.-D., Li, X.-B. & Sun, B.-W. (2008). *Acta Cryst.* **E64**, m922–m923.
- Liu, Y.-H. & Ding, M.-T. (2007). *Acta Cryst.* **E63**, m1828–m1829.
- Luo, J.-H., Huang, C.-C., Huang, X.-H. & Chen, X.-J. (2007). *Acta Cryst.* **C63**, m273–m276.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Wu, J.-Y., Chang, C.-H., Tseng, T.-W. & Lu, K.-L. (2006). *J. Mol. Struct.* **796**, 69–75.
- Yang, E., Zhang, J., Wen, Y.-H., Chen, Y.-B., Kang, Y., Li, Z.-J. & Yao, Y.-G. (2004). *Acta Cryst.* **C60**, m280–m282.
- Zhan, G. & Li, X.-H. (2010). *Acta Cryst.* **C66**, m29–m31.
- Zhao, J., Shi, D., Cheng, H., Chen, L., Ma, P. & Niu, J. (2010). *Inorg. Chem. Commun.* **13**, 822–827.

supplementary materials

Acta Cryst. (2013). E69, m273–m274 [doi:10.1107/S1600536813010246]

Poly[[diaqua(μ_4 -benzene-1,2,4,5-tetracarboxylato)tetrakis(1*H*-imidazole- κ N³)dicopper(II)] *N,N*-dimethylformamide monosolvate]

Xiao-Fei Zhou, Hong-Ping Xiao, Ya-Juan Zhao and Xin-Hua Li

Comment

In recent years, many successful implementation of crystal engineering concepts has produced a great deal of coordination and supramolecular networks (Andruh *et al.*, 2011), many of which exhibit unusual and fascinating architectures (Clarke *et al.*, 2012). The benzene-1,2,4,5-tetracarboxylate ligand as a multi-connecting ligand is also an excellent candidate for the structuring of coordination networks (Jiang *et al.*, 2008), and comparatively few examples have been reported in relation to applying it to the building of coordination polymers (Aghabozorg *et al.*, 2007; Chu *et al.*, 2001; Liu & Ding, 2007; Wu *et al.*, 2006). Here, the title complex, $\{[\text{Cu}_2(\text{idz})_4(\text{btc})(\text{H}_2\text{O})_2]\text{DMF}\}_n$ (idz=imidazole, DMF=*N,N*-dimethylformamide, btc=benzene-1,2,4,5-tetracarboxylato), (I), represents as a novel example.

As shown in Figure 1, the asymmetric unit of (I) consists of two crystallographically independent Cu^{II} atoms. Both are five-coordinate in a slight distorted tetragonal pyramid geometry, and chemical environment of them are similar. Each Cu^{II} cation is surrounded by two O atoms from two btc tetraanions, two N atoms from two monodentate idz ligands, and one water molecule (O9 or O10) which occupies the axial position. All the bond lengths fall within the typical range of Cu—N bond and Cu—O bond lengths (Luo *et al.*, 2007; Yang *et al.*, 2004).

Cu1 and Cu2 are bridged by a btc tetraanion, which acts as μ_4 -bridge, forming a two-dimensional sheet along the *a,c* axis, as shown in Figure 2. The structure can be regarded as a grid sheets with (4,4) net topology which is constructed through Cu^{II} centers bridged by btc tetraanions. Adjacent two-dimensional sheets are parallel and are linked *via* hydrogen-bonding interactions (see Table 1) through uncoordinated DMF solvent molecules to form a sandwich structure (Zhan & Li, 2010).

Experimental


A dimethylformamide (DMF) solution (20 ml) of benzene-1,2,4,5-tetracarboxylic acid (0.1 mmol, 0.0254 g) was added dropwise to an aqueous solution (10 ml) containing copper sulfate pentahydrate (0.2 mmol, 0.0498 g), *N,N*-carbonyl-diimidazole (0.3 mmol, 0.0486 g) and NaOH (0.01 mmol, 0.0004 g) at room temperature (Zhao *et al.*, 2010). The reaction mixture was filtered and the filtrate was left to stand for about three weeks until blue single crystals were obtained (yield 31%, based on Cu).

Refinement


The nitrogen H atoms were refined subject to the restraint N—H = 0.86 Å. The water H atoms were refined subject to the restraint O—H = 0.85 Å. The other H atoms were positioned geometrically and allowed to ride on their parent atoms at distances of C—H = 0.93–0.96 Å. $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C,N})$ and $1.5U_{\text{eq}}(\text{O})$.

Computing details

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT* (Bruker, 2007); program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

**Figure 1**

Structure of the (I) with the atom numbering, showing displacement ellipsoids at the 30% probability level. H atoms and DMF molecule have been omitted for clarity.


Figure 2

Perspective view of the 2-D sheet of (I).

Poly[[diaqua(μ_4 -benzene-1,2,4,5-tetracarboxylato)tetrakis(1*H*-imidazole- κ N³)dicopper(II)] *N,N*-dimethylformamide monosolvate]

Crystal data

[Cu₂(C₁₀H₂O₈)(C₃H₄N₂)₄(H₂O)₂] \cdot C₃H₇NO

$M_r = 758.65$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 8.999$ (4) Å

$b = 19.296$ (8) Å

$c = 18.926$ (7) Å

$\beta = 110.590$ (18)°

$V = 3076$ (2) Å³

$Z = 4$

$F(000) = 1552$

$D_x = 1.638$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4538 reflections

$\theta = 2.3$ – 25.1 °

$\mu = 1.46$ mm⁻¹

$T = 298$ K

Block, blue

$0.23 \times 0.21 \times 0.15$ mm

Data collection

Bruker APEXII area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2001)

$T_{\min} = 0.731$, $T_{\max} = 0.811$

17289 measured reflections

5445 independent reflections

4149 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.038$

$\theta_{\max} = 25.1$ °, $\theta_{\min} = 1.6$ °

$h = -10 \rightarrow 7$

$k = -11 \rightarrow 23$

$l = -22 \rightarrow 22$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.040$

$wR(F^2) = 0.153$

$S = 1.11$

5445 reflections

426 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.094P)^2 + 0.3347P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.51$ e Å⁻³

$\Delta\rho_{\min} = -0.60$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.49949 (5)	0.66350 (3)	0.40320 (2)	0.02190 (17)
Cu2	-0.01800 (5)	0.90670 (3)	0.40567 (2)	0.02186 (17)
O3	0.0200 (3)	0.83286 (14)	0.32205 (15)	0.0254 (6)
O4	0.1425 (4)	0.90382 (15)	0.26628 (17)	0.0308 (7)
O2	0.3393 (3)	0.77371 (15)	0.32593 (16)	0.0302 (7)
O1	0.3008 (3)	0.65963 (14)	0.31383 (15)	0.0265 (7)
O5	-0.3003 (3)	0.82712 (14)	-0.00888 (15)	0.0259 (7)
O6	-0.3577 (3)	0.71481 (15)	-0.01062 (16)	0.0289 (7)
O8	-0.0973 (4)	0.58140 (16)	0.04247 (18)	0.0432 (9)
O7	-0.0517 (3)	0.66614 (14)	-0.02611 (14)	0.0245 (6)
O10	0.0235 (3)	0.98875 (15)	0.34694 (16)	0.0313 (7)
H10A	0.0156	1.0262	0.3690	0.047*
H10B	0.1166	0.9855	0.3454	0.047*
O9	0.5490 (4)	0.54417 (17)	0.41879 (19)	0.0445 (8)
H9B	0.5993	0.5356	0.4651	0.067*
H9C	0.4612	0.5225	0.4044	0.067*
N1	0.6196 (4)	0.67888 (18)	0.33332 (19)	0.0276 (8)
N7	-0.2465 (4)	0.91606 (18)	0.34494 (19)	0.0273 (8)
N3	0.3720 (4)	0.67091 (19)	0.4723 (2)	0.0316 (9)
N5	0.2083 (4)	0.91014 (18)	0.47251 (19)	0.0265 (8)
C7	-0.2741 (4)	0.7665 (2)	0.0167 (2)	0.0227 (9)
C2	0.1268 (4)	0.7328 (2)	0.2199 (2)	0.0193 (8)
C4	0.0744 (4)	0.8489 (2)	0.2715 (2)	0.0224 (9)
C1	0.2665 (4)	0.7223 (2)	0.2915 (2)	0.0247 (9)
C10	0.0832 (4)	0.6832 (2)	0.1639 (2)	0.0220 (9)
H10	0.1380	0.6414	0.1718	0.026*
N8	-0.5031 (4)	0.8996 (2)	0.2980 (2)	0.0443 (11)
H8N	-0.5959	0.8840	0.2928	0.053*
C3	0.0416 (4)	0.7953 (2)	0.20941 (19)	0.0173 (8)
C6	-0.1278 (4)	0.7559 (2)	0.0861 (2)	0.0191 (8)
C8	-0.0412 (4)	0.6942 (2)	0.09592 (19)	0.0176 (8)
C5	-0.0858 (4)	0.8058 (2)	0.1428 (2)	0.0220 (9)
H5	-0.1438	0.8467	0.1360	0.026*
C13	0.5758 (6)	0.6644 (3)	0.2580 (3)	0.0410 (12)
H13	0.4916	0.6360	0.2308	0.049*
C17	0.3268 (5)	0.8762 (2)	0.4620 (3)	0.0358 (11)
H17	0.3161	0.8477	0.4209	0.043*

C11	0.7426 (5)	0.7202 (2)	0.3478 (3)	0.0310 (10)
H11	0.7981	0.7385	0.3953	0.037*
N6	0.4627 (4)	0.8883 (2)	0.5180 (2)	0.0450 (11)
H6N	0.5540	0.8715	0.5222	0.054*
C20	-0.3691 (5)	0.8842 (2)	0.3536 (2)	0.0336 (11)
H20	-0.3622	0.8547	0.3935	0.040*
C9	-0.0702 (4)	0.6424 (2)	0.0330 (2)	0.0239 (9)
N2	0.7785 (5)	0.7327 (2)	0.2862 (2)	0.0396 (10)
H2N	0.8543	0.7584	0.2834	0.047*
C14	0.4166 (6)	0.6577 (3)	0.5468 (3)	0.0455 (13)
H14	0.5065	0.6326	0.5745	0.055*
C19	0.2747 (5)	0.9448 (3)	0.5398 (3)	0.0404 (12)
H19	0.2200	0.9727	0.5626	0.048*
C22	-0.3115 (6)	0.9542 (3)	0.2796 (3)	0.0466 (14)
H22	-0.2548	0.9828	0.2586	0.056*
C21	-0.4691 (6)	0.9439 (3)	0.2511 (3)	0.0531 (15)
H21	-0.5405	0.9635	0.2074	0.064*
C18	0.4314 (6)	0.9319 (3)	0.5673 (3)	0.0496 (14)
H18	0.5046	0.9495	0.6116	0.060*
N4	0.2023 (5)	0.7178 (3)	0.5191 (3)	0.0549 (12)
H4N	0.1222	0.7405	0.5218	0.066*
C16	0.2400 (5)	0.7077 (3)	0.4577 (3)	0.0441 (12)
H16	0.1806	0.7246	0.4101	0.053*
C15	0.3138 (7)	0.6855 (4)	0.5755 (4)	0.0633 (17)
H15	0.3192	0.6828	0.6254	0.076*
C12	0.6730 (6)	0.6973 (3)	0.2298 (3)	0.0525 (14)
H12	0.6683	0.6960	0.1799	0.063*
O11	1.2716 (4)	0.4804 (2)	0.4224 (2)	0.0636 (12)
N9	1.0139 (5)	0.5105 (2)	0.3827 (3)	0.0555 (12)
C24	0.9597 (8)	0.4537 (4)	0.4157 (4)	0.073 (2)
H24A	0.9436	0.4139	0.3834	0.110*
H24B	0.8614	0.4660	0.4217	0.110*
H24C	1.0376	0.4431	0.4641	0.110*
C23	1.1617 (7)	0.5176 (3)	0.3867 (3)	0.0634 (17)
H23	1.1843	0.5542	0.3600	0.076*
C25	0.8863 (10)	0.5547 (4)	0.3310 (6)	0.132 (4)
H25A	0.9320	0.5877	0.3066	0.198*
H25B	0.8327	0.5788	0.3595	0.198*
H25C	0.8117	0.5261	0.2936	0.198*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.0181 (3)	0.0249 (3)	0.0169 (3)	-0.00076 (19)	-0.00107 (19)	0.0007 (2)
Cu2	0.0243 (3)	0.0221 (3)	0.0170 (3)	0.00021 (19)	0.0045 (2)	0.00088 (19)
O3	0.0329 (15)	0.0251 (17)	0.0207 (14)	-0.0055 (12)	0.0123 (12)	-0.0071 (12)
O4	0.0410 (17)	0.0231 (18)	0.0303 (16)	-0.0099 (13)	0.0149 (13)	-0.0024 (13)
O2	0.0269 (15)	0.0292 (18)	0.0246 (15)	-0.0017 (13)	-0.0034 (12)	-0.0083 (13)
O1	0.0223 (13)	0.0270 (18)	0.0223 (15)	0.0006 (12)	-0.0017 (11)	0.0024 (13)
O5	0.0264 (14)	0.0209 (17)	0.0222 (15)	0.0019 (12)	-0.0017 (11)	0.0024 (12)

O6	0.0229 (14)	0.0276 (18)	0.0291 (16)	-0.0052 (12)	0.0003 (12)	-0.0042 (13)
O8	0.074 (2)	0.0191 (19)	0.0305 (18)	-0.0097 (16)	0.0105 (16)	-0.0003 (14)
O7	0.0304 (14)	0.0242 (17)	0.0184 (14)	-0.0015 (12)	0.0081 (11)	-0.0029 (12)
O10	0.0335 (15)	0.0196 (17)	0.0401 (17)	0.0000 (13)	0.0120 (13)	-0.0002 (13)
O9	0.0472 (19)	0.030 (2)	0.053 (2)	-0.0015 (15)	0.0126 (16)	0.0056 (16)
N1	0.0276 (18)	0.026 (2)	0.0263 (19)	-0.0016 (15)	0.0055 (14)	-0.0036 (15)
N7	0.0266 (18)	0.029 (2)	0.0240 (18)	-0.0003 (15)	0.0060 (14)	0.0036 (15)
N3	0.0258 (18)	0.035 (2)	0.032 (2)	-0.0033 (15)	0.0071 (15)	0.0034 (17)
N5	0.0288 (18)	0.023 (2)	0.0249 (18)	0.0021 (15)	0.0064 (14)	0.0010 (15)
C7	0.0177 (18)	0.029 (3)	0.020 (2)	0.0013 (17)	0.0060 (15)	-0.0044 (18)
C2	0.0225 (18)	0.022 (2)	0.0096 (17)	-0.0001 (16)	0.0008 (14)	0.0012 (15)
C4	0.0200 (18)	0.020 (2)	0.022 (2)	0.0035 (16)	0.0009 (15)	-0.0021 (17)
C1	0.0188 (18)	0.029 (3)	0.024 (2)	0.0015 (17)	0.0045 (16)	-0.0004 (19)
C10	0.0226 (19)	0.019 (2)	0.023 (2)	0.0034 (16)	0.0061 (15)	-0.0001 (17)
N8	0.0246 (19)	0.055 (3)	0.046 (2)	-0.0078 (18)	0.0035 (17)	0.006 (2)
C3	0.0184 (17)	0.019 (2)	0.0125 (18)	-0.0038 (15)	0.0034 (14)	-0.0031 (15)
C6	0.0233 (18)	0.020 (2)	0.0109 (17)	-0.0009 (16)	0.0029 (14)	-0.0005 (15)
C8	0.0197 (17)	0.021 (2)	0.0102 (17)	-0.0042 (16)	0.0028 (13)	-0.0008 (15)
C5	0.0210 (18)	0.021 (2)	0.022 (2)	0.0009 (16)	0.0042 (15)	0.0005 (17)
C13	0.041 (3)	0.057 (4)	0.023 (2)	-0.012 (2)	0.0088 (19)	-0.012 (2)
C17	0.038 (2)	0.036 (3)	0.031 (2)	0.004 (2)	0.0090 (19)	-0.006 (2)
C11	0.033 (2)	0.023 (3)	0.039 (3)	-0.0059 (19)	0.0151 (19)	-0.0070 (19)
N6	0.029 (2)	0.054 (3)	0.042 (2)	0.0132 (19)	0.0003 (17)	-0.006 (2)
C20	0.035 (2)	0.035 (3)	0.027 (2)	-0.002 (2)	0.0065 (18)	0.004 (2)
C9	0.0205 (19)	0.024 (2)	0.021 (2)	0.0015 (17)	-0.0008 (15)	0.0011 (17)
N2	0.046 (2)	0.044 (3)	0.037 (2)	-0.0142 (19)	0.0242 (18)	-0.0041 (19)
C14	0.047 (3)	0.058 (4)	0.033 (3)	0.007 (2)	0.014 (2)	0.014 (2)
C19	0.039 (3)	0.041 (3)	0.030 (3)	0.009 (2)	-0.0005 (19)	-0.016 (2)
C22	0.043 (3)	0.065 (4)	0.026 (2)	-0.006 (3)	0.005 (2)	0.023 (2)
C21	0.034 (3)	0.081 (4)	0.036 (3)	0.001 (3)	0.002 (2)	0.031 (3)
C18	0.036 (3)	0.057 (4)	0.042 (3)	0.005 (2)	-0.004 (2)	-0.023 (3)
N4	0.055 (3)	0.064 (3)	0.058 (3)	0.011 (2)	0.034 (2)	-0.005 (2)
C16	0.037 (3)	0.051 (3)	0.049 (3)	0.008 (2)	0.022 (2)	0.006 (3)
C15	0.064 (4)	0.084 (5)	0.054 (4)	0.000 (3)	0.036 (3)	0.000 (3)
C12	0.059 (3)	0.069 (4)	0.036 (3)	-0.011 (3)	0.026 (2)	-0.008 (3)
O11	0.042 (2)	0.055 (3)	0.082 (3)	0.0004 (18)	0.0066 (19)	0.027 (2)
N9	0.042 (2)	0.043 (3)	0.076 (3)	0.006 (2)	0.014 (2)	-0.014 (2)
C24	0.083 (4)	0.071 (5)	0.078 (5)	-0.021 (4)	0.044 (4)	-0.027 (4)
C23	0.068 (4)	0.048 (4)	0.062 (4)	-0.014 (3)	0.008 (3)	-0.002 (3)
C25	0.091 (6)	0.083 (6)	0.173 (10)	0.048 (5)	-0.014 (6)	-0.025 (6)

Geometric parameters (Å, °)

Cu1—N1	2.003 (4)	C3—C5	1.389 (5)
Cu1—N3	2.025 (4)	C6—C5	1.393 (5)
Cu1—O1	1.984 (3)	C6—C8	1.398 (5)
Cu1—O5 ⁱ	1.985 (3)	C8—C9	1.506 (5)
Cu1—O9	2.344 (3)	C5—H5	0.9300
Cu2—N5	1.985 (3)	C13—C12	1.335 (7)
Cu2—N7	1.980 (3)	C13—H13	0.9300

Cu2—O3	2.243 (3)	C17—N6	1.327 (6)
Cu2—O7 ⁱⁱ	2.003 (3)	C17—H17	0.9300
Cu2—O10	2.043 (3)	C11—N2	1.337 (6)
O3—C4	1.257 (5)	C11—H11	0.9300
O4—C4	1.245 (5)	N6—C18	1.358 (6)
O2—C1	1.240 (5)	N6—H6N	0.8600
O1—C1	1.282 (5)	C20—H20	0.9300
O5—C7	1.256 (5)	N2—C12	1.340 (6)
O5—Cu1 ⁱⁱⁱ	1.985 (3)	N2—H2N	0.8600
O6—C7	1.247 (5)	C14—C15	1.339 (8)
O8—C9	1.228 (5)	C14—H14	0.9300
O7—C9	1.273 (5)	C19—C18	1.344 (6)
O7—Cu2 ^{iv}	2.003 (3)	C19—H19	0.9300
O10—H10A	0.8500	C22—C21	1.344 (7)
O10—H10B	0.8500	C22—H22	0.9300
O9—H9B	0.8499	C21—H21	0.9300
O9—H9C	0.8500	C18—H18	0.9300
N1—C11	1.312 (5)	N4—C16	1.335 (7)
N1—C13	1.367 (6)	N4—C15	1.335 (7)
N7—C20	1.323 (6)	N4—H4N	0.8600
N7—C22	1.380 (5)	C16—H16	0.9300
N3—C16	1.327 (6)	C15—H15	0.9300
N3—C14	1.348 (6)	C12—H12	0.9300
N5—C17	1.325 (6)	O11—C23	1.217 (7)
N5—C19	1.376 (5)	N9—C23	1.313 (8)
C7—C6	1.512 (5)	N9—C24	1.429 (8)
C2—C10	1.378 (5)	N9—C25	1.487 (8)
C2—C3	1.406 (5)	C24—H24A	0.9600
C2—C1	1.503 (5)	C24—H24B	0.9600
C4—C3	1.514 (5)	C24—H24C	0.9600
C10—C8	1.393 (5)	C23—H23	0.9300
C10—H10	0.9300	C25—H25A	0.9600
N8—C20	1.327 (6)	C25—H25B	0.9600
N8—C21	1.342 (6)	C25—H25C	0.9600
N8—H8N	0.8600		
O1—Cu1—O5 ⁱ	176.77 (12)	C3—C5—C6	120.8 (4)
O1—Cu1—N1	88.65 (13)	C3—C5—H5	119.6
O5 ⁱ —Cu1—N1	89.92 (13)	C6—C5—H5	119.6
O1—Cu1—N3	90.46 (13)	C12—C13—N1	109.2 (4)
O5 ⁱ —Cu1—N3	90.31 (13)	C12—C13—H13	125.4
N1—Cu1—N3	167.36 (15)	N1—C13—H13	125.4
O1—Cu1—O9	98.32 (12)	N5—C17—N6	111.0 (4)
O5 ⁱ —Cu1—O9	84.71 (12)	N5—C17—H17	124.5
N1—Cu1—O9	96.07 (13)	N6—C17—H17	124.5
N3—Cu1—O9	96.53 (14)	N1—C11—N2	111.8 (4)
N7—Cu2—N5	172.01 (15)	N1—C11—H11	124.1
N7—Cu2—O7 ⁱⁱ	94.38 (13)	N2—C11—H11	124.1
N5—Cu2—O7 ⁱⁱ	88.09 (13)	C17—N6—C18	107.6 (4)

N7—Cu2—O10	87.98 (13)	C17—N6—H6N	126.2
N5—Cu2—O10	88.73 (13)	C18—N6—H6N	126.2
O7 ⁱⁱ —Cu2—O10	173.16 (12)	N7—C20—N8	111.2 (4)
N7—Cu2—O3	91.92 (13)	N7—C20—H20	124.4
N5—Cu2—O3	95.39 (13)	N8—C20—H20	124.4
O7 ⁱⁱ —Cu2—O3	95.99 (11)	O8—C9—O7	124.6 (4)
O10—Cu2—O3	90.34 (12)	O8—C9—C8	120.6 (4)
C4—O3—Cu2	125.3 (3)	O7—C9—C8	114.6 (4)
C1—O1—Cu1	106.6 (2)	C11—N2—C12	106.3 (4)
C7—O5—Cu1 ⁱⁱⁱ	113.6 (2)	C11—N2—H2N	126.8
C9—O7—Cu2 ^{iv}	114.2 (3)	C12—N2—H2N	126.8
Cu2—O10—H10A	109.2	C15—C14—N3	110.8 (5)
Cu2—O10—H10B	109.3	C15—C14—H14	124.6
H10A—O10—H10B	109.5	N3—C14—H14	124.6
Cu1—O9—H9B	109.2	C18—C19—N5	108.8 (4)
Cu1—O9—H9C	109.3	C18—C19—H19	125.6
H9B—O9—H9C	109.5	N5—C19—H19	125.6
C11—N1—C13	104.8 (4)	C21—C22—N7	109.6 (4)
C11—N1—Cu1	124.3 (3)	C21—C22—H22	125.2
C13—N1—Cu1	129.3 (3)	N7—C22—H22	125.2
C20—N7—C22	104.5 (4)	N8—C21—C22	106.4 (4)
C20—N7—Cu2	128.8 (3)	N8—C21—H21	126.8
C22—N7—Cu2	126.5 (3)	C22—C21—H21	126.8
C16—N3—C14	103.6 (4)	C19—C18—N6	107.0 (4)
C16—N3—Cu1	124.9 (3)	C19—C18—H18	126.5
C14—N3—Cu1	129.4 (3)	N6—C18—H18	126.5
C17—N5—C19	105.7 (4)	C16—N4—C15	106.2 (5)
C17—N5—Cu2	126.0 (3)	C16—N4—H4N	126.9
C19—N5—Cu2	128.3 (3)	C15—N4—H4N	126.9
O6—C7—O5	125.6 (3)	N3—C16—N4	112.3 (5)
O6—C7—C6	117.8 (4)	N3—C16—H16	123.9
O5—C7—C6	116.7 (3)	N4—C16—H16	123.9
C10—C2—C3	119.5 (3)	N4—C15—C14	107.1 (5)
C10—C2—C1	121.4 (4)	N4—C15—H15	126.4
C3—C2—C1	119.0 (3)	C14—C15—H15	126.4
O4—C4—O3	127.1 (4)	C13—C12—N2	107.8 (5)
O4—C4—C3	119.1 (4)	C13—C12—H12	126.1
O3—C4—C3	113.6 (3)	N2—C12—H12	126.1
O2—C1—O1	124.0 (3)	C23—N9—C24	123.4 (5)
O2—C1—C2	119.0 (4)	C23—N9—C25	120.7 (7)
O1—C1—C2	116.9 (3)	C24—N9—C25	115.0 (6)
C2—C10—C8	121.6 (4)	N9—C24—H24A	109.5
C2—C10—H10	119.2	N9—C24—H24B	109.5
C8—C10—H10	119.2	H24A—C24—H24B	109.5
C20—N8—C21	108.2 (4)	N9—C24—H24C	109.5
C20—N8—H8N	125.9	H24A—C24—H24C	109.5
C21—N8—H8N	125.9	H24B—C24—H24C	109.5
C5—C3—C2	119.3 (3)	O11—C23—N9	125.4 (6)
C5—C3—C4	118.6 (3)	O11—C23—H23	117.3

C2—C3—C4	121.9 (3)	N9—C23—H23	117.3
C5—C6—C8	119.9 (3)	N9—C25—H25A	109.5
C5—C6—C7	119.6 (3)	N9—C25—H25B	109.5
C8—C6—C7	120.4 (3)	H25A—C25—H25B	109.5
C10—C8—C6	118.8 (3)	N9—C25—H25C	109.5
C10—C8—C9	119.3 (3)	H25A—C25—H25C	109.5
C6—C8—C9	121.7 (3)	H25B—C25—H25C	109.5
N7—Cu2—O3—C4	98.4 (3)	C1—C2—C3—C4	-7.1 (5)
N5—Cu2—O3—C4	-78.3 (3)	O4—C4—C3—C5	-78.2 (5)
O7 ⁱⁱ —Cu2—O3—C4	-167.0 (3)	O3—C4—C3—C5	98.4 (4)
O10—Cu2—O3—C4	10.4 (3)	O4—C4—C3—C2	107.7 (4)
O5 ⁱ —Cu1—O1—C1	-13 (2)	O3—C4—C3—C2	-75.7 (5)
N1—Cu1—O1—C1	-76.3 (3)	O6—C7—C6—C5	-141.6 (4)
N3—Cu1—O1—C1	91.1 (3)	O5—C7—C6—C5	38.7 (5)
O9—Cu1—O1—C1	-172.2 (3)	O6—C7—C6—C8	33.9 (5)
O1—Cu1—N1—C11	142.5 (4)	O5—C7—C6—C8	-145.8 (4)
O5 ⁱ —Cu1—N1—C11	-34.6 (3)	C2—C10—C8—C6	-3.0 (6)
N3—Cu1—N1—C11	56.4 (7)	C2—C10—C8—C9	172.2 (4)
O9—Cu1—N1—C11	-119.3 (3)	C5—C6—C8—C10	2.0 (6)
O1—Cu1—N1—C13	-20.6 (4)	C7—C6—C8—C10	-173.5 (3)
O5 ⁱ —Cu1—N1—C13	162.3 (4)	C5—C6—C8—C9	-173.0 (4)
N3—Cu1—N1—C13	-106.6 (7)	C7—C6—C8—C9	11.5 (6)
O9—Cu1—N1—C13	77.7 (4)	C2—C3—C5—C6	-1.3 (6)
N5—Cu2—N7—C20	-105.7 (10)	C4—C3—C5—C6	-175.5 (4)
O7 ⁱⁱ —Cu2—N7—C20	2.1 (4)	C8—C6—C5—C3	0.1 (6)
O10—Cu2—N7—C20	-171.5 (4)	C7—C6—C5—C3	175.6 (3)
O3—Cu2—N7—C20	98.3 (4)	C11—N1—C13—C12	-0.1 (6)
N5—Cu2—N7—C22	79.8 (11)	Cu1—N1—C13—C12	165.4 (4)
O7 ⁱⁱ —Cu2—N7—C22	-172.4 (4)	C19—N5—C17—N6	-0.8 (5)
O10—Cu2—N7—C22	14.0 (4)	Cu2—N5—C17—N6	-178.3 (3)
O3—Cu2—N7—C22	-76.2 (4)	C13—N1—C11—N2	0.2 (5)
O1—Cu1—N3—C16	-39.4 (4)	Cu1—N1—C11—N2	-166.3 (3)
O5 ⁱ —Cu1—N3—C16	137.5 (4)	N5—C17—N6—C18	0.2 (6)
N1—Cu1—N3—C16	46.5 (8)	C22—N7—C20—N8	0.7 (5)
O9—Cu1—N3—C16	-137.8 (4)	Cu2—N7—C20—N8	-174.8 (3)
O1—Cu1—N3—C14	160.2 (4)	C21—N8—C20—N7	-0.7 (6)
O5 ⁱ —Cu1—N3—C14	-23.0 (4)	Cu2 ^{iv} —O7—C9—O8	-14.0 (5)
N1—Cu1—N3—C14	-114.0 (6)	Cu2 ^{iv} —O7—C9—C8	160.5 (2)
O9—Cu1—N3—C14	61.7 (4)	C10—C8—C9—O8	58.3 (5)
N7—Cu2—N5—C17	-156.0 (9)	C6—C8—C9—O8	-126.6 (4)
O7 ⁱⁱ —Cu2—N5—C17	95.8 (4)	C10—C8—C9—O7	-116.4 (4)
O10—Cu2—N5—C17	-90.3 (4)	C6—C8—C9—O7	58.6 (5)
O3—Cu2—N5—C17	-0.1 (4)	N1—C11—N2—C12	-0.2 (6)
N7—Cu2—N5—C19	27.0 (12)	C16—N3—C14—C15	0.3 (6)
O7 ⁱⁱ —Cu2—N5—C19	-81.2 (4)	Cu1—N3—C14—C15	163.9 (4)
O10—Cu2—N5—C19	92.8 (4)	C17—N5—C19—C18	1.1 (6)
O3—Cu2—N5—C19	-177.0 (4)	Cu2—N5—C19—C18	178.6 (4)
Cu1 ⁱⁱⁱ —O5—C7—O6	3.3 (5)	C20—N7—C22—C21	-0.4 (6)

Cu1 ⁱⁱⁱ —O5—C7—C6	-177.1 (2)	Cu2—N7—C22—C21	175.2 (4)
Cu2—O3—C4—O4	13.8 (5)	C20—N8—C21—C22	0.4 (7)
Cu2—O3—C4—C3	-162.4 (2)	N7—C22—C21—N8	0.0 (7)
Cu1—O1—C1—O2	-6.3 (5)	N5—C19—C18—N6	-1.1 (7)
Cu1—O1—C1—C2	175.1 (3)	C17—N6—C18—C19	0.5 (6)
C10—C2—C1—O2	152.4 (4)	C14—N3—C16—N4	0.2 (6)
C3—C2—C1—O2	-26.0 (6)	Cu1—N3—C16—N4	-164.4 (3)
C10—C2—C1—O1	-28.9 (6)	C15—N4—C16—N3	-0.6 (7)
C3—C2—C1—O1	152.7 (4)	C16—N4—C15—C14	0.8 (7)
C3—C2—C10—C8	1.8 (6)	N3—C14—C15—N4	-0.7 (7)
C1—C2—C10—C8	-176.7 (4)	N1—C13—C12—N2	0.0 (7)
C10—C2—C3—C5	0.3 (6)	C11—N2—C12—C13	0.1 (6)
C1—C2—C3—C5	178.8 (3)	C24—N9—C23—O11	-5.1 (10)
C10—C2—C3—C4	174.4 (4)	C25—N9—C23—O11	-173.7 (6)

Symmetry codes: (i) $x+1, -y+3/2, z+1/2$; (ii) $x, -y+3/2, z+1/2$; (iii) $x-1, -y+3/2, z-1/2$; (iv) $x, -y+3/2, z-1/2$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N2—H2N \cdots O3 ^v	0.86	2.01	2.807 (5)	153
N4—H4N \cdots O7 ⁱⁱ	0.86	2.35	3.099 (5)	146
N6—H6N \cdots O6 ⁱ	0.86	2.03	2.733 (5)	138
N8—H8N \cdots O2 ^{vi}	0.86	2.35	2.952 (5)	127
N8—H8N \cdots O4 ^{vi}	0.86	2.26	3.033 (6)	149
O9—H9B \cdots O11 ^{vii}	0.85	2.06	2.909 (5)	177
O9—H9C \cdots O11 ^{vi}	0.85	2.02	2.805 (6)	153
O10—H10A \cdots O8 ^{viii}	0.85	1.90	2.655 (4)	147
O10—H10B \cdots O4	0.85	2.24	2.703 (5)	114

Symmetry codes: (i) $x+1, -y+3/2, z+1/2$; (ii) $x, -y+3/2, z+1/2$; (v) $x+1, y, z$; (vi) $x-1, y, z$; (vii) $-x+2, -y+1, -z+1$; (viii) $-x, y+1/2, -z+1/2$.